Empagliflozin PBPK/PD model: fit PK

1 Methods: Empagliflozin PBPK/PD model: fit PK

1.1 Parameters and data

11 parameters were optimized: tissue distribution emp (`ftissue_emp`), tissue/plasma partition coefficient emp (`Kp_emp`), deep tissue distribution emp (`ftissue_deep_emp`), deep tissue/plasma partition coefficient emp (`Kp_deep_emp`), rate of empagliflozin absorption (`GU__EMPABS_k`), fraction absorbed empagliflozin (`GU__F_emp_abs`), rate of the fecal transit (`GU__METEXC_k`), fed effect on the absorption rate (`GU__FED_f_absorption`), rate of the hepatic glucuronidation of empagliflozin (`EMP2EG_k`), tubular secretion of empagliflozin relative to its filtration (`KI__EMPSEC_f`) and renal clearance of EG relative to the GFR (`KI__EGEX_f`).

A fit mapping is a curve of a study, the data and the observable of the model it is compared with. The fit has 65 fit mappings of training data with 910 data points, whose residuals enter the objective; 69 fit mappings of validation data with 714 data points, which were simulated and evaluated with the fitted parameters but not fitted; and 7 fit mappings of outliers with 83 data points, data the fit dropped as not usable, which were evaluated like the validation data so that the decision to drop them can be checked. The fit was set up and run with sbmlsim [1].

1.2 Objective function

The parameters were estimated by minimizing the objective function

$$F(\vec{p}) = \frac{1}{2} \sum_k \sum_i \rho\big(w_{i,k} r_{i,k}(\vec{p})^2\big) \tag{1}$$

where k runs over the fit mappings of the training data and i over their data points, $r_{i,k}$ is the residual, $w_{i,k} = w_k w^p_{i,k}$ the weight of the data point and ρ the loss function, applied to the squared weighted residual $z = w_{i,k} r^2_{i,k}$. The loss function is linear, $\rho(z) = z$, i.e. a standard least squares problem.

The residuals are normalized by the mean $\overline{y}_k$ of the data of their curve, which makes curves of different quantities and magnitudes comparable:

$$r_{i,k} = \frac{y_{i,k} - f_{k(x_{i,k},\vec{p})}}{\overline{y}_k} \tag{2}$$

The weight of curve k is $w_k = \frac{n_k}{N_k}$, the weight n_k of its fit mapping (the weight its fit mapping collection assigns, 1 by default, or, for a collection which uses the weights of its mappings, the weight of the mapping, by default the number of subjects of its data) divided by the number N_k of its data points.

A data point is weighted with the inverse of its coefficient of variation, $w^p_{i,k} = |y_{i,k} \frac{|}{\sigma_{i,k}}$, with $\sigma_{i,k}$ the standard deviation of the data point, or its standard error for data without a standard deviation. An error of zero is missing, a missing error of a curve with errors is replaced by the largest error of the curve, and the data points of a curve without errors have the weight $w^p_{i,k} = 2$, the weight of a coefficient of variation of 0.5.

1.3 Optimization

The optimizer searched the parameters in log10 space within their bounds; bounds, values and intervals are given in the units of the fit parameters, the units of the table of the parameters.

The objective was minimized in 3 runs with differential evolution [2] of SciPy [3] within the bounds of the parameters, with the strategy `best1bin`, a population of 15 times the number of parameters, at most 1000 generations, the relative tolerance 0.01 of convergence, the recombination constant 0.7 and a mutation dithered between 0.5 and 1. The initial population of a run was drawn by latin hypercube sampling [4], with a seed of every run derived from the seed 1234 of the fit. The best member of the final population was polished with the L-BFGS-B method of SciPy.

The convergence of the runs was assessed from the traces of their cost and the waterfall plot of their final costs, and the parameter set with the lowest cost was used.

The models were simulated with libRoadRunner [5] and its integrator CVODE [6] with the relative tolerance 10^{-6} and the absolute tolerances 10^{-12} of an amount, 10^{-12} of a concentration and 10^{-6} of the other states, which for a species in a concentration is multiplied by the initial volume of its compartment. It ran with the variable step size of libRoadRunner, i.e. with an output at every step of the integrator. A data point at the time of a change after the start of a simulation, e.g. a dose or the reset of a collection, was compared with the state before the change, except the first

data point of a curve with data at later times, which starts a collection and was compared with the state after the change.

1.4 Goodness of fit

The goodness of fit is given per kind of data (training data, validation data and outliers) by metrics over its n data points with the residuals $e_i = y_i - f\left(x_i, \hat{\vec{p}}\right)$ at the fitted parameters $\hat{\vec{p}}$, the mean squared error, its root and the root mean squared error normalized by the mean $\overline{y}_{k(i)}$ of the data of the curve of a point:

$$\mathrm{MSE} = \frac{1}{n}\sum_{i=1}^{n} e_i^2, \quad \mathrm{RMSE} = \sqrt{\mathrm{MSE}}, \quad \mathrm{NRMSE} = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\left(\frac{e_i}{\overline{y}_{k(i)}}\right)^2} \tag{3}$$

The root mean square of the residuals of the objective, i.e. what the fit minimizes, and the coefficient of determination:

$$\mathrm{RMSE}_w = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\rho(w_i r_i^2)}, \quad R^2 = 1 - \frac{\sum_i e_i^2}{\sum_i \left(y_i - \overline{y}\right)^2} \tag{4}$$

The predictions of a non-linear model are no linear regression of the data, so R^2 can be negative: the prediction is worse than the mean of the data. The information criteria of Akaike [7] and Schwarz [8] with the number n_p of fitted parameters are those of a least squares fit with normally distributed residuals, up to an additive constant, so only differences between models fitted to the same data are meaningful:

$$\mathrm{AIC} = n\ln(\mathrm{MSE}) + 2n_p, \quad \mathrm{BIC} = n\ln(\mathrm{MSE}) + n_p\ln(n) \tag{5}$$

The percentage of the data points whose prediction $f_i = f\left(x_i, \hat{\vec{p}}\right)$ lies within twofold of the data is taken over the n_+ data points with positive data $y_i > 0$ and a positive prediction $f_i > 0$; the others have no ratio and are not counted, and a kind of data without such a point has no percentage:

$$P_2 = \frac{100}{n_+}\left|\left\{i : \frac{1}{2} \le \frac{f_i}{y_i} \le 2\right\}\right| \tag{6}$$

The goodness of fit plot shows the predictions against the data on logarithmic axes, with the lines of twofold $f = 2y$ and $f = \frac{y}{2}$. The Bland-Altman plot [9] shows the ratio of prediction and data as $d_i = \log_{10}\left(\frac{f_i}{y_i}\right)$ against the geometric mean $\sqrt{f_i y_i}$ for the data points with positive values, with the bias $\overline{d}$ and the limits of agreement $\overline{d} \pm 1.96 s_d$, with the sample standard deviation s_d of d_i, of the training data; they are read as the fold factors $10^{\overline{d}}$ and $10^{\overline{d} \pm 1.96 s_d}$.

1.5 Identifiability

The identifiability of the parameters was analysed with the profile likelihood [10], [11]. The profile of a parameter is the lowest objective with the parameter fixed:

$$\mathrm{PL}(p_j) = \min_{p_{i \ne j}} F(\vec{p}) \tag{7}$$

At every point of a scan the other parameters were re-optimized from the previous point with the least squares optimizer of SciPy with the relative step 0.05. A scan ran from the optimum in both directions in the space of the optimizer, with steps in decades of the parameter, starting at 0.1. A step which raised the objective by more than 0.2 of the distance between the objective at the optimum and the threshold was reduced by the factor 2 and repeated, down to 0.01, and the step after one which raised it by less than that divided by the factor was enlarged by the factor, up to 1. A scan stopped at the threshold, at the bound of the parameter, at a point which failed or after 50 points.

The confidence intervals are the values at which the profile crosses the threshold of the likelihood ratio test

$$F_{\mathrm{thr}} = F_{\mathrm{min}} + \chi^2_\alpha \frac{\mathrm{df}}{2} \tag{8}$$

with F_{min} the lowest objective of the fit and of all profiles and $\chi^2_\alpha(\mathrm{df})$ the quantile of the χ^2 distribution at the confidence level $\alpha = 0.95$ with df = 1 degrees of freedom, i.e. 1.92 above the lowest objective. The threshold has this confidence level if F is the negative log-likelihood of the data up to a constant, i.e. if the weighted residuals are independent and normally distributed with unit variance; otherwise the level is approximate. A crossing is interpolated between two points of the profile, on the logarithm of a parameter on a logarithmic scale. A parameter is identifiable if its profile reaches the threshold within the scan on both sides of the optimum, i.e. its confidence

interval is finite; practically non-identifiable towards the lower bound if its profile reaches the threshold above the optimum but not below it within the scan, which ends at the lower bound or after 50 points; practically non-identifiable towards the upper bound if its profile reaches the threshold below the optimum but not above it within the scan, which ends at the upper bound or after 50 points; practically non-identifiable if its profile reaches the threshold on neither side within the scans, which end at the bounds or after 50 points, and is not flat; and structurally non-identifiable if its profile is flat on both sides. A side of a profile is flat if it rises by less than 0.05 of the distance between the lowest objective and the threshold, and a profile with a point which failed or did not converge is not classified. The profile likelihood is preferred over the intervals of the Fisher information, its intervals are invariant under a transformation of the parameters and can be asymmetric [12].

The Fisher information is the local alternative to the profiles: it is the Gauss-Newton approximation of the curvature of the objective at the optimum, from the Jacobian J of the weighted residuals of the training data with respect to the parameters $\vec{p}$ in the space of the optimizer, computed by central finite differences with the step $h_j = \varepsilon \max(|p_j|, 1),\ \varepsilon = 10^{-6}$ of the parameter p_j:

$$I = J^\top J, \quad \mathrm{cov} = \sigma^2 I^{-1}, \quad \sigma^2 = \frac{2F}{n - n_p} \tag{9}$$

with the number n of data points of the training data and the number n_p of parameters. The information has rank 11 of 11, the number of its eigenvalues above 10^{-8} of the largest one, and the condition number 3890; the covariance of an information of a lower rank is computed with the pseudo-inverse. The Wald intervals are $\hat{p}_j \pm t_{\frac{1+\alpha}{2}, n-n_p} \mathrm{SE}_j$ with the standard errors $\mathrm{SE}_j = \sqrt{\mathrm{cov}_{jj}}$ at the confidence level $\alpha = 0.95$, transformed back into the units of the fit parameters.

1.6 Provenance

The fit PK_20261019_201013__13ddc was run on 2026-10-19 18:10:18 UTC with sbmlsim 0.10.0 [1], libRoadRunner 2.10.0 [5], SciPy 1.18.1 [3], NumPy 2.5.3 [13] and Python 3.14.4 (seed 1234). The models and data were those of commit `f43c98cc20b2368500dfcbc3ee1739eeb596f460` of `/home/mkoenig/git/pkdb_models-empagliflozin` their SHA-256 checksums are listed in Table 1.

Table 1: SHA-256 checksums of the models and data of the fit PK_20261019_201013__13ddc, their first 12 characters; the full checksums are in the provenance of the result (optimization_result.json).

File	SHA-256
pkdb_models/models/empagliflozin/data/dapagliflozin/Kim2023/.Kim2023_Fig1.tsv	faf8a67de950...
pkdb_models/models/empagliflozin/data/dapagliflozin/Kim2023/.Kim2023_Fig2.tsv	7419518c5dc2...
pkdb_models/models/empagliflozin/data/dapagliflozin/Kim2023/.Kim2023_Tab2.tsv	e4b13d62892b...
pkdb_models/models/empagliflozin/data/dapagliflozin/Kim2023/.Kim2023_Tab3.tsv	ea1325423a66...
pkdb_models/models/empagliflozin/data/dapagliflozin/Kim2023/.Kim2023_TabGroups.tsv	e6e46d776a7a...
pkdb_models/models/empagliflozin/data/dapagliflozin/Kim2023/.Kim2023_TabInterventions.tsv	00fa9049fea8...
pkdb_models/models/empagliflozin/data/dapagliflozin/Kim2023/.Kim2023_TabS2.tsv	aa635231e391...
pkdb_models/models/empagliflozin/data/dapagliflozin/vanderAartvanderBeek2020/.vanderAartvanderBeek2020_Fig3.tsv	2bd012174836...
pkdb_models/models/empagliflozin/data/dapagliflozin/vanderAartvanderBeek2020/.vanderAartvanderBeek2020_TabGroups.tsv	033515791883...
pkdb_models/models/empagliflozin/data/dapagliflozin/vanderAartvanderBeek2020/.vanderAartvanderBeek2020_TabIndividuals.tsv	77b6cfffbdb8...
pkdb_models/models/empagliflozin/data/dapagliflozin/vanderAartvanderBeek2020/.vanderAartvanderBeek2020_TabInterventions.tsv	86f743929d49...
pkdb_models/models/empagliflozin/data/empagliflozin/Ayoub2017/.Ayoub2017_Fig6.tsv	830026ab86aa...
pkdb_models/models/empagliflozin/data/empagliflozin/Ayoub2017/.Ayoub2017_Tab1.tsv	2191c2e86f41...

File	SHA-256
pkdb_models/models/empagliflozin/data/empagliflozin/Ayoub2017/.Ayoub2017_TabGroups.tsv	ab84d27300d7...
pkdb_models/models/empagliflozin/data/empagliflozin/Ayoub2017/.Ayoub2017_TabInterventions.tsv	2eebacd7400e...
pkdb_models/models/empagliflozin/data/empagliflozin/Brand2012/.Brand2012_Fig2.tsv	f1b64174e094...
pkdb_models/models/empagliflozin/data/empagliflozin/Brand2012/.Brand2012_Fig3.tsv	b8564d1b1a50...
pkdb_models/models/empagliflozin/data/empagliflozin/Brand2012/.Brand2012_Fig4.tsv	0f0f7ba3c41f...
pkdb_models/models/empagliflozin/data/empagliflozin/Brand2012/.Brand2012_Fig5.tsv	d32555572f67...
pkdb_models/models/empagliflozin/data/empagliflozin/Brand2012/.Brand2012_Tab1.tsv	edca065d782f...
pkdb_models/models/empagliflozin/data/empagliflozin/Brand2012/.Brand2012_TabGroups.tsv	ee2e1d8b121f...
pkdb_models/models/empagliflozin/data/empagliflozin/Brand2012/.Brand2012_TabInterventions.tsv	a0113deb79ef...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2015a/.Chen2015a_Fig1.tsv	57d1bcb1699e...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2015a/.Chen2015a_Fig2.tsv	46fa14430531...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2015a/.Chen2015a_Tab1.tsv	4274ba8eabce...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2015a/.Chen2015a_Tab2.tsv	e867dab9f8cb...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2015a/.Chen2015a_Tab2A.tsv	056a6134d497...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2015a/.Chen2015a_TabGroups.tsv	6996658d26dc...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2015a/.Chen2015a_TabInterventions.tsv	0a85f5e98b81...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2020/.Chen2020_Fig2.tsv	4bec8b2edb45...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2020/.Chen2020_Tab3.tsv	fe8023eb0c58...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2020/.Chen2020_TabGroups.tsv	febcf7e197e5...
pkdb_models/models/empagliflozin/data/empagliflozin/Chen2020/.Chen2020_TabInterventions.tsv	4f5508a75efa...
pkdb_models/models/empagliflozin/data/empagliflozin/ElDash2021/.ElDash2021_Fig4.tsv	23df1284387a...
pkdb_models/models/empagliflozin/data/empagliflozin/ElDash2021/.ElDash2021_Fig4A.tsv	814ff7de2b71...
pkdb_models/models/empagliflozin/data/empagliflozin/ElDash2021/.ElDash2021_Tab2.tsv	e225cf2b5de6...
pkdb_models/models/empagliflozin/data/empagliflozin/ElDash2021/.ElDash2021_Tab2A.tsv	0fd121878a39...
pkdb_models/models/empagliflozin/data/empagliflozin/ElDash2021/.ElDash2021_Tab3.tsv	75e407f845f1...
pkdb_models/models/empagliflozin/data/empagliflozin/ElDash2021/.ElDash2021_TabGroups.tsv	4e30f7c4ce3e...

File	SHA-256
pkdb_models/models/empagliflozin/data/empagliflozin/ElDash2021/.ElDash2021_TabIndividuals.tsv	56829d1f73fc...
pkdb_models/models/empagliflozin/data/empagliflozin/ElDash2021/.ElDash2021_TabInterventions.tsv	4b1eda3294a7...
pkdb_models/models/empagliflozin/data/empagliflozin/Friedrich2013/.Friedrich2013_Fig2.tsv	ce73993247d4...
pkdb_models/models/empagliflozin/data/empagliflozin/Friedrich2013/.Friedrich2013_Fig3.tsv	90be18b86d0b...
pkdb_models/models/empagliflozin/data/empagliflozin/Friedrich2013/.Friedrich2013_Fig4.tsv	7d30a98b6ffe...
pkdb_models/models/empagliflozin/data/empagliflozin/Friedrich2013/.Friedrich2013_Tab1.tsv	d1d9552db4ea...
pkdb_models/models/empagliflozin/data/empagliflozin/Friedrich2013/.Friedrich2013_TabGroups.tsv	f8ea11cf65e0...
pkdb_models/models/empagliflozin/data/empagliflozin/Friedrich2013/.Friedrich2013_TabInterventions.tsv	434be09ad515...
pkdb_models/models/empagliflozin/data/empagliflozin/Hailat2022/.Hailat2022_Fig3.tsv	139bc766d99d...
pkdb_models/models/empagliflozin/data/empagliflozin/Hailat2022/.Hailat2022_Tab1.tsv	e0b77505411f...
pkdb_models/models/empagliflozin/data/empagliflozin/Hailat2022/.Hailat2022_TabGroups.tsv	8b29e335ec42...
pkdb_models/models/empagliflozin/data/empagliflozin/Hailat2022/.Hailat2022_TabInterventions.tsv	3876a660e5fb...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013/.Heise2013_Fig1.tsv	6413d99630e7...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013/.Heise2013_Fig2.tsv	b0d17dede829...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013/.Heise2013_Fig3.tsv	ed0a136d5efc...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013/.Heise2013_Fig4.tsv	a620d58de7e6...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013/.Heise2013_Tab6.tsv	ff90e467e3d3...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013/.Heise2013_TabGroups.tsv	a9d281bb0721...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013/.Heise2013_TabInterventions.tsv	0fbec9a9334c...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013a/.Heise2013a_Fig1.tsv	d7ef90c081ca...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013a/.Heise2013a_Fig2.tsv	66764cc30622...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013a/.Heise2013a_Fig3.tsv	e992233a9aed...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013a/.Heise2013a_Fig4.tsv	f11821054eb6...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013a/.Heise2013a_Fig5.tsv	198d2c12a5e2...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013a/.Heise2013a_Tab2.tsv	6c183fd83c95...
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013a/.Heise2013a_TabGroups.tsv	31996969859e...

File	SHA-256
pkdb_models/models/empagliflozin/data/empagliflozin/Heise2013a/.Heise2013a_TabInterventions.tsv	7213cbe2ee96...
pkdb_models/models/empagliflozin/data/empagliflozin/Jiang2023b/.Jiang2023b_Fig2.tsv	bf9f8abfed43...
pkdb_models/models/empagliflozin/data/empagliflozin/Jiang2023b/.Jiang2023b_Fig3.tsv	d0e372fceddf...
pkdb_models/models/empagliflozin/data/empagliflozin/Jiang2023b/.Jiang2023b_Tab2.tsv	aee5392d871d...
pkdb_models/models/empagliflozin/data/empagliflozin/Jiang2023b/.Jiang2023b_TabGroups.tsv	c2d1464f4014...
pkdb_models/models/empagliflozin/data/empagliflozin/Jiang2023b/.Jiang2023b_TabIndividuals.tsv	659b312ed557...
pkdb_models/models/empagliflozin/data/empagliflozin/Jiang2023b/.Jiang2023b_TabInterventions.tsv	892e79317068...
pkdb_models/models/empagliflozin/data/empagliflozin/Kim2021/.Kim2021_Fig3.tsv	15c006f54d5b...
pkdb_models/models/empagliflozin/data/empagliflozin/Kim2021/.Kim2021_Fig4.tsv	b9e087186bd9...
pkdb_models/models/empagliflozin/data/empagliflozin/Kim2021/.Kim2021_Tab1.tsv	93751dfde883...
pkdb_models/models/empagliflozin/data/empagliflozin/Kim2021/.Kim2021_TabGroups.tsv	273123c26d6a...
pkdb_models/models/empagliflozin/data/empagliflozin/Kim2021/.Kim2021_TabIndividuals.tsv	0bd41eeee9de...
pkdb_models/models/empagliflozin/data/empagliflozin/Kim2021/.Kim2021_TabInterventions.tsv	4c4dad71776d...
pkdb_models/models/empagliflozin/data/empagliflozin/Laffel2018/.Laffel2018_Fig1.tsv	ecc2db7ac712...
pkdb_models/models/empagliflozin/data/empagliflozin/Laffel2018/.Laffel2018_Fig2.tsv	2b0f88f25bf3...
pkdb_models/models/empagliflozin/data/empagliflozin/Laffel2018/.Laffel2018_Fig3.tsv	b25c2a39c8f6...
pkdb_models/models/empagliflozin/data/empagliflozin/Laffel2018/.Laffel2018_Tab2.tsv	29a57164aeee...
pkdb_models/models/empagliflozin/data/empagliflozin/Laffel2018/.Laffel2018_Tab3.tsv	5dfd88cb9326...
pkdb_models/models/empagliflozin/data/empagliflozin/Laffel2018/.Laffel2018_TabGroups.tsv	0bdd74a75fb5...
pkdb_models/models/empagliflozin/data/empagliflozin/Laffel2018/.Laffel2018_TabInterventions.tsv	42e5e466779f...
pkdb_models/models/empagliflozin/data/empagliflozin/Li2020/.Li2020_Fig2.tsv	f6a0bc9ad8e0...
pkdb_models/models/empagliflozin/data/empagliflozin/Li2020/.Li2020_Fig3.tsv	13a4e31ce31c...
pkdb_models/models/empagliflozin/data/empagliflozin/Li2020/.Li2020_Tab3.tsv	f4f5e90d19a1...
pkdb_models/models/empagliflozin/data/empagliflozin/Li2020/.Li2020_TabGroups.tsv	e40e7e0fc7fc...
pkdb_models/models/empagliflozin/data/empagliflozin/Li2020/.Li2020_TabInterventions.tsv	dc30a2e7e11d...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2013e/.Macha2013e_Fig2.tsv	5eba5b683159...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2013e/.Macha2013e_Fig3.tsv	a7baa9ad84c1...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2013e/.Macha2013e_Tab1.tsv	5bba81ffec82...

File	SHA-256
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2013e/.Macha2013e_Tab3.tsv	7896cd6a0da5...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2013e/.Macha2013e_Tab4.tsv	f5441082ceda...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2013e/.Macha2013e_TabGroups.tsv	469a88aeb4e7...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2013e/.Macha2013e_TabInterventions.tsv	aeedec7c1ba5...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014/.Macha2014_Fig2.tsv	64103c1b3835...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014/.Macha2014_Tab1.tsv	482297f66665...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014/.Macha2014_Tab3.tsv	c89423f30f44...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014/.Macha2014_TabGroups.tsv	aa804cc647f2...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014/.Macha2014_TabInterventions.tsv	977f0b85a2db...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_Fig1.tsv	7984385f17ef...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_Fig2.tsv	5f6cb4a3b7a4...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_Fig3.tsv	d4cfdc0a58e0...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_Fig4.tsv	3eb1173888bf...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_Tab2.tsv	31975c12e873...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_Tab2A.tsv	bd22c15fe9b7...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_Tab3.tsv	9635c12ff03b...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_TabGroups.tsv	ed54e906b8bc...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014b/.Macha2014b_TabInterventions.tsv	603876050f2d...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_Fig1.tsv	486c6e92cd65...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_Fig2.tsv	45fb47931361...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_Fig3.tsv	06452568f722...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_Tab2.tsv	e07df4ea2ace...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_Tab2A.tsv	4818f5711ffc...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_Tab3.tsv	baa87c63714f...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_TabGroups.tsv	da12d850fb3a...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_TabIndividuals.tsv	8ad7624005d0...

File	SHA-256
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2014f/.Macha2014f_TabInterventions.tsv	c031bba350a4...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015a/.Macha2015a_Fig1.tsv	64dc19c03c72...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015a/.Macha2015a_Fig2.tsv	013016491f42...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015a/.Macha2015a_Tab1.tsv	649990a63a30...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015a/.Macha2015a_Tab3.tsv	0dbd04083a8c...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015a/.Macha2015a_TabGroups.tsv	255c9ce6f0ee...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015a/.Macha2015a_TabInterventions.tsv	f2e343c95254...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015b/.Macha2015b_Fig1.tsv	3188f476d4ff...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015b/.Macha2015b_Fig2.tsv	8c345124298b...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015b/.Macha2015b_Tab2.tsv	0cce3a48008b...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015b/.Macha2015b_Tab3.tsv	26fcd8a60d80...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015b/.Macha2015b_TabGroups.tsv	dbb37920ec4d...
pkdb_models/models/empagliflozin/data/empagliflozin/Macha2015b/.Macha2015b_TabInterventions.tsv	3aa158e0c03b...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2013/.Sarashina2013_Fig1.tsv	625ee750d33c...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2013/.Sarashina2013_Fig2.tsv	be4ca6647519...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2013/.Sarashina2013_Tab3.tsv	e7344722f126...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2013/.Sarashina2013_Tab3A.tsv	e94b66f82261...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2013/.Sarashina2013_Tab4.tsv	167a08800632...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2013/.Sarashina2013_TabGroups.tsv	b6589c4c6e2e...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2013/.Sarashina2013_TabInterventions.tsv	f34554ff6454...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Fig1.tsv	c15ac607e233...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Fig1a.tsv	95428928bacb...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Fig1b.tsv	2d77b9683cf3...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Fig2.tsv	60a1f380087e...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Fig3.tsv	ea969a16f891...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Tab2.tsv	26a04cc5bcec...

File	SHA-256
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Tab2A.tsv	28e5726006c5...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Tab3.tsv	43e1c2002719...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_Tab4.tsv	43e1c2002719...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_TabGroups.tsv	5239cd993a79...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_TabIndividuals.tsv	d3ba6b5a9c42...
pkdb_models/models/empagliflozin/data/empagliflozin/Sarashina2014/.Sarashina2014_TabInterventions.tsv	8f4432e2a943...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_Fig1.tsv	6689b6ff1369...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_Fig1B.tsv	cbfa4e727ac3...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_Fig2.tsv	26e9fb34bb19...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_Fig3.tsv	8f6f4a009c76...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_Fig4.tsv	2855782ec180...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_FigFDA.tsv	3307049f5fb2...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_FigS2.tsv	f6220577dd43...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_Tab2.tsv	58b3dae648c4...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_TabGroups.tsv	dbdbdc5278c5...
pkdb_models/models/empagliflozin/data/empagliflozin/Seman2013/.Seman2013_TabInterventions.tsv	1aa149a8084d...
pkdb_models/models/empagliflozin/data/empagliflozin/Zhao2015/.Zhao2015_Fig1.tsv	a4b3adb4b3b2...
pkdb_models/models/empagliflozin/data/empagliflozin/Zhao2015/.Zhao2015_Fig2.tsv	1e60ba0b71b2...
pkdb_models/models/empagliflozin/data/empagliflozin/Zhao2015/.Zhao2015_Fig3.tsv	22e8fc0b167e...
pkdb_models/models/empagliflozin/data/empagliflozin/Zhao2015/.Zhao2015_Tab2.tsv	f50819960480...
pkdb_models/models/empagliflozin/data/empagliflozin/Zhao2015/.Zhao2015_Tab2A.tsv	898bbb3de684...
pkdb_models/models/empagliflozin/data/empagliflozin/Zhao2015/.Zhao2015_TabGroups.tsv	10036e554ebe...
pkdb_models/models/empagliflozin/data/empagliflozin/Zhao2015/.Zhao2015_TabInterventions.tsv	580c944a9681...
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_Fig2.tsv	9a4c50ada866...
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_Fig3.tsv	2c65a0ec9ef3...
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_Fig4.tsv	8a0cb6a070e5...
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_Tab1.tsv	9af04c16cfb8...
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_Tab3.tsv	55f53bccd224...

File	SHA-256
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_Tab3A.tsv	0de829681d4f...
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_Tab4.tsv	4dbd8f06fbf0...
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_TabGroups.tsv	d77f4251d594...
pkdb_models/models/empagliflozin/data/hydrochlorothiazide/Heise2015/.Heise2015_TabInterventions.tsv	36a7fdc8e888...
pkdb_models/models/empagliflozin/data/ramipril/Macha2013/.Macha2013_Fig1.tsv	85e4de7ac42e...
pkdb_models/models/empagliflozin/data/ramipril/Macha2013/.Macha2013_Fig2.tsv	8f2eeeb33441...
pkdb_models/models/empagliflozin/data/ramipril/Macha2013/.Macha2013_Fig3.tsv	c82a126e2ac4...
pkdb_models/models/empagliflozin/data/ramipril/Macha2013/.Macha2013_Tab1.tsv	5ea0ef78b354...
pkdb_models/models/empagliflozin/data/ramipril/Macha2013/.Macha2013_Tab2.tsv	6147a2f4e148...
pkdb_models/models/empagliflozin/data/ramipril/Macha2013/.Macha2013_Tab3.tsv	621e637eef09...
pkdb_models/models/empagliflozin/data/ramipril/Macha2013/.Macha2013_TabGroups.tsv	2d2a7b78ab5e...
pkdb_models/models/empagliflozin/data/ramipril/Macha2013/.Macha2013_TabInterventions.tsv	8d1bec3da575...
pkdb_models/models/empagliflozin/models/output/models/empagliflozin_body_flat.xml	c4b7be33d3dc...

2 Results: Empagliflozin PBPK/PD model: fit PK

2.1 Parameters

Table 2: Parameters of the fit PK_20261019_201013__13ddc, the parameter set PK_20261019_201013__13ddc_0, cost 76.9, with their bounds in the optimization. The 95% confidence intervals are those of the profile likelihood, LB and UB an interval which is open towards the lower or the upper bound, and the Wald intervals of the Fisher information for a parameter without a profile. Identifiable is the classification of the profile: yes, no (LB) or no (UB) if the profile does not reach the threshold towards the lower or the upper bound, no if it reaches it on neither side, structural if it is flat.

Parameter	Description	Value [95% CI]	Unit	Bounds	Identifiable
ftissue_emp	tissue distribution emp	1.11 [0.379, UB]	l/min	[0.01, 5]	no (UB)
Kp_emp	tissue/plasma partition coefficient emp	0.238 [0.0612, 0.334]	dimensionless	[0.05, 10]	yes
ftissue_deep_emp	deep tissue distribution emp	0.113 [0.0123, 0.314]	l/min	[0.001, 0.5]	yes
Kp_deep_emp	deep tissue/plasma partition coefficient emp	0.419 [0.183, UB]	dimensionless	[0.05, 1]	no (UB)
GU__EMPABS_k	rate of empagliflozin absorption	0.0129 [0.00844, 0.0297]	1/min	[0.001, 0.1]	yes
GU__F_emp_abs	fraction absorbed empagliflozin	0.438 [0.406, 0.49]	dimensionless	[0.35, 0.85]	yes
GU__METEXC_k	rate of the fecal transit	0.000786 [0.000633, 0.00105]	1/min	[0.0001, 0.01]	yes
GU__FED_f_absorption	fed effect on the absorption rate	0.55 [0.354, 0.68]	dimensionless	[0.1, 1]	yes

Parameter	Description	Value [95% CI]	Unit	Bounds	Identifiable
EMP2EG_k	rate of the hepatic glucuronidation of empagliflozin	0.0176 [0.0151, 0.0206]	1/min	[0.001, 0.5]	yes
KI__EMPSEC_f	tubular secretion of empagliflozin relative to its filtration	1.31 [1.05, 1.55]	dimensionless	[0.05, 5]	yes
KI__EGEX_f	renal clearance of EG relative to the GFR	1.66 [1.04, 3.23]	dimensionless	[0.3, 20]	yes

2.2 Data

Table 3: Data of the fit PK_20261019_201013__13ddc: the number of fit mappings and, in parentheses, of data points per study and kind. The training data enters the cost, the validation data and the outliers are evaluated only.

Study	Training	Validation	Outlier	Total
Ayoub2017	1 (9)	-	-	1 (9)
Brand2012	2 (20)	2 (20)	-	4 (40)
Chen2015a	4 (23)	4 (46)	1 (12)	9 (81)
Chen2020	2 (36)	-	-	2 (36)
ElDash2021	1 (17)	1 (17)	-	2 (34)
Friedrich2013	2 (19)	2 (19)	-	4 (38)
Hailat2022	-	-	4 (59)	4 (59)
Heise2013	4 (147)	-	-	4 (147)
Heise2013a	3 (100)	-	-	3 (100)
Heise2015	1 (13)	2 (26)	-	3 (39)
Jiang2023b	1 (16)	-	-	1 (16)
Kim2021	1 (14)	1 (14)	-	2 (28)
Kim2023	1 (13)	1 (13)	-	2 (26)
Laffel2018	-	3 (29)	-	3 (29)
Li2020	1 (16)	1 (16)	-	2 (32)
Macha2013d	2 (31)	2 (32)	-	4 (63)
Macha2013e	1 (13)	1 (18)	-	2 (31)
Macha2014	3 (31)	4 (48)	-	7 (79)
Macha2014b	4 (27)	12 (86)	-	16 (113)
Macha2014f	2 (19)	8 (80)	-	10 (99)
Macha2015a	2 (36)	-	-	2 (36)
Macha2015b	1 (15)	1 (15)	-	2 (30)
Sarashina2013	10 (83)	-	-	10 (83)
Sarashina2014	-	8 (76)	-	8 (76)
Seman2013	16 (212)	4 (52)	-	20 (264)
vanderAartvanderBeek2020	-	4 (31)	2 (12)	6 (43)
Zhao2015	-	8 (76)	-	8 (76)
Total	65 (910)	69 (714)	7 (83)	141 (1707)

2.3 Goodness of fit

Table 4: Goodness of fit of the fit PK_20261019_201013__13ddc per kind of data, the parameter set PK_20261019_201013__13ddc_0. The cost is defined on the training data only, 2-fold is the percentage of the data points with a positive value and prediction whose prediction is within twofold of the data.

Kind	n	cost	MSE	RMSE	NRMSE	RMSE_w	R^2	2-fold [%]	AIC	BIC
training	910	76.9	1.94e-06	0.00139	0.305	0.411	0.974	91.8	−11945.1	−11892.2
validation	714	-	0.00015	0.0123	0.961	2.26	0.632	88.9	−6262.6	−6212.4
outlier	83	-	1.41e-08	0.000119	3.32	7.22	−0.848	28.4	−1478.5	−1451.9

2.4 Figures




Figure 1: Cost of the optimizers over their steps, fit PK_20261019_201013__13ddc. The cost of every optimization run over its steps. Runs which end at the same cost found the same optimum; a run which stops much higher is stuck in a local minimum.




Figure 2: Cost of the optimization runs, ordered, fit PK_20261019_201013__13ddc. The final cost of the runs, ordered. A flat plateau at the left is the global optimum found repeatedly, which is the evidence that the multistart converged; steps to the right are local minima.




Figure 3: Prediction against the measured data points, per kind of fit mapping, with the agreement of the training data, the lines of twofold and the metrics of the panel, fit PK_20261019_201013__13ddc. Within twofold of the data are 820 of the 893 training data points (91.8%), 616 of the 693 validation data points (88.9%) and 21 of the 74 outlier data points (28.4%); only the data points with a positive value and prediction are counted. Prediction against measurement, one point per data point, with a panel per kind of fit mapping. The points scatter around the diagonal when the model describes the data; a systematic deviation from it is a systematic error of the model. The band is the agreement of the training data, i.e. the same band the Bland-Altman plot draws, here as lines parallel to the diagonal. The dark red dotted lines are twofold, prediction = 2 x measurement and prediction = measurement / 2; the caption states the points between them of every kind. A point which is zero or negative is not drawn on the logarithmic axes and not counted, the title of a panel then states the points drawn of all. The validation panel shows how the fit describes data it was not fitted on, the outlier panel the data the fit dropped. The box of a panel carries its R^2, its normalized RMSE, the RMSE of the residuals of the cost, its fold errors GMFE and AFE and the percentages of its points within twofold and within two standard deviations of the data.




Figure 4: Agreement of prediction and measurement as a ratio, per kind of fit mapping, with the agreement of the training data and the agreement of the panel, fit PK_20261019_201013__13ddc. The ratio of prediction and measurement over the geometric mean of the two, i.e. the goodness of fit with the diagonal turned into the horizontal. The band is the same, the bias and the limits of agreement `bias ± 1.96 SD` of the training data as fold factors. A bias away from 1 is a systematic over- or underprediction, wide limits are a large scatter, and a trend over the mean is a model which describes the large or the small values better. The box of a panel carries the bias and the SD of its own points as fold factors and the share of them inside the limits of agreement of the training data.




Figure 5: Profile likelihood of every parameter: the cost with the parameter fixed and the other parameters optimized, the threshold of the confidence level and the confidence interval, fit PK_20261019_201013__13ddc. One profile per parameter. A profile which crosses the dashed threshold on both sides gives a finite confidence interval; one which stays below it up to a bound is practically non-identifiable; a flat profile is structurally non-identifiable, the parameter is compensated by the others.

2.5 Provenance

Table 5: Provenance of the fit PK_20261019_201013__13ddc: the versions it ran with and the revision of its models and data; the SHA-256 checksums of the models and data are listed in Table 1.

Item	Value
Fit	`PK_20261019_201013__13ddc`
Created	2026-10-19 18:10:18 UTC
Seed	1234
Python	3.14.4

Item	Value
sbmlsim	0.10.0
libRoadRunner	2.10.0
SciPy	1.18.1
NumPy	2.5.3
Repository	`/home/mkoenig/git/pkdb_models-empagliflozin`
Commit	`f43c98cc20b2368500dfcbc3ee1739eeb596f460`
Uncommitted changes	no

Bibliography

[1] M. König, "sbmlsim: SBML simulation made easy." Zenodo, 2026. doi: 10.5281/zenodo.3597770. Version 0.10.0.

[2] R. Storn and K. Price, "Differential evolution - a simple and efficient heuristic for global optimization over continuous spaces," *Journal of Global Optimization*, vol. 11, pp. 341–359, 1997, doi: 10.1023/A:1008202821328.

[3] P. Virtanen *et al.*, "SciPy 1.0: fundamental algorithms for scientific computing in Python," *Nature Methods*, vol. 17, pp. 261–272, 2020, doi: 10.1038/s41592-019-0686-2. Version 1.18.1.

[4] M. D. McKay, R. J. Beckman, and W. J. Conover, "A comparison of three methods for selecting values of input variables in the analysis of output from a computer code," *Technometrics*, vol. 21, no. 2, pp. 239–245, 1979, doi: 10.2307/1268522.

[5] C. Welsh, J. Xu, L. Smith, M. König, K. Choi, and H. M. Sauro, "libRoadRunner 2.0: a high performance SBML simulation and analysis library," *Bioinformatics*, vol. 39, no. 1, p. btac770, 2023, doi: 10.1093/bioinformatics/btac770. Version 2.10.0.

[6] A. C. Hindmarsh *et al.*, "SUNDIALS: suite of nonlinear and differential/algebraic equation solvers," *ACM Transactions on Mathematical Software*, vol. 31, no. 3, pp. 363–396, 2005, doi: 10.1145/1089014.1089020.

[7] H. Akaike, "A new look at the statistical model identification," *IEEE Transactions on Automatic Control*, vol. 19, no. 6, pp. 716–723, 1974, doi: 10.1109/TAC.1974.1100705.

[8] G. Schwarz, "Estimating the dimension of a model," *The Annals of Statistics*, vol. 6, no. 2, pp. 461–464, 1978, doi: 10.1214/aos/1176344136.

[9] J. M. Bland and D. G. Altman, "Statistical methods for assessing agreement between two methods of clinical measurement," *The Lancet*, vol. 327, no. 8476, pp. 307–310, 1986, doi: 10.1016/S0140-6736(86)90837-8.

[10] A. Raue *et al.*, "Structural and practical identifiability analysis of partially observed dynamical models by exploiting the profile likelihood," *Bioinformatics*, vol. 25, no. 15, pp. 1923–1929, 2009, doi: 10.1093/bioinformatics/btp358.

[11] C. Kreutz, A. Raue, D. Kaschek, and J. Timmer, "Profile likelihood in systems biology," *The FEBS Journal*, vol. 280, no. 11, pp. 2564–2571, 2013, doi: 10.1111/febs.12276.

[12] F.-G. Wieland, A. L. Hauber, M. Rosenblatt, C. Tönsing, and J. Timmer, "On structural and practical identifiability," *Current Opinion in Systems Biology*, vol. 25, pp. 60–69, 2021, doi: 10.1016/j.coisb.2021.03.005.

[13] C. R. Harris *et al.*, "Array programming with NumPy," *Nature*, vol. 585, pp. 357–362, 2020, doi: 10.1038/s41586-020-2649-2. Version 2.5.3.