

# Alterations in Postprandial Hepatic Glycogen Metabolism in Type 2 Diabetes

Martin Krssak,<sup>1</sup> Attila Brehm,<sup>1</sup> Elisabeth Bernroider,<sup>1</sup> Christian Anderwald,<sup>1</sup> Peter Nowotny,<sup>1</sup> Chiara Dalla Man,<sup>2</sup> Claudio Cobelli,<sup>2</sup> Gary W. Cline,<sup>3</sup> Gerald I. Shulman,<sup>3,4</sup> Werner Waldhäusl,<sup>1</sup> and Michael Roden<sup>1,5</sup>

Decreased skeletal muscle glucose disposal and increased endogenous glucose production (EGP) contribute to postprandial hyperglycemia in type 2 diabetes, but the contribution of hepatic glycogen metabolism remains uncertain. Hepatic glycogen metabolism and EGP were monitored in type 2 diabetic patients and nondiabetic volunteer control subjects (CON) after mixed meal ingestion and during hyperglycemic-hyperinsulinemic-somatostatin clamps applying <sup>13</sup>C nuclear magnetic resonance spectroscopy (NMRS) and variable infusion dual-tracer technique. Hepatocellular lipid (HCL) content was quantified by <sup>1</sup>H NMRS. Before dinner, hepatic glycogen was lower in type 2 diabetic patients (227 ± 6 vs. CON: 275 ± 10 mmol/l liver, *P* < 0.001). After meal ingestion, net synthetic rates were 0.76 ± 0.16 (type 2 diabetic patients) and 1.36 ± 0.15 mg · kg<sup>-1</sup> · min<sup>-1</sup> (CON, *P* < 0.02), resulting in peak concentrations of 283 ± 15 and 360 ± 11 mmol/l liver. Postprandial rates of EGP were ~0.3 mg · kg<sup>-1</sup> · min<sup>-1</sup> (30–170 min; *P* < 0.05 vs. CON) higher in type 2 diabetic patients. Under clamp conditions, type 2 diabetic patients featured ~54% lower (*P* < 0.03) net hepatic glycogen synthesis and ~0.5 mg · kg<sup>-1</sup> · min<sup>-1</sup> higher (*P* < 0.02) EGP. Hepatic glucose storage negatively correlated with HCL content (*R* = -0.602, *P* < 0.05). Type 2 diabetic patients exhibit 1) reduction of postprandial hepatic glycogen synthesis, 2) temporarily impaired suppression of EGP, and 3) no normalization of these defects by controlled hyperglycemic hyperinsulinemia. Thus, impaired insulin sensitivity and/or chronic glucolipotoxicity in addition to the effects of an altered insulin-to-glucagon ratio or increased free fatty acids accounts for defective hepatic glycogen metabolism in type 2 diabetic patients. *Diabetes* 53:3048–3056, 2004

From the <sup>1</sup>Department of Internal Medicine 3, Division of Endocrinology and Metabolism, Medical University of Vienna, Vienna, Austria; the <sup>2</sup>Department of Electronics and Informatics, University of Padova, Padova, Italy; the <sup>3</sup>Department of Internal Medicine, Howard Hughes Medical Institute, Yale University School of Medicine, New Haven, Connecticut; the <sup>4</sup>Cellular and Molecular Physiology Department, Howard Hughes Medical Institute, Yale University School of Medicine, New Haven, Connecticut; and the <sup>5</sup>Medical Department, Hanusch Hospital, Vienna, Austria.

Address correspondence and reprint requests to Michael Roden, MD, Medical Department, Hanusch Hospital, Heinrich Collinstrasse 30, A-1140 Vienna, Austria. E-mail: michael.roden@meduniwien.ac.at.

Received for publication 23 June 2004 and accepted in revised form 1 September 2004.

M.K. and A.B. contributed equally to this work.

EGP, endogenous glucose production; FFA, free fatty acid; HCL, hepatocellular lipid; NMRS, nuclear magnetic resonance spectroscopy.

© 2004 by the American Diabetes Association.

Under normal life conditions, meals are ingested every few hours, therefore rendering humans in the postprandial state for approximately half of the day. Knowledge of glucose metabolism after meal ingestion is central to the understanding of the pathophysiology and characterization of potential therapeutic targets in the diabetic state. Many previous studies in type 2 diabetic patients demonstrated defects in skeletal muscle glucose metabolism during experimental hyperinsulinemic hyperglycemia (1,2) and recently also after meal ingestion (3).

Nevertheless, the liver is almost exclusively responsible for endogenous glucose production (EGP), which correlates with the degree of hyperglycemia after an overnight fast (4–8) and is less suppressed in type 2 diabetic patients after meal ingestion (6,9–13). Invasive hepatic vein catheter studies suggested that splanchnic glucose uptake is decreased in type 2 diabetic patients (14,15). Impaired insulin-stimulated hepatic glucose uptake was confirmed in type 2 diabetic patients by positron emission tomography (16). Using a needle biopsy technique, Beringer and Thaler (17) demonstrated decreased hepatic glycogen concentrations in elderly diabetic patients. Noninvasive <sup>13</sup>C nuclear magnetic resonance spectroscopy (NMRS) studies found lower hepatic glycogen breakdown and increased gluconeogenesis in type 2 diabetic patients (18,19), but the contribution of liver glycogen metabolism to postprandial hyperglycemia and its sensitivity to insulin in type 2 diabetic patients remains uncertain.

Under steady-state conditions (clamp technique), the portal vein insulin, glucagon, and free fatty acid (FFA) concentrations regulate hepatic glucose metabolism in nondiabetic humans (20–25). Furthermore, increased liver fat accumulation (hepatocellular lipid [HCL] content) was found to correlate negatively with peripheral insulin sensitivity and insulin-dependent suppression of EGP (7,26). However, the impact of these factors on hepatic glycogen metabolism in type 2 diabetic patients is uncertain.

Thus, this study examined hepatic glycogen metabolism in type 2 diabetic patients and matched nondiabetic volunteers 1) after mixed meal ingestion and 2) during hyperglycemic-hyperinsulinemic-somatostatin clamps. We used noninvasive <sup>13</sup>C NMRS (18,27,28) to directly quantify net postprandial hepatic glycogen synthesis (protocol A) and variable infusion dual-tracer technique (6,27,29) to assess the postprandial time course of EGP (protocol B).




FIG. 1. Time course of ratio between plasma [6,6- $^2\text{H}_2$ ]glucose concentration and endogenous glucose concentration during the mixed-meal test in type 2 diabetic patients ( $\circ$ ) and CON ( $\blacktriangle$ ).

For evaluation of the role of hepatic insulin sensitivity in type 2 diabetic patients, we also measured flux through hepatic glycogen synthase, simultaneous flux through hepatic glycogen phosphorylase, and EGP under conditions of matched hyperglycemia, hyperinsulinemia, hypoglycagonemia, and decreased plasma FFA concentrations (clamp test, protocol C).

## RESEARCH DESIGN AND METHODS

We studied seven type 2 diabetic patients (five men/two women, age  $56 \pm 3$  years, BMI  $26.9 \pm 0.6$  kg/m $^2$ , known diabetes duration  $7 \pm 2$  years, HbA $_{1c}$   $7.1 \pm 0.3\%$ , triglycerides  $254 \pm 49$  mg/dl) and seven glucose-tolerant humans (five men/two women, age  $49 \pm 2$  years, BMI  $25.8 \pm 0.9$  kg/m $^2$ , HbA $_{1c}$   $5.2 \pm 0.1\%$ , triglycerides  $139 \pm 32$  mg/dl) during mixed-meal protocols (protocols A and B). Subsequently, we studied six type 2 diabetic patients (five men/one woman, age  $53 \pm 4$  years, BMI  $26.1 \pm 0.7$  kg/m $^2$ , known diabetes duration  $6 \pm 2$  years, HbA $_{1c}$   $7.4 \pm 0.1\%$ , triglycerides  $221 \pm 15$  mg/dl) and six healthy humans (four men/two women, age  $55 \pm 4$  years, BMI  $27.5 \pm 0.7$  kg/m $^2$ , HbA $_{1c}$   $5.4 \pm 0.1\%$ , triglycerides  $102 \pm 17$  mg/dl) during hyperglycemic-hyperinsulinemic clamp tests (protocol C). Four type 2 diabetic patients and four nondiabetic volunteer control subjects (CON) participated in all three protocols. Type 2 diabetic patients had no history of insulin or thiazolidinediones therapy and stopped their hypoglycemic agents for at least 3 days before each protocol. Female participants were either postmenopausal or were studied in the first half of their menstrual cycle. All volunteers were on an isocaloric diet and refrained from physical exercise for at least 3 days before the protocols, which were spaced by at least 3-week intervals. Informed consent to protocols, approved by the local ethics board, was obtained from all participants. **Mixed-meal tests (protocols A and B).** All volunteers were examined under identical postprandial conditions, once for liver glycogen measurement (protocol A) and again for assessing EGP (protocol B). On both occasions, they were served breakfast (231 kcal, 7:00 A.M.), lunch (208 kcal, 11:00 A.M.), and dinner (400 ml liquid meal, 652 kcal; carbohydrate/protein/fat: 83.6 g/23.2 g/24.0 g, 5:00 P.M.).

**Protocol A.** Hepatic glycogen concentrations were measured using  $^{13}\text{C}$  NMRS (28,30) before (4:00–4:30 P.M.) and after dinner (6:00–7:00 P.M., 8:00–9:00 P.M., 10:00–11:00 P.M., 12:00–1:00 A.M., and 7:00–8:00 A.M.). Liver volume measurements were performed independently on a clinical 1.5-T MR scanner (24).

**Protocol B.** At 3:00 P.M., a primed (20  $\mu\text{mol/kg}$  per 5 mmol/l) continuous (0.020  $\mu\text{mol} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ) infusion of [6,6- $^2\text{H}_2$ ]glucose (99% enriched) was started (28). At 5:00 P.M. (0 min), the test meal was ingested in which 10 g glucose were replaced by [1- $^{13}\text{C}$ ]glucose (99% enriched). After dinner, the [6,6- $^2\text{H}_2$ ]glucose infusion rate was stepwise adjusted according to a protocol developed in pilot experiments ( $n = 5$ ), which was then applied in both groups (0–3 min: 100%; 3–8 min: 60%; 8–18 min: 20%; 18–110 min: 10%; 110–180 min: 20%; 180–280 min: 30%; 280–320 min: 50%; 320–380 min: 60%; 380–480 min: 70%). This infusion protocol aimed at minimizing variations of the tracer-to-tracee ratios of endogenous glucose (Fig. 1).

The percent contribution of the direct (glucose  $\rightarrow$  glucose-6-phosphate  $\rightarrow$  UDPglucose  $\rightarrow$  glycogen) and indirect (pyruvate  $\rightarrow$  glucose-6-phosphate  $\rightarrow$  glycogen) pathways to glycogen synthesis was determined using the glucuronide probe technique with 500 mg acetaminophen given orally at 4:30 P.M. (28). From 4:45 P.M. to 1:00 A.M., 22 plasma samples were drawn for measurement of  $^{13}\text{C}$  and  $^2\text{H}$  glucose enrichments.

**Clamp test (protocol C).** Type 2 diabetic patients were admitted the evening before the study, and plasma glucose concentrations were normalized overnight using intravenous insulin infusion (Actrapid; Novo Nordisk, Vienna, Austria) (28). Nondiabetic volunteers were admitted at 6:00 A.M. the day of the study.

At 6:30 A.M. (–150 min), a primed-continuous infusion of [6,6- $^2\text{H}_2$ ]glucose was started for measurement of EGP (28). At 9:00 A.M. (0 min), a hyperglycemic-hyperinsulinemic-pancreatic clamp test was initiated by somatostatin (–5 to 300 min:  $0.1 \mu\text{g} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ; UCB Pharma, Vienna, Austria) and insulin (0–8 min:  $80 \text{ mU} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$  body surface area; 8–300 min:  $40 \text{ mU} \cdot \text{min}^{-1} \cdot \text{m}^{-2}$  body surface area; Actrapid) infusion. Plasma glucose was raised and maintained at  $\sim 180$  mg/dl by primed (0.2 g/kg) variable dextrose infusion (20% wt/vol) enriched with [1- $^{13}\text{C}$ ]glucose (20% wt/wt) and [6,6- $^2\text{H}_2$ ]glucose (2% wt/wt). For assessment of the flux through hepatic glycogen phosphorylase, the [1- $^{13}\text{C}$ ]glucose-labeled infusion was switched to natural abundance [1- $^{12}\text{C}$ ]glucose infusion at 150 min (21–23). Hepatic glycogen concentrations were measured before the clamps (–30 to 0 min), from 60 to 150 min and from 210 to 300 min during the clamps with  $^{13}\text{C}$  NMRS, as previously published (28,30). Contributions of the direct and gluconeogenic pathways to glycogen synthesis were assessed with 1,000 mg acetaminophen given orally at –30 min (27,28,31). HCL content was measured before the clamp test. During the clamps, plasma samples for measurement of  $^{13}\text{C}$  enrichments in glucose and glucuronide and  $^2\text{H}$  enrichments in glucose were drawn at 15-min intervals.

**Liver  $^{13}\text{C}$  and  $^1\text{H}$  NMRS.** Localized  $^{13}\text{C}$  NMR spectra were obtained in the 3T/80 cm Medspec (Bruker Biospin, Ettlingen, Germany) with a 10-cm circular  $^{13}\text{C}/^1\text{H}$  transmitter/receiver coil placed over the lateral aspect of the liver, applying a modified one-dimensional inversion-based sequence (28,30). Absolute glycogen concentrations were quantified by comparing the C1 glycogen peak (100.5 ppm) integral of liver spectra with that of a glycogen standard taken under identical conditions. To eliminate the contribution of [1- $^{13}\text{C}$ ]glucose to the [1- $^{13}\text{C}$ ]glycogen spectral line intensity during labeled glucose infusion in protocol C, only the left half of the [1- $^{13}\text{C}$ ]glycogen doublet in the spectra was integrated. Corrections for loading of the coil and sensitive volume of the coil were performed (28).

Localized  $^1\text{H}$  NMR spectra of the liver were acquired using the same magnetic resonance system, coil, and patient placement inside the magnet applying the breath-hold-triggered stimulated echo acquisition mode sequence (7). HCL content was quantified by integration of the  $\text{CH}_2$  and  $\text{CH}_3$  group resonances and is expressed in arbitrary units (AU) as a percentage of total  $^1\text{H}$  NMRS signals (water + lipid).

**Analytical procedures.** Plasma glucose was measured by the glucose oxidase method (Glucose Analyzer II; Beckman Instruments, Fullerton, CA). Plasma FFAs were assayed microfluorimetrically (Wako Chemicals, Richmond, VA). Plasma immunoreactive insulin, C-peptide, and glucagon were measured by commercially available radioimmunoassays (24,32,33). Atom percent excess of  $^2\text{H}$  and  $^{13}\text{C}$  in plasma glucose and  $^{13}\text{C}$  atom percent excess in plasma acetaminophen glucuronide were quantified with gas and liquid chromatography–mass spectroscopy, respectively (28).

## Calculations

**Hepatic glycogen metabolism.** Rates of net hepatic glycogen synthesis and subsequent net glycogenolysis during protocol A were calculated from linear regression of the individual glycogen concentration time curves over respective time periods of increasing (6:00 to  $\sim$ 9:00 P.M. in CON and 6:00–10:30 P.M. in type 2 diabetic patients) and decreasing (10:00 P.M. to 8:00 A.M.) glycogen concentrations (28,30). The time point of the peak glycogen concentration

was calculated from the intersection of regression lines of increasing and decreasing glycogen concentrations.

In protocol C, flux through hepatic glycogen synthase ( $V_{\text{Syn}}$ ) was assessed using previously published equations (21–23,34) from linear regression of the increments in hepatic glycogen concentration. These increments were calculated in 7-min intervals between 60 and 150 min of [ $1\text{-}^{13}\text{C}$ ]glucose infusion from the increase of the [ $1\text{-}^{13}\text{C}$ ]glycogen peak intensity, corrected by actual  $^{13}\text{C}$  plasma glucose enrichment above the natural abundance and by the contribution of the direct glycogen synthesis pathway. The percent contribution of the direct pathway to glycogen synthesis was calculated as the ratio between differences in  $^{13}\text{C}$  enrichments at the C1 and C6 positions of the glucuronide and glucose (28).

Simultaneous flux through hepatic glycogen phosphorylase ( $V_{\text{Out}}$ ) was calculated by subtracting the observed from the predicted change in [ $1\text{-}^{13}\text{C}$ ]glycogen concentrations during the chase period (210–300 min), assuming constant synthase flux and highest possible  $^{13}\text{C}$  enrichment in the C1 glycogen (21–23,34). Relative glycogen turnover (%) was calculated as  $(V_{\text{Out}} \times 100)/V_{\text{Syn}}$ . The rate of net hepatic glycogen synthesis ( $V_{\text{SynNet}}$ ) was determined as the difference between the synthase ( $V_{\text{Syn}}$ ) and simultaneous phosphorylase ( $V_{\text{Out}}$ ) flux (21–23,34). All rates describing hepatic glycogen metabolism were converted from mmol/l liver per minute into  $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  by using individual data on liver volume and body weight, which did not affect the results of the statistical analyses.

Rates of gluconeogenesis (protocols A and B, 11:00 P.M. to 1:00 A.M.) were calculated as the difference between the mean individual rates of EGP and net hepatic glycogenolysis and are given as percent contribution to EGP (30).

**EGP protocol B.** The plasma glucose concentration after a meal is the sum of exogenous (meal-derived) and endogenous (due to EGP) glucose concentrations. The exogenous glucose can be calculated on the basis of its proportionality to meal-derived [ $1\text{-}^{13}\text{C}$ ]glucose. Subtracting exogenous glucose from total plasma glucose concentration yields the endogenous glucose concentration. Rates of appearance of endogenous glucose (EGP) were quantified with a variable infusion double-tracer technique (6,27) and calculated from plasma [ $6,6\text{-}^2\text{H}_2$ ]glucose concentrations and endogenous glucose concentrations using Steele's non-steady state equation (35). EGP is derived by first expressing the disappearance rate parameter,  $k_{01}$ , from the mass balance equation of [ $6,6\text{-}^2\text{H}_2$ ]glucose tracer and then using it in the mass balance of endogenous glucose (29). The time course of [ $6,6\text{-}^2\text{H}_2$ ]glucose concentration and the ratio of [ $6,6\text{-}^2\text{H}_2$ ]glucose to endogenous glucose concentration were smoothed using an algorithm based on stochastic nonparametric deconvolution (36).

**Protocol C.** Basal rates of EGP were calculated by dividing the tracer ([ $6,6\text{-}^2\text{H}_2$ ]glucose) infusion rate times tracer enrichment by the tracer enrichment in plasma glucose and subtracting the tracer infusion rate. Rates of EGP during the clamp test were calculated for 15-min intervals using Steele's non-steady-state equation to calculate the rate of appearance of overall glucose and subtracting the mean glucose infusion rate of that 15-min interval from the overall rate of appearance of glucose (37).

**Whole-body glucose metabolism.** During protocol C, insulin-stimulated whole-body glucose disposal was calculated from glucose infusion rates and [ $6,6\text{-}^2\text{H}_2$ ]glucose dilution, both corrected for EGP (1).

**Statistical analysis.** Data are given as means  $\pm$  SE. Student's two-tailed unpaired *t* test was used for group comparison. Two-way repeated-measures ANOVA and Bonferroni/Dunnett post hoc test were used for statistical analysis of the time course EGP. Because metabolites and hormones did not differ within the groups between the protocols, individual data of every participant at each time point from both protocols were pooled for presentation. Statistical significance was considered at  $P < 0.05$ .

## RESULTS

**Mixed-meal test.** Plasma glucose concentrations were  $\sim 69\%$  higher in type 2 diabetic patients before dinner ( $150 \pm 12$  vs. CON:  $89 \pm 2$  mg/dl,  $P < 0.001$ ) (Fig. 2A) and remained elevated throughout the study. Plasma FFA concentrations were less suppressed in type 2 diabetic patients after dinner (mean 60–240 min:  $255 \pm 21$  vs. CON:  $115 \pm 14$   $\mu\text{mol/l}$ ,  $P < 0.001$ ) (Fig. 2B). In type 2 diabetic patients, plasma insulin was higher before ( $11 \pm 1$  vs. CON:  $6 \pm 1$   $\mu\text{U/ml}$ ,  $P < 0.01$ ), but was lower after the dinner (120 min;  $36 \pm 5$  vs. CON:  $93 \pm 18$   $\mu\text{U/ml}$ ,  $P < 0.01$ ), whereas plasma glucagon was higher in type 2 diabetic patients (30 min:  $108 \pm 4$  vs. CON:  $82 \pm 6$  pg/ml,  $P < 0.01$ ) (Fig. 2C and D).

Hepatic glycogen concentrations were lower in type 2 diabetic patients before dinner (type 2 diabetic patients:  $227 \pm 6$  vs. CON:  $275 \pm 10$  mmol/l,  $P < 0.001$ , Fig. 3A) and remained unchanged for 1 h after dinner (type 2 diabetic patients:  $227 \pm 10$  vs. CON:  $265 \pm 11$  mmol/l,  $P < 0.001$ ). Liver glycogen increased afterward to mean individual peak values of  $283 \pm 15$  mmol/l in type 2 diabetic patients and  $360 \pm 11$  mmol/l in CON ( $P < 0.01$ ), with a time lag of  $\sim 55$  min between the groups (calculated peak time; type 2 diabetic patients:  $302 \pm 15$  min vs. CON:  $247 \pm 17$  min;  $P < 0.05$ ). The difference between lowest individual preprandial and highest individual postprandial glycogen concentration was  $86 \pm 7$  mmol/l in CON and  $56 \pm 12$  mmol/l in type 2 diabetic patients ( $P < 0.05$ ). Rates of net glycogen synthesis (Fig. 4A) were  $\sim 44\%$  lower in type 2 diabetic patients ( $0.76 \pm 0.16$  vs. CON:  $1.36 \pm 0.15$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.05$ ). The direct pathway contributed similarly to glycogen synthesis in both study groups (type 2 diabetic patients:  $60 \pm 4\%$  vs. CON:  $56 \pm 2\%$ ,  $P = 0.514$ ). Overnight, hepatic glycogen concentrations decreased linearly, reaching values of  $\sim 215$  mmol/l in both groups in the morning (Fig. 3A). Rates of net glycogenolysis were  $\sim 50\%$  lower in type 2 diabetic patients ( $0.37 \pm 0.08$  vs. CON:  $0.75 \pm 0.09$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.01$ ) (Fig. 4A).

Before dinner, EGP was  $\sim 13\%$  higher in type 2 diabetic patients ( $1.95 \pm 0.07$  vs. CON:  $1.73 \pm 0.07$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.05$ ) (Fig. 4B). After dinner, EGP decreased in both groups but remained  $\sim 0.3$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  higher in type 2 diabetic patients from 30–170 min ( $0.98 \pm 0.11$  vs. CON:  $0.68 \pm 0.05$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.05$ ). Time course of EGP reached the nadir in CON at 60–90 min, but only at 240 min in type 2 diabetic patients, i.e., the maximum suppression of EGP occurred delayed in type 2 diabetic patients. Thereafter, EGP was similar in both groups and returned to  $\sim 60\text{--}70\%$  of predinner values at 480 min. Rates of postabsorptive gluconeogenesis in the time period of 11:00 P.M. to 1:00 A.M. represented  $67 \pm 8\%$  of EGP in type 2 diabetic patients but only  $43 \pm 4\%$  ( $P < 0.05$ ) of EGP in CON.

**Clamp tests.** During the hyperglycemic-hyperinsulinemic clamp tests, plasma glucose, FFAs, insulin, and glucagon were not different between both groups (Fig. 2E–H). Whole-body glucose uptake was  $\sim 37\%$  lower in type 2 diabetic patients (180–300 min: type 2 diabetic patients:  $8.03 \pm 0.35$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  vs. CON:  $12.73 \pm 0.48$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.05$ ).

Before the clamp test, both study groups exhibited similar hepatic glycogen concentrations (type 2 diabetic patients:  $218 \pm 10$  vs. CON:  $234 \pm 9$  mmol/l,  $P = 0.28$ , Fig. 3B). During the clamp, hepatic glycogen concentrations increased linearly in both groups (Fig. 3B), but  $V_{\text{Syn}}$  was  $\sim 46\%$  lower in type 2 diabetic patients ( $0.63 \pm 0.12$  vs. CON:  $1.17 \pm 0.15$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.05$ ), with similar contribution of the direct pathway in both groups (type 2 diabetic patients:  $60 \pm 10\%$  vs. CON:  $65 \pm 2\%$ ,  $P = 0.416$ ). Simultaneous  $V_{\text{Out}}$  (Fig. 5A) was not different (type 2 diabetic patients:  $0.21 \pm 0.07$  vs. CON:  $0.23 \pm 0.11$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P = 0.895$ ), resulting in  $\sim 54\%$  lower rates of net hepatic glycogen synthesis (Fig. 5A) in type 2 diabetic patients ( $0.42 \pm 0.10$  vs. CON:  $0.91 \pm 0.16$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.05$ ). Relative hepatic glycogen turnover was






FIG. 2. Plasma concentrations of glucose, FFAs, and major glucoregulatory hormones in type 2 diabetic patients (○) and CON (▲) after ingestion of a mixed-meal dinner (arrow) (A–D) and during hyperglycemic-hyperinsulinemic-somatostatin clamp tests (E–H). \* $P < 0.05$  for type 2 diabetic patients vs. CON.




FIG. 3. Time course hepatic glycogen concentration during a mixed-meal test (A) (protocol A, -120 to 960 min) and during the pulse period of the clamp test (B) (protocol C, -30 to 150 min) in type 2 diabetic patients (○) and CON (▲). A: \* $P < 0.05$ , type 2 diabetic patients vs. CON; § $P < 0.05$  vs. basal in CON; §§ $P < 0.05$  vs. basal in type 2 diabetic patients. B: Mean Pearson coefficient for linear regression of measured hepatic glycogen concentrations versus time was  $R = 0.96 \pm 0.01$  in CON and  $R = 0.77 \pm 0.12$  in type 2 diabetic patients.

slightly but not significantly increased in type 2 diabetic patients ( $31 \pm 11\%$  vs. CON:  $19 \pm 11\%$ ,  $P = 0.460$ ).

EGP was  $\sim 30\%$  higher in type 2 diabetic patients before the clamp (type 2 diabetic patients:  $2.38 \pm 0.10$  vs. CON:  $1.83 \pm 0.05$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.01$ ) and remained so during the clamp (120–300 min: type 2 diabetic patients:  $0.53 \pm 0.05$  vs. CON:  $0.04 \pm 0.04$   $\text{mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$ ,  $P < 0.02$ ) (Fig. 5B).

Hepatocellular lipid content was three times higher in type 2 diabetic patients ( $9.9 \pm 2.5$  AU vs. CON:  $2.8 \pm 0.7$  AU,  $P < 0.05$ ) and negatively correlated across the whole study population with rates of net hepatic glycogen synthesis ( $R = -0.602$ ,  $P < 0.05$ ) as well as with rates of whole-body glucose uptake ( $R = -0.576$ ,  $P < 0.05$ ) during the clamp test.

## DISCUSSION

This study provides the time course of cumulative glycogen storage and rates of net hepatic glycogen synthesis in type 2 diabetic patients under physiological conditions of mixed-meal ingestion as well as under that of experimen-

tal stimulation of glycogen accumulation. Postprandial glycogen synthesis is reduced in mildly overweight type 2 diabetic patients and is accompanied by a temporarily impaired EGP suppression. Because these defects of postprandial hepatic glucose metabolism of type 2 diabetic patients were observed during decreased insulin-to-glucagon ratios and impaired FFA suppression, we also assessed hepatic glucose metabolism under conditions of hyperinsulinemic hyperglycemia combined with low glucagon and FFA concentrations. Under these conditions, glycogen synthesis and suppression of EGP remained clearly impaired in type 2 diabetic patients.

**Postprandial conditions.** After dinner, glycogen accumulation rose by  $\sim 86$  mmol/l above preprandial concentration in CON in  $\sim 4.5$  h, accounting for storage of  $\sim 26$  g carbohydrates in the liver, which is similar to previous findings in nondiabetic humans (27). In type 2 diabetic patients, glycogen concentrations rose only by 56 mmol/l liver within  $\sim 5.5$  h after meal ingestion, resulting in storage of  $\sim 17$  g carbohydrates in the liver. Thus, type 2 diabetic patients exhibit reduction of net hepatic glycogen




FIG. 4. Rates of net hepatic glycogen synthesis and subsequent net hepatic glycogenolysis (A) and time course of EGP (B) in type 2 diabetic patients (T2DM) ( $\circ$ ) and CON ( $\blacktriangle$ ) after ingestion of a mixed-meal dinner at zero time. \* $P < 0.01$ , # $P < 0.05$ , type 2 diabetic patients vs. CON.

accumulation by  $\sim 35\%$ , suggesting that decreased hepatic glycogen storage contributes to their postprandial hyperglycemia.

A similar defect was observed in glucokinase-deficient maturity-onset diabetes of the young (MODY)-2 patients (38), in whom the impaired hepatic glucokinase activity is held responsible for the reduction in the contribution of glucose (the direct pathway) to hepatic glycogen synthesis (38). Of note, in the present study, the relative contribution of the direct pathway was unchanged in type 2 diabetic patients after the meal as well as during the clamp test. Previously, lower contributions of the direct pathway were reported (15,39), which could be due to the metabolic characteristics of these type 2 diabetic patients (higher  $\text{HbA}_{1c}$  and BMI) and lower insulin-to-glucagon ratios compared with the present study. In agreement with previous studies (18,19), type 2 diabetic patients also exhibited decreased net glycogenolysis and increased contribution of gluconeogenesis to glucose production during the postabsorptive period.

It is well known that EGP correlates with the degree of hyperglycemia after overnight fasting (4–8). Here we report that EGP of type 2 diabetic patients is also increased in the afternoon after a standardized breakfast and lunch, which is in contrast to normalized rates of EGP of type 2 diabetic patients after prolonged fasting (8). After dinner, EGP was rapidly suppressed in both groups but

remained  $\sim 0.3 \text{ mg} \cdot \text{kg}^{-1} \cdot \text{min}^{-1}$  higher during the first 3 h in type 2 diabetic patients, i.e., the maximum suppression of EGP occurs later in type 2 diabetic patients, providing for an additional  $\sim 5 \text{ g}$  glucose released into the circulation, thus also contributing to postprandial hyperglycemia. This time course of EGP suppression is similar to previous reports (6,10,29). Earlier studies, first describing higher rates of postprandial EGP in type 2 diabetic patients (10–12), observed more pronounced defects of EGP suppression. These differences can probably be attributed to methodological uncertainties in the assessment of metabolite fluxes under rapidly changing conditions in the classic double-tracer approach, which can be reduced by the variable infusion double-tracer technique (27,29,40).

After meal ingestion, type 2 diabetic patients had lower plasma insulin but higher glucagon and FFA concentrations than CON, as previously reported (10–12,41). Increased plasma FFAs impair EGP suppression (42) and affect autoregulation of hepatic glucose metabolism (24). The molar portal vein insulin-to-glucagon ratio, as estimated from peripheral concentrations (43,44), was  $\sim 50\%$  lower after the meal in type 2 diabetic patients. Additionally, to a possible impairment of hepatic insulin sensitivity, the lower insulin-to-glucagon ratio could have contributed to the decreased net glycogen synthesis (21,23) and EGP suppression (20,45). On the other hand, type 2 diabetic patients with markedly higher postprandial insulin secre-




FIG. 5.  $V_{\text{SynNet}}$ ,  $V_{\text{Out}}$  (A), and time course of EGP (B) in type 2 diabetic (T2DM) (○) and CON (▲) during hyperglycemic-hyperinsulinemic-somatostatin clamp tests. \* $P < 0.01$ , # $P < 0.05$ , type 2 diabetic patients vs. CON.

tion exhibit a comparable defect of EGP suppression (6). Thus, excessive postprandial hyperglycemia per se limited observed defects in hepatic glucose metabolism (46,47). Taken together, impaired net hepatic glycogen synthesis and insufficient EGP suppression resulted in decreased hepatic glucose uptake and increased amounts of glucose released into the circulation in type 2 diabetic patients.

**Hyperglycemic-hyperinsulinemic clamp conditions.** The decreased net hepatic glycogen synthesis of type 2 diabetic patients after meal ingestion can result from two distinct mechanisms: 1) decreased  $V_{\text{Syn}}$  and/or 2) simultaneously increased  $V_{\text{Out}}$  (21,23,48). Previous reports demonstrated that hyperglycemia per se does not markedly stimulate net hepatic glycogen synthesis, even in young healthy men (21,23). In addition, hyperinsulinemia per se is capable of stimulating hepatic glucose uptake in humans to some extent (23,25), and stepwise increases in the portal vein insulin concentration during hyperglycemia lead to an insulin concentration-dependent increase in net hepatic glycogen synthesis (21). Thus, we assumed that combined hyperglycemia and hyperinsulinemia are required to substantially stimulate net hepatic glycogen synthesis in the presence of hypoglucagonemia.

To evaluate hepatic glucose metabolism independently of type 2 diabetes-associated alterations in postprandial metabolite and hormonal responses, glycogen metabolism,

as well as rates of EGP and whole-body glucose disposal, were assessed under matched conditions of hyperglycemia, portal vein insulin-to-glucagon ratios, and similar plasma FFA concentrations.

$V_{\text{SynNet}}$  values observed in CON are in line with those obtained in nondiabetic humans under similar conditions (21–23). In type 2 diabetic patients, we observed ~54% reduced rates of net hepatic glycogen synthesis. Because the  $V_{\text{Out}}$  was comparable between type 2 diabetic patients and CON, the lower  $V_{\text{SynNet}}$  values are entirely explained by ~46% reduced  $V_{\text{Syn}}$  in type 2 diabetic patients, which under clamp conditions, cannot be explained by short-term differences in the metabolic or hormonal environment between type 2 diabetic patients and CON (Fig. 1). The comparable phosphorylase flux supports previous findings of similar hepatic glucagon sensitivity in type 2 diabetic patients and nondiabetic humans (49,50). During the hyperglycemic-hyperinsulinemic pancreatic clamp test, rates of EGP also remained higher in type 2 diabetic patients, which is in line with persistent decreased insulin sensitivity at the level of the liver (4,10).

In line with previous reports (7,26,51), HCL content was increased in type 2 diabetic patients and inversely correlated to insulin-stimulated whole-body glucose uptake during the clamp test. Furthermore, HCL content inversely correlated with  $V_{\text{SynNet}}$ , suggesting that HCL content can

serve as a marker not only of peripheral but also of hepatic insensitivity to insulin action.

During the clamp test, the ~40% reduction in rates of whole-body glucose uptake indicated peripheral insulin resistance in type 2 diabetic patients. Of note,  $V_{\text{SynNet}}$  contributed by ~6% and ~8% to whole-body glucose disposal in type 2 diabetic patients and CON, respectively, supporting the contention that skeletal muscle accounts for ~90% of whole-body glucose disposal under clamp conditions (1). Because both glucose disposal and net hepatic glycogen synthesis were similarly decreased by 40–50% in type 2 diabetic patients of our study and skeletal muscle glycogen synthesis is reduced by 55% in type 2 diabetic patients under clamp conditions (1), glycogen synthesis seems to be equally insensitive to insulin in liver and skeletal muscle.

In contrast to current knowledge of the importance of skeletal muscle insulin resistance in the development of overt type 2 diabetic patients, there is still debate on the causality between the defects in hepatic glucose metabolism and the onset of type 2 diabetic patients. There are some arguments in favor of the hypothesis that the defects in hepatic glycogen metabolism are secondary to type 2 diabetic patients and could result from effects of acute and/or chronic hyperglycemia, increased availability of lipids, and alterations of secretion of major glucoregulatory hormones. Both the blunted rise of insulin-to-glucagon ratios and higher plasma FFA concentrations likely contributed to the impaired net hepatic glycogen synthesis in type 2 diabetic patients during the mixed-meal study. When insulin, glucagon, glucose, and FFA concentrations were successfully matched during the clamp protocol, hepatic glycogen synthesis was also impaired, indicating an additional defect aside from the effects of metabolic and hormonal differences. Moreover, the impact of chronic changes in glycemia and lipidemia, frequently termed “glucolipotoxicity,” on liver metabolism cannot be completely excluded from the results of the clamp test.

Chronically increased availability of both circulating (FFAs and triglycerides) and intracellular (HCL) lipids are frequently observed in different nondiabetic insulin-resistant populations as well as in type 2 diabetic patients (52,53) and are associated with defects in hepatic and/or whole-body glucose metabolism (26,52–55). Thus, increased HCL content in type 2 diabetic patients and its correlation with net hepatic glycogen synthesis as observed in our study indicates that fat accumulation in liver could play a role in the development of defects of hepatic glycogen metabolism in type 2 diabetic patients.

In conclusion, we show that type 2 diabetic patients exhibit a marked reduction of net hepatic glycogen synthesis after mixed-meal ingestion. Furthermore, even stimulation of glycogen synthesis by combined hyperinsulinemic hyperglycemia is unable to overcome this defect of hepatic glycogen metabolism. Thus, impaired insulin sensitivity and/or chronic glucolipotoxicity in addition to any effects of altered insulin-to-glucagon ratio or increased FFA accounts for defective hepatic glucose metabolism in type 2 diabetic patients.

#### ACKNOWLEDGMENTS

The study was supported by grants from the Austrian Science Foundation (FWF; P13722-MED, P15656), the Eu-

ropean Foundation for the Study of Diabetes (EFSD and Novo-Nordisk Type 2 Program Focused Research Grant), the Herzfelder'sche Familienstiftung (to M.R.), Novo-Nordisk (to W.W.), and the U.S. Public Health Service (ROI DK-49230, P30 DK-45735) (to G.I.S.).

We gratefully acknowledge the cooperation of A. Hofer, H. Lentner, and G. Hofstetter (Metabolic Research Unit and Endocrine Laboratory, Department of Internal Medicine III, University of Vienna) and Professors E. Moser and S. Trattnig (MR Center of Excellence, University of Vienna).

#### REFERENCES

- Shulman GI, Rothman DL, Jue T, Stein P, DeFronzo RA, Shulman RG: Quantitation of muscle glycogen synthesis in normal subjects and subjects with non-insulin-dependent diabetes by  $^{13}\text{C}$  nuclear magnetic resonance spectroscopy. *N Engl J Med* 322:223–228, 1990
- Cline GW, Petersen KF, Krssak M, Shen J, Hundal RS, Trajanoski Z, Inzucchi S, Dresner A, Rothman DL, Shulman GI: Impaired glucose transport as a cause of decreased insulin-stimulated muscle glycogen synthesis in type 2 diabetes. *N Engl J Med* 341:240–246, 1999
- Carey PE, Halliday J, Snaar JE, Morris P, Taylor R: Direct assessment of muscle glycogen storage after mixed meals in normal and type 2 diabetic subjects. *Am J Physiol Endocrinol Metab* 284:E688–E694, 2002
- DeFronzo RA: Pathogenesis of type 2 diabetes: metabolic and molecular implications for identifying diabetes genes. *Diabetes Reviews* 5:177–269, 1997
- Boden G, Chen X, Stein TP: Gluconeogenesis in moderately and severely hyperglycemic patients with type 2 diabetes mellitus. *Am J Physiol Endocrinol Metab* 280:E23–E30, 2001
- Singhal P, Caumo A, Carey PE, Cobelli C, Taylor R: Regulation of endogenous glucose production after a mixed meal in type 2 diabetes. *Am J Physiol Endocrinol Metab* 283:E275–E283, 2002
- Anderwald C, Bernroider E, Krssak M, Stingl H, Brehm A, Bischof MG, Nowotny P, Roden M, Waldhäusl W: Effects of insulin treatment in type 2 diabetic patients on intracellular lipid content in liver and skeletal muscle. *Diabetes* 51:3025–3032, 2002
- Radziuk J, Pye S: Production and metabolic clearance of glucose under basal conditions in type II (non-insulin-dependent) diabetes mellitus. *Diabetologia* 44:983–991, 2001
- Ferrannini E, Simonson DC, Katz LD, Reichard G Jr, Bevilacqua S, Barrett EJ, Olsson M, DeFronzo RA: The disposal of an oral glucose load in patients with non-insulin-dependent diabetes. *Metabolism* 37:79–85, 1988
- Firth RG, Bell PM, Marsh HM, Hansen I, Rizza RA: Postprandial hyperglycemia in patients with noninsulin-dependent diabetes mellitus: role of hepatic and extrahepatic tissues. *J Clin Invest* 77:1525–1532, 1986
- Kelley D, Mookan M, Veneman T: Impaired postprandial glucose utilization in non-insulin-dependent diabetes mellitus. *Metabolism* 43:1549–1557, 1994
- McMahon M, Marsh HM, Rizza RA: Effects of basal insulin supplementation on disposition of mixed meal in obese patients with NIDDM. *Diabetes* 38:291–303, 1989
- Mitrakou A, Kelley D, Veneman T, Jenssen T, Pangburn T, Reilly J, Gerich J: Contribution of abnormal muscle and liver glucose metabolism to postprandial hyperglycemia in NIDDM. *Diabetes* 39:1381–1390, 1990
- Felig P, Wahren J, Hendler R: Influence of maturity-onset diabetes on splanchnic glucose balance after oral glucose ingestion. *Diabetes* 27:121–126, 1978
- Basu A, Basu R, Shah P, Vella A, Johnson CM, Jensen M, Nair KS, Schwenk WF, Rizza RA: Type 2 diabetes impairs splanchnic uptake of glucose but does not alter intestinal glucose absorption during enteral glucose feeding: additional evidence for a defect in hepatic glucokinase activity. *Diabetes* 50:1351–1362, 2001
- Iozzo P, Hallsten K, Oikonen V, Virtanen KA, Kempainen J, Solin O, Ferrannini E, Knuuti J, Nuutila P: Insulin-mediated hepatic glucose uptake is impaired in type 2 diabetes: evidence for a relationship with glycemic control. *J Clin Endocrinol Metab* 88:2055–2060, 2003
- Beringer A, Thaler H: [On quantitative analyses of the glycogen content of the liver in normal and sick persons.] *Wien Klin Wochenschr* 26:509–513, 1963
- Magnusson I, Rothman DL, Katz LD, Shulman RG, Shulman GI: Increased rate of gluconeogenesis in type II diabetes mellitus: a  $^{13}\text{C}$  nuclear magnetic resonance study. *J Clin Invest* 90:1323–1327, 1992



19. Hundal RS, Krssak M, Dufour S, Laurent D, Lebon V, Chandramouli V, Inzucchi SE, Schumann WC, Petersen KF, Landau BR, Shulman GI: Mechanism by which metformin reduces glucose production in type 2 diabetes. *Diabetes* 49:2063–2069, 2000
20. Cherrington AD: Banting Lecture 1997: Control of glucose uptake and release by the liver in vivo. *Diabetes* 48:1198–1214, 1999
21. Roden M, Perseghin G, Petersen KF, Hwang JH, Cline GW, Gerow K, Rothman DL, Shulman GI: The roles of insulin and glucagon in the regulation of hepatic glycogen synthesis and turnover in humans. *J Clin Invest* 97:642–648, 1996
22. Cline GW, Rothman DL, Magnusson I, Katz LD, Shulman GI: <sup>13</sup>C-nuclear magnetic resonance spectroscopy studies of hepatic glucose metabolism in normal subjects and subjects with insulin-dependent diabetes mellitus. *J Clin Invest* 94:2369–2376, 1994
23. Petersen KF, Laurent D, Rothman DL, Cline GW, Shulman GI: Mechanism by which glucose and insulin inhibit net hepatic glycogenolysis in humans. *J Clin Invest* 101:1203–1209, 1998
24. Stingl H, Krssak M, Krebs M, Bischof MG, Nowotny P, Fornsinn C, Shulman GI, Waldhäusl W, Roden M: Lipid-dependent control of hepatic glycogen stores in healthy humans. *Diabetologia* 44:48–54, 2001
25. Iozzo P, Geisler F, Oikonen V, Maki M, Takala T, Solin O, Ferrannini E, Knuuti J, Nuutila P: Insulin stimulates liver glucose uptake in humans: an <sup>18</sup>F-FDG PET Study. *J Nucl Med* 44:682–689, 2003
26. Ryysy L, Hakkinen AM, Goto T, Vehkavaara S, Westerbacka J, Halavaara J, Yki-Jarvinen H: Hepatic fat content and insulin action on free fatty acids and glucose metabolism rather than insulin absorption are associated with insulin requirements during insulin therapy in type 2 diabetic patients. *Diabetes* 49:749–758, 2000
27. Taylor R, Magnusson I, Rothman DL, Cline GW, Caumo A, Cobelli C, Shulman GI: Direct assessment of liver glycogen storage by <sup>13</sup>C nuclear magnetic resonance spectroscopy and regulation of glucose homeostasis after a mixed meal in normal subjects. *J Clin Invest* 97:126–132, 1996
28. Bischof MG, Bernroider E, Krssak M, Krebs M, Stingl H, Nowotny P, Yu C, Shulman GI, Waldhäusl W, Roden M: Hepatic glycogen metabolism in type 1 diabetes after long-term near normoglycemia. *Diabetes* 51:49–54, 2002
29. Basu R, Di Camillo B, Toffolo G, Basu A, Shah P, Vella A, Rizza R, Cobelli C: Use of a novel triple-tracer approach to assess postprandial glucose metabolism. *Am J Physiol Endocrinol Metab* 284:E55–E69, 2003
30. Rothman DL, Magnusson I, Katz LD, Shulman RG, Shulman GI: Quantitation of hepatic glycogenolysis and gluconeogenesis in fasting humans with <sup>13</sup>C NMR. *Science* 254:573–576, 1991
31. Hwang JH, Perseghin G, Rothman DL, Cline GW, Magnusson I, Petersen KF, Shulman GI: Impaired net hepatic glycogen synthesis in insulin-dependent diabetic subjects during mixed meal ingestion: a <sup>13</sup>C nuclear magnetic resonance spectroscopy study. *J Clin Invest* 95:783–787, 1995
32. Roden M, Krssak M, Stingl H, Gruber S, Hofer A, Fornsinn C, Moser E, Waldhäusl W: Rapid impairment of skeletal muscle glucose transport/phosphorylation by free fatty acids in humans. *Diabetes* 48:358–364, 1999
33. Krebs M, Stingl H, Nowotny P, Weghuber D, Bischof M, Waldhäusl W, Roden M: Prevention of in vitro lipolysis by tetrahydrolipstatin. *Clin Chem* 46:950–954, 2000
34. Magnusson I, Rothman DL, Jucker B, Cline GW, Shulman RG, Shulman GI: Liver glycogen turnover in fed and fasted humans. *Am J Physiol* 266:E796–E803, 1994
35. Steele R: Influences of glucose loading and of injected insulin on hepatic glucose output. *Ann NY Acad Sci* 82:420–430, 1959
36. Sparacino G, Pillonetto G, Capello M, De Nicolao G, Cobelli C: WINSTO-DEC: a stochastic deconvolution interactive program for physiological and pharmacokinetic systems. *Comput Methods Programs Biomed* 67:67–77, 2002
37. Krebs M, Krssak M, Bernroider E, Anderwald C, Brehm A, Meyerspeer M, Nowotny P, Roth E, Waldhäusl W, Roden M: Mechanism of amino acid-induced skeletal muscle insulin resistance in humans. *Diabetes* 51:599–605, 2002
38. Velho G, Petersen KF, Perseghin G, Hwang JH, Rothman DL, Pueyo ME, Cline GW, Froguel P, Shulman GI: Impaired hepatic glycogen synthesis in glucokinase-deficient (MODY-2) subjects. *J Clin Invest* 98:1755–1761, 1996
39. Basu A, Basu R, Shah P, Vella A, Johnson CM, Nair KS, Jensen MD, Schwenk WF, Rizza RA: Effects of type 2 diabetes on the ability of insulin and glucose to regulate splanchnic and muscle glucose metabolism: evidence for a defect in hepatic glucokinase activity. *Diabetes* 49:272–283, 2000
40. Cobelli C, Mari A, Ferrannini E: Non-steady state: error analysis of Steele's model and developments for glucose kinetics. *Am J Physiol* 252:E679–E689, 1987
41. Waldhäusl W, Bratusch-Marrain P, Gasic S, Korn A, Nowotny P: Insulin production rate, hepatic insulin retention and splanchnic carbohydrate metabolism after oral glucose ingestion in hyperinsulinaemic type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia* 23:6–15, 1982
42. Boden G, Chen X, Capulong E, Mozzoli M: Effects of free fatty acids on gluconeogenesis and autoregulation of glucose production in type 2 diabetes. *Diabetes* 50:810–816, 2001
43. Waldhäusl W, Bratusch-Marrain P, Gasic S, Korn A, Nowotny P: Insulin production rate following glucose ingestion estimated by splanchnic C-peptide output in normal man. *Diabetologia* 17:221–227, 1979
44. Bjorkman O, Eriksson LS, Nyberg B, Wahren J: Gut exchange of glucose and lactate in basal state and after oral glucose ingestion in postoperative patients. *Diabetes* 39:747–751, 1990
45. Unger RH: Role of glucagon in the pathogenesis of diabetes: the status of the controversy. *Metabolism* 27:1691–1709, 1978
46. Vella A, Reed AS, Charkoudian N, Shah P, Basu R, Basu A, Joyner MJ, Rizza RA: Glucose-induced suppression of endogenous glucose production: dynamic response to differing glucose profiles. *Am J Physiol Endocrinol Metab* 285:E25–E30, 2003
47. Hother-Nielsen O, Vaag A, Skott P, Beck-Nielsen H: Effect of hyperglycemia per se on glucose turnover rates in patients with insulin-dependent diabetes. *Metabolism* 42:86–93, 1993
48. Landau BR: Methods for measuring glycogen cycling. *Am J Physiol Endocrinol Metab* 281:E413–E419, 2001
49. Matsuda M, DeFronzo RA, Glass L, Consoli A, Giordano M, Bressler P, Delprato S: Glucagon dose-response curve for hepatic glucose production and glucose disposal in type 2 diabetic patients and normal individuals. *Metabolism* 51:1111–1119, 2002
50. Nielsen MF, Wise S, Dinneen SF, Schwenk WF, Basu A, Rizza RA: Assessment of hepatic sensitivity to glucagon in NIDDM: use as a tool to estimate the contribution of the indirect pathway to nocturnal glycogen synthesis. *Diabetes* 46:2007–2016, 1997
51. Mayerson AB, Hundal RS, Dufour S, Lebon V, Befroy D, Cline GW, Enocksson S, Inzucchi SE, Shulman GI, Petersen KF: The effects of rosiglitazone on insulin sensitivity, lipolysis, and hepatic and skeletal muscle triglyceride content in patients with type 2 diabetes. *Diabetes* 51:797–802, 2002
52. Petersen KF, Befroy D, Dufour S, Dziura J, Ariyan C, Rothman DL, DiPietro L, Cline GW, Shulman GI: Mitochondrial dysfunction in the elderly: possible role in insulin resistance. *Science* 300:1140–1142, 2003
53. Seppala-Lindroos A, Vehkavaara S, Hakkinen AM, Goto T, Westerbacka J, Sovijarvi A, Halavaara J, Yki-Jarvinen H: Fat accumulation in the liver is associated with defects in insulin suppression of glucose production and serum free fatty acids independent of obesity in normal men. *J Clin Endocrinol Metab* 87:3023–3028, 2002
54. Tiikkainen M, Tamminen M, Hakkinen AM, Bergholm R, Vehkavaara S, Halavaara J, Teramo K, Rissanen A, Yki-Jarvinen H: Liver-fat accumulation and insulin resistance in obese women with previous gestational diabetes. *Obes Res* 10:859–867, 2002
55. Marchesini G, Brizi M, Bianchi G, Tomassetti S, Bugianesi E, Lenzi M, McCullough AJ, Natale S, Forlani G, Melchionda N: Nonalcoholic fatty liver disease: a feature of the metabolic syndrome. *Diabetes* 50:1844–1850, 2001