

Effect of quinolones on caffeine disposition

Six healthy volunteers received a single caffeine dose after pretreatment with norfloxacin, pipemidic acid, or placebo in a crossover, randomized, single-blind clinical trial. Quinolones altered the pharmacokinetics of caffeine, with a significant increase in the AUCs and a decrease in plasma clearance. The elimination half-life increased significantly with pipemidic acid. The apparent volume of distribution, mean renal clearance, and time to reach maximum caffeine concentrations remained unaltered. There was a decline in caffeine metabolite levels in the 24-hour urine samples for both quinolone treatments, suggesting that pipemidic acid and, to a lesser degree, norfloxacin inhibit metabolism of the *N*-demethylation pathways of caffeine. The practical consequence of this observation could be caffeine accumulation during repeated intake of coffee. In two additional healthy volunteers under a controlled multiple-dose regimen of caffeine ingestion, administration of pipemidic acid for 2 days caused a fourfold increase in the plasma concentrations of caffeine. (CLIN PHARMACOL THER 1989;45:234-40.)

Marcel·li Carbó, MD, Jordi Segura, PhD, Rafael De la Torre, PhD,
Josep M. Badenas, MD, Jordi Camí, MD *Barcelona, Spain*

The primary pathways of caffeine metabolism are *N*-demethylations,¹ which take place in the hepatic microsomal oxidative system (cytochrome P-450) mediated by isozymes, also related to the biotransformation of polycyclic aromatic hydrocarbons.² Factors such as smoking habit,³ pregnancy,^{4,5} genetic inheritance,⁶ liver disorders,⁷ and interaction with other drugs like cimetidine,^{8,9} disulfiram,¹⁰ oral contraceptives,^{11,12} and mexiletine¹³ can moderately affect metabolic clearance of caffeine. Intense metabolic inhibition of caffeine *N*-demethylation by idroclamide,¹⁴ furafylline,¹⁵ and methoxsalen¹⁶ has been detected in human beings. Studies using animal models¹⁷⁻¹⁹ confirmed these interactions and demonstrated that the phenomenon is shared by other compounds. Recent findings have shown that antibiotics of the quinolone group can affect the clearance of substances with a caffeine-like structure such as the-

ophylline.²⁰⁻²⁴ Side effects probably caused by caffeine accumulation have been observed in patients treated with enoxacin^{25,26} and pipemidic acid.²⁷ Alterations in caffeine disposition as a result of treatment with enoxacin and ciprofloxacin have also been described.^{28,29} To study how the quinolones alter caffeine disposition, a clinical trial has been carried out to evaluate the variations of the pharmacokinetic parameters and the metabolic profile of caffeine caused by treatment with pipemidic acid or norfloxacin. Caffeine accumulation after multiple doses in the presence of concomitant treatment with multiple doses of pipemidic acid was also investigated.

SUBJECTS AND METHODS

Subjects. Eight healthy male volunteers (age range 20 to 37 years) were selected after evaluation of their clinical history, physical examination, ECG, and complete blood and urine analyses. Seven subjects were regular smokers (10 to 40 cigarettes per day) and volunteer No. 8 was a nonsmoker. The subjects had not taken drugs during the 4 weeks before the trial. They were informed of extent, nature, and objectives of the study and gave written consent. All the studies were approved by the Hospital del Mar Ethical Committee and authorized by the Dirección General de Farmacia y Productos Sanitarios (Nos. 87/260). Drugs were administered orally with 120 ml water after a 10-hour fast. The volunteers were requested to abstain from the

From the Department of Pharmacology and Toxicology, Institut Municipal d'Investigació Mèdica.

Supported by the Comisión Asesora de Investigación Científica y Técnica (Ministerio de Educación y Ciencia) CAICYT 188/84 and the Institut Municipal d'Investigació Mèdica.

Presented in preliminary form at the Third European Congress of Biopharmaceutics and Pharmacokinetics Friburg, Federal Republic of Germany, April 20-24, 1987.

Received for publication April 7, 1988; accepted July 29, 1988.

Reprint requests: Jordi Segura, PhD, Department of Pharmacology and Toxicology, Institut Municipal d'Investigació Mèdica, Passeig Marítim 25-29, 08003 Barcelona, Spain.




Fig. 1. Caffeine plasma concentrations for six volunteers (mean \pm SE) after administration of a single dose of caffeine and treatment with pipemidic acid (black triangles), norfloxacin (white triangles), or placebo (black circles).




Fig. 2. Concentrations ratios (mean \pm SE) of *N*-demethylated caffeine (CF) metabolites in 24-hour urine samples after administration of a single dose of caffeine and treatment with pipemidic acid (black bars), norfloxacin (stripped bars), or placebo (white bars). The *N*-demethylations affected are indicated at the bottom of the figure. Ratios are indicated as follows: paraxanthine/caffeine (PX/CF), theophylline/caffeine (TF/CF), theobromine/caffeine (TB/CF), 1-methylxanthine/theophylline (1-MX/TF), 1-methylxanthine/paraxanthine (1-MX/PX), and 1-methylxanthine/caffeine (1-MX/CF). (* 0.02 < *p* < 0.05 vs placebo; ** 0.005 < *p* < 0.02 vs placebo; *** *p* < 0.005 vs placebo.)

Table I. Pharmacokinetic parameters for caffeine after single dose and during treatment with pipemidic acid, norfloxacin, or placebo

Volunteer			Placebo		
No.	Age (yr)	Weight (kg)	C_{max} ($\mu\text{g/ml}$)	t_{max} (hr)	$t_{1/2}$ (hr)
1	23	74.5	6.5	1.0	3.53
2	23	61.7	8.11	1.0	3.71
3	22	59.4	5.87	2.0	1.02
4	19	60.5	8.17	1.0	13.47
5	21	60.5	6.36	1.0	2.77
6	22	69.5	7.35	2.0	3.40
$\bar{X} \pm \text{SE}$	21.6 ± 0.6	64.3 ± 2.5	7.06 ± 0.39	1.3 ± 0.2	4.65 ± 1.81
p Value					

Norfloxacin							
C_{max} ($\mu\text{g/ml}$)	t_{max} (hr)	$t_{1/2}$ (hr)	$AUC_{(0-24)}$ ($\text{mg} \cdot \text{hr/L}$)	$AUC_{0-\infty}$ ($\text{mg} \cdot \text{hr/L}$)	V_{area} (L/kg)	CL (L/hr)	CL_R^* (L/hr)
4.66	1.0	5.41	45.88	45.88	0.80	7.63	0.041
6.53	1.0	4.11	50.08	50.08	0.67	6.99	0.093
7.86	1.0	2.03	32.18	32.18	0.54	10.88	—
9.92	2.0	12.33	116.22	161.06	0.64	2.17	0.260
8.27	1.0	2.68	48.93	48.93	0.46	7.15	0.134
10.52	2.0	7.63	108.30	124.59	0.45	2.81	0.070
7.96 ± 0.88	1.3 ± 0.2	5.70 ± 1.56	66.93 ± 14.60	77.12 ± 21.46	0.59 ± 0.05	6.27 ± 1.33	0.120 ± 0.038
NS	NS	NS	<0.05	NS	NS	<0.01	NS

NS, not significant differences vs placebo.

* $n = 5$ (loss of urine volume of volunteer 3).

consumption of alcohol and drinks and foods that contained methylxanthines for 48 hours before the onset of each phase of treatment.

Single caffeine dose pharmacokinetic study. Six volunteers (Nos. 1 to 6) were subjected to a Latin square design with single-blind administration according to previous random assignment. There were three phases of treatment, each lasting 2 days and separated by a 7-day drug-free interval. During the first day of each phase, volunteers received one of the following treatments: pipemidic acid, 800 mg b.i.d., norfloxacin, 800 mg b.i.d., or placebo b.i.d. On the second day of each phase each volunteer was given a single dose of caffeine (350 mg) together with the first daily dose of the assigned drug, the antecubital vein was catheterized, and a 10 ml blood sample was drawn before caffeine administration ($t = 0$) and 1, 2, 3, 4, 6, 8, 10, 12, 14, and 24 hours after administration. Twenty-four-hour urine samples were also collected. Plasma and urine samples were stored at -20°C until analysis. Vital signs were recorded immediately before blood extractions and self-reported side effects were assessed at 0, 14, and 24 hours.

Multiple caffeine dose study. Volunteers 7 and 8

were submitted to controlled caffeine intake for 48 hours before pipemidic acid administration (days 1 and 2 of this protocol), which was continued until day 5. Doses were 100 mg q.i.d. for volunteer 7 and 100 mg t.i.d. for volunteer 8. On days 3 and 4 they also received 400 mg b.i.d. pipemidic acid. Blood samples were obtained throughout the study and handled as described above.

Analysis of biologic samples. Plasma samples were taken at room temperature and subjected to an ultrasound bath for 30 minutes. More than 250 μL plasma was dispensed in 500 μL of methanolic solution containing ethyltheophylline (10 $\mu\text{g/ml}$) as internal standard. The sample was vortexed for 20 seconds and then centrifuged at 3000 rpm for 15 minutes. The methanolic supernatant was collected and 20 μL injected into an HPLC system. The HPLC mobile phase was a mixture of 2% tetrahydrofuran in 10 mmol/L sodium sulfate adjusted to pH 6.5 with phosphoric acid/acetonitrile/methanol (80:10:10). Spherisorb ODS-1 column (Phase Separation Ltd., Queenserry, U.K.), 20 by 0.46 cm, particle size 10 μm , and detection wavelength of 280 nm were used. The assay sensitivity for caffeine was 0.3 $\mu\text{g/ml}$. The interassay coefficient of variation

Placebo							
$AUC_{(0-24)}$ (mg · hr/L)		$AUC_{0 \rightarrow \infty}$ (mg · hr/L)	V_{area} (L/kg)	CL (L/hr)	CL_R^* (L/hr)		
30.91		30.91	0.78	11.32	0.113		
42.71		42.71	0.71	8.19	0.175		
20.09		20.09	0.43	17.42	—		
86.67		86.67	1.29	4.03	0.113		
34.24		34.24	0.67	10.22	0.093		
49.87		49.87	0.49	7.01	0.063		
44.08 ± 9.48		44.08 ± 9.48	0.73 ± 0.124	9.70 ± 1.86	0.111 ± 0.018		
Pipemidic acid							
C_{max} (µg/ml)	t_{max} (hr)	$t_{1/2}$ (hr)	$AUC_{(0-24)}$ (mg · hr/L)	$AUC_{0 \rightarrow \infty}$ (mg · hr/L)	V_{area} (L/kg)	CL (L/hr)	CL_R^* (L/hr)
6.38	1.0	7.16	68.30	76.15	0.64	4.6	0.076
7.72	2.0	10.32	90.75	116.66	0.72	3.00	0.290
7.41	1.0	3.86	45.52	45.52	0.72	7.68	—
11.89	4.0	19.73	141.91	274.58	0.60	1.27	0.157
7.13	3.0	9.08	82.95	101.82	0.74	3.43	0.038
8.24	3.0	19.58	150.03	254.29	0.56	1.38	0.043
8.13 ± 0.79	2.3 ± 0.5	11.62 ± 2.69	96.58 ± 16.87	144.84 ± 39.17	0.66 ± 0.03	3.56 ± 0.97	0.121 ± 0.05
NS	NS	<0.02	<0.005	<0.05	NS	<0.005	NS

was 7% at 1 µg/ml and 5.4% at 10 µg/ml ($n = 8$). Intrassay coefficient of variation was 4.6% and 3.2% ($n = 8$) at the same concentrations.

Caffeine and its metabolites were analyzed in urine samples according to the method developed by Grant et al.³⁰ with the following modifications. More than 0.5 ml urine was dispensed in 0.2 ml of methanolic solution containing 8-chlorotheophylline as internal standard (10 µg/ml). Six milliliters of chloroform/isopropanol (85:15) was added and the mixture was shaken for 30 seconds and centrifuged for 5 minutes at 2500 rpm. The organic phase was transferred and evaporated; the residue was reconstituted in 400 µl mobile phase and 20 µl was injected into an HPLC system. Assay sensitivity was 0.1 µg/ml for caffeine and metabolites; the corresponding interassay coefficients of variation were 5.5% at 1 µg/ml and 3.1% at 10 µg/ml ($n = 10$). Intraassay coefficients of variation were 4.1% and 2% ($n = 10$).

Data analysis. Pharmacokinetic parameters are described in Table I. Experimental values are the maximum plasma concentrations (C_{max}), time to reach this value (t_{max}), and $AUC_{0 \rightarrow 24}$.

Calculations were carried out to determine the best

least-squares fit of the individual curves to a mono-compartmental model. The resulting plasma elimination half-life ($t_{1/2}$), apparent volume of distribution (V_{area}), and area under the curve from zero to infinity ($AUC_{0 \rightarrow \infty}$) are also reported together with the total plasma clearance (CL) and mean renal clearance (CL_R), calculated as total caffeine excreted in 24-hour urine divided by $AUC_{0 \rightarrow 24}$. Complete absorption was assumed.³¹

The paired Student t test was used for each volunteer (single dose study) to evaluate the effects of pipemidic acid and norfloxacin on pharmacokinetic parameters compared with placebo values.

RESULTS

Quinolone effects on pharmacokinetics of caffeine.

Fig. 1 shows the plasma caffeine concentrations of volunteers in the caffeine single-dose study and Table I presents the individual pharmacokinetic parameters. C_{max} was usually attained 1 to 3 hours after caffeine administration. Plasma caffeine levels were significantly greater in the pipemidic acid treatment group than in the placebo group at times from 3 hours on. At 8 and 24 hours, caffeine levels of 5.7 ± 2.33




Fig. 3. Caffeine plasma concentrations in volunteers 8 (nonsmoker) and 7 (smoker) after several caffeine doses per day. Arrows indicate pipemidic acid administration.

µg/ml and 2.0 ± 1.78 µg/ml were reached with pipemidic acid compared, respectively, with 2.2 ± 1.82 µg/ml and undetectable in the placebo group. Average levels were always higher for the norfloxacin group than for the placebo group, although there were significant differences only between 3 and 4 hours after caffeine administration.

The elimination $t_{1/2}$ of caffeine was 248% greater ($p < 0.02$) in the pipemidic acid group. Plasma clearance decreased for both the pipemidic acid ($p < 0.005$) and norfloxacin ($p < 0.01$) groups. $AUC_{0 \rightarrow 24}$ increased for both pipemidic acid ($p < 0.005$) and norfloxacin ($p < 0.05$). Apparent V_{area} , mean CL_R , C_{max} , and t_{max} did not vary significantly.

Ratios of the urinary concentrations of caffeine metabolites to their respective substrates along the metabolic pathways (Fig. 2) showed that paraxanthine/caffeine, theophylline/caffeine, theobromine/caffeine, 1-methylxanthine/caffeine, and 1-methylxanthine/theophylline decreased with pipemidic acid administration. After norfloxacin administration, only paraxanthine/caffeine and 1-methylxanthine/caffeine showed significant differences with regard to the placebo. The ratio 1-methylxanthine/paraxanthine did not vary significantly as a result either of pipemidic acid or norfloxacin pretreatments. However, independent of the statistically significant differences, all mean values of the ratios were always lower after the quinolone than after placebo treatments.

Quinolone effects on caffeine steady-state plasma caffeine levels. After repeated administration of caf-

feine (Fig. 3), plasma caffeine concentrations in volunteer 8 stabilized at about 3 to 5 µg/ml and then increased after pipemidic acid intake, reaching levels of about 15 µg/ml. The plasma levels of volunteer 7 stabilized at about 0.1 to 1.5 µg/ml and increased to 5 µg/ml after pipemidic acid treatment. In both cases there was a fourfold increase in plasma concentrations after 2 days of pipemidic acid treatment. No adverse effects were observed in either volunteer.

DISCUSSION

The results show that caffeine elimination is inhibited by both pipemidic acid and norfloxacin. The differences in pharmacokinetic parameters suggest that the alterations in caffeine clearance induced by pipemidic acid and norfloxacin are quantitatively different, pipemidic acid inhibiting caffeine elimination more than norfloxacin. These differences are not surprising, because recent studies involving other quinolones (i.e., enoxacin and ciprofloxacin^{28,29}) have also shown a difference in the degree of effects on plasma caffeine clearance.

The absence of significant differences with respect to the placebo in both the apparent V_{area} and the mean CL_R suggests that the alteration in caffeine disposition is not related to changes in the distribution or renal elimination of the unaltered drug. Caffeine absorption was not affected by pretreatment with the quinolones, because neither t_{max} nor C_{max} varied for either drug. This fact is in agreement with the assumption of complete bioavailability of caffeine after oral administration,^{31,32}

These findings suggest altered metabolic elimination

as the most probable mode of interaction. Primary metabolic degradation of caffeine takes place in the hepatic mixed-function oxidase system (cytochrome P-450, forms c and d) by *N*-demethylation, *N*³-demethylation being the principal route of biotransformation.² This metabolic pathway seems to be the one most affected by quinolone treatment, because of the relative reduction in the paraxanthine/caffeine ratio in 24-hour urine samples (60% and 39% for the pipemidic acid and norfloxacin groups, respectively). The important decrease in the 1-methylxanthine/theophylline ratio is also related to inhibition of the *N*³-demethylation. The differences in the theobromine/caffeine, theophylline/caffeine, and 1-methylxanthine/paraxanthine ratios suggest that pipemidic acid inhibits not only *N*³-demethylation but also to some extent *N*⁷- and *N*¹-demethylation pathways. Obviously, those metabolites subjected to consecutive metabolic pathways potentially affected show a significant inhibition (e.g., 1-methylxanthine/caffeine; Fig. 2).

The caffeine multiple-dose study confirmed the alterations in caffeine disposition, because plasma concentrations increased after pipemidic acid administration. Rapid accumulation was induced even under conditions of normal caffeine intake. The plasma levels of volunteers 7 and 8, like those in the single-dose study, did not produce adverse effects. This is probably due in part to the fact that all volunteers except number 8 were regular cigarettes smokers. It is known that smoking produces enzymatic induction of caffeine biotransformation,³³ which blunts the effects of quinolone inhibition. Volunteer 8 (nonsmoker) probably would have attained caffeine plasma concentrations sufficient to produce symptoms of caffeine intoxication if the regimen of pipemidic acid administration had continued. Those situations in which alteration in caffeine disposition was caused by other enzymatic inhibitors (oral contraceptives, cimetidine, disulfiram, methoxalen, etc.) should be given special attention when treatment with antibiotics of the quinolone group is considered.

References

1. Tang-liu D, Williams RL, Riegelman S. Disposition of caffeine and its metabolites in man. *J Pharmacol Exp Ther* 1983;224:180-5.
2. Kotake AN, Schoeller DA, Lambert GH, Baker AL, Schaffer DD, Josephs H. The caffeine CO₂ breath test: dose response and route of *N*-demethylation in smokers and nonsmokers. *CLIN PHARMACOL THER* 1982;32:261-9.
3. Parsons WD, Neims AH. Effect of smoking in caffeine clearance. *CLIN PHARMACOL THER* 1978;24:40-5.
4. Scott NR, Chakraborty J, Marks V. Urinary metabolites of caffeine in pregnant women. *Br J Clin Pharmacol* 1986;22:475-8.
5. Cummings AJ. A survey of pharmacokinetic data from pregnant women. *Clin Pharmacokinet* 1983;8:344-5.
6. Kalow W. Genetics of drug transformation. *Clin Biochem* 1986;19:76-82.
7. Wang T, Kleber G, Stellard F, Paumgartner G. Caffeine elimination: a test of liver function. *Klin Wochenschr* 1985;63:1124-8.
8. May DC, Jarboe CH, Van Bakel AB, Williams WM. Effects of cimetidine on caffeine disposition in smokers and nonsmokers. *CLIN PHARMACOL THER* 1982;31:656-61.
9. Desmond PV, Patwardhan R, Parker R, Shenker S, Speeg KV. Effect of cimetidine and other antihistaminics on the elimination of aminopyrine, phenacetin and caffeine. *Life Sci* 1980;26:1261-8.
10. Beach CA, Mays DC, Guiler RC, Jacober CH, Gerber N. Inhibition of elimination of caffeine by disulfiram in normal subjects and recovering alcoholics. *CLIN PHARMACOL THER* 1986;39:265-70.
11. Abernethy DR, Todd EL. Impairment of caffeine clearance by chronic use of low-dose oestrogen-containing oral contraceptives. *Eur J Clin Pharmacol* 1985;28:425-8.
12. Patwardhan RV, Desmond PV, Johnson RF, Schenker S. Impaired elimination of caffeine by oral contraceptive steroids. *J Lab Clin Med* 1980;95:603-8.
13. Joeres R, Richter E. Mexiletine and caffeine elimination. *N Engl J Med* 1987;317:117.
14. Brazier JL, Descotes J, Lery N, Ollagnier M, Evreux JCI. Inhibition by idroclamide of the disposition of caffeine. *Eur J Clin Pharmacol* 1980;17:37-43.
15. Tarrús E, Camí J, Roberts RG, Spickett RGW, Celdran E, Segura J. Accumulation of caffeine in healthy volunteers treated with furafylline. *Br J Clin Pharmacol* 1987;23:9-18.
16. Mays DC, Camisa C, Cheney P, Pacula CM, Nawoot S, Gerber N. Methoxsalen is a potent inhibitor of metabolism of caffeine in humans. *CLIN PHARMACOL THER* 1987;42:621-6.
17. Tarrús E, García I, Segura J. An animal model to detect the inhibition of the metabolism of caffeine. *Methods Find Exp Clin Pharmacol* 1987;9:311-6.
18. Segura J, Tarrús E. Severe inhibition of caffeine metabolism by xanthine derivatives. Proceedings of the Second European Congress of Biopharmaceutics and Pharmacokinetics. *Exp Pharmacokinet* 1984;2:520-8.
19. Letteron P, Descatoire V, Larey D, Tinel M, Geneve J, Pessayre D. Inactivation and induction of cytochrome P-450 by various psoralen derivatives in rats. *J Pharmacol Exp Ther* 1986;238:685-92.
20. Maesen FPV, Teengs JP, Baur C, Davies BI. Quinolones and raised plasma concentrations of theophylline. *Lancet* 1984;ii:530.

21. Frech K, Shah PM. Drug interaction: theophylline and quinolones. *Quinolones Bull* 1984; i:3.
22. Khan F, Raoof S. Ciprofloxacin in the treatment of respiratory tract infections. *Ciprofloxacin Symposium, Leberkussen. Proceedings Excerpta Medica* 1985:252-6.
23. Davies BI, Maesen FPV, Baur C. Ciprofloxacin in the treatment of acute exacerbations of chronic bronchitis. *Eur J Clin Microbiol* 1986;i:226-31.
24. Wijnands WJA, Vree TB, Van Herwaarden LA. The influence of quinolone derivatives on theophylline clearance. *Br J Clin Pharmacol* 1986;22:677-83.
25. Simpson KJ, Brodie MJ. Convulsions related to enoxacin. *Lancet* 1985;ii:161.
26. Wijnands WJA, Van Herwaarden LA. Enoxacin raises plasma theophylline concentrations. *Lancet* 1984;ii:108.
27. San José Valverde L, Domínguez Lázaro AR, García Cubero A, Rodríguez Mosquera M, Blanco Carmona JG. Efectos del ácido pipemídico en el aclaramiento de teofilinas. *Rev Esp Alergología Inmunol Clin* 1986;1:14-6.
28. Staib AH, Harper P, Mieke S, Beer S, Stille W, Shah P. Gyrase inhibitors impair caffeine elimination in man. *Methods Find Exp Clin Pharmacol* 1987;9:193-8.
29. Shah PM. Drug interaction: caffeine and quinolones. *Quinolones Bull* 1986;2:18.
30. Grant DM, Tang BK, Kalow W. A simple test for acetylator phenotype using caffeine. *Br J Pharmacol* 1984; 19:459-64.
31. Blanchard J, Sawers SJA. The absolute bioavailability of caffeine. *Eur J Clin Pharmacol* 1983;24:93-8.
32. Bonati M, Latini R, Galletti F, Young JF, Tognoni G, Garattini S. Caffeine disposition after oral doses. *CLIN PHARMACOL THER* 1982;32:98-106.
33. Brodie MJ, Boobis AR, Bulpitt CJ, Davies DS. Influence of liver disease and environmental factors on hepatic monooxygenase activity in vitro. *Eur J Clin Pharmacol* 1981;31:39-46.