



Precision dosing in clinical medicine: present and future

Thomas M. Polasek, Sepehr Shakib & Amin Rostami-Hodjegan

To cite this article: Thomas M. Polasek, Sepehr Shakib & Amin Rostami-Hodjegan (2018) Precision dosing in clinical medicine: present and future, Expert Review of Clinical Pharmacology, 11:8, 743-746, DOI: [10.1080/17512433.2018.1501271](https://doi.org/10.1080/17512433.2018.1501271)

To link to this article: <https://doi.org/10.1080/17512433.2018.1501271>



Accepted author version posted online: 16 Jul 2018.
Published online: 20 Jul 2018.



Submit your article to this journal [↗](#)



Article views: 1028



View Crossmark data [↗](#)

EDITORIAL



Precision dosing in clinical medicine: present and future

Thomas M. Polasek^{a,b}, Sepehr Shakib^c and Amin Rostami-Hodjegan^{a,d}

^aCertara, Princeton, NJ, USA; ^bCentre for Medicines Use and Safety, Monash University, Melbourne, Australia; ^cDepartment of Clinical Pharmacology, University of Adelaide, Adelaide, Australia; ^dCentre for Applied Pharmacokinetic Research, University of Manchester, Manchester, UK

ARTICLE HISTORY Received 2 June 2018; Accepted 13 July 2018

KEYWORDS Precision dosing; prescribing; medication safety; physiologically based pharmacokinetics; model-informed precision dosing

1. The ‘problem’ of variability in drug response

There are four potential outcomes when a patient commences a new drug and its effects are assessed at the correct time. First, the new drug provides no clinical benefits and gives the patient intolerable adverse effects (negative benefit: risk). Second, clinical benefits occur but the adverse effects are so severe that the drug must be ceased (negative benefit: risk). Third, there are no clinical benefits or adverse effects because meaningful pharmacological activity is absent in that patient, including any interaction with the immune system (zero benefit: risk). Finally, the patient experiences clinical benefits with little or no adverse effects and treatment continues, ideally with ongoing monitoring of efficacy and vigilance for potential toxicities (positive benefit: risk). Unfortunately, failed drug therapy occurs widely in clinical medicine, i.e. outcomes 1–3. The financial costs to healthcare are considerable, with a recent analysis by the World Health Organization estimating that the cost of medication-related harm globally is about US \$42 billion per annum [1]. No data are available to quantify the cost of ineffective treatment with medication, which is likely to be considerable. This ‘problem’ exists because drug response is highly variable between different patients who take the same dose (inter-individual variability), and because drug response changes in the same patient taking the same dose over time (intra-individual variability). Assuming a correct diagnosis and complete adherence by the patient to the drug regimen, the causes of drug response variability are either pharmacokinetic and/or pharmacodynamic. Pharmacokinetic variability occurs when drug concentration at the site of action differs between patients or changes with time resulting in different responses. Pharmacodynamic variability occurs when pharmacological activity following molecular target binding differs between patients or changes with time at the same drug concentration [2].

There is incongruity between drug development and clinical medicine in how to tackle variability in drug response. Drug development has a one-dose-fits-all culture because it is impractical and cost prohibitive to study too many different doses and too many different types of patients. There are no regulatory requirements to find the best dose, rather, only a safe and effective dose at the study population (not the individual) level. Clinical trials therefore limit the number of doses studied. Participation in clinical trials is also homogenized by tight inclusion criteria. This decreases

response variability and improves the chances of success. For example, many cancer patients are excluded from clinical trials due to age (>65 years old), significant organ dysfunction, prior or concurrent malignancies, comorbidities, and potentially interacting concomitant medications [3]. Clinical trials therefore limit the types of patients studied. Drug therapy fails in the ‘real world’ because limited doses are approved, and because drugs are given to patient types for which established efficacy and safety data are lacking, i.e. a ‘black hole’ in Professional Drug Information (PI)/Summary of Product Characteristics. Prescribers attempt to counteract these difficulties by selecting drugs with fewer or less severe adverse effects, by selecting drugs with lower variability in response, and hence more predictable clinical outcomes, and by individualizing dose. The latter option is ‘precision dosing,’ defined as dose selection by a prescriber for an individual patient at a given time. Importantly, there are examples where the approved dose is no longer used in clinical practice, with safer and more effective dosing having evolved with real world experience, e.g. sunitinib.

2. Current precision dosing

Precision dosing in clinical medicine is needed for drugs with a narrow therapeutic index (TI) and for patients who belong to ‘special populations.’ Therapeutic classes with narrow TI drugs include the antiarrhythmics, anticoagulants, antiepileptics, antineoplastics, aminoglycoside antibiotics, and immunosuppressants. Patients in the special populations are at increased risk of medication-related harm compared to other groups – pediatrics, the elderly, those with renal or hepatic impairment, the infant who is breastfeeding, and patients taking concomitant medications that cause pharmacokinetic or pharmacodynamic drug–drug interactions [4]. The most compelling cases for precision dosing are when a narrow TI drug is essential for a patient in one or more of the special populations.

2.1. Professional drug information/summary of product characteristics and drug monographs

The PI is a regulatory approved document from the manufacturer. Drug monographs (DMs) are contained in commercial and independent drug information resources [5]. Both are used for precision dosing when no robust biomarker is

available to allow empirical dosing (see below), e.g. direct oral anticoagulants. Recommended doses come directly from clinical studies (e.g. a dedicated renal impairment study), or indirectly via population pharmacokinetic \pm pharmacodynamic analyses (pop PK \pm PD) of large clinical datasets. These approaches identify covariates driving inter- and intra-individual variability in exposure or response, including age, gender, weight, body surface area, ethnicity, and renal and hepatic functions (note that some of these covariates correspond to the special populations listed above). The recommended dose is dependent on the value of one, sometimes two, of these covariates for a given patient [6]. For example, the recommended dose of apixaban for stroke prevention in non-valvular atrial fibrillation is halved to 2.5 mg twice daily in patients with at least two of the following characteristics: age ≥ 80 years, total body weight ≤ 60 kg, or serum creatinine ≥ 1.5 mg/dL. However, PI/DM are difficult to apply to patients who have multiple covariates known to influence drug exposure or response, i.e. the unique combination of their covariates has not been studied in clinical trials or addressed by pop PK \pm PD analyses so a 'black hole' exists in PI/DM. Sticking with the apixaban example, it is difficult to choose a dose for an 85-year-old patient who has renal impairment (serum creatinine = 1.7 mg/dL), ischemic heart disease, hypertension, and osteoarthritis, and is taking verapamil, a drug that increases apixaban exposure by inhibiting cytochrome P4503A4 and p-glycoprotein, and ibuprofen, a drug with anti-platelet action that increases the risk of bleeding [7,8].

2.2. Empirical approach

An initial dose is selected and then adjusted based on exposure or response (as measured clinically or through a biomarker, e.g. warfarin and the international normalization ratio, INR). Since there is prescriber feedback and room to move with dose, prescribing requires greater thought than with PI/DMs that stipulate a fixed dose based on a patient covariate. With the empirical approach, the intensity of monitoring and frequency of dose-titration depends on the nature of the medical condition (acute vs. chronic), the TI of the drug (narrow or wide), and the robustness of the biomarker(s) of exposure or response (correlation to clinical outcomes). This is well done in acute settings, such as immunosuppression following whole organ transplantation, and is improving in primary care with automatic reminders to monitor biomarkers of chronic diseases, such as blood HbA1c concentration in patients on anti-diabetics. Whilst the empirical approach allows some dose flexibility, potentially useful drugs are ceased often prematurely due to limited dose options. Indeed, there may be legal ramifications if things go wrong on a dose outside the approved range, and this range may be narrow from the clinical trials.

2.3. Genetic testing (pharmacogenomics)

Germ line and somatic mutations are increasingly important in explaining variability in drug response. The main utility of pharmacogenomics, however, is for drug selection. Examples include the choice of targeted pharmacotherapy

after molecular diagnoses in oncology (e.g. dabrafenib in patients with malignant melanoma harboring *BRAF V600* mutations [9]), or to warn prescribers about unfavorable risks of toxicity (e.g. abacavir hypersensitivity in HIV patients with the *HLA-B*5701* genotype [10]). The case for pharmacogenomics is less clear for dose selection. Some practical guidance is available for polymorphisms in drug metabolizing enzymes and transporters (DMET) [11,12]. For example, a dose reduction of 50% is recommended when initiating several selective serotonin re-uptake inhibitor antidepressants in patients who are cytochrome P4502C19 and cytochrome P4502D6 poor metabolizers [13]. There are early indications that DMET testing has economic value in high-risk patients (e.g. the elderly on polypharmacy [14]), and several large prospective studies of preemptive DMET genotyping on health and economic outcomes are underway in the United States and Europe [15].

2.4. Model-informed

Biosimulation in 'high-impact' cases predicts the dose for a given patient that is most likely to improve efficacy and/or lower toxicity based on their unique values of covariates known to influence drug response [6]. This niche is limited to academic-hospital centers with strong clinical pharmacology expertise, and to trained healthcare providers with access to specialized software and guidelines that typically require a therapeutic drug monitoring service (TDM). Algorithms based on pop PK \pm PD models are available for the initiation of warfarin, although empirical dosing via the INR takes over once treatment has started. For other narrow TI drugs, biosimulation is applied after initiating drug treatment, with a Bayesian approach using pop PK \pm PD models to predict subsequent doses from personalized inputs to the model that include a recent drug concentration. This approach is more powerful than dose recommendations in PI/DMs because it combines the flexibility of empirical dosing via biomarker feedback whilst accounting for multiple patient covariates that determine variability in drug response [16]. Examples include model-informed precision dosing (MIPD) of antibiotics in the critically ill [17], the use of models to suggest metformin dose in patients with renal impairment [18], and MIPD of immunosuppressants and chemotherapy in serious pediatric illnesses [19,20].

3. Future precision dosing

Precision dosing is set to become more sophisticated. Affordable omics technologies (genomics, transcriptomics, proteomics, and metabolomics), improved medical imaging, rapid pathology testing, characterization of the gut microbiome, superior analysis of biological samples, and powerful computational tools to analyze 'big data' are all important factors in advancing precision dosing. Our understanding of inter-individual and intra-individual variability in drug response now goes beyond simple patient covariates (e.g. age, gender, weight, etc.) to include the 'hidden' physiological and molecular determinates of pharmacokinetics and pharmacodynamics – organ sizes, blood flows, DMET activities, inflammatory status, gut microbiome,

genetics of molecular targets, etc. [21,22]. An improved understanding of disease heterogeneity as a source of pharmacodynamic variability is also increasingly recognized, e.g. doses of the IgE-specific monoclonal antibody omalizumab are determined by age, weight, and pretreatment serum plasma IgE concentration in moderate to severe allergic asthma [23].

Capitalizing on the complexity of inter-related patient information to predict dose is only possible with biosimulation. In addition to pop PK \pm PD models linked to TDM, physiologically based (PB) PK \pm PD models and quantitative systems pharmacology (QSP) approaches will become important tools for precision dosing. These allow generation of a patient's virtual twin based on their unique physiological and molecular characteristics, including pharmacogenomics. In theory, prescribers could predict drug response prior to treatment, as concomitant drugs change, and as medical conditions progress or improve [16]. Model-informed approaches are now revolutionizing drug development [24,25], and efforts are underway to repurpose PBPK \pm PD and QSP for clinical application [26–30]. Importantly, the model-informed approach is a way to distill large quantities of drug-related and patient-related information into relatively simple dosing instructions. For example, a recent study identified abundance of hepatic cytochrome P4503A4 as the major driver of axitinib exposure, accounting for >90% of inter-individual variability (Dr Andrew Rowland, personal communication). When the technology becomes available, a blood test result of this single covariate may be sufficient to predict an axitinib dose that improves the chances of survival for a patient with metastatic renal cell carcinoma prior to treatment and/or when TDM is unavailable [31].

Economic incentives that encourage precision dosing in drug development will improve precision dosing in clinical medicine. Value-based or outcome-based pricing is when a drug is paid for only if it works, i.e. outcome 4. Compared to fixed-pricing, there would be greater incentive for manufacturers to find the best dose and to find the types of patients that benefit the most. Exposure-controlled clinical trials and response-controlled clinical trials may be needed to generate this evidence, although these are expensive and logistically more challenging than current trial designs. However, the provision of lower doses for certain patient populations would lower manufacturing costs, whilst patients who fail drug therapy could be treated successfully at a higher dose [2].

In conclusion, this editorial outlines why variability in drug response occurs in clinical medicine and describes the current precision dosing strategies to address the problem. Model-informed approaches that incorporate the unique physiological and molecular characteristics of individual patients will become increasingly important for precision dosing in the era of precision medicine.

Funding

This manuscript was not funded but was supported by Certara, the makers of biosimulation tools to aide drug development.

Declaration of interest

TM Polasek and A Rostami-Hodjegan are both employees of Certara. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

Reviewer disclosures

Peer reviewers on this manuscript have no relevant financial or other relationships to disclose.

References

Papers of special note have been highlighted as either of interest (*) or of considerable interest (**) to readers.

1. World Health Organization. Medication without harm – global patient safety challenge on medication safety. Geneva: World Health Organization; 2017.
2. Peck RW. Precision medicine is not just genomics: the right dose for every patient. *Annu Rev Pharmacol Toxicol*. 2018;58:105–122.
**** An excellent review of the sources of inter-individual variability in drug response and the ways to improve the personalization of drug therapy.**
3. Masters JC, Wiernik PH. Are we ready to include organ-impaired patients in oncology trials? A clinical pharmacology perspective on recent recommendations. *J Clin Pharmacol*. 2018;58(6):701–703.
4. Snyder BD, Polasek TM, Doogue MP. Drug interactions: principles and practice. *Australian Prescriber*. 2012;35(3):85–88.
5. Rossi S, Hurley E, Sutcliffe A. Australian medicines handbook. 19 ed. Adelaide: The Royal Australian College of General Practitioners, the Australasian Society of Clinical and Experimental Pharmacologists and Toxicologists and the Pharmaceutical Society of Australia; 2018.
6. Darwich AS, Ogungbenro K, Vinks AA, et al. Why has model-informed precision dosing not yet become common clinical reality? Lessons from the past and a roadmap for the future. *Clin Pharmacol Ther*. 2017;101(5):646–656.
**** The first 'white paper' for model-informed precision dosing, describing the clinical reality and the ways forward to address challenges.**
7. Forbes HL, Polasek TM. Potential drug–drug interactions with direct oral anticoagulants in elderly hospitalized patients. *Ther Adv Drug Saf*. 2017;8(10):319–328.
8. Polasek TM, Lin FPY, Miners JO, et al. Perpetrators of pharmacokinetic drug–drug interactions from altered cytochrome P450 activity: a criteria-based assessment. *Br J Clin Pharmacol*. 2011;71(5):727–736.
9. Polasek TM, Ambler K, Scott HS, et al. Targeted pharmacotherapy after somatic cancer mutation screening. *F1000Res*. 2016;5:1551.
10. Martin MA, Klein TE, Dong BJ, et al. Clinical pharmacogenetics implementation consortium guidelines for HLA-B genotype and abacavir dosing. *Clin Pharmacol Ther*. 2012;91(4):734–738.
11. Relling MV, Klein TE. CPIC: clinical pharmacogenetics implementation consortium of the pharmacogenomics research network. *Clin Pharmacol Ther*. 2011;89(3):464–467.
• Paper introducing the structure, aims and importance of the US Clinical Pharmacogenomics Implementation Consortium.
12. Swen JJ, Nijenhuis M, De Boer A, et al. Pharmacogenetics: from bench to byte-an update of guidelines. *Clin Pharmacol Ther*. 2011;89(5):662–673.
• This paper describes clinically actionable pharmacogenomic information as per the Dutch Pharmacogenomics Working Group.
13. Hicks JK, Bishop JR, Sangkuhl K, et al. Clinical pharmacogenetics implementation consortium (CPIC) guideline for CYP2D6 and

- CYP2C19 genotypes and dosing of selective serotonin reuptake inhibitors. *Clin Pharmacol Ther.* **2015**;98(2):127–134.
14. Brixner D, Biltaji E, Bress A, et al. The effect of pharmacogenetic profiling with a clinical decision support tool on healthcare resource utilization and estimated costs in the elderly exposed to polypharmacy. *J Med Econ.* **2016**;19(3):213–228.
 15. Van Der Wouden CH, Cambon-Thomsen A, Cecchin E, et al. Implementing pharmacogenomics in Europe: design and implementation strategy of the ubiquitous pharmacogenomics consortium. *Clin Pharmacol Ther.* **2017**;101(3):341–358.
 16. Tucker GT. Personalized drug dosage – closing the loop. *Pharm Res.* **2017**;34:1539–1543.
 17. Tängdén T, Ramos Martín V, Felton TW, et al. The role of infection models and PK/PD modelling for optimising care of critically ill patients with severe infections. *Intensive Care Med.* **2017**;43(7):1021–1032.
 18. Duong JK, Kroonen MYAM, Kumar SS, et al. A dosing algorithm for metformin based on the relationships between exposure and renal clearance of metformin in patients with varying degrees of kidney function. *Eur J Clin Pharmacol.* **2017**;73(8):981–990.
 19. Vinks AA, Emoto C, Fukuda T. Modeling and simulation in pediatric drug therapy: application of pharmacometrics to define the right dose for children. *Clin Pharmacol Ther.* **2015**;98(3):298–308.
 20. Darwich AS, Ogungbenro K, Hatley OJ, et al. Role of pharmacokinetic modeling and simulation in precision dosing of anticancer drugs. *Transl Cancer Res.* **2017**;6(S10):S1512–S1529.
 21. Hauser, AS, Chavali S, Masuho I, et al. Pharmacogenomics of GPCR drug targets. *Cell.* **2017**;172:41–59.
 22. Morgan ET, Dempsey JL, Mimche SM, et al. Physiological regulation of drug metabolism and transport: pregnancy, microbiome, inflammation, infection and fasting. *Drug Metab Dispos.* **2018**;46:503–513.
 23. Peck RW. The right dose for every patient: a key step for precision medicine. *Nat Rev Drug Discov.* **2016**;15(3):145–146.
 24. Allerheiligen SR. Impact of modeling and simulation: myth or fact? *Clin Pharmacol Ther.* **2014**;96(4):413–415.
 - **A nice perspective piece describing how modeling and simulation saves the Pharmaceutical Industry time and money.**
 25. Milligan PA, Brown MJ, Marchant B, et al. Model-based drug development: a rational approach to efficiently accelerate drug development. *Clin Pharmacol Ther.* **2013**;93(6):502–514.
 26. Rowland A, Van Dyk M, Hopkins AM, et al. Physiologically based pharmacokinetic modeling to identify physiological and molecular characteristics driving variability in drug exposure. *Clin Pharmacol Ther.* **2018**. doi:10.1002/cpt.1076. [Epub ahead of print]
 - **The first paper to show that Simcyp® can be used to identify covariates that influence drug pharmacokinetics.**
 27. Marsousi N, Samer CF, Fontana P, et al. Coadministration of ticagrelor and ritonavir: toward prospective dose adjustment to maintain an optimal platelet inhibition using the PBPK approach. *Clin Pharmacol Ther.* **2016**;100(3):295–304.
 28. Patel N, Wiśniowska B, Jamei M, et al. Real patient and its virtual twin: application of quantitative systems toxicology modelling in the cardiac safety assessment of citalopram. *AAPS J.* **2017**;20(1):6.
 29. Polasek TM, Tucker GT, Sorich MJ, et al. Prediction of olanzapine exposure in individual patients using physiologically based pharmacokinetic modelling and simulation. *Br J Clin Pharmacol.* **2018**;84(3):462–476.
 - **This paper describes the first use of physiological-based pharmacokinetic modeling and simulation to predict drug exposure in patients rather than in a drug development setting.**
 30. Dinh JC, Black BT, Matzuka BJ, et al. Prospective validation of a population pharmacokinetic model derived dosing algorithm of atomoxetine, a CYP2D6 model substrate. *Clin Pharmacol Ther.* **2018**;103(S1):S29–S30.
 31. Rini BI, Garrett M, Poland B, et al. Axitinib in metastatic renal cell carcinoma: results of a pharmacokinetic and pharmacodynamic analysis. *J Clin Pharmacol.* **2013**;53(5):491–504.