Rivaroxaban PBPK/PD model: fit PD

1 Methods: Rivaroxaban PBPK/PD model: fit PD

1.1 Parameters and data

5 parameters were optimized: relative increase of the PT per rivaroxaban concentration (`PT_slope`), maximum effect aPTT (`Emax_aPTT`), EC50 rivaroxaban aPTT (`EC50_riv_aPTT`), maximum effect Xa inhibition (`Emax_Xa`) and EC50 rivaroxaban Xa inhibition (`EC50_riv_Xa`). The parameters of the adopted fit PK_20261018_075718__82026 were fixed at their fitted values.

A fit mapping is a curve of a study, the data and the observable of the model it is compared with. The fit has 64 fit mappings of training data with 823 data points, whose residuals enter the objective; 39 fit mappings of validation data with 1043 data points, which were simulated and evaluated with the fitted parameters but not fitted; and 3 fit mappings of outliers with 45 data points, data the fit dropped as not usable, which were evaluated like the validation data so that the decision to drop them can be checked. The fit was set up and run with sbmlsim [1].

1.2 Objective function

The parameters were estimated by minimizing the objective function

$$F(\vec{p}) = \frac{1}{2} \sum_k \sum_i \rho\left(w_{i,k} r_{i,k}(\vec{p})^2\right) \tag{1}$$

where k runs over the fit mappings of the training data and i over their data points, $r_{i,k}$ is the residual, $w_{i,k} = w_k w_{i,k}^p$ the weight of the data point and ρ the loss function, applied to the squared weighted residual $z = w_{i,k} r_{i,k}^2$. The loss function is linear, $\rho(z) = z$, i.e. a standard least squares problem.

The residuals are normalized by the mean $\overline{y}_k$ of the data of their curve, which makes curves of different quantities and magnitudes comparable:

$$r_{i,k} = \frac{y_{i,k} - f_{k(x_{i,k},\vec{p})}}{\overline{y}_k} \tag{2}$$

The weight of curve k is $w_k = \frac{n_k}{N_k}$, the weight n_k of its fit mapping (the weight its fit mapping collection assigns, 1 by default, or, for a collection which uses the weights of its mappings, the weight of the mapping, by default the number of subjects of its data) divided by the number N_k of its data points.

A data point is weighted with the inverse of its coefficient of variation, $w_{i,k}^p = |y_{i,k} \frac{|}{\sigma_{i,k}}$, with $\sigma_{i,k}$ the standard deviation of the data point, or its standard error for data without a standard deviation. An error of zero is missing, a missing error of a curve with errors is replaced by the largest error of the curve, and the data points of a curve without errors have the weight $w_{i,k}^p = 2$, the weight of a coefficient of variation of 0.5.

1.3 Optimization

The optimizer searched the parameters in log10 space within their bounds; bounds, values and intervals are given in the units of the fit parameters, the units of the table of the parameters.

The objective was minimized in 3 runs with differential evolution [2] of SciPy [3] within the bounds of the parameters, with the strategy `best1bin`, a population of 15 times the number of parameters, at most 1000 generations, the relative tolerance 0.01 of convergence, the recombination constant 0.7 and a mutation dithered between 0.5 and 1. The initial population of a run was drawn by latin hypercube sampling [4], with a seed of every run derived from the seed 1234 of the fit. The best member of the final population was polished with the L-BFGS-B method of SciPy.

The convergence of the runs was assessed from the traces of their cost and the waterfall plot of their final costs, and the parameter set with the lowest cost was used.

The models were simulated with libRoadRunner [5] and its integrator CVODE [6] with the relative tolerance 10^{-6} and the absolute tolerances 10^{-13} of an amount, 10^{-13} of a concentration and 10^{-6} of the other states, which for a species in a concentration is multiplied by the initial volume of its compartment. It ran with the variable step size of libRoadRunner, i.e. with an output at every step of the integrator. A data point at the time of a change after the start of a simulation, e.g. a dose or the reset of a collection, was compared with the state before the change, except the first data point of a curve with data at later times, which starts a collection and was compared with the state after the change.

1.4 Goodness of fit

The goodness of fit is given per kind of data (training data, validation data and outliers) by metrics over its n data points with the residuals $e_i = y_i - f\left(x_i, \hat{\vec{p}}\right)$ at the fitted parameters $\hat{\vec{p}}$, the mean squared error, its root and the root mean squared error normalized by the mean $\overline{y}_{k(i)}$ of the data of the curve of a point:

$$\mathrm{MSE} = \frac{1}{n}\sum_{i=1}^{n} e_i^2, \quad \mathrm{RMSE} = \sqrt{\mathrm{MSE}}, \quad \mathrm{NRMSE} = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\left(\frac{e_i}{\overline{y}_{k(i)}}\right)^2} \tag{3}$$

The root mean square of the residuals of the objective, i.e. what the fit minimizes, and the coefficient of determination:

$$\mathrm{RMSE}_w = \sqrt{\frac{1}{n}\sum_{i=1}^{n}\rho(w_i r_i^2)}, \quad R^2 = 1 - \frac{\sum_i e_i^2}{\sum_i \left(y_i - \overline{y}\right)^2} \tag{4}$$

The predictions of a non-linear model are no linear regression of the data, so R^2 can be negative: the prediction is worse than the mean of the data. The information criteria of Akaike [7] and Schwarz [8] with the number n_p of fitted parameters are those of a least squares fit with normally distributed residuals, up to an additive constant, so only differences between models fitted to the same data are meaningful:

$$\mathrm{AIC} = n\ln(\mathrm{MSE}) + 2n_p, \quad \mathrm{BIC} = n\ln(\mathrm{MSE}) + n_p\ln(n) \tag{5}$$

The percentage of the data points whose prediction $f_i = f\left(x_i, \hat{\vec{p}}\right)$ lies within twofold of the data is taken over the n_+ data points with positive data $y_i > 0$ and a positive prediction $f_i > 0$; the others have no ratio and are not counted, and a kind of data without such a point has no percentage:

$$P_2 = \frac{100}{n_+}\left|\left\{i : \frac{1}{2} \le \frac{f_i}{y_i} \le 2\right\}\right| \tag{6}$$

The goodness of fit plot shows the predictions against the data on logarithmic axes, with the lines of twofold $f = 2y$ and $f = \frac{y}{2}$. The Bland-Altman plot [9] shows the ratio of prediction and data as $d_i = \log_{10}\left(\frac{f_i}{y_i}\right)$ against the geometric mean $\sqrt{f_i y_i}$ for the data points with positive values, with the bias $\overline{d}$ and the limits of agreement $\overline{d} \pm 1.96 s_d$, with the sample standard deviation s_d of d_i, of the training data; they are read as the fold factors $10^{\overline{d}}$ and $10^{\overline{d} \pm 1.96 s_d}$.

1.5 Identifiability

The identifiability of the parameters was analysed with the profile likelihood [10], [11]. The profile of a parameter is the lowest objective with the parameter fixed:

$$\mathrm{PL}(p_j) = \min_{p_{l \ne j}} F(\vec{p}) \tag{7}$$

At every point of a scan the other parameters were re-optimized from the previous point with the least squares optimizer of SciPy with the relative step 0.05. A scan ran from the optimum in both directions in the space of the optimizer, with steps in decades of the parameter, starting at 0.1. A step which raised the objective by more than 0.2 of the distance between the objective at the optimum and the threshold was reduced by the factor 2 and repeated, down to 0.01, and the step after one which raised it by less than that divided by the factor was enlarged by the factor, up to 1. A scan stopped at the threshold, at the bound of the parameter, at a point which failed or after 50 points.

The confidence intervals are the values at which the profile crosses the threshold of the likelihood ratio test

$$F_{\mathrm{thr}} = F_{\mathrm{min}} + \chi^2_\alpha \frac{\mathrm{df}}{2} \tag{8}$$

with F_{min} the lowest objective of the fit and of all profiles and $\chi^2_\alpha(\mathrm{df})$ the quantile of the χ^2 distribution at the confidence level $\alpha = 0.95$ with df = 1 degrees of freedom, i.e. 1.92 above the lowest objective. The threshold has this confidence level if F is the negative log-likelihood of the data up to a constant, i.e. if the weighted residuals are independent and normally distributed with unit variance; otherwise the level is approximate. A crossing is interpolated between two points of the profile, on the logarithm of a parameter on a logarithmic scale. A parameter is identifiable if its profile reaches the threshold within the scan on both sides of the optimum, i.e. its confidence interval is finite; practically non-identifiable towards the lower bound if its profile reaches the threshold above the optimum but not below it within the scan, which ends at the lower bound or after 50 points; practically non-

identifiable towards the upper bound if its profile reaches the threshold below the optimum but not above it within the scan, which ends at the upper bound or after 50 points; practically non-identifiable if its profile reaches the threshold on neither side within the scans, which end at the bounds or after 50 points, and is not flat; and structurally non-identifiable if its profile is flat on both sides. A side of a profile is flat if it rises by less than 0.05 of the distance between the lowest objective and the threshold, and a profile with a point which failed or did not converge is not classified. The profile likelihood is preferred over the intervals of the Fisher information, its intervals are invariant under a transformation of the parameters and can be asymmetric [12].

The Fisher information is the local alternative to the profiles: it is the Gauss-Newton approximation of the curvature of the objective at the optimum, from the Jacobian J of the weighted residuals of the training data with respect to the parameters $\vec{p}$ in the space of the optimizer, computed by central finite differences with the step $h_j = \varepsilon \max\left(|p_j|, 1\right),\ \varepsilon = 10^{-6}$ of the parameter p_j:

$$I = J^{\top} J, \quad \mathrm{cov} = \sigma^2 I^{-1}, \quad \sigma^2 = \frac{2F}{n - n_p} \tag{9}$$

with the number n of data points of the training data and the number n_p of parameters. The information has rank 5 of 5, the number of its eigenvalues above 10^{-8} of the largest one, and the condition number 2420; the covariance of an information of a lower rank is computed with the pseudo-inverse. The Wald intervals are $\hat{p}_j \pm t_{\frac{1+\alpha}{2}, n-n_p} \mathrm{SE}_j$ with the standard errors $\mathrm{SE}_j = \sqrt{\mathrm{cov}_{jj}}$ at the confidence level $\alpha = 0.95$, transformed back into the units of the fit parameters.

1.6 Provenance

The fit PD_20261018_082509__94e87 was run on 2026-10-18 06:25:12 UTC with sbmlsim 0.10.0 [1], libRoadRunner 2.10.0 [5], SciPy 1.18.1 [3], NumPy 2.5.3 [13] and Python 3.14.4 (seed 1234). The models and data were those of commit `b15d1ca073221f8e25d56c4caef1fef784205bf2` of `/home/mkoenig/git/pkdb_models-rivaroxaban` their SHA-256 checksums are listed in Table 1. The fit kept the values `GU__RIVABS_k` = 0.0818778, `GU__Ka_emptying_riv` = 11.7785, `GU__Vmax_dis_riv` = 4.03659, `GU__ktransit_riv` = 0.5, `GU__FED_f_solubility` = 1.60754, `GU__FED_f_emptying` = 0.202834, `LI__RIV2RX_k` = 0.0482107, `KI__RIVEX_f` = 6.39515, `Kp_riv` = 0.407069, `ftissue_riv` = 5.80765 and `LI__RXEX_BI_k` = 0.179531 of the model fixed.

Table 1: SHA-256 checksums of the models and data of the fit PD_20261018_082509__94e87, their first 12 characters; the full checksums are in the provenance of the result (optimization_result.json).

File	SHA-256
pkdb_models/models/rivaroxaban/data/apixaban/Frost2014/.Frost2014_Fig2.tsv	df47c941fd84...
pkdb_models/models/rivaroxaban/data/apixaban/Frost2014/.Frost2014_Fig3.tsv	10b5b148e803...
pkdb_models/models/rivaroxaban/data/apixaban/Frost2014/.Frost2014_Fig4.tsv	de5f58673bdc...
pkdb_models/models/rivaroxaban/data/apixaban/Frost2014/.Frost2014_Fig4Mean.tsv	20c5a8eeb2f7...
pkdb_models/models/rivaroxaban/data/apixaban/Frost2014/.Frost2014_Tab2.tsv	73d67b04ee04...
pkdb_models/models/rivaroxaban/data/apixaban/Frost2014/.Frost2014_Tab3.tsv	dc6374a9c6a9...
pkdb_models/models/rivaroxaban/data/apixaban/Frost2014/.Frost2014_TabGroups.tsv	dfaab6207c67...
pkdb_models/models/rivaroxaban/data/apixaban/ Frost2014/.Frost2014_TabIndividuals.tsv	6cef70f6b764...
pkdb_models/models/rivaroxaban/data/apixaban/ Frost2014/.Frost2014_TabInterventions.tsv	a2285172067a...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_Fig2.tsv	94243c590984...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_Fig3.tsv	216b03d0818a...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_Fig4.tsv	6a6c7f642285...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_Fig6.tsv	6d31b7e2bebe...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_FigS1.tsv	56fb1bd401d4...
pkdb_models/models/rivaroxaban/data/apixaban/ Kreutz2017/.Kreutz2017_FigS1Mean.tsv	e40ddaad8a2b...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_FigS3.tsv	f2b6ba261408...

File	SHA-256
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_Tab2.tsv	1bf835d0716b...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_TabGroups.tsv	e9c0b77517ec...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_TabInterventions.tsv	926a1221fc08...
pkdb_models/models/rivaroxaban/data/apixaban/Kreutz2017/.Kreutz2017_TabS2.tsv	4dd36fd908e6...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2024/.Lenard2024_Fig2.tsv	6ef59c8303ca...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2024/.Lenard2024_Tab1.tsv	e22229078327...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2024/.Lenard2024_Tab2.tsv	7c2cf210c32d...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2024/.Lenard2024_TabGroups.tsv	1563a66c5472...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2024/.Lenard2024_TabInterventions.tsv	028d640d560c...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2024/.Lenard2024_Tab_Supplement.tsv	3ba8777b45fe...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2025/.Lenard2025_Fig2.tsv	2c0e1a6bc114...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2025/.Lenard2025_Fig3.tsv	4ad80c096bcc...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2025/.Lenard2025_Fig4.tsv	f3dac8691767...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2025/.Lenard2025_Tab1.tsv	0651a8e8e14e...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2025/.Lenard2025_Tab2.tsv	8dae4e8812a1...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2025/.Lenard2025_Tab3.tsv	f7b8f4279c5e...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2025/.Lenard2025_TabGroups.tsv	1e834066fe0e...
pkdb_models/models/rivaroxaban/data/edoxaban/Lenard2025/.Lenard2025_TabInterventions.tsv	2477fb8a8c7e...
pkdb_models/models/rivaroxaban/data/edoxaban/Metze2021/.Metze2021_Fig1.tsv	914a8f2bc513...
pkdb_models/models/rivaroxaban/data/edoxaban/Metze2021/.Metze2021_Fig1A.tsv	74b4252228da...
pkdb_models/models/rivaroxaban/data/edoxaban/Metze2021/.Metze2021_Tab2.tsv	9c4de7b2dd65...
pkdb_models/models/rivaroxaban/data/edoxaban/Metze2021/.Metze2021_TabGroups.tsv	5096f3d68a1f...
pkdb_models/models/rivaroxaban/data/edoxaban/Metze2021/.Metze2021_TabIndividuals.tsv	632b916218ea...
pkdb_models/models/rivaroxaban/data/edoxaban/Metze2021/.Metze2021_TabInterventions.tsv	5abff06c2e57...
pkdb_models/models/rivaroxaban/data/edoxaban/Rohr2024/.Rohr2024_Fig2.tsv	124cfd04b982...
pkdb_models/models/rivaroxaban/data/edoxaban/Rohr2024/.Rohr2024_Tab1.tsv	50ef32d48f1e...
pkdb_models/models/rivaroxaban/data/edoxaban/Rohr2024/.Rohr2024_Tab1A.tsv	f939a77c6620...
pkdb_models/models/rivaroxaban/data/edoxaban/Rohr2024/.Rohr2024_TabGroups.tsv	5ef7ccc355e2...
pkdb_models/models/rivaroxaban/data/edoxaban/Rohr2024/.Rohr2024_TabInterventions.tsv	544c2fd7c13c...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_Fig2.tsv	3c26410b1cd4...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_Fig3.tsv	37e53911132b...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_Fig4a.tsv	fa7acdee3717...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_Fig4b.tsv	2cca7875276e...

File	SHA-256
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_Tab2.tsv	e224637202b5...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_Tab3.tsv	7c2ab22d384b...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_Tab4.tsv	233b5f2d3074...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_TabGroups.tsv	949a47df1174...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Alalawneh2024/.Alalawneh2024_TabInterventions.tsv	b0ee0b24568f...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Ding2020/.Ding2020_Fig1.tsv	eb8f2e57e283...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Ding2020/.Ding2020_Fig2.tsv	c03d27128ae5...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Ding2020/.Ding2020_Tab2.tsv	ac3b885f1eb2...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Ding2020/.Ding2020_Tab3.tsv	592fb9c52218...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Ding2020/.Ding2020_TabGroups.tsv	179f5dc32ec3...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Ding2020/.Ding2020_TabInterventions.tsv	ac8688a556ee...
pkdb_models/models/rivaroxaban/data/rivaroxaban/GenisNajera2022/.GenisNajera2022_Fig1.tsv	55d0d4190d89...
pkdb_models/models/rivaroxaban/data/rivaroxaban/GenisNajera2022/.GenisNajera2022_Tab2.tsv	c569910d6980...
pkdb_models/models/rivaroxaban/data/rivaroxaban/GenisNajera2022/.GenisNajera2022_TabGroups.tsv	70d18910fdfb...
pkdb_models/models/rivaroxaban/data/rivaroxaban/GenisNajera2022/.GenisNajera2022_TabInterventions.tsv	7d4551bc61ac...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Graff2007/.Graff2007_Fig3A.tsv	32632877eec7...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Graff2007/.Graff2007_Fig3B.tsv	c6bc2b44d774...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Graff2007/.Graff2007_Fig5.tsv	97e80b0c5b30...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Graff2007/.Graff2007_TabGroups.tsv	a1797cbf760c...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Graff2007/.Graff2007_TabInterventions.tsv	c58e8c0ae6a9...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Greenblatt2018/.Greenblatt2018_Fig1.tsv	cb6ed3dca65a...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Greenblatt2018/.Greenblatt2018_Fig2.tsv	18fad2493c38...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Greenblatt2018/.Greenblatt2018_Fig5.tsv	3245e202448a...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Greenblatt2018/.Greenblatt2018_Tab2.tsv	34a28fc41330...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Greenblatt2018/.Greenblatt2018_TabGroups.tsv	d4944dea98ee...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Greenblatt2018/.Greenblatt2018_TabInterventions.tsv	a8f2bb4fcc08...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Jiang2010/.Jiang2010_Fig1.tsv	68be58ef66b3...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Jiang2010/.Jiang2010_Fig2.tsv	2c1a23e24ff0...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Jiang2010/.Jiang2010_Fig3.tsv	8f9cba4cdda3...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Jiang2010/.Jiang2010_Tab2.tsv	3f2c8f70e56c...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Jiang2010/.Jiang2010_Tab2A.tsv	0b0b6f9851cd...

File	SHA-256
pkdb_models/models/rivaroxaban/data/rivaroxaban/Jiang2010/.Jiang2010_Tab3.tsv	9ab42405a779...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Jiang2010/.Jiang2010_TabGroups.tsv	3a4a37cc707f...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Jiang2010/.Jiang2010_TabInterventions.tsv	dbcc5ef00640...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig1A.tsv	00051f41af0b...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig1B.tsv	e4b6cb7b3962...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig1C.tsv	08c6a286b869...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig1D.tsv	0b79b41a13de...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig2A.tsv	4465f37e7977...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig2B.tsv	c75a5bfe2b89...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig2C.tsv	aaf9d0d9cd66...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig2D.tsv	23a9024da761...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Fig3.tsv	bf13808f993c...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_Tab2.tsv	2addafd9ab39...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_TabGroups.tsv	09932960d40e...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005/.Kubitza2005_TabInterventions.tsv	056f18775973...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Fig3.tsv	aa0a412f0292...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Fig3A.tsv	ab2b7ad64927...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Fig3B.tsv	1df0fd85175c...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Fig3C.tsv	6c187f419416...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Fig3D.tsv	bb19174927d0...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Fig3E.tsv	c4c82400bd26...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Fig3F.tsv	10d2a7c9784c...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Fig5.tsv	d2e64e78579c...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_Tab2.tsv	94e9646fc2fa...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_TabGroups.tsv	486d17d8d092...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2005a/.Kubitza2005a_TabInterventions.tsv	e021d6953849...

File	SHA-256
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_Fig2.tsv	a52fb7b96bb0...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_Fig3.tsv	a4387993e608...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_Fig4.tsv	127065a8ec75...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_Fig5.tsv	cb3d7c2451db...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_Tab2.tsv	72b919a49462...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_Tab4.tsv	b796365cb6aa...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_Tab6.tsv	653d687daf1a...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_TabGroups.tsv	8c1256d9b1b5...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2006/.Kubitza2006_TabInterventions.tsv	5938603d330e...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2013/.Kubitza2013_Fig1.tsv	cba1051845d9...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2013/.Kubitza2013_Fig2A.tsv	86661263a3dd...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2013/.Kubitza2013_Fig2B.tsv	d18d61ef6558...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2013/.Kubitza2013_Tab2.tsv	335176718043...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2013/.Kubitza2013_TabGroups.tsv	16e939f1ee8a...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Kubitza2013/.Kubitza2013_TabInterventions.tsv	bd327d70b8e6...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Liu2024/.Liu2024_Fig2.tsv	707729a78861...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Liu2024/.Liu2024_Fig3.tsv	4fa439064439...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Liu2024/.Liu2024_Fig3b.tsv	e6ce95ed8237...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Liu2024/.Liu2024_Tab2.tsv	d782e58bf3ae...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Liu2024/.Liu2024_TabGroups.tsv	a03376f1e3f8...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Liu2024/.Liu2024_TabInterventions.tsv	4364bca90d4e...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Moore2014/.Moore2014_Fig2.tsv	f4fd44a21f12...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Moore2014/.Moore2014_Tab2.tsv	37b38e946068...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Moore2014/.Moore2014_Tab23.tsv	fe7a6b854ead...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Moore2014/.Moore2014_Tab2A.tsv	348c8aea5cae...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Moore2014/.Moore2014_TabGroups.tsv	ef8e9d1470fe...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Moore2014/.Moore2014_TabInterventions.tsv	2efb0800398f...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Santos2024/.Santos2024_Fig1.tsv	870de5cb4686...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Santos2024/.Santos2024_Tab2.tsv	d4fb84005d2f...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Santos2024/.Santos2024_TabGroups.tsv	195e010da909...

File	SHA-256
pkdb_models/models/rivaroxaban/data/rivaroxaban/Santos2024/.Santos2024_TabInterventions.tsv	0e6d426eaffd...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Tang2025/.Tang2025_Fig1.tsv	3957a7c25ac5...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Tang2025/.Tang2025_Tab1.tsv	c33102eef944...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Tang2025/.Tang2025_TabGroups.tsv	67a5d0f66a72...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Tang2025/.Tang2025_TabInterventions.tsv	5549a37c6b41...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Weinz2009/.Weinz2009_Fig2.tsv	4f3041d1089f...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Weinz2009/.Weinz2009_Fig3.tsv	bfd9c8edab22...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Weinz2009/.Weinz2009_Tab2.tsv	ba3448d628fb...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Weinz2009/.Weinz2009_Tab3.tsv	e9bbf4bc2e0b...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Weinz2009/.Weinz2009_Tab3_4.tsv	8558c48b6689...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Weinz2009/.Weinz2009_Tab4.tsv	ee7ada59e487...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Weinz2009/.Weinz2009_TabGroups.tsv	ea8bd11147b3...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Weinz2009/.Weinz2009_TabInterventions.tsv	dc31b3dadf20...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Zhao2009/.Zhao2009_Fig1A_2A.tsv	2360e79c86d7...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Zhao2009/.Zhao2009_Fig1B_2B.tsv	0b0bcd996ca2...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Zhao2009/.Zhao2009_Fig2C.tsv	52f245dc89e0...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Zhao2009/.Zhao2009_Tab2_4.tsv	69eca24f723c...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Zhao2009/.Zhao2009_Tab2_4U.tsv	e23964826282...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Zhao2009/.Zhao2009_TabGroups.tsv	dadc31a4fdc6...
pkdb_models/models/rivaroxaban/data/rivaroxaban/Zhao2009/.Zhao2009_TabInterventions.tsv	fe4ba4604d30...
pkdb_models/models/rivaroxaban/data/rivaroxaban/bland_altman_Fig3b.tsv	b149f817f4e0...
pkdb_models/models/rivaroxaban/models/output/models/rivaroxaban_body_flat.xml	c4a224f59919...

2 Results: Rivaroxaban PBPK/PD model: fit PD

2.1 Parameters

Table 2: Parameters of the fit PD_20261018_082509__94e87, the parameter set PD_20261018_082509__94e87_0, cost 68.7, with their bounds in the optimization. The 95% confidence intervals are those of the profile likelihood, LB and UB an interval which is open towards the lower or the upper bound, and the Wald intervals of the Fisher information for a parameter without a profile. Identifiable is the classification of the profile: yes, no (LB) or no (UB) if the profile does not reach the threshold towards the lower or the upper bound, no if it reaches it on neither side, structural if it is flat.

Parameter	Description	Value [95% CI]	Unit	Bounds	Identifiable
PT_slope	relative increase of the PT per rivaroxaban concentration	628 [453, 803]	1/(mmol/l)	[100, 1e+04]	yes
Emax_aPTT	maximum effect aPTT	1.6 [0.538, UB]	dimensionless	[0.1, 5]	no (UB)
EC50_riv_aPTT	EC50 rivaroxaban aPTT	0.000845 [0.000133, 0.0044]	mmol/l	[1e-05, 0.01]	yes
Emax_Xa	maximum effect Xa inhibition	1 [0.81, UB]	dimensionless	[0.5, 1]	no (UB)
EC50_riv_Xa	EC50 rivaroxaban Xa inhibition	0.000558 [0.00041, 0.000623]	mmol/l	[1e-05, 0.01]	yes

2.2 Data

Table 3: Data of the fit PD_20261018_082509__94e87: the number of fit mappings and, in parentheses, of data points per study and kind. The training data enters the cost, the validation data and the outliers are evaluated only.

Study	Training	Validation	Outlier	Total
Alalawneh2024	-	4 (36)	-	4 (36)
Graff2007	4 (32)	-	-	4 (32)
Jiang2010	11 (136)	-	1 (13)	12 (149)
Kreutz2017	-	2 (24)	-	2 (24)
Kubitza2005	-	20 (702)	-	20 (702)
Kubitza2005a	20 (366)	-	1 (19)	21 (385)
Kubitza2006	4 (75)	-	-	4 (75)
Kubitza2013	2 (23)	4 (47)	-	6 (70)
Liu2024	12 (48)	-	-	12 (48)
Zhao2009	11 (143)	9 (234)	1 (13)	21 (390)
Total	64 (823)	39 (1043)	3 (45)	106 (1911)

2.3 Goodness of fit

Table 4: Goodness of fit of the fit PD_20261018_082509__94e87 per kind of data, the parameter set PD_20261018_082509__94e87_0. The cost is defined on the training data only, 2-fold is the percentage of the data points with a positive value and prediction whose prediction is within twofold of the data.

Kind	n	cost	MSE	RMSE	NRMSE	RMSE_w	R^2	2-fold [%]	AIC	BIC
training	823	68.7	0.0304	0.174	0.325	0.408	0.895	92.7	−2864.6	−2841.1
validation	1043	-	1.48	1.22	0.274	0.26	0.947	94.1	420.2	444.9
outlier	45	-	0.00122	0.035	1.22	1.61	−0.0557	-	−291.8	−282.8

2.4 Figures




Figure 1: Cost of the optimizers over their steps, fit PD_20261018_082509__94e87. The cost of every optimization run over its steps. Runs which end at the same cost found the same optimum; a run which stops much higher is stuck in a local minimum.




Figure 2: Cost of the optimization runs, ordered, fit PD_20261018_082509__94e87. The final cost of the runs, ordered. A flat plateau at the left is the global optimum found repeatedly, which is the evidence that the multistart converged; steps to the right are local minima.




Figure 3: Prediction against the measured data points, per kind of fit mapping, with the agreement of the training data, the lines of twofold and the metrics of the panel, fit PD_20261018_082509__94e87. Within twofold of the data are 732 of the 790 training data points (92.7%) and 953 of the 1013 validation data points (94.1%); only the data points with a positive value and prediction are counted. Prediction against measurement, one point per data point, with a panel per kind of fit mapping. The points scatter around the diagonal when the model describes the data; a systematic deviation from it is a systematic error of the model. The band is the agreement of the training data, i.e. the same band the Bland-Altman plot draws, here as lines parallel to the diagonal. The dark red dotted lines are twofold, prediction = 2 x measurement and prediction = measurement / 2; the caption states the points between them of every kind. A point which is zero or negative is not drawn on the logarithmic axes and not counted, the title of a panel then states the points drawn of all. The validation panel shows how the fit describes data it was not fitted on, the outlier panel the data the fit dropped. The box of a panel carries its R^2, its normalized RMSE, the RMSE of the residuals of the cost, its fold errors GMFE and AFE and the percentages of its points within twofold and within two standard deviations of the data.




Figure 4: Agreement of prediction and measurement as a ratio, per kind of fit mapping, with the agreement of the training data and the agreement of the panel, fit PD_20261018_082509__94e87. The ratio of prediction and measurement over the geometric mean of the two, i.e. the goodness of fit with the diagonal turned into the horizontal. The band is the same, the bias and the limits of agreement `bias ± 1.96 SD` of the training data as fold factors. A bias away from 1 is a systematic over- or underprediction, wide limits are a large scatter, and a trend over the mean is a model which describes the large or the small values better. The box of a panel carries the bias and the SD of its own points as fold factors and the share of them inside the limits of agreement of the training data.




Figure 5: Profile likelihood of every parameter: the cost with the parameter fixed and the other parameters optimized, the threshold of the confidence level and the confidence interval, fit PD_20261018_082509__94e87. One profile per parameter. A profile which crosses the dashed threshold on both sides gives a finite confidence interval; one which stays below it up to a bound is practically non-identifiable; a flat profile is structurally non-identifiable, the parameter is compensated by the others.

2.5 Provenance

Table 5: Provenance of the fit PD_20261018_082509__94e87: the versions it ran with and the revision of its models and data; the SHA-256 checksums of the models and data are listed in Table 1.

Item	Value
Fit	`PD_20261018_082509__94e87`
Created	2026-10-18 06:25:12 UTC
Seed	1234
Python	3.14.4
sbmlsim	0.10.0
libRoadRunner	2.10.0
SciPy	1.18.1
NumPy	2.5.3
Repository	`/home/mkoenig/git/pkdb_models-rivaroxaban`
Commit	`b15d1ca073221f8e25d56c4caef1fef784205bf2`
Uncommitted changes	no

Bibliography

[1] M. König, “sbmlsim: SBML simulation made easy.” Zenodo, 2026. doi: 10.5281/zenodo.3597770. Version 0.10.0.

[2] R. Storn and K. Price, “Differential evolution - a simple and efficient heuristic for global optimization over continuous spaces,” *Journal of Global Optimization*, vol. 11, pp. 341–359, 1997, doi: 10.1023/A:1008202821328.

[3] P. Virtanen *et al.*, “SciPy 1.0: fundamental algorithms for scientific computing in Python,” *Nature Methods*, vol. 17, pp. 261–272, 2020, doi: 10.1038/s41592-019-0686-2. Version 1.18.1.

[4] M. D. McKay, R. J. Beckman, and W. J. Conover, “A comparison of three methods for selecting values of input variables in the analysis of output from a computer code,” *Technometrics*, vol. 21, no. 2, pp. 239–245, 1979, doi: 10.2307/1268522.

[5] C. Welsh, J. Xu, L. Smith, M. König, K. Choi, and H. M. Sauro, "libRoadRunner 2.0: a high performance SBML simulation and analysis library," *Bioinformatics*, vol. 39, no. 1, p. btac770, 2023, doi: 10.1093/bioinformatics/btac770. Version 2.10.0.

[6] A. C. Hindmarsh *et al.*, "SUNDIALS: suite of nonlinear and differential/algebraic equation solvers," *ACM Transactions on Mathematical Software*, vol. 31, no. 3, pp. 363–396, 2005, doi: 10.1145/1089014.1089020.

[7] H. Akaike, "A new look at the statistical model identification," *IEEE Transactions on Automatic Control*, vol. 19, no. 6, pp. 716–723, 1974, doi: 10.1109/TAC.1974.1100705.

[8] G. Schwarz, "Estimating the dimension of a model," *The Annals of Statistics*, vol. 6, no. 2, pp. 461–464, 1978, doi: 10.1214/aos/1176344136.

[9] J. M. Bland and D. G. Altman, "Statistical methods for assessing agreement between two methods of clinical measurement," *The Lancet*, vol. 327, no. 8476, pp. 307–310, 1986, doi: 10.1016/S0140-6736(86)90837-8.

[10] A. Raue *et al.*, "Structural and practical identifiability analysis of partially observed dynamical models by exploiting the profile likelihood," *Bioinformatics*, vol. 25, no. 15, pp. 1923–1929, 2009, doi: 10.1093/bioinformatics/btp358.

[11] C. Kreutz, A. Raue, D. Kaschek, and J. Timmer, "Profile likelihood in systems biology," *The FEBS Journal*, vol. 280, no. 11, pp. 2564–2571, 2013, doi: 10.1111/febs.12276.

[12] F.-G. Wieland, A. L. Hauber, M. Rosenblatt, C. Tönsing, and J. Timmer, "On structural and practical identifiability," *Current Opinion in Systems Biology*, vol. 25, pp. 60–69, 2021, doi: 10.1016/j.coisb.2021.03.005.

[13] C. R. Harris *et al.*, "Array programming with NumPy," *Nature*, vol. 585, pp. 357–362, 2020, doi: 10.1038/s41586-020-2649-2. Version 2.5.3.