

Celiac Disease Triggers

Using Mendelian Randomization to infer causality

Matthijs Knigge¹, Serena Sanna²

1. Hanze University of Groningen, Institute for Life Science & Technology, Bioinformatics, University of Medical Center Groningen, Department of Genetics, Groningen, The Netherlands.

2. University of Groningen, University of Medical Center Groningen, Department of Genetics, Groningen, The Netherlands.

INTRODUCTION


Celiac disorder affects up to two percent of the European population, and it is hereditary. The inheritance model of celiac disease is oligogenic.^[1] In this model, a genetic variant in the Major Histocompatibility Complex (MHC) region, the HLA-DQ2 haplotype, accounts for 40% of the inheritable risk, but it is not sufficient enough to develop the disease. 30% of the Europeans are in fact carriers of the variant, but only 3% express celiac disease. Other 42 non-HLA variants have been identified, but still remains little understanding on what triggers the disease. We aimed to systematically assess multiple clinical parameters and molecular mechanisms to identify causal and protective factors, that can explain what is triggering disease manifestation in HLA-DQ2 carriers.^[2]

MATERIALS AND METHODS

We investigated over 500.000 clinical parameters and molecular phenotypes, for which summary statistics from genome-wide association studies (GWAS) were available. We used the concept of Mendelian Randomization^[3] (MR) approach, and applied two-samples MR methods: Inverse Variance Weighted^[4] (IVW) and MR-Egger method^[5] to infer causality links by combining summary statistics from GWAS. For our analyses, we discarded variants located in the MHC region to avoid over-estimation of causality for factors that also influenced by the same or other HLA haplotypes.

RESULTS

We identified 1133 significant (FDR < 0.05) clinical parameters that cause or protect for celiac disease, and others that show feedback mechanisms. For example autoimmune diseases such as Type 1 Diabetes (T1D), have been largely described to co-occur with celiac disease.^[6,7] We also confirmed a causative effect of eosinophils (and protective of lymphocytes) in increasing the risk of celiac disease, which was recently described using the same approach^[8]. Here we also observed that there is feedback mechanisms, with the disease also affecting, in turn, the number of eosinophils and lymphocytes. Finally, in addition to parameters that are supported by previous epidemiological studies, we found an interesting causative role for infection diseases and allergies.



OTHER (NON-AUTOIMMUNE) DISEASE OF DIGESTION SYSTEM

HYPERTENSION

PBMC

NATURAL KILLER CELLS, CD4+

WHITE BLOODCELLS
EOSINOPHILS, LYMPHOCYTES

INFECTIONS,
ALLERGIES

OTHER AUTO-
IMMUNE DISEASES

ABNORMAL
ERYTHROPOIESIS
ANEMIA

BONE-RELATED
DISORDERS

SKIN
DISORDERS

NEUROLOGICAL
DISORDERS

REDUCED BODY
FAT/MASS

THYROID
ABNORMALITIES

THROMBOCYTOSIS

CELIAC DISEASE

CONCLUSION & DISCUSSION

We have carried out a systematic application of Mendelian Randomization to hundred of thousands of phenotypes, making this study the largest of its kind. We have identified causal and protective factors for celiac disease, recapitulating knowledge from epidemiological studies and also highlighting new key players. We expect that application to other diseases can bring new insights in the understanding of pathophysiology as we well as of biological mechanism, especially as more genome-wide association studies on -omics measurements will be available in the future.

REFERENCES. ¹Robert Di Niro, et al. Nature Medicine (2012); ²Gujral, N. et al. World journal of Gastroenterology (2012); ³David M. Evans, et al. Human Genetics (2014); ⁴Debbie A. Lawlor. Journal of Epidemiology (2016); ⁵Stephan Burgess, et al. Eur J Epidemiol (2017); ⁶William Hagopian, et al. Pediatrics (2017); ⁷Festen EAM, et al. PloS Genetics (2011); ⁸William J. Astle, et al. Cell (2016).