

1 Jacobian-vector products

Consider the parameterized ODE initial value problem

$$\dot{y} = f(t, y, a), \quad y(0, a) = y_0(a), \quad (1)$$

by which we mean

$$\partial_0 y(t, a)[1] = f(t, y(t, a), a), \quad y(0, a) = y_0(a), \quad (2)$$

for all t and a in some domains. We want to understand how the solution to the ODE changes (e.g. at particular values of t) for small perturbations of a . That is, we want to be able to compute the Jacobian-vector product

$$(a, v) \mapsto \partial_1 y(t, a)[v] \quad (3)$$

at any particular values of t and a , where v can be interpreted as a small perturbation to the value of a .

Since the ODE holds true for all values of a (or at least those close to a particular a_0 in which we are interested), we can view both sides as functions of a , and assuming differentiability we can differentiate both sides with respect to a to find a new equation that must be satisfied:

$$\partial_1((t, a) \mapsto \partial_0 y(t, a)[1]) = \partial_2 f(t, y(t, a), a) + \partial_1 f(t, y(t, a), a) \circ \partial_1 y(t, a). \quad (4)$$

Applying both sides to a particular perturbation vector v and using the fact that partial derivatives commute, we have

$$\partial_0((t, a) \mapsto \partial_1 y(t, a)[v])[1] = \partial_2 f(t, y(t, a), a)[v] + \partial_1 f(t, y(t, a), a)[\partial_1 y(t, a)[v]].$$

We can identify $z(t, a) \triangleq \partial_1 y(t, a)[v]$ as a new state vector to write a joint ODE system

$$\begin{bmatrix} \dot{y} \\ \dot{z} \end{bmatrix} = \begin{bmatrix} f(t, y, a) \\ g(t, y, z, a) \end{bmatrix}, \quad \begin{bmatrix} y(0, a) \\ z(0, a) \end{bmatrix} = \begin{bmatrix} y_0(a) \\ \partial y_0(a)[v] \end{bmatrix}, \quad (5)$$

$$g(t, y, z, a) = \partial_1 f(t, y, a)[z] + \partial_2 f(t, y, a)[v]. \quad (6)$$

Notice that the dynamics on the z component are linear/affine in z (and v !).

2 Vector-Jacobian products

Consider the parameterized linear/affine ODE IVP

$$\dot{z}(t) = A(t)z(t) + B(t)v, \quad z(0) = Cv. \quad (7)$$

The implicit mapping $\mathcal{T}_1 : v \mapsto z$ is linear, and so for any linear functional on solution functions $\mathcal{T}_2 : z \mapsto \mathbb{R}$ there is linear function on perturbations v defined by $\mathcal{T}_2 \circ \mathcal{T}_1 : v \mapsto \mathbb{R}$. Given such a linear functional \mathcal{T}_2 , we wish to find an explicit representer vector for $\mathcal{T}_2 \circ \mathcal{T}_1$.

Concretely, consider linear functionals of the form

$$\mathcal{T}_2[z] = w^T z(T) + \int_0^T w(t)^T z(t) dt. \quad (8)$$

We wish to find a function $\lambda(t)$ such that

$$w^T z(T) + \int_0^T w(t)^T z(t) dt = \lambda(0)^T C v + \int_0^T \lambda(t)^T B(t)[v] dt, \quad (9)$$

for all v , when z is a solution to (7).

Consider first the special case when $B \equiv 0$, so that we have the ODE

$$\dot{z}(t) = A(t)z(t), \quad z(0) = C v. \quad (10)$$

Moreover consider the special case of a weighted evaluation functional

$$\mathcal{T}_2[z] = w^T z(T). \quad (11)$$

We wish to find a representer vector λ such that

$$\lambda^T z(0) = w^T z(T). \quad (12)$$

Since the particular time $t = 0$ is arbitrary, a more general problem would be to find a function $\lambda(t)$ such that

$$\lambda(t)^T z(t) = w^T z(T). \quad (13)$$

That is, we can fix $\lambda(T) = w$ and ensure that the value of $\lambda(t)^T z(t)$ does not change with time:

$$0 = \partial(t \mapsto \lambda(t)^T z(t)) = \dot{\lambda}(t)^T z(t) + \lambda(t)^T \dot{z}(t) \quad (14)$$

$$= \dot{\lambda}(t)^T z(t) + \lambda(t)^T A(t)z(t), \quad (15)$$

where on the last line we have used the ODE (10). To satisfy this equation for all t and nonzero solutions $z(t)$, we can choose

$$\dot{\lambda}(t) = -A(t)^T \lambda(t), \quad \lambda(T) = w. \quad (16)$$

This gives us a means of computing a representer for the linear functional \mathcal{T}_2 by solving another ODE IVP, integrating backward in time from $t = T$ to $t = 0$ to compute $\lambda(0)$. (Notice that by linearity we can handle a linear functional that is a linear combination of such weighted evaluation functionals, say at times $0 < T_1 < T_2$, by pulling back the functional at time $t = T_2$ to a representer at time $t = T_1$ and summing before pulling back the sum to $t = 0$.)