

Sample BEAMER Presentation

Matt Kinsinger¹



ARIZONA STATE UNIVERSITY
SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES

May 9, 2017

¹mkinsing@asu.edu

Introduction

- Heat as a function of 1-dimensional space and time: $u(x, t)$
- Notation for u evaluated at discretized points in space and time

$$u(x_i, t_n) = u_i^n$$



- The heat equation, a second order partial differential equation



$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}$$

- Euler's method approximation

$$u_i^{n+1} \approx u_i^n + \frac{k\Delta t}{\Delta x^2} [u_{i+1}^n - 2u_i^n + u_{i-1}^n]$$

- We need the initial temperature profile of the rod and the temperatures for all points in time of the endpoints of the rod
 - We will give these values in our MatLab code
- Example 1
 - Length of rod, L : 1 meter
 - Number of points: 50
 - Total time, t_f : 5 seconds
 - Number of time points: 2000
 - Diffusivity constant: k
 - $r = k \frac{\Delta t}{\Delta x^2} \quad r < 0.50$ required
 - $u(1, :) = \sin\left(\frac{2\pi}{t_f} t\right)$ Left endpoint boundary conditions
 - $u(N, :) = \cos\left(\frac{2\pi}{t_f} t\right)$ Right endpoint boundary conditions
 - $u(:, 1) = \sin\left(\frac{2\pi}{L} x\right)$ Initial conditions





- A better approximation method: Second order Runge-Katta
 - Approximate the slopes at the endpoints of each time interval
 - \tilde{K}_{i+1} is found using an approximation for u_i^{n+1}
 - Use the average of these slopes to linearly approximate the solution over the interval

$$u_i^{n+1} = u_i^n + \Delta t \left[\frac{K_i + \tilde{K}_{i+1}}{2} \right]$$

- $K_i \approx \text{slope at } u_i^n$
- $\tilde{K}_{i+1} \approx \text{slope at } u_i^{n+1}$

- Pieces to the approximation



- $K_i = \frac{u_i^{n+1} - u_i^n}{\Delta t} = \frac{k}{\Delta x^2} [u_{i-1}^n - 2u_i^n + u_{i+1}^n]$



- $\tilde{u}_i^{n+1} = u_i^n + \frac{k\Delta t}{\Delta x^2} [u_{i-1}^n - 2u_i^n + u_{i+1}^n]$

- $\tilde{K}_{i+1} = \frac{u_i^{n+2} - u_i^{n+1}}{\Delta t} = \frac{k}{\Delta x^2} [\tilde{u}_{i-1}^{n+1} - 2\tilde{u}_i^{n+1} + \tilde{u}_{i+1}^{n+1}]$

• Example 2

- All parameters the same as in example 1
- Still had the same limiting diffusivity constant, $k \approx 0.083$





Implicit Time Stepping

- Benefits
 - Increased stability of approximation
 - Able to handle larger diffusivity constants
- Drawbacks
 - Need to solve a linear system $Ax = b$

- Setting up the approximation

- Use the Taylor's series expansion, but the RHS of our final approximation expression is in terms of $u_x^{t_{n+1}}$ rather than $u_x^{t_n}$.

$$\frac{u_i^{n+1} - u_i^n}{\Delta t} = \frac{k}{\Delta x^2} \left[u_{i-1}^{n+1} - 2u_i^{n+1} + u_{i+1}^{n+1} \right]$$

$$u_i^n = u_i^{n+1} - \frac{k\Delta t}{\Delta x^2} \left[u_{i-1}^{n+1} - 2u_i^{n+1} + u_{i+1}^{n+1} \right]$$

$$u_i^n = u_i^{n+1} \left[1 + 2r \right] - r \left[u_{i-1}^{n+1} + u_{i+1}^{n+1} \right] \quad , r = \frac{k\Delta t}{\Delta x^2}$$

- We are using the value at the *next* time step

- Fix n (time) and let i (space) float over all of the interior points of our rod
 - Boundary conditions $u(x_1, t)$ and $u(x_N, t)$ are known for all t
 - Interior points

$$u_2^n = u_2^{n+1} [1 + 2r] - r [u_1^{n+1} + u_3^{n+1}]$$

$$u_3^n = u_3^{n+1} [1 + 2r] - r [u_2^{n+1} + u_4^{n+1}]$$

$$u_4^n = u_4^{n+1} [1 + 2r] - r [u_3^{n+1} + u_5^{n+1}]$$

⋮

⋮




$$u_{N-2}^n = u_{N-2}^{n+1} [1 + 2r] - r [u_{N-3}^{n+1} + u_{N-1}^{n+1}]$$



$$u_{N-1}^n = u_{N-1}^{n+1} [1 + 2r] - r [u_{N-2}^{n+1} + u_N^{n+1}]$$

We move our known terms u_1^{n+1} and u_N^{n+1} to the RHS.



Recognizing the pattern we build the linear system:



$$\begin{pmatrix} 1+2r & -r & 0 & 0 & \cdot & \cdot & 0 \\ -r & 1+2r & -r & 0 & 0 & \cdot & 0 \\ 0 & -r & 1+2r & -r & 0 & \cdot & 0 \\ \cdot & & \cdot & \cdot & \cdot & & \cdot \\ \cdot & & \cdot & \cdot & \cdot & & \cdot \\ \cdot & & \cdot & \cdot & \cdot & & \cdot \\ 0 & 0 & \cdot & \cdot & -r & 1+2r & -r \\ 0 & 0 & \cdot & \cdot & 0 & -r & 1+2r \end{pmatrix} \begin{pmatrix} u_2^{n+1} \\ u_3^{n+1} \\ u_4^{n+1} \\ \cdot \\ \cdot \\ \cdot \\ u_{N-2}^{n+1} \\ u_{N-1}^{n+1} \end{pmatrix} = \begin{pmatrix} u_2^n + ru_1^{n+1} \\ u_3^n \\ u_4^n \\ \cdot \\ \cdot \\ \cdot \\ u_{N-2}^n \\ u_{N-1}^n + ru_N^{n+1} \end{pmatrix}$$

*Euler's**Improved Euler's**Implicit Time Stepping*



We can greatly increase the diffusivity constant and the approximation remains stable





Exploring Error

- Simplify IC and Boundary conditions so the we can solve for an explicit solution

- IC

$$u(x, 0) = \sin(\pi x) + 0.2 \sin(10\pi x)$$

- Boundary conditions


$$u(0, t) = u(1, t) = 0$$

- Guess a solution to $u_t = ku_{xx}$


$$u(t, x) = e^{-\pi^2 kt} \sin(\pi x) + 0.2e^{-(10\pi)^2 kt} \sin(10\pi x)$$

$$\begin{aligned}u_t &= \frac{\partial}{\partial t} [u(x, t)] \\&= \frac{\partial}{\partial t} [e^{-\pi^2 kt} \sin(\pi x) + 0.2e^{-(10\pi)^2 kt} \sin(10\pi x)] \\&= -\pi^2 k e^{-\pi^2 kt} \sin(\pi x) + 0.2 [-(10\pi)^2 k] e^{-(10\pi)^2 kt} \sin(10\pi x) \\&= k \left[-\pi^2 e^{-\pi^2 kt} \sin(\pi x) + 0.2 [-(10\pi)^2 k] e^{-(10\pi)^2 kt} \sin(10\pi x) \right] \\&= k \frac{\partial^2 u}{\partial x^2} [e^{-\pi^2 kt} \sin(\pi x) + 0.2e^{-(10\pi)^2 kt} \sin(10\pi x)] \\&= k \frac{\partial^2 u}{\partial x^2} [u(x, t)] \\&= ku_{xx}\end{aligned}$$


Plots with $M = 2000, N = 200, k = 0.10 \rightarrow r = 0.2972$



Exact solution



Euler's method




Error


At which point in time should we sample the error across the rod?

- t large and $t \approx 0$ the error is nearly zero
- Error appears non-trivial at $t \approx 0.10$ seconds


Improved Euler's



Exact solution




Improved Euler's method




Error


Implicit time stepping




Exact solution






Implicit time step method






Error



Euler's method at $t = .1$ seconds



Improved Euler's method at $t = .1$ seconds



Remarks

Acknowledgments