

Exploring Approximations for the Diffusion Equation

Matt Kinsinger¹



ARIZONA STATE UNIVERSITY
SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES

May 12, 2017

¹mkinsing@asu.edu

Introduction

- Heat as a function of 1-dimensional space and time: $u(x, t)$
- Notation for u evaluated at discretized points in space and time

$$u(x_i, t_n) = u_i^n$$

- The heat equation, a second order partial differential equation

$$\frac{\partial u}{\partial t} = k \frac{\partial^2 u}{\partial x^2}$$

- Three approximation strategies
 - Euler's method
 - Improved Euler's method
 - Implicit Time Stepping

Euler's Method

$$u_i^{n+1} \approx u_i^n + \frac{k\Delta t}{\Delta x^2} [u_{i+1}^n - 2u_i^n + u_{i-1}^n]$$

- This can be derived using Taylor series expansions in each variable

Improved Euler's method

- Approximate the slopes at the endpoints of each time interval
 - \tilde{K}_{i+1} is found using an approximation for u_i^{n+1}
- Use the average of these slopes to linearly approximate the solution over the interval

$$u_i^{n+1} = u_i^n + \Delta t \left[\frac{K_i + \tilde{K}_{i+1}}{2} \right]$$

- $K_i \approx$ slope at u_i^n
- $\tilde{K}_{i+1} \approx$ slope at u_i^{n+1}

- Pieces to the approximation

- $K_i = \frac{u_i^{n+1} - u_i^n}{\Delta t} = \frac{k}{\Delta x^2} [u_{i-1}^n - 2u_i^n + u_{i+1}^n]$

- $\tilde{u}_i^{n+1} = u_i^n + \frac{k\Delta t}{\Delta x^2} [u_{i-1}^n - 2u_i^n + u_{i+1}^n]$

- $\tilde{K}_{i+1} = \frac{u_i^{n+2} - u_i^{n+1}}{\Delta t} = \frac{k}{\Delta x^2} [\tilde{u}_{i-1}^{n+1} - 2\tilde{u}_i^{n+1} + \tilde{u}_{i+1}^{n+1}]$

Implicit Time Stepping

- Benefits
 - Increased stability of approximation
 - Able to handle larger diffusivity constants
- Drawbacks
 - Need to solve a linear system $Ax = b$

- Setting up the approximation

- Use the Taylor's series expansion, but the RHS of our final approximation expression is in terms of u_x^{n+1} rather than u_x^n .

$$\frac{u_i^{n+1} - u_i^n}{\Delta t} = \frac{k}{\Delta x^2} \left[u_{i-1}^{n+1} - 2u_i^{n+1} + u_{i+1}^{n+1} \right]$$

$$u_i^n = u_i^{n+1} - \frac{k\Delta t}{\Delta x^2} \left[u_{i-1}^{n+1} - 2u_i^{n+1} + u_{i+1}^{n+1} \right]$$

$$u_i^n = u_i^{n+1} \left[1 + 2r \right] - r \left[u_{i-1}^{n+1} + u_{i+1}^{n+1} \right] \quad , r = \frac{k\Delta t}{\Delta x^2}$$

- We are using the value at the *next* time step

- Fix n (time) and let i (space) float over all of the interior points of our rod
 - Boundary conditions $u(x_1, t)$ and $u(x_N, t)$ are known for all t
 - Interior points

$$u_2^n = u_2^{n+1} [1 + 2r] - r [u_1^{n+1} + u_3^{n+1}]$$

$$u_3^n = u_3^{n+1} [1 + 2r] - r [u_2^{n+1} + u_4^{n+1}]$$

$$u_4^n = u_4^{n+1} [1 + 2r] - r [u_3^{n+1} + u_5^{n+1}]$$

⋮

⋮

$$u_{N-2}^n = u_{N-2}^{n+1} [1 + 2r] - r [u_{N-3}^{n+1} + u_{N-1}^{n+1}]$$

$$u_{N-1}^n = u_{N-1}^{n+1} [1 + 2r] - r [u_{N-2}^{n+1} + u_N^{n+1}]$$


We move our known terms u_1^{n+1} and u_N^{n+1} to the RHS.

Recognizing the pattern we build the linear system:


$$\begin{pmatrix} 1+2r & -r & 0 & 0 & \cdot & \cdot & 0 \\ -r & 1+2r & -r & 0 & 0 & \cdot & 0 \\ 0 & -r & 1+2r & -r & 0 & \cdot & 0 \\ \cdot & & \cdot & \cdot & \cdot & & \cdot \\ \cdot & & \cdot & \cdot & \cdot & & \cdot \\ \cdot & & \cdot & \cdot & \cdot & & \cdot \\ 0 & 0 & \cdot & \cdot & -r & 1+2r & -r \\ 0 & 0 & \cdot & \cdot & 0 & -r & 1+2r \end{pmatrix} \begin{pmatrix} u_2^{n+1} \\ u_3^{n+1} \\ u_4^{n+1} \\ \cdot \\ \cdot \\ \cdot \\ u_{N-2}^{n+1} \\ u_{N-1}^{n+1} \end{pmatrix} = \begin{pmatrix} u_2^n + ru_1^{n+1} \\ u_3^n \\ u_4^n \\ \cdot \\ \cdot \\ \cdot \\ u_{N-2}^n \\ u_{N-1}^n + ru_N^{n+1} \end{pmatrix}$$

Comparing plots


- We need the initial temperature profile of the rod and the temperatures for all points in time of the endpoints of the rod
 - We will give these values in our MatLab code
- Example 1
 - Length of rod, L : 1 meter
 - Number of points: 50
 - Total time, t_f : 5 seconds
 - Number of time points: 2000
 - Diffusivity constant: k
 - $r = k \frac{\Delta t}{\Delta x^2}$ $r < 0.50$ required
 - $u(1, :) = \sin\left(\frac{2\pi}{t_f} t\right)$ Left endpoint boundary conditions
 - $u(N, :) = \cos\left(\frac{2\pi}{t_f} t\right)$ Right endpoint boundary conditions
 - $u(:, 1) = \sin\left(\frac{2\pi}{L} x\right)$ Initial conditions



Euler's



Improved Euler's



Implicit Time Stepping

Exploring Error

- Simplify IC and Boundary conditions so the we can solve for an explicit solution

- IC

$$u(x, 0) = \sin(\pi x) + 0.2 \sin(10\pi x)$$

- Boundary conditions

$$u(0, t) = u(1, t) = 0$$

- Guess a solution to $u_t = ku_{xx}$




$$u(t, x) = e^{-\pi^2 kt} \sin(\pi x) + 0.2e^{-(10\pi)^2 kt} \sin(10\pi x)$$

$$\begin{aligned}
 u_t &= \frac{\partial}{\partial t} [u(x, t)] \\
 &= \frac{\partial}{\partial t} [e^{-\pi^2 kt} \sin(\pi x) + 0.2e^{-(10\pi)^2 kt} \sin(10\pi x)] \\
 &= -\pi^2 k e^{-\pi^2 kt} \sin(\pi x) + 0.2 [-(10\pi)^2 k] e^{-(10\pi)^2 kt} \sin(10\pi x) \\
 &= k \left[-\pi^2 e^{-\pi^2 kt} \sin(\pi x) + 0.2 [-(10\pi)^2 k] e^{-(10\pi)^2 kt} \sin(10\pi x) \right] \\
 &= k \frac{\partial^2 u}{\partial x^2} [e^{-\pi^2 kt} \sin(\pi x) + 0.2e^{-(10\pi)^2 kt} \sin(10\pi x)] \\
 &= k \frac{\partial^2 u}{\partial x^2} [u(x, t)] \\
 &= ku_{xx}
 \end{aligned}$$



It is a solution! We can use this to measure the error in our approximations!

Plots with $M = 2000, N = 200, k = 0.01 \rightarrow r = 0.2972$



Exact solution



Euler's









Improved Euler's



Implicit







At which point in time should we sample the error across the rod?

- t large and $t \approx 0$ the error is nearly zero
- Error appears non-trivial at $t \approx 0.10$ seconds
- Experiment:
 - A given number of sample points in space
 - Range through different numbers of sample points in time
 - $r = \frac{k\Delta t}{\Delta x^2} \downarrow$ as $\Delta t \downarrow$



Implicit time stepping: The code ran forever

Log plot to get a closer look at Euler's method approximation



At some point just above $r = 0.10$ we see the error becoming very small!

Intersting...Why?

$$U_t = kU_{xx}$$

Approximation:
$$\frac{U^{n+1} - U^n}{\Delta t} = k \frac{U_{j+1} - 2U_j - U_{j-1}}{\Delta x^2}$$

Recall the Taylor series expansion in one variable

$$u(x_j \pm \Delta x) = \sum_{k=0}^N (\pm \Delta x)^k f^{(k)}(x_j) + \textit{error term}.$$

$$u(t^n \pm \Delta t) = \sum_{k=0}^N (\pm \Delta t)^k f^{(k)}(t^n) + \textit{error term}.$$

x first...

$$\begin{aligned}
 U_{j+1} &= u(x_j + \Delta x) \\
 &= u(x_j) + \Delta x U_x(x_j) + \frac{\Delta x^2}{2!} U_{xx}(x_j) + \frac{\Delta x^3}{3!} U_{xxx}(x_j) + \frac{\Delta x^4}{4!} U_{4x}(x_j) + \dots
 \end{aligned}$$

$$\begin{aligned}
 U_{j-1} &= u(x_j - \Delta x) \\
 &= u(x_j) - \Delta x U_x(x_j) + \frac{\Delta x^2}{2!} U_{xx}(x_j) - \frac{\Delta x^3}{3!} U_{xxx}(x_j) + \frac{\Delta x^4}{4!} U_{4x}(x_j) + \dots
 \end{aligned}$$

$$\frac{U_{j+1} - 2U_j + U_{j-1}}{\Delta x^2} = U_{xx}(x_j) + 2\frac{\Delta x^2}{4!} U_{4x}(x_j) + \dots \Delta x^{large} \quad (1)$$

Now t...

$$U^{n+1} = u(t^n + \Delta t) = u(t^n) + \Delta t U_t(t^n) + \frac{\Delta t^2}{2!} U_{tt}(t^n) + \frac{\Delta t^3}{3!} U_{ttt}(t^n) + \dots \Delta t^{large}$$

$$\frac{U^{n+1} - U^n}{\Delta t} = U_t(t^n) + \frac{\Delta t}{2!} U_{tt}(t^n) + \frac{\Delta t^2}{3!} U_{ttt}(t^n) + \dots \Delta t^{large} \quad (2)$$

$$(2) = k(1)$$

$$\frac{U^{n+1} - U^n}{\Delta t} = k \left[\frac{U_{j+1} - 2U_j + U_{j-1}}{\Delta x^2} \right]$$

$$U_t(t^n) + \frac{\Delta t}{2!} U_{tt}(t^n) + \frac{\Delta t^2}{3!} U_{ttt}(t^n) + \dots = k \left[U_{xx}(x_j) + 2 \frac{\Delta x^2}{4!} U_{4x}(x_j) + \dots \right]$$

$$U_t = kU_{xx} - \frac{\Delta t}{2} U_{tt} - \frac{\Delta t^2}{6} U_{3t} + k \frac{\Delta x^2}{12} U_{4x} + \dots \quad (3)$$

$$U_t = kU_{xx} - \text{error}$$

$$\frac{\partial}{\partial t} \left[U_t = kU_{xx} \right] \longrightarrow U_{tt} = kU_{xxt}$$

$$\frac{\partial^2}{\partial x^2} \left[kU_{xx} = U_t \right] \longrightarrow kU_{4x} = U_{txx}$$

So,

$$U_{tt} = k^2 U_{4x}$$

$$U_t = kU_{xx} - \frac{\Delta t}{2} \left[k^2 U_{4x} \right] - \frac{\Delta t^2}{6} U_{3t} + k \frac{\Delta x^2}{12} U_{4x} + \dots$$

$$U_t = kU_{xx} + U_{4x} \left[k \frac{\Delta x^2}{12} - k^2 \frac{\Delta t}{2} \right] - \frac{\Delta t^2}{6} U_{3t} + \dots$$







$$0 = k \frac{\Delta x^2}{12} - k^2 \frac{\Delta t}{2}$$




$$0 = \frac{\Delta x^2}{6} - k \Delta t$$

$$k \Delta t = \frac{\Delta x^2}{6}$$

$$\frac{k \Delta t}{\Delta x^2} = \frac{1}{6}$$

$$r = \frac{1}{6}.$$



Error with different r values $M = 24400$  $M = 14400$  $M = 8000$

Remarks

Thank you Juan

Thank you Professor Platte

Acknowledgments