

## ORIGINAL ARTICLE

**Correspondence:**

Clare Turnbull, Division of Genetics and Epidemiology, The Institute of Cancer Research, London, SM2 5NG, UK.  
E-mail: clare.turnbull@icr.ac.uk

\*Denotes equal author contribution.

**Keywords:**

genome-wide association studies, germ cell tumour, growth, height, testicular cancer

Received: 14-Mar-2017

Accepted: 4-May-2017

doi: 10.1111/andr.12388

# Mendelian randomisation analysis provides no evidence for a relationship between adult height and testicular cancer risk

<sup>1,\*</sup>M. Levy, <sup>1,\*</sup>D. Hall, <sup>1</sup>A. Sud, <sup>1</sup>P. Law, <sup>1</sup>K. Litchfield, <sup>1</sup>D. Dudakia, <sup>2</sup>T. B. Haugen, <sup>3</sup>R. Karlsson, <sup>4</sup>A. Reid, <sup>4,5</sup>R. A. Huddart, <sup>6</sup>T. Grotmol, <sup>3</sup>F. Wiklund, <sup>1</sup>R. S. Houlston and <sup>1,7,8</sup>C. Turnbull

<sup>1</sup>Division of Genetics & Epidemiology, The Institute of Cancer Research, London, UK, <sup>2</sup>Faculty of Health Sciences, Oslo and Akershus University College of Applied Sciences, Oslo, Norway, <sup>3</sup>Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden, <sup>4</sup>Academic Radiotherapy Unit, Institute of Cancer Research, <sup>5</sup>Academic Uro-oncology Unit, The Royal Marsden NHS Foundation Trust, Sutton, Surrey, UK, <sup>6</sup>Department of Research, Cancer Registry of Norway, Oslo, Norway, <sup>7</sup>William Harvey Research Institute, Queen Mary University, and <sup>8</sup>Department of Clinical Genetics, Guy's and St Thomas' NHS Trust, London, UK

**SUMMARY**

Observational studies have suggested anthropometric traits, particularly increased height are associated with an elevated risk of testicular cancer (testicular germ cell tumour). However, there is an inconsistency between study findings, suggesting the possibility of the influence of confounding factors. To examine the association between anthropometric traits and testicular germ cell tumour using an unbiased approach, we performed a Mendelian randomisation study. We used genotype data from genome wide association studies of testicular germ cell tumour totalling 5518 cases and 19,055 controls. Externally weighted polygenic risk scores were created and used to evaluate associations with testicular germ cell tumour risk per one standard deviation (s.d) increase in genetically-defined adult height, adult BMI, adult waist hip ratio adjusted for BMI (WHRadjBMI), adult hip circumference adjusted for BMI (HIPadjBMI), adult waist circumference adjusted for BMI (WCadjBMI), birth weight (BW) and childhood obesity. Mendelian randomisation analysis did not demonstrate an association between any anthropometric trait and testicular germ cell tumour risk. In particular, despite good power, there was no global evidence for association between height and testicular germ cell tumour. However, three SNPs for adult height individually showed association with testicular germ cell tumour (rs4624820: OR = 1.47, 95% CI: 1.41–1.55,  $p = 2.7 \times 10^{-57}$ ; rs12228415: OR = 1.17, 95% CI: 1.11–1.22,  $p = 3.1 \times 10^{-10}$ ; rs7568069: OR = 1.13, 95% CI: 1.07–1.18,  $p = 1.1 \times 10^{-6}$ ). This Mendelian randomisation analysis, based on the largest testicular germ cell tumour genome wide association dataset to date, does not support a causal etiological association between anthropometric traits and testicular germ cell tumour aetiology. Our findings are more compatible with confounding by shared environmental factors, possibly related to prenatal growth with exposure to these risk factors occurring *in utero*.

**INTRODUCTION**

Testicular cancer (testicular germ cell tumour, TGCT) has doubled in incidence over the last 30 years in Western Countries and has an unusual incidence rate in comparison to many other neoplasms with rates peaking at ~30 years and declining rapidly thereafter (Bray *et al.*, 2006; Le Cornet *et al.*, 2014; Ruf *et al.*, 2014). Tumours with this kind of age-incidence profile are likely to have onset early in life, and previous studies have therefore focussed on perinatal and adolescent stages of development as well as adult anthropometric traits (Akre *et al.*, 2000; Dieckmann & Pichlmeier, 2002; Garner *et al.*, 2003; Rasmussen *et al.*, 2003; Richiardi *et al.*, 2003; Bjørge *et al.*, 2006; McGlynn *et al.*, 2007;

Dieckmann *et al.*, 2009). Although there is disagreement as to the effect of adult body mass index (BMI) on TGCT risk many, but not all, studies have reported increased height as a risk factor for the disease (Akre *et al.*, 2000; Dieckmann & Pichlmeier, 2002; Rasmussen *et al.*, 2003; Richiardi *et al.*, 2003; Bjørge *et al.*, 2006; McGlynn *et al.*, 2007). However, none of these studies have established causal pathways linking traits such as height with TGCT, with some studies suggesting that confounding environmental factors such as childhood nutrition could play a greater role.

An alternative approach to a traditional epidemiology observational study is the Mendelian randomisation (MR) approach.

The strategy of MR uses as instrumental variables (IV) genetic markers known to be associated with a factor in the assessment of its effect on another trait (Palmer *et al.*, 2012). This methodology permits the nature of the relation between the trait and the factor to be assessed and, importantly, establishes whether an association is causal. A further attribute of MR is the avoidance of the influence of factors whose effect may be time sensitive, for example temporal trends in calorific intake. Therefore, a genetically defined IV has the potential to more accurately assess lifetime exposure when compared to measurements of potential risk factors recorded in an observational study.

Recent genome-wide association studies (GWAS) have identified single nucleotide polymorphisms (SNPs) at multiple loci that are robustly associated with height and other anthropometric traits (Wood *et al.*, 2014; Locke *et al.*, 2015). Given the unexplained recent rapid increase in TGCT incidence, the paucity of established risk factors for TGCT and the mixed observational epidemiology on the association of anthropometric traits with TCGT risk, we performed a MR study by analysing genotype data from three GWAS conducted on TGCT (Turnbull *et al.*, 2010; Kristiansen *et al.*, 2015). The traits we considered were adult height, adult BMI, adult waist-hip ratio adjusted for BMI (WHRadjBMI), adult waist circumference adjusted for BMI (WCadjBMI), adult hip circumference adjusted for BMI (HIPadjBMI), childhood obesity and birth weight (BW).

## MATERIALS AND METHODS

### Testicular cancer GWAS data sources

Our MR analysis was based on data from three previously reported GWAS of TGCT totalling 5518 cases and 19,055 controls. Briefly, these GWAS were based on individuals with European ancestry and comprised: UK GWAS1 (985 cases, 4946 controls), UK Oncoarray (3206 cases, 7422 controls), and Scandinavian GWAS1 (1327 cases, 6687 controls). Details of the genotyping, quality control and imputation of untyped SNPs have been previously published (Turnbull *et al.*, 2010; Kristiansen *et al.*, 2015). Briefly, we excluded SNPs with a minor allele frequency of <1%, a call rate <95%, those SNPs violating Hardy-Weinberg equilibrium, and individuals with non-European ancestry as assessed using data from HapMap v2 (International HapMap *et al.*, 2007). IMPUTEV2 software (Howie *et al.*, 2012) was used to recover untyped SNP genotypes using a merged reference panel consisting of UK10K and 1000 Genomes Project data. Poorly imputed SNPs, defined by an INFO score of <0.9, were excluded. Summary statistics from these GWAS were used to calculate the ratio estimates for the anthropometric-related SNPs.

### Instrumental variables

For each anthropometric trait we used data from recent GWAS of individuals of European-descent. Specifically for height, we used data from the Genetic Investigation of Anthropometric Traits (GIANT) consortium, which was based on an analysis of up to 253,288 individuals (Wood *et al.*, 2014). For adult BMI, data was obtained from a consortium meta-analysis comprising 339,224 individuals from 125 studies (Locke *et al.*, 2015). Data for WHRadjBMI, WCadjBMI and HIPadjBMI were obtained from related consortium meta-analyses comprising up to 224,459 individuals (Shungin *et al.*, 2015). Data reported by the Early Growth Genetics (EGG) consortium of a meta-analysis of 5530

cases and 8318 controls was used for childhood obesity (Bradfield *et al.*, 2012). Birth weight data was obtained from a meta-analysis of up to 69,308 individuals from 43 studies (Horikoshi *et al.*, 2013).

To calculate the genetic scores we considered only SNPs that were genome-wide significant ( $p < 5 \times 10^{-8}$ ) for each anthropometric trait and were present in all three GWAS TCGT datasets. To avoid co-linearity between SNPs we imposed a threshold  $R^2$  value of  $\geq 0.8$  for linkage disequilibrium (LD) including only the SNPs with the strongest effect on the trait in the genetic risk score. After imposing these metrics we analysed 379 SNPs for adult height, 69 SNPs for adult BMI, 44 SNPs for WHRadjBMI, 25 SNPs for WCadjBMI and 10 SNPs for HIPadjBMI, seven SNPs for birth weight and seven SNPs for childhood obesity.

We also examined male-specific risk-associated SNPs. However, because of a paucity of data, we relaxed the significance threshold and considered all SNPs at  $p < 1.0 \times 10^{-7}$ . Imposing this criteria provided seven male-specific SNPs for WHRadjBMI, four for HIPadjBMI and nine for WCadjBMI (Randall *et al.*, 2013). Fifteen SNPs displaying significant male dimorphism for adult height were also considered (Randall *et al.*, 2013). For all SNPs used, we recovered the chromosome position, risk alleles, association estimates and standard errors. None of the GWAS that comprised our dataset had reported non-additive effects of the SNPs, hence per allele effects were considered additive. For each SNP, we extracted the respective effect estimates and  $p$ -values from the TGCT GWAS.

### Statistical analysis

A central assumption of MR analysis is that there is no pleiotropism between the genetic instruments influencing TGCT and the trait being tested. Prior to performing MR, we used linkage disequilibrium regression (LDR) to test for global pleiotropy and used MR-Egger and inverse-variance weighted (IVW) regression to examine standard IV assumptions (Bowden *et al.*, 2015). LDR was performed for TGCT against each trait to estimate genetic covariance with TGCT (Bulik-Sullivan *et al.*, 2015).

We performed MR analysis to assess the association between each anthropometric trait and TGCT using summary statistics (Burgess *et al.*, 2015). Using this method, the combined ratio estimate ( $\hat{\beta}$ ) of all SNPs associated with the particular risk factor on TGCT was calculated under a fixed effects model:

$$\hat{\beta} = \frac{\sum_k X_k Y_k \sigma_{Y_k}^{-2}}{\sum_k X_k^2 \sigma_{Y_k}^{-2}}$$

where  $X_k$  represents the anthropometric trait effect size for SNP  $k$ ,  $Y_k$  represents the TGCT risk effect size for SNP  $k$  and  $\sigma_{Y_k}$  denotes the standard error in the value of  $Y_k$ . The standard error of the TGCT risk effect size, denoted by  $\text{se}(\hat{\beta})$ , is given by:

$$\text{se}(\hat{\beta}) = \sqrt{\frac{1}{\sum_k X_k^2 \sigma_{Y_k}^{-2}}}$$

With the statistics generated by following these calculations on the three different cohorts in the TGCT data, we performed a meta-analysis under a fixed-effects model to derive the final odds ratios and confidence intervals.

For each statistical test we considered a global nominal significance level of  $p \leq 0.05$  as satisfactory to derive conclusions. A Bonferroni correction was applied to significance levels to

compensate for multiple testing. All analyses were undertaken using R software (Version 2.14.1).

To gain biological insight into individual SNPs associated with both TGCT and anthropometric traits, we investigated the effects of significantly associated individual SNPs on gene expression using the GTEx resource, examining association with expression of all genes within 1 MB of the associated SNP (Lonsdale *et al.*, 2013).

Finally, we estimated the power of our MR analysis to demonstrate an association between TGCT and anthropometric traits tested over a range of effect sizes (Brion *et al.*, 2013).

## RESULTS

Using LDR we found no evidence for global pleiotropism between TGCT and any of the anthropometric traits after correction for multiple testing (Table S1). Of the seven tested anthropometric traits and four male-specific components of traits, the strongest correlation was found between TGCT and childhood obesity (Genetic Correlation (GC) = 0.14, 95% confidence interval (CI) −0.26, −0.012,  $p = 0.032$ ). MR-Egger and IVW regression tests provided no evidence of infringement of the standard IV assumptions (Figure S1–S11, Tables S2 and S3). All genetic variants used were therefore considered valid IVs and the following MR analysis valid.

MR analysis found no evidence for a genetic relationship between any of the seven anthropometric traits and TGCT risk (Table 1). To explore the effect of male dimorphism we additionally performed MR using IVs generated from male-specific SNPs for four of the anthropometric traits. This restricted analysis found no relationship between any IV and TGCT risk (Table 1).

Although in the LDR analysis we found no evidence for a global pleiotropism between height and TGCT, three of the IV SNPs for height showed a strong association with TGCT risk, even after applying a Bonferroni correction for testing 379 SNPs (rs4624820 (5q31.3): OR 1.47, 95% CI 1.41–1.55,  $p = 2.7 \times 10^{-57}$ ; rs12228415 (12p13.1): OR 1.17, 95% CI 1.11–1.22,  $p = 3.1 \times 10^{-10}$ ; and rs7568069 (2p13.3): OR 1.13, 95% CI 1.07–1.18,  $p = 1.1 \times 10^{-6}$ ). Only rs7568069 showed a significant eQTL effect in testicular tissue (for *DYSF*,  $p = 5.8 \times 10^{-6}$ ). No other trait tested had any SNP associated with TGCT risk.

Analyses of the power afforded by these MR analyses for each anthropometric trait analysed are shown in Figures S12–S15. Since the IV SNPs for adult height explain ~20% of variance of the trait (Wood *et al.*, 2014), our study has 90% power to detect

an OR  $\geq 1.1$ . In contrast, the SNPs used as IVs for adult BMI only account for 3% of the variation (Locke *et al.*, 2015), and hence we had <30% power to detect an OR  $\geq 1.1$ . Similarly, we had <20% power to demonstrate associations for WHRadjBMI and birth weight for effect sizes of OR  $\geq 1.1$ .

## DISCUSSION

Following up upon associations reported in observational epidemiological studies between anthropometric traits and TGCT, our MR analysis using a TGCT GWAS dataset did not find an association between any anthropometric trait tested and risk. Although based on large datasets, aside from adult height, the SNP sets used as IVs only account for a modest proportion of the genetic variance underlying the anthropometric traits analysed. Hence we cannot exclude the possibility that null findings were simply a consequence of limited study power.

The primary motivation for this study was to further explore the epidemiological evidence for a positive relationship between adult height and TGCT, which has been reproduced across a number of studies (Akre *et al.*, 2000; Dieckmann & Pichlmeier, 2002; Rasmussen *et al.*, 2003; Richiardi *et al.*, 2003; Bjørge *et al.*, 2006; McGlynn *et al.*, 2007). Despite 90% power to detect an OR  $> 1.1$ , in our MR analysis we did not find global genetic evidence supporting a role of height in TGCT aetiology. These results suggest that the observed association between height and TGCT, if true, may be due to the influence of confounding due to shared environmental factors (Dieckmann *et al.*, 2008).

Although we found no evidence of global pleiotropism between height and TGCT through LDR, our results, however, do not discount pleiotropic influences of individual SNPs on both adult height and TGCT development. Interestingly, three SNPs associated with increased height also showed a strong association with TGCT risk. The strongest of these SNPs, rs4624820 (5q31.3), is within the previously well-established TGCT-associated locus containing *SPRY4* (Kanetsky *et al.*, 2009; Rapley *et al.*, 2009; Turnbull *et al.*, 2010). Another of these SNPs, rs7568069 (2p13.3), showed a significant eQTL effect in testicular tissue of *DYSF* (Dystrophy-Associated Fer-1-Like 1), which has been implicated in both muscular growth and developmental processes including sperm activation during fertilization and early embryogenesis (Smith & Wakimoto, 2007). While these results are consistent with overlapping pleiotropic pathways for these variants, these three SNPs are among 379 instrumental variables for height and contribute very modestly individually or in combination to each trait (Cole, 2000; Le Cornet *et al.*, 2014).

A number of studies have suggested that BMI plays a role in TGCT risk (Akre *et al.*, 2000; Garner *et al.*, 2003; Dieckmann *et al.*, 2009), although others have found no significant positive or negative association (Dieckmann & Pichlmeier, 2002; Rasmussen *et al.*, 2003; Bjørge *et al.*, 2006). Physical activity was found in one study to have a preventative effect on TGCT risk (Gallagher *et al.*, 1995). There is some evidence for a negative association of BMI from a dataset of 500,000 Norwegian men (Akre *et al.*, 2000); examining our Scandinavian cohort alone, waist circumference adjusted for BMI displayed a nominally significant ( $p = 0.04$ ) preventative association with TGCT risk but there were no other associations of note. Globally we did not find genetic evidence for a role in TGCT of BMI, waist circumference, or any other related traits. However, we acknowledge our power to detect an association of BMI and related

**Table 1** Meta-analysis results of Mendelian Randomisation (MR) for anthropometric phenotypes against TGCT risk. MR was performed for each phenotype against three cohorts

Phenotype	OR (95% CI)	<i>p</i> -value	<i>I</i> <sup>2</sup>	<i>P</i> <sub>het</sub>
Adult Height	1.066 (0.968–1.173)	0.197	25.70	0.260
Adult Height (Male only)	0.922 (0.600–1.416)	0.710	62.89	0.068
Adult BMI	0.848 (0.658–1.093)	0.203	9.79	0.330
WHRadjBMI	0.956 (0.712–1.284)	0.767	0	0.649
WHRadjBMI (Male only)	1.388 (0.755–2.551)	0.292	0	0.579
WCadjBMI	1.163 (0.787–1.719)	0.449	58.83	0.088
WCadjBMI (Male only)	1.245 (0.720–2.150)	0.433	27.46	0.252
HIPadjBMI	1.664 (0.936–2.957)	0.083	53.83	0.115
HIPadjBMI (Male only)	0.996 (0.498–1.994)	0.991	0	0.507
Childhood Obesity	0.863 (0.673–1.107)	0.246	0	0.517
Birth Weight	1.149 (0.783–1.687)	0.478	0	0.599

anthropometric traits with TGCT using MR was limited. Accordingly, our null results therefore do not further clarify whether the reported association between BMI and related anthropometric traits with TGCT are true or not, and if true, whether causal or due to the confounding influence of common environmental factors. Similarly, MR analyses for association of both birth weight and childhood obesity on TGCT risk were subject to low power. To be more informative, further MR analysis of these traits would require significantly larger SNP sets capturing a higher proportion of phenotypic variation.

In summary, in our MR analysis we found no evidence for influence of adult height, BMI or other anthropometric traits in TGCT aetiology. Our MR analysis of height with TGCT was well-powered, and our null result suggests that the previously reported positive association between height and TGCT, if a true association, is unlikely to be driven by a common aetiology but instead is more likely to reflect confounding. Several potential mechanisms underlying the association between height and TGCT have previously been postulated, such as individual variation in the insulin-like growth factor I system (Zavos *et al.*, 2004), age at puberty (Akre *et al.*, 2000), and childhood nutrition (Dieckmann *et al.*, 2008). Molecular and epidemiological data support a model whereby initial neoplastic transformation into the TGCT precursor lesion arises during the foetal period with the pre-invasive TGCT lesion lying dormant through childhood into adolescence and malignant growth finally occurring in early adulthood (Skakkebaek *et al.*, 1987; Jacobsen & Henriques, 1992; Oosterhuis & Looijenga, 2005; Rajpert-De Meyts, 2006; Kristensen *et al.*, 2008). This model of prenatal determination of TGCT would therefore be consistent with proposed exposure to an environmental factor common to adult height and TGCT risk occurring *in utero*. It is therefore plausible that foetal nutrition or another maternal or *in utero* exposure is positively associated with each of adult height and TGCT, thus potentially driving the observed association between these traits. Although much epidemiological study has examined association of maternal and *in utero* exposures with TGCT, to date no compelling evidence that implicates any specific pre-natal risk factors in the aetiology of TGCT has emerged from these studies (McGlynn & Cook, 2009).

In conclusion, although the genomic basis of TGCTs is becoming increasingly well understood (Litchfield *et al.*, 2016), the environmental factors underlying the rapid increase in incidence of TGCT over the last four decades remain unclear. From the MR analyses we have presented, it is likely that if any of the associations of TGCT with anthropometric variables are true, they are a product of confounding influence from early-life environmental exposures which are yet to be elucidated.

## ACKNOWLEDGEMENTS

We thank the patients and all clinicians forming part of the UK Testicular Cancer Collaboration (UKTCC) for their participation in the studies from which the TGCT GWAS datasets were generated. Dominic Hall was supported by Wellcome Trust. K. Litchfield and A. Sud are supported by a PhD fellowship from Cancer Research UK. R.S.H. is supported by Cancer Research UK (C1298/A8362 Bobby Moore Fund for Cancer Research UK). We acknowledge National Health Service funding to the National Institute for Health Research Biomedical Research Centre. This study was supported by the Movember Foundation and the Institute of Cancer Research.

## DISCLOSURE

The authors declare no conflict of interest.

## AUTHOR CONTRIBUTIONS

C.T. and R.S.H. designed the study. C.T. and R.H. lead patient recruitment for UK GWAS1 and UK Oncoarray, for which D.D. coordinated assembly and tracking of samples. T.B.H., R.K., T.G. and F.W. supplied GWAS data. D.H. conducted bioinformatic and statistical analyses with assistance from A.S., P.L. and K.L.. M.L. and D.H. drafted the manuscript with assistance from C.T. and R.S.H. All authors reviewed and contributed to the manuscript.

## REFERENCES

- Akre O, Ekbom A, Sparén P & Tretli S. (2000) Body size and testicular cancer. *J Natl Cancer Inst* 92, 1093–1096.
- Björge T, Tretli S, Lie AK & Engeland A. (2006) The impact of height and body mass index on the risk of testicular cancer in 600,000 Norwegian men. *Cancer Causes Control* 17, 983–987.
- Bowden J, Davey Smith G & Burgess S. (2015) Mendelian randomization with invalid instruments: effect estimation and bias detection through Egger regression. *Int J Epidemiol* 44, 512–525.
- Bradfield JP, Taal HR, Timpson NJ, Scherag A, Lecoeur C, Warrington NM, Hyppönen E, Holst C, Valcarcel B, Thiering E, Salem RM, Schumacher FR, Cousminer DL, Sleiman PM, Zhao J, Berkowitz RI, Vimalaswaran KS, Jarick I, Pennell CE, Evans DM, St Pourcain B, Berry DJ, Mook-Kanamori DO, Hofman A, Rivadeneira F, Uitterlinden AG, van Duijn CM, van der Valk RJ, de Jongste JC, Postma DS, Boomsma DI, Gauderman WJ, Hassanein MT, Lindgren CM, Magi R, Boreham CA, Neville CE, Moreno LA, Elliott P, Pouta A, Hartikainen AL, Li M, Raitakari O, Lehtimäki T, Eriksson JG, Palotie A, Dallongeville J, Das S, Deloukas P, McMahon G, Ring SM, Kemp JP, Buxton JL, Blakemore AI, Bustamante M, Guxens M, Hirschhorn JN, Gillman MW, Kreiner-Möller E, Bisgaard H, Gilliland FD, Heinrich J, Wheeler E, Barroso I, O'Rahilly S, Meirhaeghe A, Sorensen TI, Power C, Palmer LJ, Hinney A, Widen E, Farooqi IS, McCarthy MI, Froguel P, Meyre D, Hebebrand J, Jarvelin MR, Jaddoe VW, Smith GD, Hakonarson H, Grant SF & Early Growth Genetics C (2012) A genome-wide association meta-analysis identifies new childhood obesity loci. *Nat Genet* 44, 526–531.
- Bray F, Ferlay J, Devesa SS, McGlynn KA & Moller H. (2006) Interpreting the international trends in testicular seminoma and nonseminoma incidence. *Nature clinical practice. Urology* 3, 532–543.
- Brion M-JA, Shakhbazov K & Visscher PM. (2013) Calculating statistical power in Mendelian randomization studies. *Int J Epidemiol* 42, 1497–1501.
- Bulik-Sullivan B, Finucane HK, Anttila V, Gusev A, Day FR, Loh P-R; ReproGen Consortium, Psychiatric Genomics Consortium, Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3, Duncan L, Perry JRB, Patterson N, Robinson EB, Daly MJ, Price AL & Neale BM (2015) An atlas of genetic correlations across human diseases and traits. *Nat Genet* 47, 1236–1241.
- Burgess S, Scott RA, Timpson NJ, Davey Smith G, Thompson SG & Consortium E-I. (2015) Using published data in Mendelian randomization: a blueprint for efficient identification of causal risk factors. *Eur J Epidemiol* 30, 543–552.
- Cole TJ. (2000) Secular trends in growth. *Proc Nutr Soc* 59, 317–324.
- Dieckmann K-P & Pichlmeier U. (2002) Is risk of testicular cancer related to body size? *Eur Urol* 42, 564–569.
- Dieckmann KP, Hartmann JT, Classen J, Ludde R, Diederichs M & Pichlmeier U. (2008) Tallness is associated with risk of testicular cancer: evidence for the nutrition hypothesis. *Br J Cancer* 99, 1517–1521.
- Dieckmann KP, Hartmann JT, Classen J, Diederichs M & Pichlmeier U. (2009) Is increased body mass index associated with the incidence of testicular germ cell cancer? *J Cancer Res Clin Oncol* 135, 731–738.

- Gallagher RP, Huchcroft S, Phillips N, Hill GB, Coldman AJ, Coppin C & Lee T. (1995) Physical activity, medical history, and risk of testicular cancer (Alberta and British Columbia, Canada). *Cancer Causes Control* 6, 398–406.
- Garner MJ, Birkett NJ, Johnson KC, Shatenstein B, Ghadirian P & Krewski D; Canadian Cancer Registries Epidemiology Research Group (2003) Dietary risk factors for testicular carcinoma. *Int J Cancer* 106, 934–941.
- Horikoshi M, Yaghootkar H, Mook-Kanamori DO, Sovio U, Taal HR, Hennig BJ, Bradfield JP, St Pourcain B, Evans DM, Charoen P, Kaakinen M, Cousminer DL, Lehtimäki T, Kreiner-Møller E, Warrington NM, Bustamante M, Feenstra B, Berry DJ, Thiering E, Pfab T, Barton SJ, Shields BM, Kerkhof M, van Leeuwen EM, Fulford AJ, Kutalik Z, Zhao JH, den Hoed M, Mahajan A, Lindi V, Goh LK, Hottenga JJ, Wu Y, Raitakari OT, Harder MN, Meirhaeghe A, Ntalla I, Salem RM, Jameson KA, Zhou K, Monies DM, Lagou V, Kirin M, Heikkinen J, Adair LS, Alkuraya FS, Al-Odaib A, Amouyel P, Andersson EA, Bennett AJ, Blakemore AI, Buxton JL, Dallongeville J, Das S, de Geus EJ, Estivill X, Flexeder C, Froguel P, Geller F, Godfrey KM, Gottrand F, Groves CJ, Hansen T, Hirschhorn JN, Hofman A, Hollegaard MV, Hougaard DM, Hyppönen E, Inskip HM, Isaacs A, Jorgensen T, Kanaka-Gantenbein C, Kemp JP, Kiess W, Kilpeläinen TO, Klopp N, Knight BA, Kuzawa CW, McMahon G, Newnham JP, Niinikoski H, Oostra BA, Pedersen L, Postma DS, Ring SM, Rivadeneira F, Robertson NR, Sebert S, Simell O, Slowinski T, Tiesler CM, Tonjes A, Vaag A, Viikari JS, Vink JM, Vissing NH, Wareham NJ, Willemssen G, Witte DR, Zhang H, Zhao J, Meta-Analyses of G, Insulin-related traits C, Wilson JF, Stumvoll M, Prentice AM, Meyer BF, Pearson ER, Boreham CA, Cooper C, Gillman MW, Dedoussis GV, Moreno LA, Pedersen O, Saarinen M, Mohlke KL, Boomsma DI, Saw SM, Lakka TA, Korner A, Loos RJ, Ong KK, Vollenweider P, van Duijn CM, Koppelman GH, Hattersley AT, Holloway JW, Hoche B, Heinrich J, Power C, Melbye M, Guxens M, Pennell CE, Bonnelykke K, Bisgaard H, Eriksson JG, Widen E, Hakonarson H, Uitterlinden AG, Pouta A, Lawlor DA, Smith GD, Frayling TM, McCarthy MI, Grant SF, Jaddoe VW, Jarvelin MR, Timpson NJ, Prokopenko I, Freathy RM & Early Growth Genetics C (2013) New loci associated with birth weight identify genetic links between intrauterine growth and adult height and metabolism. *Nat Genet* 45, 76–82.
- Howe B, Fuchsberger C, Stephens M, Marchini J & Abecasis GR. (2012) Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat Genet* 44, 955–959.
- International HapMap C, Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, Gibbs RA, Belmont JW, Boudreau A, Hardenbol P, Leal SM, Pasternak S, Wheeler DA, Willis TD, Yu F, Yang H, Zeng C, Gao Y, Hu H, Hu W, Li C, Lin W, Liu S, Pan H, Tang X, Wang J, Wang W, Yu J, Zhang B, Zhang Q, Zhao H, Zhou J, Gabriel SB, Barry R, Blumenstiel B, Camargo A, Defelice M, Faggart M, Goyette M, Gupta S, Moore J, Nguyen H, Onofrio RC, Parkin M, Roy J, Stahl E, Winchester E, Ziaugra L, Altshuler D, Shen Y, Yao Z, Huang W, Chu X, He Y, Jin L, Liu Y, Shen Y, Sun W, Wang H, Wang Y, Wang Y, Xiong X, Xu L, Wayne MM, Tsui SK, Xue H, Wong JT, Galver LM, Fan JB, Gunderson K, Murray SS, Oliphant AR, Chee MS, Montpetit A, Chagnon F, Ferretti V, Leboeuf M, Olivier JF, Phillips MS, Roumy S, Sallee C, Verner A, Hudson TJ, Kwok PY, Cai D, Koboldt DC, Miller RD, Pawlikowska L, Taillon-Miller P, Xiao M, Tsui LC, Mak W, Song YQ, Tam PK, Nakamura Y, Kawaguchi T, Kitamoto T, Morizono T, Nagashima A, Ohnishi Y, Sekine A, Tanaka T, Tsunoda T, Deloukas P, Bird CP, Delgado M, Dermitzakis ET, Gwilliam R, Hunt S, Morrison J, Powell D, Stranger BE, Whittaker P, Bentley DR, Daly MJ, de Bakker PI, Barrett J, Chretien JR, Maller J, McCarrroll S, Patterson N, Pe'er I, Price A, Purcell S, Richter DY, Sabeti P, Saxena R, Schaffner SF, Sham PC, Varilly P, Altshuler D, Stein LD, Krishnan L, Smith AV, Tello-Ruiz MK, Thorisson GA, Chakravarti A, Chen PE, Cutler DJ, Kashuk CS, Lin S, Abecasis GR, Guan W, Li Y, Munro HM, Qin ZS, Thomas DJ, McVean G, Auton A, Bottolo L, Cardin N, Eyheramendy S, Freeman C, Marchini J, Myers S, Spencer C, Stephens M, Donnelly P, Cardon LR, Clarke G, Evans DM, Morris AP, Weir BS, Tsunoda T, Mullikin JC, Sherry ST, Feolo M, Skol A, Zhang H, Zeng C, Zhao H, Matsuda I, Fukushima Y, Macer DR, Suda E, Rotimi CN, Adebamowo CA, Ajayi I, Aniagwu T, Marshall PA, Nkwdimma C, Royal CD, Leppert MF, Dixon M, Peiffer A, Qiu R, Kent A, Kato K, Niikawa N, Adewole IF, Knoppers BM, Foster MW, Clayton EW, Watkin J, Gibbs RA, Belmont JW, Muzny D, Nazareth L, Sodergren E, Weinstock GM, Wheeler DA, Yakub I, Gabriel SB, Onofrio RC, Richter DJ, Ziaugra L, Birren BW, Daly MJ, Altshuler D, Wilson RK, Fulton LL, Rogers J, Burton J, Carter NP, Clee CM, Griffiths M, Jones MC, McLay K, Plumb RW, Ross MT, Sims SK, Willey DL, Chen Z, Han H, Kang L, Godbout M, Wallenburg JC, L'Archeveque P, Bellemare G, Saeki K, Wang H, An D, Fu H, Li Q, Wang Z, Wang R, Holden AL, Brooks LD, McEwen JE, Guyer MS, Wang VO, Peterson JL, Shi M, Spiegel J, Sung LM, Zacharia LF, Collins FS, Kennedy K, Jamieson R & Stewart J (2007) A second generation human haplotype map of over 3.1 million SNPs. *Nature* 449, 851–861.
- Jacobsen GK & Henriques UV. (1992) A fetal testis with intratubular germ cell neoplasia (ITGCN). *Mod Pathol* 5, 547–549.
- Kanetsky PA, Mitra N, Vardhanabhuti S, Li M, Vaughn DJ, Letrero R, Ciosek SL, Doody DR, Smith LM, Weaver J, Albano A, Chen C, Starr JR, Rader DJ, Godwin AK, Reilly MP, Hakonarson H, Schwartz SM & Nathanson KL. (2009) Common variation in KITLG and at 5q31.3 predisposes to testicular germ cell cancer. *Nat Genet* 41, 811–815.
- Kristensen DM, Sonne SB, Ottesen AM, Perrett RM, Nielsen JE, Almstrup K, Skakkebaek NE, Leffers H & Rajpert-De Meyts E. (2008) Origin of pluripotent germ cell tumours: the role of microenvironment during embryonic development. *Mol Cell Endocrinol* 288, 111–118.
- Kristiansen W, Karlsson R, Rounge TB, Whittington T, Andreassen BK, Magnusson PK, Fossa SD, Adami HO, Turnbull C, Haugen TB, Grotmol T & Wiklund F. (2015) Two new loci and gene sets related to sex determination and cancer progression are associated with susceptibility to testicular germ cell tumor. *Hum Mol Genet* 24, 4138–4146.
- Le Cornet C, Lortet-Tieulent J, Forman D, Beranger R, Flechon A, Fervers B, Schuz J & Bray F. (2014) Testicular cancer incidence to rise by 25% by 2025 in Europe? Model-based predictions in 40 countries using population-based registry data. *Eur J Cancer* 50, 831–839.
- Litchfield K, Levy M, Huddart RA, Shipley J & Turnbull C. (2016) The genomic landscape of testicular germ cell tumours: from susceptibility to treatment. *Nature reviews. Urology* 13, 409–419.
- Locke AE, Kahali B, Berndt SI, Justice AE, Pers TH, Day FR, Powell C, Vedantam S, Buchkovich ML, Yang J, Croteau-Chonka DC, Esko T, Fall T, Ferreira T, Gustafsson S, Kutalik Z, Luan J, Magi R, Randall JC, Winkler TW, Wood AR, Workalemahu T, Faul JD, Smith JA, Hua Zhao J, Zhao W, Chen J, Fehrmann R, Hedman AK, Karjalainen J, Schmidt EM, Absher D, Amin N, Anderson D, Beekman M, Bolton JL, Bragg-Gresham JL, Buyske S, Demirkan A, Deng G, Ehret GB, Feenstra B, Feitosa MF, Fischer K, Goel A, Gong J, Jackson AU, Kanoni S, Kleber ME, Kristiansson K, Lim U, Lotay V, Mangino M, Mateo Leach I, Medina-Gomez C, Medland SE, Nalls MA, Palmer CD, Pasko D, Pechlivanis S, Peters MJ, Prokopenko I, Shungin D, Stancakova A, Strawbridge RJ, Ju Sung Y, Tanaka T, Teumer A, Trompet S, van der Laan SW, van Setten J, Van Vliet-Ostaptchouk JV, Wang Z, Yengo L, Zhang W, Isaacs A, Albrecht E, Arnlöv J, Arscott GM, Attwood AP, Bandinelli S, Barrett A, Bas IN, Bellis C, Bennett AJ, Berne C, Blagieva R, Bluher M, Bohringer S, Bonnycastle LL, Bottcher Y, Boyd HA, Bruinenberg M, Caspersen IH, Ida Chen Y-D, Clarke R, Warwick Daw E, de Craen AJM, Delgado G, Dimitriou M, Doney ASF, Eklund N, Estrada K, Eury E, Folkersen L, Fraser RM, Garcia ME, Geller F, Giedraitis V, Gigante B, Go AS, Golay A, Goodall AH, Gordon SD, Gorski M, Grabe H-J, Grallert H, Grammer TB, Graszer J, Gronberg H, Groves CJ, Gusto G, Haessler J, Hall P, Haller T, Hallmans G, Hartman CA, Hassinen M, Hayward C, Heard-Costa NL, Helmer Q, Hengstenberg C, Holmen O, Hottenga J-J, James AL, Jeff JM, Johansson A, Jolley J, Juliusdottir T, Kinnunen L, Koenig W, Koskenvuo

- M, Kratzer W, Laitinen J, Lamina C, Leander K, Lee NR, Lichtner P, Lind L, Lindstrom J, Sin Lo K, Lobbens S, Lorbeer R, Lu Y, Mach F, Magnusson PKE, Mahajan A, McArdle WL, McLachlan S, Menni C, Merger S, Mihailov E, Milani L, Moayyeri A, Monda KL, Morken MA, Mulas A, Muller G, Muller-Nurasyid M, Musk AW, Nagaraja R, Nothen MM, Nolte IM, Pilz S, Rayner NW, Renstrom F, Rettig R, Ried JS, Ripke S, Robertson NR, Rose LM, Sanna S, Scharnagl H, Scholtens S, Schumacher FR, Scott WR, Seufferlein T, Shi J, Vernon Smith A, Smolonska J, Stanton AV, Steinhorsdottir V, Stirrups K, Stringham HM, Sundstrom J, Swertz MA, Swift AJ, Syvanen A-C, Tan S-T, Tayo BO, Thorand B, Thorleifsson G, Tyrer JP, Uh H-W, Vandenput L, Verhulst FC, Vermeulen SHGusto G, Haessler J, Hall P, Haller T, Hallmans G, Hartman CA, Hassinen M, Hayward C, Heard-Costa NL, Helmer Q, Hengstenberg C, Holmen O, Hottenga J-J, James AL, Jeff JM, Johansson A, Jolley J, Juliusdottir T, Kinnunen L, Koenig W, Koskenvuo M, Kratzer W, Laitinen J, Lamina C, Leander K, Lee NR, Lichtner P, Lind L, Lindstrom J, Sin Lo K, Lobbens S, Lorbeer R, Lu Y, Mach F, Magnusson PKE, Mahajan A, McArdle WL, McLachlan S, Menni C, Merger S, Mihailov E, Milani L, Moayyeri A, Monda KL, Morken MA, Mulas A, Muller G, Muller-Nurasyid M, Musk AW, Nagaraja R, Nothen MM, Nolte IM, Pilz S, Rayner NW, Renstrom F, Rettig R, Ried JS, Ripke S, Robertson NR, Rose LM, Sanna S, Scharnagl H, Scholtens S, Schumacher FR, Scott WR, Seufferlein T, Shi J, Vernon Smith A, Smolonska J, Stanton AV, Steinhorsdottir V, Stirrups K, Stringham HM, Sundstrom J, Swertz MA, Swift AJ, Syvanen A-C, Tan S-T, Tayo BO, Thorand B, Thorleifsson G, Tyrer JP, Uh H-W, Vandenput L, Verhulst FC, Vermeulen SH, Verweij N, Vonk JM, Waite LL, Warren HR, Waterworth D, Weedon MN, Wilkens LR, Willenborg C, Wilsgaard T, Wojczynski MK, Wong A, Wright AF, Zhang Q, The LifeLines Cohort S, Brennan EP, Choi M, Dastani Z, Drong AW, Eriksson P, Franco-Cereceda A, Gadin JR, Gharavi AG, Goddard ME, Handsaker RE, Huang J, Karpe F, Kathiresan S, Keildson S, Kiryluk K, Kubo M, Lee J-Y, Liang L, Lifton RP, Ma B, McCarroll SA, McKnight AJ, Min JL, Moffatt MF, Montgomery GW, Murabito JM, Nicholson G, Nyholt DR, Okada Y, Perry JRB, Dorajoo R, Reinmaa E, Salem RM, Sandholm N, Scott RA, Stolk L, Takahashi A, Tanaka T, van't Hooft FM, Vinkhuysen AAE, Westra H-J, Zheng W, Zondervan KT, The AC, The A-BMIWG, The CDC, The CC, The G, The I, The MI, The Mu TC, The MC, The PC, The ReproGen C, The GC, The International Endogene C, Heath AC, Arveiler D, Bakker SJL, Beilby J, Bergman RN, Blangero J, Bovet P, Campbell H, Caulfield MJ, Cesana G, Chakravarti A, Chasman DI, Chines PS, Collins FS, Crawford DC, Adrienne Cupples L, Cusi D, Danesh J, de Faire U, den Ruijter HM, Dominiczak AF, Erbel R, Erdmann J, Eriksson JG, Farrall M, Felix SB, Ferrannini E, Ferrieres J, Ford I, Forouhi NG, Forrester T, Franco OH, Gansevoort RT, Gejman PV, Gieger C, Gottesman O, Gudnason V, Gyllenstein U, Hall AS, Harris TB, Hattersley AT, Hicks AA, Hindorf LA, Hingorani AD, Hofman A, Homuth G, Kees Hovingh G, Humphries SE, Hunt SC, Hypponen E, Illig T, Jacobs KB, Jarvelin M-R, Jockel K-H, Johansen B, Jousilahti P, Wouter Jukema J, Julia AM, Kaprio J, Kastelein JJP, Keinänen-Kiukaanniemi SM, Kiemeny LA, Knekt P, Kooner JS, Kooperberg C, Kovacs P, Kraja AT, Kumari M, Kuusisto J, Lakka TA, Langenberg C, Le Marchand L, Lehtimäki T, Lyssenko V, Mannisto S, Marette A, Matise TC, McKenzie CA, McKnight B, Moll FL, Morris AD, Morris AP, Murray JC, Nelis M, Ohlsson C, Oldehinkel AJ, Ong KK, Madden PAF, Pasterkamp G, Peden JF, Peters A, Postma DS, Pramstaller PP, Price JF, Qi L, Raitakari OT, Rankinen T, Rao DC, Rice TK, Ridker PM, Rioux JD, Ritchie MD, Rudan I, Salomaa V, Samani NJ, Saramies J, Sarzynski MA, Schunkert H, Schwarz PEH, Sever P, Shuldiner AR, Sinisalo J, Stolk RP, Strauch K, Tonjes A, Tregouet D-A, Tremblay A, Tremoli E, Virtamo J, Vohl M-C, Volker U, Waeber G, Willemsen G, Witterman JC, Carola Zillikens M, Adair LS, Amouyel P, Asselbergs FW, Assimes TL, Bochud M, Boehm BO, Boerwinkle E, Bornstein SR, Bottinger EP, Bouchard C, Cauchi S, Chambers JC, Chanock SJ, Cooper RS, de Bakker PIW, Dedoussis G, Ferrucci L, Franks PW, Froguel P, Groop LC, Haiman CA, Hamsten A, Hui J, Hunter DJ, Hveem K, Kaplan RC, Kivimäki M, Kuh D, Laakso M, Liu Y, Martin NG, Marz W, Melbye M, Metspalu A, Moebus S, Munroe PB, Njolstad I, Oostra BA, Palmer CNA, Pedersen NL, Perola M, Perusse L, Peters U, Power C, Quertermous T, Rauramaa R, Rivadeneira F, Saaristo TE, Saleheen D, Sattar N, Schadt EE, Schlessinger D, Eline Slagboom P, Snieder H, Spector TD, Thorsteinsdottir U, Stumvoll M, Tuomilehto J, Uitterlinden AG, Uusitupa M, van der Harst P, Walker M, Wallaschofski H, Wareham NJ, Watkins H, Weir DR, Wichmann HE, Wilson JF, Zanen P, Borecki IB, Deloukas P, Fox CS, Heid IM, O'Connell JR, Strachan DP, Stefansson K, van Duijn CM, Abecasis GR, Franke L, Frayling TM, McCarthy MI, Visscher PM, Scherag A, Willer CJ, Boehnke M, Mohlke KL, Lindgren CM, Beckmann JS, Barroso I, North KE, Ingelsson E, Hirschhorn JN, Loos RJJ & Speliotes EK (2015) Genetic studies of body mass index yield new insights for obesity biology. *Nature* 518, 197–206.
- Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, Hasz R, Walters G, Garcia F, Young N, Foster B, Moser M, Karasik E, Gillard B, Ramsey K, Sullivan S, Bridge J, Magazine H, Syron J, Fleming J, Siminoff L, Traino H, Mosavel M, Barker L, Jewell S, Rohrer D, Maxim D, Filkins D, Harbach P, Cortadillo E, Berghuis B, Turner L, Hudson E, Feenstra K, Sobin L, Robb J, Branton P, Korzeniewski G, Shive C, Tabor D, Qi L, Groch K, Nampally S, Buia S, Zimmerman A, Smith A, Burges R, Robinson K, Valentino K, Bradbury D, Cosentino M, Diaz-Mayoral N, Kennedy M, Engel T, Williams P, Erickson K, Ardlie K, Winckler W, Getz G, DeLuca D, MacArthur D, Kellis M, Thomson A, Young T, Gelfand E, Donovan M, Meng Y, Grant G, Mash D, Marcus Y, Basile M, Liu J, Zhu J, Tu Z, Cox NJ, Nicolae DL, Gamazon ER, Im HK, Konkashbaev A, Pritchard J, Stevens M, Fluttre T, Wen X, Dermitzakis ET, Lappalainen T, Guigo R, Monlong J, Sammeth M, Koller D, Battle A, Mostafavi S, McCarthy M, Rivas M, Maller J, Rusyn I, Nobel A, Wright F, Shabalin A, Feolo M, Sharopova N, Sturcke A, Paschal J, Anderson JM, Wilder EL, Derr LK, Green ED, Struwing JP, Temple G, Volpi S, Boyer JT, Thomson EJ, Guyer MS, Ng C, Abdallah A, Colantuoni D, Insel TR, Koester SE, Little AR, Bender PK, Lehner T, Yao Y, Compton CC, Vaught JB, Sawyer S, Lockhart NC, Demchok J & Moore HF (2013) The Genotype-Tissue Expression (GTEx) project. *Nat Genet* 45, 580–585.
- McGlynn KA & Cook MB. (2009) Etiologic factors in testicular germ-cell tumors. *Future Oncol* 5, 1389–1402.
- McGlynn KA, Sakoda LC, Rubertone MV, Sesterhenn IA, Lyu C, Graubard BI & Erickson RL. (2007) Body size, dairy consumption, puberty, and risk of testicular germ cell tumors. *Am J Epidemiol* 165, 355–363.
- Oosterhuis JW & Looijenga LH. (2005) Testicular germ-cell tumours in a broader perspective. *Nat Rev Cancer* 5, 210–222.
- Palmer TM, Lawlor DA, Harbord RM, Sheehan NA, Tobias JH, Timpson NJ, Davey Smith G & Sterne JA. (2012) Using multiple genetic variants as instrumental variables for modifiable risk factors. *Stat Methods Med Res* 21, 223–242.
- Rajpert-De Meyts E. (2006) Developmental model for the pathogenesis of testicular carcinoma in situ: genetic and environmental aspects. *Human Reprod Update* 12, 303–323.
- Randall JC, Winkler TW, Kutalik Z, Berndt SI, Jackson AU, Monda KL, Kilpeläinen TO, Esko T, Mägi R, Li S, Workalemahu T, Feitosa MF, Croteau-Chonka DC, Day FR, Fall T, Ferreira T, Gustafsson S, Locke AE, Mathieson I, Scherag A, Vedantam S, Wood AR, Liang L, Steinhorsdottir V, Thorleifsson G, Dermitzakis ET, Dimas AS, Karpe F, Min JL, Nicholson G, Clegg DJ, Person T, Krohn JP, Bauer S, Buechler C, Eisinger K, Bonnetfond A, Froguel P, Hottenga J-J, Prokopenko I, Waite LL, Harris TB, Smith AV, Shuldiner AR, McArdle WL, Caulfield MJ, Munroe PB, Grönberg H, Chen Y-DI, Li G, Beckmann JS, Johnson T, Thorsteinsdottir U, Teder-Laving M, Khaw K-T, Wareham NJ, Zhao JH, Amin N, Oostra BA, Kraja AT, Province MA, Cupples LA, Heard-Costa NL, Kaprio J, Ripatti S, Surakka I, Collins FS, Saramies J, Tuomilehto J, Julia A, Salomaa V, Erdmann J, Hengstenberg C, Loley C, Schunkert H, Lamina C, Wichmann HE, Albrecht E, Gieger C, Hicks

- AA, Johansson Å, Pramstaller PP, Kathiresan S, Speliotes EK, Penninx B, Hartikainen A-L, Jarvelin M-R, Gyllenstein U, Boomsma DI, Campbell H, Wilson JF, Chanock SJ, Farrall M, Goel A, Medina-Gomez C, Rivadeneira F, Estrada K, Uitterlinden AG, Hofman A, Zillikens MC, den Heijer M, Kiemeny LA, Maschio A, Hall P, Tyrer J, Teumer A, Völzke H, Kovacs P, Tönjes A, Mangino M, Spector TD, Hayward C, Rudan I, Hall AS, Samani NJ, Attwood AP, Sambrook JG, Hung J, Palmer LJ, Lokki M-L, Sinisalo J, Boucher G, Huikuri H, Lorentzon M, Ohlsson C, Eklund N, Eriksson JG, Barlassina C, Rivolta C, Nolte IM, Snieder H, Van der Klauw MM, Van Vliet-Ostaptchouk JV, Gejman PV, Shi J, Jacobs KB, Wang Z, Bakker SJL, Mateo Leach I, Navis G, van der Harst P, Martin NG, Medland SE, Montgomery GW, Yang J, Chasman DI, Ridker PM, Rose LM, Lehtimäki T, Raitakari O, Absher D, Iribarren C, Basart H, Hovingh KG, Hyppönen E, Power C, Anderson D, Beilby JP, Hui J, Jolley J, Sager H, Bornstein SR, Schwarz PEH, Kristiansson K, Perola M, Lindström J, Swift AJ, Uusitupa M, Atalay M, Lakka TA, Rauramaa R, Bolton JL, Fowkes G, Fraser RM, Price JF, Fischer K, KrjutAikov K, Metspalu A, Mihailov E, Langenberg C, Luan J, Ong KK, Chines PS, Keinanen-Kiukkaanniemi SM, Saaristo TE, Edkins S, Franks PW, Hallmans G, Shungin D, Morris AD, Palmer CNA, Erbel R, Moebus S, Nöthen MM, Pechlivanis S, Hveem K, Narisu N, Hamsten A, Humphries SE, Strawbridge RJ, Tremoli E, Grallert H, Thorand B, Illig T, Koenig W, Müller-Nurasyid M, Peters A, Boehm BO, Kleber ME, März W, Winkelmann BR, Kuusisto J, Laakso M, Arveiler D, Cesana G, Kuulasmaa K, Virtamo J, Yarnell JWG, Kuh D, Wong A, Lind L, de Faire U, Gigante B, Magnusson PKE, Pedersen NL, Dedoussis G, Dimitriou M, Kolovou G, Kanoni S, Stirrups K, Bonnycastle LL, Njolstad I, Wilsgaard T, Ganna A, Rehnberg E, Hingorani A, Kivimäki M, Kumari M, Assimes TL, Barroso I, Boehnke M, Borecki IB, Deloukas P, Fox CS, Frayling T, Groop LC, Haritunians T, Hunter D, Ingelsson E, Kaplan R, Mohlke KL, O'Connell JR, Schlessinger D, Strachan DP, Stefansson K, van Duijn CM, Abecasis GR, McCarthy MI, Hirschhorn JN, Qi L, Loos RJJ, Lindgren CM, North KE, Heid IM, Consortium D & Investigators M (2013) Sex-stratified Genome-wide Association Studies Including 270,000 Individuals Show Sexual Dimorphism in Genetic Loci for Anthropometric Traits. *PLoS Genet* 9, e1003500.
- Rapley EA, Turnbull C, Al Olama AA, Dermitzakis ET, Linger R, Huddart RA, Renwick A, Hughes D, Hines S, Seal S, Morrison J, Nsengimana J, Deloukas P, Testicular Cancer Collaboration UK, Rahman N, Bishop DT, Easton DF & Stratton MR. (2009) A genome-wide association study of testicular germ cell tumor. *Nat Genet* 41, 807–810.
- Rasmussen F, Gunnell D, Ekblom A, Hallqvist J & Tynelius P. (2003) Birth weight, adult height, and testicular cancer: cohort study of 337,249 Swedish young men. *Cancer Causes Control* 14, 595–598.
- Richiardi L, Askling J, Granath F & Akre O. (2003) Body size at birth and adulthood and the risk for germ-cell testicular cancer. *Cancer Epidemiol Biomark Prev* 12, 669–673.
- Ruf CG, Isbarn H, Wagner W, Fisch M, Matthies C & Dieckmann KP. (2014) Changes in epidemiologic features of testicular germ cell cancer: age at diagnosis and relative frequency of seminoma are constantly and significantly increasing. *Urol Oncol* 32, 33 e31–36.
- Shungin D, Winkler TW, Croteau-Chonka DC, Ferreira T, Locke AE, Magi R, Strawbridge RJ, Pers TH, Fischer K, Justice AE, Workalemahu T, Wu JM, Buchkovich ML, Heard-Costa NL, Roman TS, Drong AW, Song C, Gustafsson S, Day FR, Esko T, Fall T, Kutalik Z, Luan J, Randall JC, Scherag A, Vedantam S, Wood AR, Chen J, Fehrmann R, Karjalainen J, Kahali B, Liu CT, Schmidt EM, Absher D, Amin N, Anderson D, Beekman M, Bragg-Gresham JL, Buyske S, Demirkan A, Ehret GB, Feitosa MF, Goel A, Jackson AU, Johnson T, Kleber ME, Kristiansson K, Mangino M, Mateo Leach I, Medina-Gomez C, Palmer CD, Pasko D, Pechlivanis S, Peters MJ, Prokopenko I, Stancakova A, Ju Sung Y, Tanaka T, Teumer A, Van Vliet-Ostaptchouk JV, Yengo L, Zhang W, Albrecht E, Arnlöv J, Arscott GM, Bandinelli S, Barrett A, Bellis C, Bennett AJ, Berne C, Bluher M, Bohringer S, Bonnet F, Bottcher Y, Bruinenberg M, Carba DB, Caspersen IH, Clarke R, Daw EW, Deelen J, Deelman E, Delgado G, Doney AS, Eklund N, Erdos MR, Estrada K, Eury E, Friedrich N, Garcia ME, Giedraitis V, Gigante B, Go AS, Golay A, Grallert H, Grammer TB, Grasser J, Grewal J, Groves CJ, Haller T, Hallmans G, Hartman CA, Hassinen M, Hayward C, Heikkilä K, Herzig KH, Helmer Q, Hillege HL, Holmen O, Hunt SC, Isaacs A, Itermann T, James AL, Johansson I, Juliusdottir T, Kalafati IP, Kinnunen L, Koenig W, Kooner IK, Kratzer W, Lamina C, Leander K, Lee NR, Lichtner P, Lind L, Lindstrom J, Lobbens S, Lorentzon M, Mach F, Magnusson PK, Mahajan A, McArdle WL, Menni C, Merger S, Mihailov E, Milani L, Mills R, Moayyeri A, Monda KL, Mooijart SP, Muhleisen TW, Mulas A, Muller G, Muller-Nurasyid M, Nagaraja R, Nalls MA, Narisu N, Glorioso N, Nolte IM, Olden M, Rayner NW, Renstrom F, Ried JS, Robertson NR, Rose LM, Sanna S, Schernagl H, Scholtens S, Sennblad B, Seufferlein T, Sitlani CM, Vernon Smith A, Stirrups K, Stringham HM, Sundstrom J, Swertz MA, Swift AJ, Syvanen AC, Tayo BO, Thorand B, Thorleifsson G, Tomaschitz A, Troffa C, van Oort FV, Verweij N, Vonk JM, Waite LL, Wennauer R, Wilsgaard T, Wojczynski MK, Wong A, Zhang Q, Hua Zhao J, Brennan EP, Choi M, Eriksson P, Folkersen L, Franco-Cereceda A, Gharavi AG, Hedman AK, Hivert MF, Huang J, Kanoni S, Karpe F, Keildson S, Kiryluk K, Liang L, Lifton RP, Ma B, McKnight AJ, McPherson R, Metspalu A, Min JL, Moffatt MF, Montgomery GW, Murabito JM, Nicholson G, Nyholt DR, Olsson C, Perry JR, Reinmaa E, Salem RM, Sandholm N, Schadt EE, Scott RA, Stolk L, Vallejo EE, Westra HJ, Zondervan KT, Consortium AD, Consortium CAD, Consortium CK, Consortium G, Consortium G, Glgc, Icbp, International Endogene C, LifeLines Cohort S, Investigators M, Mu TC, Consortium P, ReproGen C, Amouyel P, Arveiler D, Bakker SJ, Beilby J, Bergman RN, Blangero J, Brown MJ, Burnier M, Campbell H, Chakravarti A, Chines PS, Claudi-Boehm S, Collins FS, Crawford DC, Danesh J, de Faire U, de Geus EJ, Dorr M, Erbel R, Eriksson JG, Farrall M, Ferrannini E, Ferrieres J, Forouhi NG, Forrester T, Franco OH, Gansevoort RT, Gieger C, Gudnason V, Haiman CA, Harris TB, Hattersley AT, Heliovaara M, Hicks AA, Hingorani AD, Hoffmann W, Hofman A, Homuth G, Humphries SE, Hyppönen E, Illig T, Jarvelin MR, Johansen B, Jousilahti P, Julia AM, Kaprio J, Kee F, Keinanen-Kiukkaanniemi SM, Kooner JS, Kooperberg C, Kovacs P, Kraja AT, Kumari M, Kuulasmaa K, Kuusisto J, Lakka TA, Langenberg C, Le Marchand L, Lehtimäki T, Lyssenko V, Mannisto S, Marette A, Matise TC, McKenzie CA, McKnight B, Musk AW, Mohlenkamp S, Morris AD, Nelis M, Ohlsson C, Oldehinkel AJ, Ong KK, Palmer LJ, Penninx BW, Peters A, Pramstaller PP, Raitakari OT, Rankinen T, Rao DC, Rice TK, Ridker PM, Ritchie MD, Rudan I, Salomaa V, Samani NJ, Saramies J, Sarzynski MA, Schwarz PE, Shuldiner AR, Staessen JA, Steinthorsdottir V, Stolk RP, Strauch K, Tonjes A, Tremblay A, Tremoli E, Vohl MC, Volker U, Vollenweider P, Wilson JF, Witteman JC, Adair LS, Bochud M, Boehm BO, Bornstein SR, Bouchard C, Cauchi S, Caulfield MJ, Chambers JC, Chasman DI, Cooper RS, Dedoussis G, Ferrucci L, Froguel P, Grabe HJ, Hamsten A, Hui J, Hveem K, Jockel KH, Kivimäki M, Kuh D, Laakso M, Liu Y, März W, Munroe PB, Njolstad I, Oostra BA, Palmer CN, Pedersen NL, Perola M, Perusse L, Peters U, Power C, Quertermous T, Rauramaa R, Rivadeneira F, Saaristo TE, Saleheen D, Sinisalo J, Slagboom PE, Snieder H, Spector TD, Thorsteinsdottir U, Stumvoll M, Tuomilehto J, Uitterlinden AG, Uusitupa M, van der Harst P, Veronesi G, Walker M, Wareham NJ, Watkins H, Wichmann HE, Abecasis GR, Assimes TL, Berndt SI, Boehnke M, Borecki IB, Deloukas P, Franke L, Frayling TM, Groop LC, Hunter DJ, Kaplan RC, O'Connell JR, Qi L, Schlessinger D, Strachan DP, Stefansson K, van Duijn CM, Willer CJ, Visscher PM, Yang J, Hirschhorn JN, Zillikens MC, McCarthy MI, Speliotes EK, North KE, Fox CS, Barroso I, Franks PW, Ingelsson E, Heid IM, Loos RJ, Cupples LA, Morris AP, Lindgren CM & Mohlke KL (2015) New genetic loci link adipose and insulin biology to body fat distribution. *Nature* 518, 187–196.
- Skakkebaek NE, Berthelsen JG, Giwercman A & Muller J. (1987) Carcinoma-in-situ of the testis: possible origin from gonocytes and



- precursor of all types of germ cell tumours except spermatocytoma. *Int J Androl* 10, 19–28.
- Smith MK & Wakimoto BT. (2007) Complex regulation and multiple developmental functions of misfire, the *Drosophila melanogaster* *ferlin* gene. *BMC Dev Biol* 7, 21.
- Turnbull C, Rapley EA, Seal S, Pernet D, Renwick A, Hughes D, Ricketts M, Linger R, Nsengimana J, Deloukas P, Huddart RA, Bishop DT, Easton DF, Stratton MR, Rahman N & Testicular Cancer Collaboration UK. (2010) Variants near DMRT1, TERT and ATF7IP are associated with testicular germ cell cancer. *Nat Genet* 42, 604–607.
- Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S, Chu AY, Estrada K, Luan J, Kutalik Z, Amin N, Buchkovich ML, Croteau-Chonka DC, Day FR, Duan Y, Fall T, Fehrmann R, Ferreira T, Jackson AU, Karjalainen J, Lo KS, Locke AE, Magi R, Mihailov E, Porcu E, Randall JC, Scherag A, Vinkhuyzen AA, Westra HJ, Winkler TW, Workalemahu T, Zhao JH, Absher D, Albrecht E, Anderson D, Baron J, Beekman M, Demirkan A, Ehret GB, Feenstra B, Feitosa MF, Fischer K, Fraser RM, Goel A, Gong J, Justice AE, Kanoni S, Kleber ME, Kristiansson K, Lim U, Lotay V, Lui JC, Mangino M, Mateo Leach I, Medina-Gomez C, Nalls MA, Nyholt DR, Palmer CD, Pasko D, Pechlivanis S, Prokopenko I, Ried JS, Ripke S, Shungin D, Stancakova A, Strawbridge RJ, Sung YJ, Tanaka T, Teumer A, Trompet S, van der Laan SW, van Setten J, Van Vliet-Ostaptchouk JV, Wang Z, Yengo L, Zhang W, Afzal U, Arnlöv J, Arscott GM, Bandinelli S, Barrett A, Bellis C, Bennett AJ, Berne C, Bluher M, Bolton JL, Bottcher Y, Boyd HA, Bruinenberg M, Buckley BM, Buyske S, Caspersen IH, Chines PS, Clarke R, Claudi-Boehm S, Cooper M, Daw EW, De Jong PA, Deelen J, Delgado G, Denny JC, Dhonukshe-Rutten R, Dimitriou M, Doney AS, Dorr M, Eklund N, Eury E, Folkersen L, Garcia ME, Geller F, Giedraitis V, Go AS, Grallert H, Grammer TB, Grassler J, Gronberg H, de Groot LC, Groves CJ, Haessler J, Hall P, Haller T, Hallmans G, Hannemann A, Hartman CA, Hassinen M, Hayward C, Heard-Costa NL, Helmer Q, Hemani G, Henders AK, Hillege HL, Hlatky MA, Hoffmann W, Hoffmann P, Holmen O, Houwing-Duistermaat JJ, Illig T, Isaacs A, James AL, Jeff J, Johansen B, Johansson A, Jolley J, Juliusdottir T, Junttila J, Kho AN, Kinnunen L, Klopp N, Kocher T, Kratzer W, Lichtner P, Lind L, Lindstrom J, Lobbens S, Lorentzon M, Lu Y, Lyssenko V, Magnusson PK, Mahajan A, Maillard M, McArdle WL, McKenzie CA, McLachlan S, McLaren PJ, Menni C, Merger S, Milani L, Moayyeri A, Monda KL, Morken MA, Muller G, Muller-Nurasyid M, Musk AW, Narisu N, Nauck M, Nolte IM, Nothen MM, Oozageer L, Pilz S, Rayner NW, Renstrom F, Robertson NR, Rose LM, Roussel R, Sanna S, Scharnagl H, Scholtens S, Schumacher FR, Schunkert H, Scott RA, Sehmi J, Seufferlein T, Shi J, Silventoinen K, Smit JH, Smith AV, Smolonska J, Stanton AV, Stirrups K, Stott DJ, Stringham HM, Sundstrom J, Swertz MA, Syvanen AC, Tayo BO, Thorleifsson G, Tyrer JP, van Dijk S, van Schoor NM, van der Velde N, van Heemst D, van Oort FV, Vermeulen SH, Verweij N, Vonk JM, Waite LL, Waldenberger M, Wennauer R, Wilkens LR, Willenborg C, Wilsgaard T, Wojczynski MK, Wong A, Wright AF, Zhang Q, Arveiler D, Bakker SJ, Beilby J, Bergman RN, Bergmann S, Biffar R, Blangero J, Boomsma DI, Bornstein SR, Bovet P, Brambilla P, Brown MJ, Campbell H, Caulfield MJ, Chakravarti A, Collins R, Collins FS, Crawford DC, Cupples LA, Danesh J, de Faire U, den Ruijter HM, Erbel R, Erdmann J, Eriksson JG, Farrall M, Ferrannini E, Ferrieres J, Ford I, Forouhi NG, Forrester T, Gansevoort RT, Gejman PV, Gieger C, Golay A, Gottesman O, Gudnason V, Gyllenstein U, Haas DW, Hall AS, Harris TB, Hattersley AT, Heath AC, Hengstenberg C, Hicks AA, Hindorf LA, Hingorani AD, Hofman A, Hovingh GK, Humphries SE, Hunt SC, Hypponen E, Jacobs KB, Jarvelin MR, Jousilahti P, Jula AM, Kaprio J, Kastelein JJ, Kayser M, Kee F, Keinanen-Kiukkaanniemi SM, Kiemeny LA, Kooner JS, Kooperberg C, Kosken S, Kovacs P, Kraja AT, Kumari M, Kuusisto J, Lakka TA, Langenberg C, Le Marchand L, Lehtimäki T, Lupoli S, Madden PA, Mannisto S, Manunta P, Marette A, Matise TC, McKnight B, Meitinger T, Moll FL, Montgomery GW, Morris AD, Morris AP, Murray JC, Nelis M, Ohlsson C, Oldehinkel AJ, Ong KK, Ouwehand WH, Pasterkamp G, Peters A, Pramstaller PP, Price JF, Qi L, Raitakari OT, Rankinen T, Rao DC, Rice TK, Ritchie M, Rudan I, Salomaa V, Samani NJ, Saramies J, Sarzynski MA, Schwarz PE, Sebert S, Sever P, Shuldiner AR, Sinisalo J, Steinthorsdottir V, Stolk RP, Tardif JC, Tonjes A, Tremblay A, Tremoli E, Virtamo J, Vohl MC, Electronic Medical R, Genomics C, Consortium MI, Consortium P, LifeLines Cohort S, Amouyel P, Asselbergs FW, Assimes TL, Bochud M, Boehm BO, Boerwinkle E, Bottinger EP, Bouchard C, Cauchi S, Chambers JC, Chanock SJ, Cooper RS, de Bakker PI, Dedoussis G, Ferrucci L, Franks PW, Froguel P, Groop LC, Haiman CA, Hamsten A, Hayes MG, Hui J, Hunter DJ, Hveem K, Jukema JW, Kaplan RC, Kivimäki M, Kuh D, Laakso M, Liu Y, Martin NG, Marz W, Melbye M, Moebus S, Munroe PB, Njolstad I, Oostra BA, Palmer CN, Pedersen NL, Perola M, Perusse L, Peters U, Powell JE, Power C, Quertermous T, Rauramaa R, Reinmaa E, Ridker PM, Rivadeneira F, Rotter JJ, Saaristo TE, Saleheen D, Schlessinger D, Slagboom PE, Snieder H, Spector TD, Strauch K, Stumvoll M, Tuomilehto J, Uusitupa M, van der Harst P, Volzke H, Walker M, Wareham NJ, Watkins H, Wichmann HE, Wilson JF, Zanen P, Deloukas P, Heid IM, Lindgren CM, Mohlke KL, Speliotes EK, Thorsteinsdottir U, Barroso I, Fox CS, North KE, Strachan DP, Beckmann JS, Berndt SI, Boehnke M & Borecki IB, McCarthy MI, Metspalu A, Stefansson K, Uitterlinden AG, van Duijn CM, Franke L, Willer CJ, Price AL, Lettre G, Loos RJ, Weedon MN, Ingelsson E, O'Connell JR, Abecasis GR, Chasman DI, Goddard ME, Visscher PM, Hirschhorn JN & Frayling TM (2014) Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat Genet* 46, 1173–1186.
- Zavos C, Andreadis C, Diamantopoulos N & Mouratidou D. (2004) A hypothesis on the role of insulin-like growth factor I in testicular germ cell tumours. *Med Hypotheses* 63, 511–514.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

**Figure S1** TGCT and adult height effect sizes (and 95% CI) for 379 adult height risk SNPs.

**Figure S2** TGCT and BMI effect sizes (and 95% CI) for 69 BMI risk SNPs.

**Figure S3** TGCT and HIPadjBMI effect sizes (and 95% CI) for ten HIPadjBMI risk SNPs.

**Figure S4** TGCT and WCadjBMI effect sizes (and 95% CI) for 25 WCadjBMI risk SNPs.

**Figure S5** TGCT and WHRadjBMI effect sizes (and 95% CI) for 44 WHRadjBMI risk SNPs.

**Figure S6** TGCT and birth weight effect sizes (and 95% CI) for seven birth weight risk SNPs.

**Figure S7** TGCT and childhood obesity effect sizes (and 95% CI) for seven childhood obesity risk SNPs.

**Figure S8** TGCT and height (male dimorphism) effect sizes (and 95% CI) for 15 height (male dimorphism) risk SNPs.

**Figure S9** TGCT and HIPadjBMI (male specific) effect sizes (and 95% CI) for four HIPadjBMI (male-specific) risk SNPs.

**Figure S10** TGCT and WCadjBMI (male specific) effect sizes (and 95% CI) for nine WCadjBMI (male specific) risk SNPs.

**Figure S11** TGCT and WHRadjBMI (male specific) effect sizes (and 95% CI) for seven WHRadjBMI (male specific) risk SNPs.

**Figure S12** Estimated power of the present study to detect a range of odds ratios of height and TGCT risk. SNPs used here are estimated to explain ~20% of the phenotypic variation in height (Wood *et al.* 2014).



**Figure S13** Estimated power of the present study to detect a range of odds ratios of BMI and TGCT risk. SNPs used here are estimated to explain ~3% of the phenotypic variation in BMI (Locke *et al.* 2015).

**Figure S14** Estimated power of the present study to detect a range of odds ratios of WHRadjBMI and TGCT risk. SNPs used here are estimated to explain ~1.36% of the phenotypic variation of WHRadjBMI (Shungin *et al.* 2015).

**Figure S15** Estimated power of the present study to detect a range of odds ratios of birth weight and TGCT risk. SNPs used here are estimated

to explain ~0.76% of the phenotypic variation of birth weight (Horikoshi *et al.* 2013).

**Table S1** Results of linkage disequilibrium regression analysis for seven anthropometric traits against TGCT, as well as male-specific components for four of these traits.

**Table S2** Egger regression results for each of the traits for which Mendelian randomisation was performed.

**Table S3** Inverse weighted variance results for each of the traits for which Mendelian randomisation was performed.