

Title tbd

Matthew A. Niemiro

July 8, 2023

Contents

0.1	December 2022	3
0.1.1	(11/28) Ok, let's give this a try	3
0.1.2	(12/1) Why simplicial sets, simplicial complexes	3
0.1.3	(12/4) Basic structure in \mathbf{sSet}	3
0.1.4	(12/6) Colimits in/functors out of \mathbf{sSet}	4
0.1.5	(12/23) The singular complex and geometric realization	5
0.2	January 2023	7
0.2.1	(1/23) Plans have changed, nerves of categories	7
0.2.2	(1/26) Spines	7
0.2.3	(1/30) Inner Horns	8
0.3	February 2023	9
0.3.1	(2/4) Quasicategories	9
0.3.2	(2/6) Sub, opposite quasicategories	9
0.3.3	(2/7) Examples of ∞ -categories	10
0.3.4	(2/8) The fundamental category of a simplicial set	11
0.3.5	(2/9) Homotopy for ∞ -categories	11
0.3.6	(2/12) About composition in ∞ -categories	12
0.3.7	(2/13) Isomorphisms and inverses in ∞ -categories	12
0.3.8	(2/13) ∞ -groupoids, cores, and Kan complexes	13
0.3.9	(2/15) The functor quasicategory	13
0.3.10	(2/16) Lifting properties time — weakly saturated classes	14
0.3.11	(2/16) Classes of horns, anodyne morphisms	14
0.3.12	(2/16) Lifting calculus	15
0.3.13	(2/17) Inner fibrations	15
0.3.14	(2/20) Factorizations	16
0.3.15	(2/21) Factorization systems and unique lifts	16
0.3.16	(2/23) Degenerate cells	17
0.3.17	(2/23) The skeletal filtration	18
0.3.18	(2/25) Pushout-products, pullback-homs	19
0.4	March 2023	20
0.4.1	(3/5) Pullback-hom as an enriched lifting problem	20
0.5	April 2023	21
0.6	May 2023	22
0.6.1	(5/3) Operads for (Peter) May	22
0.6.2	(5/5) Basic examples of operads	24
0.6.3	(5/6) Warm-up: monoids are $\mathcal{A}\text{ssoc}$ -algebras	24
0.6.4	(5/9) Associativity up to homotopy, Stasheff associahedra, and A_∞ -operads	25
0.6.5	(5/11) A_n -operad stuff	26
0.6.6	(5/13) Rings via operads?	27
0.6.7	The rest of May	28
0.7	June 2023	29
0.7.1	(6/13) June activities, monoidal categories	29
0.7.2	(6/15) Initial, final objects	29
0.7.3	(6/18) Stable ∞ -categories	30
0.7.4	(6/20) Idempotents	32
0.7.5	(6/21) I hate idempotents today. K -theory?	33
0.7.6	(6/26) Structure of $\text{Hom}_{\mathcal{C}}(-, -)$ for (various adjectives) categories	34

0.7.7	(6/28) Anima and other examples of ∞ -categories via N^c	35
0.8	July 2023	37
0.8.1	(7/5) Derived ∞ -categories	37

0.1 December 2022

0.1.1 (11/28) Ok, let's give this a try

I want to read *Higher Topos Theory* (HTT). That book is > 700 pages, and fairly dense ones, so it's a bit of a project. On top of that, HTT isn't really a self-contained read (and that's sort of the nature of the subject). So I'll be drawing on lots of additional material. The subject seems well worth learning, even unavoidable at times (e.g., in the areas I am interested in), so this should be a productive little activity.

...

I need to cover some ground before actually opening HTT. First I need to think about *simplicial sets*. Their role in higher category theory is ubiquitous. They give us a combinatorial model for the homotopy theory of spaces, and also a model for ∞ -categories. (On the list of things to do is make precise sense of those statements.) Some references are [Rie], [Mat], [Fri08], and kerodon.net.

Definition 0.1.1. Denote by Δ the *simplex category*, defined to have...

- As objects, the ordered set $[n] := \{0 < 1 < \dots < n\}$ for each $n \geq 0$; and
- As maps, the weakly order-preserving set maps.

Definition 0.1.2. A *simplicial set* is a contravariant functor $\Delta \rightarrow \text{Set}$. The *category of simplicial sets*, denoted \mathbf{sSet} , is the functor category $\text{Fun}(\Delta^{op}, \text{Set})$.

Notation 0.1.1. Let $X : \Delta^{op} \rightarrow \text{Set}$ be a simplicial set. We may write it X_\bullet , and denote by X_n the set $X([n])$. We call the elements of X_n the *n -simplices* of X .

Notation 0.1.2. We may write $\langle f_0, \dots, f_n \rangle$ to denote the function $[n] \rightarrow [m]$ given by $n \mapsto f_n$.

Simplicial sets are not just simplices. They carry additional structure, that arising from morphisms in Δ . We can give a simple description of Δ . This in turn gives some intuition for what a simplicial set "is."

Proposition 0.1.3 (The structure of Δ). *For each $n \geq 0$ and $0 \leq i \leq n$, define the*

$$\begin{aligned} & i\text{-th face map } d^i : [n-1] \rightarrow [n] \text{ as } \langle 0, \dots, \hat{i}, \dots, n \rangle, \text{ and the} \\ & i\text{-th degeneracy map } s^i : [n+1] \rightarrow [n] \text{ as } \langle 0, \dots, i, i, \dots, n \rangle. \end{aligned}$$

Every morphism in Δ may be written as a composition of face and degeneracy maps. (Also, the face/degeneracy maps satisfy various relations, the simplicial identities; in fact Δ is the category generated by those maps, subject to these identities.)

Thus a simplicial set X_\bullet can be described as a collection of sets X_n (n -cells) together with face and degeneracy maps which satisfy the "simplicial identities." I should write more about how this notion arises from topology, in particular the singular complex. That in turn would be a good time to relate all this back to topology (nerves, geometric realization, ...) which is important.

0.1.2 (12/1) Why simplicial sets, simplicial complexes

I had stuff written here. But it was incomplete, and the "story" here is an aside I want to write about a bit more carefully at some point. I'm leaving this day blank for the time being.

0.1.3 (12/4) Basic structure in \mathbf{sSet}

We need to make some terminology regarding / record examples of simplicial sets.

Definition 0.1.4. The *standard n -simplex* Δ^n is the simplicial set represented by $[n]$, i.e. $\Delta^n := \text{Hom}_\Delta(-, [n])$.

Definition 0.1.5. Let X_\bullet, Y_\bullet be simplicial sets. We say Y_\bullet is a *simplicial subset* of X_\bullet if $Y_n \subseteq X_n$ and $Xf|_{Y_n} = Yf$ for every $n \geq 0$ and simplicial operator f . In other words, the action of operators on Y is the restriction of their action on X . In other words, Y_\bullet is a subfunctor of X_\bullet .

Proposition 0.1.6. Let X_\bullet be a simplicial set. The Yoneda lemma asserts a bijection $\text{Hom}_{\text{sSet}}(\Delta^n, X_\bullet) \cong X_n$. Under this bijection, each n -cell $a \in X_n$ corresponds to a map $f_a : \Delta^n \rightarrow X_\bullet$ satisfying $f_a(\text{id}_{[n]}) = a$.

Definition 0.1.7. Let X_\bullet be a simplicial set. By the above, we may identify its n -cells with maps $\Delta^n \rightarrow X_\bullet$. Call a cell $a \in X_n$ *degenerate* if it factors as $\Delta^n \rightarrow \Delta^m \rightarrow X_\bullet$ for some $m < n$. (See [Lur22, Tag 0011] for equivalent conditions.)

Proposition 0.1.8. The standard simplex Δ^n has a unique non-degenerate n -simplex, that arising from $\text{id}_{[n]}$. We may call this the *generator* of Δ^n .

Definition 0.1.9 (Boundary of Δ^n). Define a simplicial subset $\partial\Delta^n$, the *boundary of Δ^n* , by

$$(\partial\Delta^n)_k := \{\text{non-surjective maps } [k] \rightarrow [n]\} \subseteq \text{Hom}_{\Delta^{op}}([k], [n]).$$

Proposition 0.1.10. The boundary of Δ^n is the maximal proper simplicial subset of Δ^n .

Definition 0.1.11 (Horns in Δ^n). For $0 \leq i \leq n$, define a simplicial subset Λ_i^n , the *i -th horn in Δ^n* , by

$$(\Lambda_i^n)_k := \{f \in \text{Hom}_{\Delta^{op}}([k], [n]) : f([k]) \cup \{i\} \neq [n]\}.$$

In other words, its cells are those maps “missing something besides i .” A horn Λ_i^n is called *outer* if $i \neq 0, n$ and *inner* otherwise.

Any simplicial operator $f : [m] \rightarrow [n]$ factors through its image, i.e. we can uniquely write $f = f^{inj} f^{surj}$, a surjection followed by an injection. Furthermore, this is unique. We get the following.

Proposition 0.1.12. Let $\sigma : \Delta^n \rightarrow X_\bullet$ be an n -cell of X_\bullet . Then σ factors uniquely as

$$\Delta^n \xrightarrow{\alpha} \Delta^m \xrightarrow{\tau} X_\bullet,$$

Where α represents a surjection $[n] \rightarrow [m]$ and τ is not degenerate. Call m the *dimension* of the cell σ . (My notation, maybe poor, not that important.)

So, degenerate n -simplices are just non-degenerate simplices in a lower dimension (their “dimension”), trying to bite off more than they can chew.

Definition 0.1.13 (Skeleta). Let X_\bullet be a simplicial set. For $k \geq -1$, define a simplicial subset $\text{sk}_k(X_\bullet)$, the *k -skeleton* of X_\bullet , by

$$(\text{sk}_k(X_\bullet))_n := \{n\text{-simplices of } X_\bullet \text{ with dimension at most } k\}.$$

Remark 0.1.14. The face maps $d^i : [n-1] \rightarrow [n]$ induce maps $d^i : \Delta^{n-1} \rightarrow \Delta^n$ via post-composition. Now, consider an n -cell $a \in X_n$ and its representation $a : \Delta^n \rightarrow X_\bullet$. We have that $d_i(a) \in X_{n-1}$ is represented by ad^i .

0.1.4 (12/6) Colimits in/functors out of sSet

Today I want to understand part of Akhil’s notes, about functors out of sSet. This is closely related to understanding colimits in sSet, by general theory for presheaf categories. So we also want to understand colimits in sSet. (And we should want to understand these regardless.) Let’s go over this.

Here’s a standard structure result for presheaf categories.

Proposition 0.1.15. If a category C is small, then every presheaf on C is canonically the colimit of representable presheaves. In particular, every simplicial set is canonically the colimit of standard simplices.

Proof. This is written out in Akhil’s notes. I’ll give the idea. Also see [Lur22, Remark 00X5].

Consider a presheaf $F : C^{op} \rightarrow \text{Set}$. We associate to F the category D_F with

- Objects: morphisms from represented presheaves to F , i.e. arrows $[-, X] \rightarrow F$; and
- Morphisms: morphisms between represented presheaves such that the obvious triangle commutes.

There is a functor $\phi_F : D_F \rightarrow \text{PShv}(\mathcal{C})$ which sends objects $[-, X] \rightarrow F$ to $[-, X]$. By construction, for each object $c \in D_F$, there is a morphism $\phi_F(c) \rightarrow F$, and the diagram described by ϕ_F together with these morphisms commutes. We therefore have a distinguished morphism

$$\lim_{D_F} \phi_F \rightarrow F.$$

This map turns out to be an isomorphism. \square

Hereafter, denote by $\overline{\mathcal{C}}$ the category of presheaves on \mathcal{C} .

Suppose D is cocomplete. We want to understand functors $\overline{F} : \overline{\mathcal{C}} \rightarrow D$. The previous proposition says that objects in $\overline{\mathcal{C}}$ are colimits of representables. So, if \overline{F} preserves colimits, then \overline{F} is determined by $\overline{F}|_{\mathcal{C}}$, i.e. what it does to \mathcal{C} (embedded via Yoneda). We've described an injection of sets

$$\text{Fun}'(\overline{\mathcal{C}}, D) \hookrightarrow \text{Fun}(\mathcal{C}, D). \quad (16)$$

Here, Fun' denotes the set of colimit-preserving functors.

Conversely, suppose given a functor $F : \mathcal{C} \rightarrow D$. Does it extend along the Yoneda embedding to a functor $\overline{F} : \overline{\mathcal{C}} \rightarrow D$? We can do something here, let me write it out:

- (1) As above, for each presheaf $G : \mathcal{C}^{op} \rightarrow \text{Set}$, consider it as a colimit of $\phi_G : D_G \rightarrow \mathcal{C}$. (We can do this because it lands in represented functors.)
- (2) This is 'functorial' in the following sense: a morphism $G \rightarrow H$ induces a functor $D_G \rightarrow D_H$ such that the obvious triangle commutes.
- (3) Define a functor $\overline{F} : \overline{\mathcal{C}} \rightarrow D$ by

$$\overline{F}(G) := \text{colim}_{D_G} F \circ \phi_G.$$

This is a functor because of (2).

This functor \overline{F} really extends F , i.e. the obvious diagram commutes. For suppose $G = [-, c]$; then D_G has a final object $[-, c] \rightarrow [-, c]$, therefore $\overline{F}(G) = \text{colim}_{D_G} F \circ \phi_G = F(G)$.

Proposition 0.1.17. *Suppose given a functor $F : \mathcal{C} \rightarrow D$ to a cocomplete category. Then the associated functor $\overline{F} : \overline{\mathcal{C}} \rightarrow D$ constructed above preserves all colimits. In fact, \overline{F} is a left adjoint. The right adjoint to \overline{F} is the functor defined by*

$$D \ni d \mapsto (c \mapsto \text{Hom}_D(Fc, d)) \in \overline{\mathcal{C}}.$$

Proposition 0.1.18. *Suppose given a functor $F : \mathcal{C} \rightarrow D$ to a cocomplete category. Then the mapping $F \mapsto \overline{F}$ describes a bijection of sets*

$$\text{Fun}(\mathcal{C}, D) \xrightarrow{\sim} \text{LeftAdjoints}(\overline{\mathcal{C}}, D).$$

The proofs are short and formal.

Corollary 0.1.19. *If a functor $\overline{F} : \overline{\mathcal{C}}^{op} \rightarrow \text{Set}$ takes colimits to limits, then it is representable.*

Proof. Suppose as given \overline{F} . By the above, it is left adjoint to some $\overline{G} : \text{Set}^{op} \rightarrow \mathcal{C}$. Define $f := \overline{G}(\{pt\})$, the image of the terminal object in Set^{op} . I claim that f represents F . (Insert short, formal proof; it's in Akhil's notes.) \square

0.1.5 (12/23) The singular complex and geometric realization

Finals are over and I've had some time to wind down at home. Last time I worked through part of Akhil's notes about functors out of sSet . (Emily Riehl talks about something similar in her notes, but I have not gotten through those, so let me say nothing about that right now.)

Next I want to relate Top , Cat , and sSet . This is the backdrop for the idea that higher categories "bridge" topology/homotopy theory and ordinary categories. Today I'll go over the relation of sSet to Top , by which I mean the adjunction

the geometric realization functor \dashv the total singular complex functor.

There are a few ways to introduce this adjunction. Lurie, Charles, and Akhil each do it differently. As a matter of taste, I prefer Akhil's approach. (Possibly related: [Lur22, Tag 002D].) Lurie's approach has some important ideas behind it too, I think, but that is overruled because I am feeling sleepy today.

Definition 0.1.20. Define a functor $|-| : \Delta \rightarrow \mathbf{Top}$ as follows.

- Each object $[n]$ is sent to the *topological n -simplex* $\Delta_{top}^n \subseteq \mathbb{R}^{n+1}$, defined as those (t_0, \dots, t_n) satisfying $t_i \geq 0$ and $\sum t_i = 1$ and given the subspace topology.
- Each morphism $f : [m] \rightarrow [n]$ is sent to the map

$$(t_0, \dots, t_n) \mapsto (u_j), \quad u_j = \sum_{i: f(i)=j} t_i.$$

Definition 0.1.21 (Geometric realization). Since \mathbf{Top} is cocomplete, according to Proposition 0.1.17 and 0.1.18 the functor of Definition 0.1.20 extends uniquely to a left adjoint $|-| : \mathbf{sSet} \rightarrow \mathbf{Top}$. We call this *geometric realization*.

In fact, as in Proposition 0.1.18, we know the right adjoint to geometric realization. It sends a space X to the simplicial set $[n] \mapsto \mathrm{Hom}_{\mathbf{Top}}([n], X) = \Delta_{top}^n(X)$. This is an important construction I maybe should have defined earlier.

Definition 0.1.22 (Singular complex). Let X be a space. Denote by $\mathrm{Sing}(X)_\bullet$ the simplicial set given as follows.

- The n -cells are the continuous maps $\Delta_{top}^n \rightarrow X$, and
- Each simplicial operator $f : [m] \rightarrow [n]$ acts by precomposing with the continuous map

$$\Delta_{top}^m \rightarrow \Delta_{top}^n, \quad (t_j) \mapsto (u_j = \sum_{f(i)=j} t_i).$$

We call $\mathrm{Sing}(X)_\bullet$ the *singular complex* of X . We define a functor $\mathrm{Sing}(-)_\bullet : \mathbf{Top} \rightarrow \mathbf{sSet}$ in the obvious way.

Proposition 0.1.23. *Prior discussion tells us that geometric realization $|-| : \mathbf{sSet} \rightarrow \mathbf{Top}$ is left adjoint to the singular complex functor.*

Proposition 0.1.24. *Since geometric realization is a left adjoint, it commutes with colimits. Furthermore, geometric realization commutes with finite limits of compactly generated spaces.*

We will see later that this adjunction is homotopically well-behaved.

I next want to describe geometric realization. We already have the general construction laid out for us by Proposition 0.1.17 and the preceding discussion. Given X_\bullet , we will form the *category of simplices*, also called its *category of elements*, whose elements are the morphisms $\Delta^n \rightarrow X_\bullet$ (i.e., the cells of X_\bullet), and we take the colimit of $|-|$ restricted to this subcategory. (This is not circular since we are really applying the “baby” geometric realization to the simplex category, Yoneda embedded.)

Definition 0.1.25. Given $X_\bullet \in \mathbf{sSet}$, its *category of simplices* or *category of elements* has as objects all morphisms $\Delta^n \rightarrow X_\bullet$ for every n , and morphisms the maps $\Delta^m \rightarrow \Delta^n$ making the obvious diagram commute. We write this category $\mathrm{el}(X)$.

This category of elements/simplices $\mathrm{el}(X)$ is precisely the category D_X described on (12/6) with $C = \Delta$. (Lurie writes this Δ_X .) Also as noted there, there is a natural functor $\phi_X : \mathrm{el}(X) \rightarrow \mathbf{sSet}$. Geometric realization is by definition the colimit

$$|X| \cong \mathrm{co} \lim_{\mathrm{el}(X)} |-| \circ \phi_X.$$

Here, we are thinking of the “baby” geometric realization defined only on Δ .

Remark 0.1.26. General machinery gave us geometric realization. I think there are a few things worth saying about this, but I don't totally know what. I'll leave this remark here as a “to-do.” Some possibly related keywords and references: “Grothendieck construction,” “Kan extension,” nLab, [Rie, §4], and Subsection 01Q7.

0.2 January 2023

0.2.1 (1/23) Plans have changed, nerves of categories

I've gone radio silent for a month. One big reason why is that I am busy this semester. Another is that some mutuals want to organize a reading group/seminar similar, but not identical to, what I've been trying to do here, and I may join them. Maybe the biggest difference is that they want to focus on Charles' quasicategory notes (under Charles' supervision).

This will probably mean repeating myself a bit while I change tracks to Charles' notes.

In any case, I want to talk about the nerve of a category. This is part of the basic "Spaces, categories, and simplicial sets" picture. In particular, the nerve of a category is a simplicial set encoding that category.

Definition 0.2.1. Let C be a category. Define a simplicial set NC , the *nerve* of C to have as cells $(NC)_n := \text{Hom}_{\text{Cat}}([n], C)$ and so that operators $f : [m] \rightarrow [n]$ act by precomposition. This defines the *nerve functor* $N : \text{Cat} \rightarrow \text{sSet}$.

Here's a feel for the structure of a nerve. The n -cells of NC may be canonically identified with the set of length n tuples of composable arrows in C . The 0-cells in particular may be identified with objects of C . An operator $f : [m] \rightarrow [n]$, or in Charles' notation $\langle f_1, \dots, f_m \rangle$, acts by taking n -strings of composable arrows and "collapsing edges" by composing the arrows, those collapsed edges being determined by the f_i . (At least that's how I try to think about it. I think that's correct. UPDATE: Yes this is correct, see Charles' notes, Proposition 4.4.) For instance, $\langle 0, 2 \rangle^*$ takes a pair of composable arrows $(f, g) \in (NC)_2$ and sends them to their composite $gf \in (NC)_1$. See also Charles notes, p. 13.

Now we ask a nascent question: can we characterize nerves of categories?

Proposition 0.2.2. Let X be a simplicial set. For $n \geq 2$, consider the function

$$\phi_n : X_n \rightarrow \{(g_i) \in (X_1)^n : g_j \langle 1 \rangle = g_{j+1} \langle 0 \rangle \text{ for all } i\}$$

(The latter set being the collection of " n -paths" of 1-cells in X) Which acts by $a \mapsto (a \langle 0, 1 \rangle, \dots, a \langle n-1, n \rangle)$. These ϕ_n are bijections for all $n \geq 2$ if and only if X is the nerve of a category.

Maybe a lazy way to digest this is that "a simplicial set is the nerve of a category iff, thinking of n -cells as length n strings of arrows, their 1-dimensional structure exactly reflects the structure that should arise from the existence and uniqueness of composites."

Proposition 0.2.3. The nerve functor $N : \text{Cat} \rightarrow \text{sSet}$ is fully-faithful. That is, morphisms of nerves $NC \rightarrow ND$ correspond exactly to functors $C \rightarrow D$.

0.2.2 (1/26) Spines

We characterized nerves as those simplicial sets whose n -cells were exactly determined by the collection of length n strings of "composable" 1-cells, in the obvious way. This captures the existence and uniqueness of composites for morphisms in a category. We can go about this characterization a bit more systematically.

Definition 0.2.4. Let $n \geq 0$. The *spine* of the standard n -simplex Δ^n is the simplicial subset defined by

$$(\text{Spine}^n)_k := \{\langle f_0, \dots, f_k \rangle : [k] \rightarrow [n] : f_k \leq f_0 + 1\} \subseteq \Delta_k^n.$$

Informally, the spine is the set of vertices of Δ^n together with the arrows between adjacent vertices (considered with their total ordering).

Proposition 0.2.5. Let X be a simplicial set. For every $n \geq 0$, the map

$$\text{Hom}(\text{Spine}^n, X) \rightarrow \{(a_i) \in (X_1)^n : a_i \langle 1 \rangle = a_{i+1} \langle 0 \rangle\} \quad (6)$$

Given by $f \mapsto (f \langle 0, 1 \rangle, \dots, f \langle n-1, n \rangle)$ is a bijection.

Pictorially, I think this is obvious. Here's a clean proof.

Proof. One point we need: we previously talked about colimits in \mathbf{sSet} . Or at least I intended to. Here's the main fact: *a colimit of simplicial sets X_α exists and has as its n -cells the colimit of the n -cells of the X_α .* This is true for presheaves in general; we say their (co)limits are "computed objectwise."

Another point we need: here's a definition. Suppose S is a totally ordered set. We denote by Δ^S the simplicial set having $(\Delta^S)_n := \{\text{order-preserving maps } [n] \rightarrow S\}$. If S is finite and nonempty, there is a unique isomorphism $\Delta^{|S|-1} \cong \Delta^S$. In the case that $S \subseteq [n]$, this is a good way to notate subcomplexes of Δ^n .

Here's a fact I won't prove: *given a subcomplex $K \subseteq \Delta^n$, writing A for the poset of $S \subseteq [n]$ such that $\Delta^S \subseteq K$, the canonical map $\text{colim}_{S \in A} \Delta^S \rightarrow K$ is an isomorphism.*

Finally, our proposition: in the case that $K = \text{Spine}^n$, the poset A consists of sets of the form $\{j\}$ and $\{j+1\}$, and we have that $\text{colim}_{S \in A} \Delta^S \cong \text{Spine}^n$. Now:

$$\text{Hom}(\text{Spine}^n, X) \cong \text{Hom}(\text{colim}_{S \in A} \Delta^S, X) \cong \lim_A \text{Hom}(\Delta^S, X).$$

See that the latter set is precisely the RHS of (6). □

Maybe the key observation is that Δ^n is "generated" precisely by the arrows of Spine^n . (Make this formal? Say this better? Well, this is how I think about it.)

Proposition 0.2.7. *A simplicial set X is the nerve of some category if and only if for each $n \geq 2$, every morphism $f : \text{Spine}^n \rightarrow X$ extends uniquely along the inclusion $\text{Spine}^n \hookrightarrow \Delta^n$.*

Proof. The unique extension condition is equivalent to bijectivity of the restriction $\text{Hom}(\Delta^n, X) \rightarrow \text{Hom}(\text{Spine}^n, X)$. By Proposition 0.2.5, the latter set is isomorphic to

$$\{(a_i) \in (X_1)^n : a_i \langle 1 \rangle = a_{i+1} \langle 0 \rangle\}.$$

Then the desired result is immediate considering Proposition 0.3.34. □

0.2.3 (1/30) Inner Horns

Recall that for each n and $0 \leq i \leq n$ we defined the i -th horn $\Lambda_i^n \subseteq \Delta^n$ to have as k -cells those cells $f : [k] \rightarrow [n]$ of Δ^n which "miss something other than i ," i.e. those satisfying $f([n]) \cup \{i\} \neq [n]$. If $j \neq 0, n$ then we called Λ_j^n an *inner horn*.

Drawing some pictures of horns and thinking of 1-cells as arrows, we think of inner horns as those collections of arrows that "should be composable." Similar to how we handled spines, we may think to characterize nerves as those simplicial sets whose inner horns have unique extensions. By our analogy comparing inner horns to composable arrows, this unique extension condition is analogous the existence and uniqueness of composites.

Theorem 0.2.8. *A simplicial set X is the nerve of some category if and only if for every $n \geq 2$ and inner horn Λ_j^n , every morphism $\Lambda_j^n \rightarrow X$ extends uniquely along $\Lambda_j^n \hookrightarrow \Delta^n$.*

Proof. Charles gives a full proof on p. 21 of his notes. The "only if" direction is not complicated. For the "if" direction:

- We can construct a category C whose nerve realizes X explicitly. The objects and morphisms are specified by X_0 and X_1 .
- Existence and uniqueness of fillers are necessary for the existence and uniqueness of composites.
- Existence of a filler for Λ_1^3 or Λ_2^3 is necessary for the associative law to hold.
- Then one exhibits an isomorphism $X \rightarrow NC$.

□

0.3 February 2023

0.3.1 (2/4) Quasicategories

A *quasicategory* or ∞ -category is a simplicial set X such that every inner horn $\Lambda_j^n \rightarrow X$ has a filler (i.e. an extension along $\Lambda_j^n \hookrightarrow \Delta^n$.) We have shown that ordinary categories are precisely those quasicategories with *unique* horn extensions.

Definition 0.3.1. Some terminology. Let X, Y be quasicategories.

- *Objects* of $X := X_0$.
- *Morphisms* of $X := X_1$.
- *Identity morphism* of $x \in X_0 := x\langle 0, 0 \rangle$.
- *Products of quasicategories* are just their products as simplicial sets.
- *Coroducts of quasicategories* are just their products as simplicial sets.
- A *morphism of quasicategories* $X \rightarrow Y$ is a map of simplicial sets.
- A *Natural transformation* $f_0 \implies f_1$ of functors $f_0, f_1 : X \rightarrow Y$ is a map of simplicial sets $\phi : X \times \Delta^1 \rightarrow Y$ such that $\phi|_{X \times \{i\}} = f_i$.

It's a fact to be proven that the (co)product of quasicategories is again a quasicategory. Here's more terminology.

Definition 0.3.2. Let X be a simplicial set. Let \sim denote the equivalence relation on the set $\coprod X_n$ of cells of X generated by the relation which identifies a cell a with any other cell of the form af for some simplicial operator f . A *connected component* of X is an equivalence class of \sim . We write $\pi_0 X$ for the set of equivalence classes. A simplicial set is called *connected* if $\pi_0 X$ is a singleton.

Proposition 0.3.3. Let X be a simplicial set and suppose x, y are cells in the same connected component of X , i.e. $x = yf$ for some $f : [m] \rightarrow [n]$. If $F : X \rightarrow Y$ is a map of simplicial sets, then $F(yf) = F(y)Y(Xf)$. The latter is $F(x)$ by hypothesis, thus $F(x) \sim F(y)$. So morphisms $F : X \rightarrow Y$ induce maps $\pi_0 X \rightarrow \pi_0 Y$ on connected components.

Proposition 0.3.4. The induced map $\pi_0(X \times Y) \rightarrow \pi_0(X) \times \pi_0(Y)$ is a bijection.

0.3.2 (2/6) Sub, opposite quasicategories

Definition 0.3.5. Let C be a quasicategory. A subcomplex $C' \subseteq C$ is called a *subcategory* of C if for all $n \geq 2$ and $0 < k < n$, every $f : \Lambda_k^n \rightarrow C$ such that $f(\Lambda_k^n) \subseteq C'$ extends "into C' ," i.e. extends to a map $f : \Delta^n \rightarrow C'$.

It is clear that subcategories of quasicategories are quasicategories.

Definition 0.3.6. Let C be a simplicial set. A simplicial subset $C' \subseteq C$ is called *full* if

- For every cell $\sigma : \Delta^n \rightarrow C$ such that for every $0 \leq i \leq n$ the vertex $\sigma(i) \in C$ belongs in C' , the cell σ belongs in C' .

If C is a quasicategory and C' is a full subcomplex, then it is a quasicategory. In this case we say C' is a *full subcategory*.

Next we can define the opposite of a quasicategory. In ordinary categories, we do this by reversing composition. We can do something similar once we identify an involution on the simplex category Δ .

Definition 0.3.7. Define an involution functor $\text{op} : \Delta \rightarrow \Delta$ as follows.

- It acts as the identity on objects.
- It sends the morphism $\langle f_0, \dots, f_m \rangle : [m] \rightarrow [n]$ to its "reverse" $\langle n - f_m, \dots, n - f_0 \rangle : [m] \rightarrow [n]$.

Definition 0.3.8. Let $X : \Delta^{\text{op}} \rightarrow \text{Set}$ be a simplicial set. We define the *opposite simplicial set* as $X^{\text{op}} := X \circ \text{op}$.

One sees that $(\Delta_j^n)^{\text{op}} \cong \Delta_{n-j}^n$ and that $(NC)^{\text{op}} = N(C^{\text{op}})$. The former fact ensures that opposites of quasicategories are quasicategories. The latter ensures that this notion of opposites restricts to the usual 1-categorical notion.

0.3.3 (2/7) Examples of ∞ -categories

Example 0.3.1. The nerve NC of a category C is a quasicategory. This is immediate by our characterization of nerves (Theorem 0.2.8).

Example 0.3.2. The singular complex $\text{Sing}(X)$ of a space X is a quasicategory. In fact, we can say a little bit more. Denote by $(\Lambda_j^n)_{\text{top}}$ the *topological horn*, defined as you might expect:

$$(\Lambda_j^n)_{\text{top}} := \{t \in \Delta^n : t_i = 0 \text{ for some } i \in [n] \setminus \{j\}\}.$$

It is clear that for any j , the simplex Δ^n_{top} retracts onto $(\Lambda_j^n)_{\text{top}}$. Thus by precomposing with the retract we get, for every j , an inverse to the restriction

$$\text{Hom}(\Delta^n, \text{Sing}X) \rightarrow \text{Hom}(\Lambda_j^n, \text{Sing}X).$$

In other words, we can fill every $\Lambda_j^n \rightarrow \text{Sing}X$ via the retract. This shows that $\text{Sing}(X)$ is a quasicategory. In fact, we've shown all horns fill, not just inner horns. We call such simplicial sets *Kan complexes*.

Example 0.3.3. Let A be an abelian group and $d \geq 0$ an integer. In spaces, the *Eilenberg-MacLane spaces* $K(A, d)$ represent $H^d(-; A)$. We will define an analogous simplicial set $K = K(A, d)$, the *Eilenberg-MacLane objects* in \mathbf{sSet} , like so.

- An element of K_n is a collection $(a_\delta \in A)_\delta$, where the index δ occurs over all operators $\delta : [d] \rightarrow [n]$, so that
 - If δ is not injective, $a_\delta = 0$; and
 - For each operator $\gamma : [d+1] \rightarrow [n]$ we have $\sum_{j=0}^{d+1} (-1)^j a_{\delta d j} = 0$.
- For each operator $f : [m] \rightarrow [n]$ and $a \in K_n$, we define $(af)_\delta := a_{f\delta}$.

These $K(A, d)$'s are ∞ -categories. In fact, they are simplicial abelian groups, which are always Kan complexes. In \mathbf{sSet} , they represent *normalized d -cocycles with values in A* . (See Charles' notes, p. 29.)

Example 0.3.4. There is a simplicial set of ordinary categories, denoted \mathbf{Cat}_1 . We define it like so.

- Each n -cell $(\mathbf{Cat}_1)_n$ is the data of $(C_i, F_{ij}, \zeta_{ijk})$ where
 - For each $i \in [n]$, C_i is a category,
 - For each $i \leq j$ in $[n]$, $F_{ij} : C_i \rightarrow C_j$ is a functor, and
 - For each $i \leq j \leq k$ in $[n]$, $\zeta_{ijk} : F_{jk}F_{ij} \rightarrow F_{ik}$ is a natural isomorphism,
 - And furthermore, these data are subject to certain basic properties (e.g. $F_{ii} = \text{id}_{C_i}$).
- Each operator $f : [m] \rightarrow [n]$ acts on an n -cell $(C_i, F_{ij}, \zeta_{ijk})$ by composing with the indices.

The simplicial set \mathbf{Cat}_1 is an ∞ -category. Let's discuss fillers.

- A 2-horn $\Lambda_1^2 \rightarrow \mathbf{Cat}_1$ is the data of functors $C_0 \xrightarrow{F_{01}} C_1 \xrightarrow{F_{12}} C_2$. An extension is the data of a functor $F_{02} : C_0 \rightarrow C_2$ and a natural isomorphism $\zeta_{012} : F_{12}F_{01} \Rightarrow F_{02}$. An obvious *but not necessarily unique* candidate is $F_{02} := F_{12}F_{01}$.
- The data of a 3-horn $\Lambda_1^3 \rightarrow \mathbf{Cat}_1$ is a bit of a picture. A filler amounts to finding a natural isomorphism to fill the following diagram.

$$\begin{array}{ccc} F_{03} & \xrightarrow{\zeta_{013}} & F_{13}F_{01} \\ \downarrow & & \downarrow \zeta_{123}F_{01} \\ F_{23}F_{02} & \xrightarrow{F_{23}\zeta_{012}} & F_{23}F_{12}F_{01} \end{array}$$

We can always find this and it is unique, since we required the ζ 's to be natural *isomorphisms*.

0.3.4 (2/8) The fundamental category of a simplicial set

The fundamental groupoid $\pi_{\leq 1}X$ of a space X can be recovered from its singular complex $\text{Sing}(X)$. We will recast this construction $\pi_{\leq 1}X$ for any ∞ -category. The result will no longer be a groupoid in general (it will only be so for Kan complexes, I think). Let's see how far we get.

First we will look at a certain construction for all simplicial sets. By its definition, it's essentially a left adjoint to the nerve functor.

Definition 0.3.9. Let X be a simplicial set. A *fundamental category of X* is a category hX and a map $\alpha : X \rightarrow N(hX)$ such that for every nerve NC , the restriction

$$\alpha^* : \text{Hom}(N(hX), NC) \rightarrow \text{Hom}(X, NC)$$

Is a bijection. This characterizes hX up to unique isomorphism, if it exists. (It always does.)

Proposition 0.3.10. Every simplicial set has a fundamental category.

Proof. Charles sketches this on p. 30. The objects of our category are X_0 . The morphisms are (those generated by) the edges X_1 , where we identify composites according to the 2-cells of X . So we turn X into a category in the most obvious way, flattening the higher-categorical structure in the process. The map $\alpha : X \rightarrow N(hX)$ is the one you'd expect. \square

Proposition 0.3.11. The fundamental category describes a functor $h : \text{sSet} \rightarrow \text{Cat}$, and this functor is left adjoint to the nerve functor N .

0.3.5 (2/9) Homotopy for ∞ -categories

Now we start down a long, dark path. Neither of these adjectives matter up-to-homotopy, however.

Definition 0.3.12. Let C denote an ∞ -category and let $f, g : x \rightarrow y$ be two morphisms between objects x, y in C . A *homotopy from f to g* is a 2-cell $a \in C_2$ such that $a_{01} = f$, $a_{12} = \text{id}_y$, and $a_{02} = g$.

Proposition 0.3.13. The homotopy relation is an equivalence relation on $\text{Hom}_C(x, y)$, i.e. the set of edges i with $i_0 = x$ and $i_1 = y$. Thus we may unambiguously say maps are (or are not) *homotopic* and speak of *homotopy classes*.

Remark 0.3.14. The existence of inner horn extensions is necessary for this relation to be symmetric and transitive. So quasicategories stand out amongst simplicial sets as those having a good notion of homotopy.

Proposition 0.3.15. Maps f, g are homotopic in C if and only if they are homotopic in C^{op} .

It's maybe a little weird that " f homotopic to g " is a slightly asymmetric definition, in that even if f is homotopic to g , the data of a homotopy does not suffice to get a homotopy from f to g in C^{op} . I don't think this matters much, in light of the previous proposition. Lurie also gives an alternate, symmetric notion of homotopy to address this point [Lur22, Tag 00V0].

Suppose as given $f \in \text{Hom}_C(x, y)$ and $g \in \text{Hom}_C(y, z)$. We say an edge $h \in \text{Hom}_C(x, z)$ is a *composite* of (g, f) if there exists a 2-cell a such that (what you expect).

Proposition 0.3.16. Composition respects the homotopy relation on morphisms. Thus, composites are unique up to homotopy.

Definition 0.3.17. Let C be an ∞ -category. Its *homotopy category hC* is the category having as objects C_0 and as morphisms the homotopy classes of morphisms of C .

Now we have defined the *fundamental category of a simplicial set* and the *homotopy category of an ∞ -category*. The fundamental category is supposed to be the homotopy category to some extent, so we should compare these two notions where they are both defined (∞ -categories).

Definition 0.3.18. Let C be an ∞ -category. There is a natural map $\pi : C \rightarrow N(hC)$ that "passes to homotopy." It acts like so.

- An object is sent to itself (note $C_0 = (hC)_0 = N(hC)_0$).

- A morphism f is sent to its homotopy class.
- An n -cell $a \in C_n$ is sent to the unique $\pi(a) \in N(hC)_n$ that satisfies $\pi(a)_{i-1,i} = \pi(a_{i-1,i})$ for all i . (See also [Lur22, Construction 004G].)

This map $\pi : C \rightarrow N(hC)$ is compatible with simplicial operators, in the sense that given a cell $a \in C_n$, one has $[a_{01}] \circ \cdots \circ [a_{n-1,n}] = [a_{0n}]$.

Proposition 0.3.19. *If C is an ordinary category, then $f \simeq g$ iff $f = g$. Thus if an ∞ -category C is isomorphic to a nerve of a 1-category, then $\pi : C \rightarrow N(hC)$ is an isomorphism, so it must be isomorphic to the nerve of its homotopy category.*

Proposition 0.3.20 (Universal property of homotopy category). *Let C be an ∞ -category and D a small category. If $f : C \rightarrow ND$ is a map of simplicial sets, then there exists a unique map $g : N(hC) \rightarrow ND$ such that $f = g \circ \pi$.*

Proof. We will construct g . We will do so by constructing a functor $g : hC \rightarrow D$. On objects $c \in ob(hC) = C_0$, we define $g(c) := f(c) \in (ND)_0 = ob(D)$. On morphisms, we define $g([h]) := f(h)$. This is well-defined, for if $h \simeq h'$ exhibited by some $a \in C_2$, then $\phi(a) \in (ND)_2$ exhibits the identity $f(h') = id \circ h$. \square

Corollary 0.3.21. *The homotopy category construction is left adjoint to the nerve functor:*

$$h : \mathbf{qCat} \rightleftarrows \mathbf{Cat} : N$$

(Easy to-do: homotopy category of products.)

0.3.6 (2/12) About composition in ∞ -categories

Let C be an ∞ -category. Let $x, y, z \in C_0$ be objects and let $f \in \text{Hom}_C(x, y)$ and $g \in \text{Hom}_C(y, z)$ be morphisms (i.e. 1-cells starting/ending at their domains/targets.) Last time we defined a *composite of morphisms f, g in C* to be any $h \in \text{Hom}_C(x, z)$ such that there exists a 2-cell $a \in C_2$ such that $a_{01} = f$, $a_{12} = g$, and $a_{02} = h$. Composites exist and are unique up-to-homotopy (thus we can compose homotopy classes), but are not uniquely determined in general. We may ask whether every representative of a homotopy class of a composite can be realized on-the-nose as the extension of its (compositees?) The answer is yes.

Proposition 0.3.22. *If $f : x \rightarrow y, g : y \rightarrow z$, and $h : x \rightarrow z$ are morphisms in an ∞ -category C , then $h \in [g] \circ [f]$ if and only if h is a composite of f with g , i.e. there exists $u \in C_2$ satisfying*

$$u|_{\Delta^{0,1}} = f, \quad u|_{\Delta^{1,2}} = g, \quad u|_{\Delta^{0,2}} = h.$$

The proof is nice. I'd reproduce it here but I don't feel like making that diagram right now.

0.3.7 (2/13) Isomorphisms and inverses in ∞ -categories

Denote by C an ∞ -category and $f : x \rightarrow y$ a morphism in C . We say f is an *isomorphism* or an *equivalence* if $[f]$ is an isomorphism in hC . Unwinding a bit, this is equivalent to the existence of a $g : y \rightarrow x$ such that $[f] \circ [g] = [1_y]$ and $[g] \circ [f] = [1_x]$. The property of being an isomorphism is related to inverses; the following is elementary.

Proposition 0.3.23. *Let $f : x \rightarrow y$ be a morphism in an ∞ -category C . A morphism $g : y \rightarrow x$ is called a *preinverse* to f if $[f] \circ [g] = [id_y]$, and a *postinverse* if $[g] \circ [f] = [id_x]$. If g is both, we call it an *inverse*. TFAE.*

1. f is an isomorphism.
2. f has an inverse.
3. f has a preinverse and a postinverse.
4. f has a preinverse with a preinverse.
5. f has a postinverse with a postinverse.

As for composites, inverses are not generally unique, but are so up-to-homotopy.

Proposition 0.3.24. *If $F : C \rightarrow D$ is a map of quasicategories, then F sends isomorphisms to isomorphisms.*

Proof. Suppose that $f : x \rightarrow y$ is an isomorphism in C . By Proposition 0.3.23, f admits an inverse $g : y \rightarrow x$. By definition, it satisfies $[f] \circ [g] = [\text{id}_y]$, so by Proposition 0.3.22 there exists $u \in C_2$ witnessing $\text{id}_y = f \circ g$. By this, I mean the obvious identities with simplicial operators hold. Morphisms of simplicial sets commute with operators, hence $F(u)$ witnesses $\text{id}_{F(y)} = F(f) \circ F(g)$. This shows that $F(g)$ is a preinverse to $F(f)$. By an identical argument, one sees that $F(g)$ is a postinverse. So $F(g)$ is an inverse and we are done. \square

0.3.8 (2/13) ∞ -groupoids, cores, and Kan complexes

Here are some definitions.

- An ∞ -category C is called a *quasigroupoid* or *∞ -groupoid* if hC is a groupoid, i.e. if every morphism is an isomorphism.
- For an ordinary category C , its *core* C^{core} is the subcategory with the same objects and only the isomorphisms of C .
- For an ∞ -category C , its *core* C^{core} is the simplicial subset consisting of all cells of C whose edges are all isomorphisms.
- Recall that a simplicial set is called a *Kan complex* if all horns (not necessarily inner) have extensions.

Note that $N(C^{\text{core}}) = (NC)^{\text{core}}$, so our terminology is justified. Now here are some facts.

Proposition 0.3.25. *If C is an ∞ -category, then $\pi_0 C^{\text{core}} = \{\text{objects of } C\} / \cong$.*

Proposition 0.3.26. *If C is an ∞ -category, then C^{core} is a subcategory and an ∞ -groupoid. Furthermore, every sub- ∞ -groupoid is contained in C^{core} . In other words, C^{core} is the maximal subcategory which is also an ∞ -groupoid.*

Proposition 0.3.27. *Every Kan complex is an ∞ -groupoid.*

Proof. Suppose that K is a Kan complex and $f : x \rightarrow y$ is a morphism in K . Consider the horn $u : \Lambda_0^2 \rightarrow K$ with $u_{01} = f$ and $u_{02} = \text{id}_x$. Extending this horn gives us $g := u_{12}$ with $[g] \circ [f] = [\text{id}_x]$. Thus, every morphism admits a preinverse, and this turns out to be sufficient for an ∞ -category to be an ∞ -groupoid. \square

Proposition 0.3.28 (Joyal's theorem; harder). *Every ∞ -groupoid is a Kan complex.*

Definition 0.3.29. Since $\text{Sing } X$ is a Kan complex, the proposition allows us to define the *fundamental ∞ -groupoid* of a space X as $\text{Sing } X$.

0.3.9 (2/15) The functor quasicategory

The nerve functor $N : \text{Cat} \rightarrow \text{sSet}$ is fully faithful, so functors of categories correspond to morphisms of their nerves. Maybe this suggests that if we want a “mapping space,” or really a “mapping ∞ -category,” it's 0- and 1-categorical structure should consist of morphisms $X \rightarrow Y$ of simplicial sets and natural transformations between them. Morphisms are the same thing as maps $X \times \Delta^0 \rightarrow Y$. A natural transformation is a suitable map $X \times \Delta^1 \rightarrow Y$. This suggests the following.

Definition 0.3.30. Let X, Y be simplicial sets. Their *function complex* is the simplicial set $\text{Fun}(X, Y)$ with

$$(\text{Fun}(X, Y))_n := \text{Hom}_{\text{sSet}}(\Delta^n \times X, Y).$$

Proposition 0.3.31. *There is a natural bijection*

$$\text{Hom}(X \times Y, Z) \xrightarrow{\sim} \text{Hom}(X, \text{Fun}(Y, Z)).$$

Proposition 0.3.32. *For ordinary C and D , one has $N(\text{Fun}(C, D)) \cong \text{Fun}(NC, ND)$.*

Eventually we will see that *function complexes to an ∞ -category are ∞ -categories*. This is what we want. We can't prove this yet.

0.3.10 (2/16) Lifting properties time — weakly saturated classes

My assessment to become a personal fitness trainer is approaching, so I'm shifting focus to that for a little while. I'll probably be leaving more to-do's than I should be.

Quasicategories are defined in terms of lifting properties. Now we will take some time to generally study lifting properties, which will be useful for studying quasicategories.

Definition 0.3.33. Let C be a category admitting small colimits. A class of morphisms \mathcal{A} is called a *weakly saturated class* if it

1. Contains all isomorphisms,
2. Is closed under cobase change (also called pushouts), composition, transfinite composition, coproducts, and retracts. (See Charles p. 38 for the definitions.)

Given any class of morphisms S , its *weak saturation* \bar{S} is the smallest weakly saturated class containing S .

Example 0.3.5. Take $C = \text{Set}$. The weak saturation of $\{\{0, 1\} \rightarrow \{1\}\}$ is the class of surjective maps. The weak saturation of $\{\emptyset \rightarrow \{1\}\}$ is the class of injective maps.

Example 0.3.6. Take $C = \text{Set}$. The surjections/injections also arise as weak saturations. Of what?

Proposition 0.3.34. Let S be a category with small colimits and let C be a class of objects. Let \mathcal{A} be the class of maps with the following lifting property: if $i : A \rightarrow B$ is in \mathcal{A} , then for every $f : A \rightarrow C$ to an object of C , we can fill the following diagram:

$$\begin{array}{ccc} A & \xrightarrow{f} & C \\ \downarrow i & \nearrow & \\ B & & \end{array}$$

Then \mathcal{A} is a weakly saturated class. (Example: $S = \text{sSet}$, $C = \{\infty\text{-categories}\}$).

Proof. To-do. (Worked out in meeting.) □

0.3.11 (2/16) Classes of horns, anodyne morphisms

As indicated, we are interested in ∞ -categories, so we ought to study lifting properties of horns in particular. We make some definitions for this.

Definition 0.3.35. We define the following sets of horns.

$$\begin{aligned} \text{InnHorn} &:= \{\Lambda_k^n \hookrightarrow \Delta^n : 0 < k < n, n \geq 2\}, \\ \text{Horn} &:= \{\Lambda_k^n \hookrightarrow \Delta^n : 0 \leq k \leq n, n \geq 1\}, \\ \text{RHorn} &:= \{\Lambda_k^n \hookrightarrow \Delta^n : 0 < k \leq n, n \geq 1\}, \\ \text{LHorn} &:= \{\Lambda_k^n \hookrightarrow \Delta^n : 0 \leq k < n, n \geq 1\}. \end{aligned}$$

We call their weak saturations in sSet the *(inner, right, left) anodyne morphisms*.

Proposition 0.3.36. Monomorphisms of simplicial sets form a weakly saturated class. Therefore, since Horn consists of monomorphisms, its weak saturation must too. So (inner, right, left) anodyne maps are always monomorphisms.

Proposition 0.3.37. Let C be an ∞ -category. If $A \hookrightarrow B$ is an inner anodyne inclusion, then every $f : A \rightarrow C$ extends to B .

Proof. Let \mathcal{A} denote the set of maps of simplicial sets $X \rightarrow Y$ which extend along every map $X \rightarrow C$. (C is fixed here.) Since C is a quasicategory, we have $\text{InnHorn} \subseteq \mathcal{A}$. By Proposition 0.3.34, the class \mathcal{A} is weakly saturated, so $\overline{\text{InnHorn}} \subseteq \mathcal{A}$. That is what we wanted to show. □

Proposition 0.3.38. (To-do: Prop 16.10.)

Example 0.3.7. Here are some examples of inner anodyne morphisms. (Important to-do: finish.)

- The inclusions of spines $I^n \hookrightarrow \Delta^n$ are inner anodyne for every n . In particular, if C is an ∞ -category, every map $I^n \rightarrow C$ extends to a $\Delta^n \rightarrow C$.
- \vdots

0.3.12 (2/16) Lifting calculus

Here's a definition I expected a bit earlier in Charles' notes.

Definition 0.3.39. Say an object X *satisfies the extension property* for $f : A \rightarrow B$ if for every $u : A \rightarrow X$ we can find an extension $B \rightarrow X$.

Definition 0.3.40. Suppose as given maps $f : A \rightarrow B$ and $g : X \rightarrow Y$. A *lifting problem* for (f, g) is a pair of maps $u : A \rightarrow X$ and $v : B \rightarrow Y$ making a commutative square. A *lift* for the lifting problem is a fill s to the obvious diagram:

$$\begin{array}{ccc} A & \xrightarrow{u} & X \\ f \downarrow & \nearrow s & \downarrow g \\ B & \xrightarrow{v} & Y \end{array}$$

Definition 0.3.41. Let f, g be morphisms in a category. We write $f \sqsubset g$ if every lifting problem for (f, g) admits a lift. We call this the *lifting relation* on morphisms. If $f \sqsubset g$, we say:

- f has the left lifting property rel. to g , or
- g has the right lifting property rel. to f , or
- f lifts against g .

Definition 0.3.42. Let A be a class of morphisms. We define the *right complement* $A^\sqsupset := \{g : a \sqsubset g, \forall a \in A\}$. We define the *left complement* ${}^\sqsubset A$ similarly.

Proposition 0.3.43. Let \mathcal{A} be any class of morphisms in a category with small colimits. The left complement ${}^\sqsubset \mathcal{A}$ is weakly saturated, and the right complement \mathcal{A}^\sqsupset is weakly cosaturated.

Proof. (To-do: prove. Also, Charles' related exercises.) □

Example 0.3.8. Let C be an abelian category, let $\mathcal{P} = \{0 \rightarrow P : P \text{ projective}\}$, and let \mathcal{B} be the class of epimorphisms in C . By the definition of projective objects, we have $\mathcal{P} \sqsubset \mathcal{B}$. Thus $\mathcal{B} \subseteq \mathcal{P}^\sqsupset$.
(To-do: show converse?)

0.3.13 (2/17) Inner fibrations

A map p of simplicial sets is called an *inner fibration* if $\text{InnHorn} \sqsubset p$. Thus, $\text{InnFib} = \text{InnHorn}^\sqsupset$.

Proposition 0.3.44. A simplicial set C is an ∞ -category iff $C \rightarrow *$ is an inner fibration.

Proposition 0.3.45. InnFib is defined as a right complement, thus InnFib is weakly cosaturated. This implies, for instance, that if C is an ∞ -category and $D \rightarrow C$ is an inner fibration, then D is an ∞ -category.

Proposition 0.3.46 (. kerodon] If X is a simplicial set, then a morphism $X \rightarrow ND$ is an inner fibration iff X is an ∞ -category.

Proof. (To-do.) □

0.3.14 (2/20) Factorizations


Recall that we defined *inner fibrations* InnFib as the right complement of InnHorn . This tells us something—maybe this is why we call it a “complement.”

Proposition 0.3.47 (Small object argument). *Let S be a set of morphisms in \mathbf{sSet} . Then every map f of simplicial sets admits a factorization $f = p \circ j$ with $p \in S^\square$ and $j \in \bar{S}$.*


Corollary 0.3.48. *If S is any set of morphisms in \mathbf{sSet} , then $\bar{S} = {}^\square(S^\square)$.*

Proof. Since ${}^\square(S^\square)$ is a left complement, it is weakly saturated (Proposition 0.3.43), thus $\bar{S} \subseteq {}^\square(S^\square)$.

Now suppose that $f \in S^\square$. By the previous proposition, we may write $f = pj$ for $p \in S^\square$ and $j \in \bar{S}$, and by assumption f admits a lift in the following diagram.



Thus, we get the following commutative diagram.



This exhibits f as a retract of j . Since $j \in \bar{S}$ and weak saturations are closed under retracts, we have $f \in \bar{S}$. \square

Corollary 0.3.49. *Every map f of simplicial sets can be factored $f = pj$ with p an inner fibration and j inner anodyne.*

0.3.15 (2/21) Factorization systems and unique lifts

Last time, we proved that for a class S of maps in \mathbf{sSet} , we have $\bar{S} = {}^\square(S^\square)$ and we can factor every map as a composite of one map from \bar{S} and S^\square . Starting with $S = \text{InnHorn}$, we got a factorization of an arbitrary map as an inner anodyne followed by an inner fibration. We study such systems in general.


Definition 0.3.50. A *weak factorization system* in a category is a pair of classes of maps $(\mathcal{L}, \mathcal{R})$ with the following properties.

1. Every morphism factors as rl for $l \in \mathcal{L}$ and $r \in \mathcal{R}$; and
2. $\mathcal{L} = {}^\square \mathcal{R}$ and $\mathcal{R} = \mathcal{L}^\square$.

Example 0.3.9. The pair $(\overline{\text{InnHorn}}, \text{InnHorn}^\square)$ is a weak factorization system.

We would like to understand lifting problems with unique solutions.

Definition 0.3.51. In a category with coproducts, let $f : A \rightarrow B$ be a morphism. We define the *fold* of f , denoted $f^\wedge : B \amalg_A B \rightarrow B$, as the unique map making the following diagram commute.



Proposition 0.3.52. *Let f, g be morphisms. The following are equivalent.*

1. *We have $f \sqsubset g$ and $f^\vee \sqsubset g$.*
2. *The solution to any lifting problem for (f, g) exists and is unique.*

Proof. (2) \implies (1) is obvious. For (1) \implies (2), existence is assumed, so we need only show uniqueness. Wait, what does uniqueness mean in an arbitrary category? \square

Definition 0.3.53. A weak factorization system $(\mathcal{L}, \mathcal{R})$ is called an *orthogonal factorization system* if $\mathcal{L} = {}^\square \mathcal{R}$ and $\mathcal{R} = \mathcal{L}^\square$ are realized by *unique* lifts.

Proposition 0.3.54. *The factorization $f = rl$ is unique up to unique isomorphism in an orthogonal factorization system.*

Proposition 0.3.55. $(\{\text{surjections}\}, \{\text{injections}\})$ form an orthogonal factorization system in Set . (Proof is obvious.)

Proposition 0.3.56. *For any class of simplicial maps S , the pair $(\overline{S \cup S^\vee}, (S \cup S^\vee)^\square)$ is an orthogonal system.*

0.3.16 (2/23) Degenerate cells

We want to concretely understand monomorphisms of simplicial sets. For this, recall that we defined the *boundary* of Δ^n as the subcomplex of Δ^n whose k -cells are the non-surjective maps $[k] \rightarrow [n]$. Write Cell for the class of inclusions $\partial\Delta^n \hookrightarrow \Delta^n$ and $\text{InnFib} := \text{Cell}^\square$. Since Cell consists of monos, we know $\overline{\text{Cell}}$ contains all monomorphisms. Our main theorem is the converse.

Proposition 0.3.57. *The class $\overline{\text{Cell}}$ is exactly the class of monomorphisms of simplicial sets.*

We'll prove this (Proof 0.3.17) once we've set some stuff up.

Toward proving this, recall the notion of degenerate cells: a cell $\sigma : \Delta^n \rightarrow X$ is called *degenerate* if there exists a non-injective operator $f : [m] \rightarrow [n]$ such that $\sigma = \tau f$. Since every simplicial operator factors uniquely as $f = f^{\text{inj}} f^{\text{surj}}$, we see that σ is degenerate if and only if there is some non-identity surjective f such that $\sigma = \tau f$. A cell which is not degenerate is called *non-degenerate*. We write $X_n = X_n^{\text{deg}} \coprod X_n^{\text{nd}}$ for the decomposition of X_n into (non)-degenerate cells. Neither assemble to a subcomplex.

Proposition 0.3.58. *Here are some straightforward facts about degenerate cells.*

1. *If $f : X \rightarrow Y$ is a map of simplicial sets, then $f(X_n^{\text{deg}}) \subseteq Y_n^{\text{deg}}$.*
2. *If $f : X \rightarrow Y$ is a map of simplicial sets, then $f^{-1}(Y_n^{\text{nd}}) \subseteq X_n^{\text{nd}}$.*
3. *If $A \hookrightarrow X$ is a subcomplex, then*

$$\begin{aligned} A_n^{\text{nd}} &= X_n^{\text{nd}} \cap A_n, \text{ and} \\ A_n^{\text{deg}} &= X_n^{\text{deg}} \cap A_n. \end{aligned}$$

4. *The elements of $(\Delta^n)_k^{\text{nd}}$ are in bijection with the subsets of $[n]$ of size k .*
5. *The *simplicial n -sphere* $\Delta^n / \partial\Delta^n$, defined as the pushout of $\Delta^n \leftarrow \partial\Delta^n \rightarrow \Delta^0$, has exactly two nondegenerate cells: its unique vertex and the generator $\langle 0, 1, \dots, n \rangle$. In other words, the image of $\Delta^0 \rightarrow \Delta^n / \partial\Delta^n$ and of the generator in $\Delta^n \rightarrow \Delta^n / \partial\Delta^n$ in the pushout square:*

$$\begin{array}{ccc} \partial\Delta^n & \xrightarrow{\quad} & \Delta^n \\ \downarrow & & \downarrow \\ \Delta^0 & \xrightarrow{\quad} & \Delta^n / \partial\Delta^n \end{array}$$

The [Eilenberg-Zilber lemma](#) says that every cell a of X occurs *uniquely* as $a = b\sigma$ for a nondegenerate b and surjective operator σ . (This is not too complicated; the nontrivial part is uniqueness.) Let's state this in a slightly different, stronger form.

Proposition 0.3.59. *If X is a simplicial set, then for every n the map*

$$\coprod_{j \geq 0} X_j^{nd} \times \text{Hom}_{\Delta_{surj}}([n], [j]) \rightarrow X_n$$

Given by $(j, a, \sigma) \mapsto a\sigma$ is a bijection. Furthermore, this map is natural with respect to surjective operators $[n'] \rightarrow [n]$ and with respect to monomorphisms of simplicial sets $X \rightarrow X'$.

There are various things to be said now. I think I will move on, and refer back to these things when they are needed.

0.3.17 (2/23) The skeletal filtration

If $\sigma : \Delta^n \rightarrow X$ is an n -cell, it uniquely factors as $\Delta^n \rightarrow \Delta^m \rightarrow X$ where the first map is surjective and the second is nondegenerate. So σ is “really” an m -cell, for some $m \leq n$. Now, we want a notion of the k -skeleton of X . Its n -cells should be the n -cells of X which are “really” j -cells for some $j \leq k$.

Definition 0.3.60. Let X be a simplicial set. The k -skeleton of X , written $Sk_k X$, is the smallest subcomplex containing all cells of dimension $\leq k$. Thus, we have

$$(Sk_k X)_n = \bigcup_{0 \leq j \leq k} \{yf : y \in X_j \text{ and } f : [n] \rightarrow [j]\}.$$

A nondegenerate cell $\Delta^k \rightarrow X$ determines a cell $\Delta^k \rightarrow Sk_k X$. This map carries $\partial\Delta^{k-1}$ to $Sk_{k-1} X$.

Proposition 0.3.61. *The evident square*

$$\begin{array}{ccc} \coprod_{i \in X_k^{nd}} \partial\Delta^k & \longrightarrow & Sk_{k-1} X \\ \downarrow & & \downarrow \\ \coprod_{i \in X_k^{nd}} \Delta^k & \longrightarrow & Sk_k X \end{array}$$

Is a pushout square. More generally, if $A \subseteq X$ is a subcomplex, the following is a pushout square.

$$\begin{array}{ccc} \coprod_{i \in X_k^{nd}/A_k^{nd}} \partial\Delta^k & \longrightarrow & A \cup Sk_{k-1} X \\ \downarrow & & \downarrow \\ \coprod_{i \in X_k^{nd}/A_k^{nd}} \Delta^k & \longrightarrow & A \cup Sk_k X \end{array}$$

It is in this sense that simplicial sets are built out of standard simplices: a simplicial set X is filtered by $X_0 = Sk_0 X \subseteq Sk_1 X \subseteq Sk_2 X \subseteq \dots$, and each Sk_n is obtained from Sk_{n-1} by attaching copies of Δ^n as in Proposition 0.3.61.

Now we are ready to prove our characterization of monomorphisms (Proposition 0.3.57).

Proof. A monomorphism of simplicial sets is isomorphic to an inclusion $A \hookrightarrow X$. It is clear that $X \cong \text{colim}_{\rightarrow k} A \cup Sk_k X$. But see that, by the above proposition, the maps $A \cup Sk_{k-1} X \rightarrow A \cup Sk_k X$ arise via cobase change from coproducts of maps in Cell . Then the inclusion is exhibited as a countable composition(?) of maps in $\overline{\text{Cell}}$, thus is in $\overline{\text{Cell}}$. \square

And so we have some handle on monomorphisms in sSet now.

Corollary 0.3.62 (Geometric realizations are CW). Recall that we constructed the geometric realization functor $|-| : \mathbf{sSet} \rightarrow \mathbf{Top}$ as a left adjoint. Left adjoints preserve colimits, hence we have a pushout diagram

$$\begin{array}{ccc} \coprod_{a \in X_k^{nd}} \partial \Delta_{top}^k & \longrightarrow & |Sk_{k-1}X| \\ \downarrow & & \downarrow \\ \coprod_{a \in X_k^{nd}} \Delta_{top}^k & \longrightarrow & |Sk_kX| \end{array}$$

Additionally, we have $|X| = \lim_{\rightarrow} |Sk_kX|$. This describes a canonical CW structure on the geometric realization $|X|$ of a simplicial set. Evidently, cells of $|X|$ correspond to nondegenerate simplices of X .

0.3.18 (2/25) Pushout-products, pullback-homs

Let $f : A \rightarrow B$ and $g : K \rightarrow L$ be morphisms in \mathbf{sSet} . We define the *pushout-product* of f and g , denoted $f \square g$, as the unique dotted map making the following pushout square diagram commute.

$$\begin{array}{ccc} A \times K & \xrightarrow{\text{id} \times g} & A \times L \\ \downarrow f \times \text{id} & & \downarrow f \times \text{id} \\ B \times K & \xrightarrow{\quad \quad} & (B \times K) \amalg_{A \times K} (A \times L) \\ & \searrow \text{id} \times g & \nearrow f \times \text{id} \\ & & B \times L \end{array}$$

$f \square g$ (dotted arrow from $B \times K$ to $B \times L$)

Dually, we define the *pullback-hom* of f and g , denoted $f \square^* g$, to be the unique dotted map making the following pullback square diagram commute.

$$\begin{array}{ccccc} \text{Fun}(L, A) & & \xrightarrow{\text{Fun}(g, \text{id})} & & \text{Fun}(K, A) \\ & \searrow f \square^* g & & \nearrow \text{Fun}(\text{id}, f) & \\ & \text{Fun}(K, A) \times_{\text{Fun}(K, B)} \text{Fun}(L, B) & \xrightarrow{\quad \quad} & & \text{Fun}(K, A) \\ & \downarrow \text{Fun}(\text{id}, f) & & \downarrow \text{Fun}(\text{id}, f) & \\ & \text{Fun}(L, B) & \xrightarrow{\text{Fun}(g, \text{id})} & & \text{Fun}(K, B) \end{array}$$

0.4 March 2023

0.4.1 (3/5) Pullback-hom as an enriched lifting problem

Suppose given $g : K \rightarrow L$ and $h : X \rightarrow Y$. We have the pullback-hom

$$h^{\square g} : \text{Fun}(L, X) \rightarrow \text{Fun}(K, X) \times_{\text{Fun}(K, Y)} \text{Fun}(L, Y).$$

The vertices of $\text{Fun}(L, X)$ are morphisms $L \rightarrow X$ in \mathbf{sSet} . The vertices of $\text{Fun}(K, X) \times_{\text{Fun}(K, Y)} \text{Fun}(L, Y)$ are those pairs of morphisms $(s : K \rightarrow X, t : L \rightarrow Y)$ such that $hs = tg$, i.e. lifting problems for (g, h) . The pullback-hom $h^{\square g}$ takes a morphism $w : L \rightarrow X$ and composes it to (wg, gh) . This gives us a lifting problem for (g, h) that is solvable. Then the following is clear.

Proposition 0.4.1. *The pullback-hom $h^{\square g}$ is surjective on vertices iff $g \sqsupseteq h$.*

In this sense, $h^{\square g}$ encodes an “enriched” lifting problem for (g, h) . The target $\text{Fun}(K, X) \times_{\text{Fun}(K, Y)} \text{Fun}(L, Y)$ parametrizes lifting problems for (g, h) while the source $\text{Fun}(L, X)$ parametrizes families of lifting problems together with a chosen lift.

Also, let’s talk about so-called *adjunctions of lifting problems*. The product and function complex constructions are adjoint. Ultimately, this leads to the following.

Proposition 0.4.2. *One has $(f \square g) \sqsupseteq h$ if and only if $f \sqsupseteq h^{\square g}$.*

Here’s a special case. Take $K = \emptyset$ and $Y = *$. Then the proposition gives us that

$$(f \times \text{id}_L) \sqsupseteq (X \rightarrow *) \iff f \sqsupseteq (\text{Fun}(L, X) \rightarrow *).$$

0.5 April 2023

0.6 May 2023

0.6.1 (5/3) Operads for (Peter) May

For May, I will learn about operads and monads. Here are my motives.

- (1) Peter May coined the term *operad*.
- (2) They're interesting and fit into the ∞ -categorical framework, eventually.
- (3) I read some of Moerdijk-Weiss's *Dendroidal sets* for Charles' Kan Seminar and thought it was exciting.
- (4) Peter May tipped me off that monads and operads would make a big appearance in his talks for the 2023 UChicago REU. (Probably related to his recent work with Ruoyi Zhang and Foling Zou.)

Here are some potential references.

- (1) May, *The Geometry of Iterated Loop Spaces* (1971)
 - (2) Markl-Shnider-Stasheff, *Operads in Algebra, Topology, and Physics* (2000)
 - (3) Heuts, *Simplicial and Dendroidal Homotopy Theory* (2022)
 - (4) Markl, *Operads and PROPs* (2006)
 - (5) Lawson, *E_n -ring spectra and Dyer-Lashof operations*
-

Let me say what I *think* operads are supposed to do/be before I dive into it: An operad abstracts away the structure of “an operation on a structure its identities/coherences.” We'll see this worked out later, for the first time in Example 0.6.4. Here's a vague indication as to why this is useful: say an algebraic thing X has operations which “play nice” with its algebraic structure. If this occurs, it can have useful consequences. It's natural that we then (1) find and study objects for which this occurs, and (2) study them in concert. But this has problems: (A) it may require *lots* of data to verify or realize that X 's operations “play nice” with its structure (*coherence data*), especially for complicated X , and/or especially if we're thinking up-to-homotopy, and (B) if we want to study such objects relative to each other, we'll have to compare several of these huge packages of data. Operads do the work for us: the idea is to say, “let \mathcal{C} be the operad codifying the possession of coherent operations.” Then, given an object X , a choice (if one exists) of such structure on X amounts to a morphism $\mathcal{C} \rightarrow \mathcal{E}nd_X$, the latter being a canonical “endomorphism operad” associated to X . That morphism essentially says, “we can interpret the structure within \mathcal{C} as some class of operations on X .” In other words, for an algebraic structure X , an operad classifies “coherent” operations on X , the details (e.g., how coherent?) dependent on which operad you're considering.

Maybe another way of putting it is that operads represent the formal algebraic theory, while we're interested in “instantiations” of these theories, i.e. their representations—I think we call these “algebras over the operads.”

OK, let me actually learn what these are now.

Definition 0.6.1. An *operad*, *symmetric operad*, or *classical operad* \mathcal{C} is a collection of sets $(\mathcal{C}(i))_{i \geq 0}$ with a distinguished operation $1_{\mathcal{C}} \in \mathcal{C}(1)$ and functions $\gamma : \mathcal{C}(n) \times \mathcal{C}(k_1) \times \cdots \times \mathcal{C}(k_n) \rightarrow \mathcal{C}(\sum k_s)$, which we regard as *operations*, the *unit operation*, and as *composition*, respectively, and in this regard we require that these structures are suitably associative, unital, and equivariant up to reordering of inputs.

Concretely, \mathcal{C} is the data of

- (Operations) For each $i \geq 0$, a set $\mathcal{C}(i)$ called *i -ary operations*; and
- (Composites) For each $n \geq 0$ and $k_1, \dots, k_n \geq 0$, a *composition* map $\gamma : \mathcal{C}(n) \times \mathcal{C}(k_1) \times \cdots \times \mathcal{C}(k_n) \rightarrow \mathcal{C}(\sum k_s)$; and
- (Unit) A distinguished *identity* operation $1_{\mathcal{C}} \in \mathcal{C}(1)$; and

- (Symmetries) For each $i \geq 0$, an action $\Sigma_i \rightarrow \text{Aut}\mathcal{C}(i)$

satisfying the following conditions.

- (Unitality) For every operation $d \in \mathcal{C}(j)$, we have $\gamma(1; d) = d$ and $\gamma(d; 1^{\times j}) = d$.
- (Associativity) Blah blah blah
- (Σ_n -Equivariance) For each $\Sigma_1^k j_s = j$, $c \in \mathcal{C}(k)$, $d_s \in \mathcal{C}(j_s)$, $\sigma \in \Sigma_k$, and $\tau_s \in \Sigma_{j_s}$, we have

$$\gamma(c\sigma; (d_i)) = \gamma(c; (d_{\sigma^{-1}i})) \cdot \sigma(j_1, \dots, j_s), \quad \text{and}$$

$$\gamma(c; (d_i\tau_i)) = \gamma(c; (d_i))(\tau_1 \oplus \dots \oplus \tau_k).$$


Here, $\sigma(j_1, \dots, j_s) :=$ the permutation of j letters given by permuting the k blocks of letters determined by the partition $j = \Sigma j_s$ according to σ .

Definition 0.6.2. An operad with no Σ_i -actions or equivariance is called *plain* or *non- Σ* or *non-symmetric*.

Definition 0.6.3. Above, we defined an operad in Set . We can make an analogous definition in any bicomplete symmetric monoidal category $(\mathcal{C}, \otimes, 1)$. In this case, the unit/identity is a distinguished morphism $1 \rightarrow \mathcal{C}(1)$. The symmetries become maps $\Sigma_i \rightarrow \text{Iso}(\mathcal{C}(i), \mathcal{C}(i))$. Such a thing is called an *operad in \mathcal{C}* . This is like “enriching” an operad over a category.

Definition 0.6.4. A *morphism of operads* $\mathcal{C} \rightarrow \mathcal{C}'$ is a collection of maps $f_i : \mathcal{C}(i) \rightarrow \mathcal{C}'(i)$ such that $f_1(1) = 1$ and (equivariance, compatibility with composition).

Remark 0.6.5. As indicated, we think of $\mathcal{C}(i)$ as a set of i -ary operations, and the functions $\gamma : \mathcal{C}(k) \times \mathcal{C}(n_1) \times \dots \times \mathcal{C}(n_k) \rightarrow \mathcal{C}(\sum n_s)$ as taking a k -ary operation c and plugging in k other operations (d_i) . The Σ_n -equivariance demands that if we tamper with the inputs for c then plug in the (d_i) , that is the same as plugging in the (d_i) in a different order then tampering with the order of their inputs. See the little picture.



An operad’s purpose in life is to help define algebras over an operad. Such a thing establishes an “algebraic structure representing the operad” upon an object. Here is the definition.

Definition 0.6.6. Let \mathcal{C} be an operad in a symmetrical monoidal \mathcal{C} . A *\mathcal{C} -algebra* A is an object A and maps $\mathcal{C}(i) \otimes A^{\otimes i} \rightarrow A$ that are suitably associative, unital, and equivariant. (We take $A^{\otimes 0} = 1_{\mathcal{C}}$.)

I’ll go over many examples soon. Some of these will let us reinterpret some of the above structures. But for the rest of today, I’ll just make a little remark.

Remark 0.6.7 (Classical operads generalize monoids). Let \mathcal{C} be a classical operad. (A non- Σ operad works too.) There is an associated category $j_!\mathcal{C}$ with one object and morphisms given by $\mathcal{C}(1)$, the unary operations. Given $f, g \in \mathcal{C}(1)$, their composite $g \circ f :=$ their image in $\mathcal{C}(1) \times \mathcal{C}(1) \rightarrow \mathcal{C}(1)$, and the unit is the identity operation $1_{\mathcal{C}} \in \mathcal{C}(1)$. This checks out thanks to the unitality and associativity axioms.

Conversely, given a one-object category M , i.e. a monoid, we may form an operad j^*M with solely unitary operations, given by $(j^*M)(1) := \text{Hom}_M(*, *)$. The unit and composition functions are obvious, and the Σ_i -actions are trivial.

Altogether we get an *adjunction*

$$\text{Monoids} \begin{array}{c} \xrightarrow{j_!} \\ \xleftarrow{j^*} \end{array} \text{Operads}$$

0.6.2 (5/5) Basic examples of operads

In what follows, let $(C, \otimes, 1)$ denote a symmetric monoidal category.

Example 0.6.1. Let A be an object of C . If C is closed,¹ we denote by End_A the *endomorphism operad* of A , defined by $(\text{End}_A)(i) := \text{Hom}(A^{\otimes i}, A)$. The unit is $\text{id}_A \in \text{End}_A(1)$ and the compositions are given by composing tensor product maps. The right Σ_i -action is given by the left Σ_i -action on tensor powers.

Proposition 0.6.8. Let $A \in C$ and let \mathcal{C} be an operad in C . Via the tensor-Hom adjunction, a \mathcal{C} -algebra structure on A is “the same thing as” a morphism $\mathcal{C} \rightarrow \text{End}_A$.

Example 0.6.2. We denote by Comm the *commutative operad* in Set . It is defined to have a single operation $\text{Comm}(i) := \{*\}$ for every i .

Example 0.6.3. We denote by Assoc the *associative operad* in Set . It is defined to have $\text{Assoc}(i) := \Sigma_i$ for every i . The unit and Σ_i -action are obvious. The maps $\gamma : \Sigma_n \times \Sigma_{k_1} \times \cdots \times \Sigma_{k_n} \rightarrow \Sigma_{k_1 + \cdots + k_n}$ are defined as follows: given $\sigma \in \Sigma_n$ and $\tau_j \in \Sigma_{k_j}$ regarded as matrices, one inserts τ_j in place of the 1 in the j -th column of σ , for each $1 \leq j \leq n$.

Proposition 0.6.9. In Set , Assoc -algebras (resp. Comm -algebras) are precisely monoids (resp. commutative monoids).

In fact, we can encode monoids in the arbitrary C with operad actions. If $(C, \otimes, 1)$ has finite coproducts, for a finite set S let $1[S]$ denote the coproduct $\coprod_S 1$.

Definition 0.6.10. We denote by Comm the *commutative operad* in C . It is defined to have $\text{Comm}(i) := 1$.

Definition 0.6.11. If C has finite coproducts, we denote by Assoc the *associative operad* in C . It is defined to have $\text{Assoc}(i) := 1[\Sigma_i]$. The rest of its structure is mostly obvious.

Proposition 0.6.12. In C , the Comm -algebras are precisely the monoids in C . If C has finite coproducts, then the Assoc -algebras are precisely the commutative monoids in C .

Remark 0.6.13 (Algebras over symmetric vs. plain operads). Above, we are regarding Assoc and Comm as *symmetric* operads, and this is manifest in the structure of algebras over them. We can instead skip any mention of Σ_n ’s and consider Assoc , Comm as *plain* operads. If we do, then Comm -algebras become precisely monoids in C . And Assoc -algebras become...something? Maybe this indicates we should avoid plain operads if possible.

0.6.3 (5/6) Warm-up: monoids are Assoc -algebras

Let me work out a concrete example of how monoids are “the same thing as” Assoc -algebras.

Example 0.6.4 (How do Assoc -algebras encode monoids?). Let X be a set. Suppose it is a monoid, i.e. that we have chosen a unital, associative product $\mu : X \times X \rightarrow X$. As I said yesterday, the monoid (X, μ) is “the same thing as” an Assoc -algebra which is “the same thing as” a choice of morphism $f : \text{Assoc} \rightarrow \text{End}_X$.

I’ll describe the morphism f . Let me write e_n for the identity in Σ_n .

- f sends $\text{Assoc}(1) = \{*\}$ to the identity $\text{id}_X \in \text{End}_X(1)$.
- f sends $e_2 \in \text{Assoc}(2) = \Sigma_2$ to $\mu \in \text{End}_X(2)$.
- Where does f send $\sigma \in \text{Assoc}(2) = \Sigma_2$? Equivariance demands that $f(\sigma) = f(e_2)\sigma$ = “swap inputs then do $f(e_2) = \mu$.”
- Similarly, the value of f on $\text{Assoc}(3)$ is already totally determined by its value $f(e_2) = \mu$. Here’s why. In Assoc , the composition $\gamma : \text{Assoc}(2) \times \text{Assoc}(2) \times \text{Assoc}(1) \rightarrow \text{Assoc}(3)$ satisfies $\gamma(e_2, e_2, e_1) = e_3$. Since f respects composition, it is determined at e_3 :

$$f(e_3) = \gamma(f(e_2), f(e_2), f(e_1)) = \gamma(\mu, \mu, e) = ((a, b, c) \mapsto \mu(\mu(a, b), c)).$$

¹I.e., if it has an internal Hom

Moreover, notice that $\gamma(e_2, e_1, e_2) = e_3$. Then by the same argument, we find that

$$f(e_3) = ((a, b, c) \mapsto \mu(a, \mu(b, c))).$$

So, $(ab)c = a(bc)$, where we're suppressing μ from notation. Associativity! Similarly, we could take into account the Σ_3 -action and show the other associativity identities hold, e.g. $a(cb) = (ac)b$. So, the composition and equivariance conditions gave us associativity (with three inputs)!

- We could repeat the above to see that f 's values on $\mathcal{A}\text{ssoc}(n)$ are forced by $f(e_2) = \mu$. Again, the various equalities $\gamma(e_k, e_{n_1}, \dots, e_{n_k}) = e_{n_1 + \dots + n_k}$, the equivariance equalities, and the fact that f must respect these force all the associativity laws.

Remark 0.6.14. The above example was nice in that after we specified $f(e_2 \in \mathcal{A}\text{ssoc}(2)) = \mu$, we could determine the rest of f based on composition laws, equivariance, and the fact that f must respect those. This is because in some sense, $\mathcal{A}\text{ssoc}$ is “generated by” the element $e_2 \in \mathcal{A}\text{ssoc}(2)$.

So there's an example of how we can use an operad to describe an algebraic operation with coherence. In this case, coherence was strict associativity: we had equalities such as

$$(ab)c = a(bc), a(cb) = (ac)b, \text{ and } a((bc)d) = ((ab)c)d.$$

These equalities were present “formally” in $\mathcal{A}\text{ssoc}$, and since $\mu : X \times X \rightarrow X$ was associative, we could find a corresponding $f : \mathcal{A}\text{ssoc} \rightarrow \text{End}_X$ implementing μ .


That example was overkill. We “just know” what the “coherence” is—it's associativity. The utility of operads arises when the coherences are more complicated. Next time I will look at an example wherein we'll replace the set X with a space, μ with a continuous map, and strict associativity with “associativity up to specified homotopy.” In this case, we'll have e.g. $(ab)c \simeq a(bc)$ and the data of a homotopy realizing this equivalence.

0.6.4 (5/9) Associativity up to homotopy, Stasheff associahedra, and A_∞ -operads

Let Y denote a based space. If $X = \Omega Y$, then X has a multiplication (loop concatenation). Let's parameterize it like so: for $x, y \in X$, define $xy : [0, 1] \rightarrow Y$ as the loop “do y over the first half of the interval, then x over the other.”

Generally, we have $(ab)c \neq a(bc)$, thus X is not an $\mathcal{A}\text{ssoc}$ -algebra. Loop concatenation is not “strictly associative.” But clearly $(ab)c \simeq a(bc)$, realized by (say) a linear reparametrization of $[0, 1]$. That would be a map $[0, 1] \times I \rightarrow X$ starting at $a(bc)$ and landing at $(ab)c$. In $\text{Fun}(X^3, X)$, in which $(ab)c$ and $a(bc)$ reside,² such a homotopy is a path from $(ab)c$ to $a(bc)$.

Now say we're concatenating four loops. There are five ways to do this (by reordering parentheses). Say we want a *choice* of homotopy between each reordering. This is the same as taking the pentagon below and sending it to $\text{Fun}(X^4, X)$, each vertex going to the labeled 4-ary operation. **This pentagon is not filled in.**



Again, we think of e.g. $((ab)c)d$ and $(ab)(cd)$ as points in $\text{Fun}(X^4, X)$ and the image in $\text{Fun}(X^4, X)$ of the edge between them as a choice of homotopy equivalence $((ab)c)d \simeq (ab)(cd)$.

Actually, if we chose a path in $\text{Fun}(X^3, X)$ from $a(bc)$ to $(ab)c$ (i.e., if we've made a choice of homotopy equivalence realizing $(ab)c \simeq a(bc)$), that already gives us the five homotopy equivalences above—that is, where to send the pentagon above in $\text{Fun}(X^4, X)$.

Notice that there are determined TWO homotopy equivalences between e.g. $a(b(cd))$ and $(ab)(cd)$. One follows the path in $\text{Fun}(X^4, X)$ from $a(b(cd))$ to $(ab)(cd)$ given by traversing the pentagon clockwise,

²That is, the maps $X^3 \rightarrow X$ given by “concatenate (a, b, c) to $(ab)c$, or to $a(bc)$.”

the other counterclockwise. **Since X is a loop space, it turns out that there is a higher homotopy equivalence between these two homotopies.** (In general, this need not be the case, since $\pi_1 \text{Fun}(X^4, X)$ is not generally trivial.) Therefore, **there is determined a continuous map from the solid pentagon to $\text{Fun}(X^4, X)$.** This is like a “higher” level of associativity, a “higher” level of coherence.

In fact, since X is a loop space, its multiplication is associative up to *all* higher homotopy coherences. Let’s define an operad whose algebras are spaces with a multiplication that is associative up to all higher homotopy coherences.

Definition 0.6.15. Denote by \mathcal{K} the non- Σ *Stasheff operad*. It is defined to have $\mathcal{K}(n) :=$ the convex $(n - 2)$ -dimensional polygon with a vertex for each parenthetization of n ordered letters. (The composition maps can be defined if we use a more explicit description; I won’t give that.)

It is somewhat clear (to me, maybe everyone) that the Stasheff operad \mathcal{K} works largely because $\mathcal{K}(n)$ is contractible for each n .

Definition 0.6.16. Let \mathcal{C} be a non- Σ operad in Top . Say it is an A_∞ -operad if each $\mathcal{C}(n)$ is contractible. Say that a space is an A_∞ -space if it an algebra over an A_∞ -operad.

Here’s the main thing.

Theorem 0.6.17. *Up to weak equivalence,*

1. A_∞ -spaces are precisely the \mathcal{K} -algebras, and
2. Loop spaces are precisely the grouplike \mathcal{K} -algebras.

0.6.5 (5/11) A_n -operad stuff

Let X be a space. Given an operation $\mu : X^2 \rightarrow X$, we may ask if it is...

- (1) Associative *up to first homotopy*, i.e. we can choose an equivalence $\mu(\mu(-, -), -) \simeq \mu(-, \mu(-, -))$. A choice is the same data as a path $\mathcal{K}(3) = I \rightarrow \text{Fun}(X^3, X)$ from $(ab)c$ to $a(bc)$.
- (2) (Harder) Associative *up to second homotopy*, i.e. not only can we choose an equivalence $\mu(\mu(-, -), -) \simeq \mu(-, \mu(-, -))$, but can do so in such a way that the resulting homotopy equivalences between e.g. $\mu(\mu(\mu(-, -), -), -)$ and $\mu(-, \mu(\mu(-, -), -))$ are themselves realted by a higher homotopy equivalence. (A “second order” homotopy equivalence.) Such a choice amounts to (A) the structure described in (2), plus (B) a “compatible” map from the solid pentagon $\mathcal{K}(4)$ to $\text{Fun}(X^4, X)$ which sends the vertices to the various parenthetizations of $abcd$. See (5/9).
- \vdots
- (∞) (Even harder) Associative *up to all higher homotopies*, i.e. we can choose a morphism $\mathcal{K} \rightarrow \text{End}_X$ such that the path $\mathcal{K}(2) = I \rightarrow \text{End}_X$ connects $(ab)c$ and $a(bc)$.
- (?) (Too hard) *Strictly* associative, i.e. $\mu(\mu(-, -), -) = \mu(-, \mu(-, -))$.

The structure of (∞) on X is intuitively captured by a choice of \mathcal{K} -algebra structure on X (\iff a choice of A_∞ -algebra structure). We saw that if $X \simeq \Omega Y$ and we take $\mu =$ loop concatenation, then X has a \mathcal{K} -algebra structure. Moreover, up to (some notion of equivalence between \mathcal{K} -algebras?), all grouplike \mathcal{K} -algebras arise from a loop space and loop concatenation. That’s the $n = 1$ case of May’s recognition principle.

We’re often thinking about the case where μ is only associative up to n -th homotopy for some $n < \infty$. For example, *homotopy associative H -spaces* are precisely those with an operation that is associative up to first homotopy.³ (And, in my notation, there is something which might be called “associative up to no homotopies,” whose algebras are precisely not-necessarily-homotopy-associative H -spaces.)

We want to characterize these structures with operads too. This means taking the data of an A_∞ -space and truncating its coherence data at the n -th level. So for example, an “ A_3 -space” should be a based space X , a homotopy monoidal structure $\mu : X^2 \rightarrow X$, and a homotopy equivalence $\mathcal{K}(3) = I \rightarrow$

³Actually, this is wrong as I’ve described “associativity up to n -th homotopy,” but I’m just trying to give some intuition for A_n -algebra structures, so I’ll gloss over this point as it’ll all work when I actually get to A_n -operads.

$\text{Fun}(X^3, X)$ from $(ab)c$ to $a(bc)$. Then an “ A_n -operad” should be an operad whose algebras are precisely A_n spaces, modulo weak homotopy equivalence.

Our flagship A_∞ -operad is the Stasheff operad \mathcal{K} . Each associahedron $\mathcal{K}(i)$ is a CW complex (in fact a simplicial complex) in an obvious way. My first thought was, “maybe taking the $(n - 2)$ -skeleton of every $\mathcal{K}(i)$ will produce an A_n -operad.” After all, taking e.g. $n = 3$, if we take the 1-skeleton of \mathcal{K} , call it \mathcal{K}_3 , then an algebra over this operad will have a homotopy monoidal structure (specified by where we send $\mathcal{K}_3(2) = *$) that is homotopy associative (specified by where we send $\mathcal{K}_3(3) = I$), but it will NOT be “associative up to second homotopy” since $\mathcal{K}_3(4)$ is the *hollow* pentagon. (The image of its interior is what *would have* specified the “second-order homotopy coherence” for associativity.)

BUT, this thing \mathcal{K}_3 is NOT an operad. For a simple reason: if $\mathcal{K}_3(4)$ contains intervals, then composition necessitates that $\mathcal{K}_3(5)$ contains products of intervals. (We haven’t precisely defined composition, but no matter.) But this proposal for \mathcal{K}_3 is such that $\mathcal{K}_3(5)$ has no products of intervals, so it doesn’t work.

I don’t think we’re far off, though. There’s a *free operad* construction on “collections of objects” (yet-undefined) that I suspect will recover an operad from an appropriate n -truncation of \mathcal{K} which should be rightfully called an A_n -operad.

0.6.6 (5/13) Rings via operads?

Tangent today. So far, I’ve thought of operads as devices for studying the structure of “an operation with coherences” on a given object. We also care about *rings* and *ringlike* structures. These have TWO operations, plus coherence data for identities that may involve BOTH operations at once. (Distributivity.)

Question 0.6.18. Can we use operads to capture the structure of rings?

The answer depends on the base category! We cannot use operads to characterize rings “all at once,” i.e. in Set . We can instead try changing our base category to “handle” some of one operation first, though—we’ll find that we can recover rings as algebras over operads in Ab .

Let’s start in Set . We want an operad \mathcal{R} such that \mathcal{R} -algebras are precisely rings. Or commutative rings—it won’t matter, since \mathcal{R} does not exist in either case. I’ll give two proofs why.

The first proof is moral, not a real proof. Suppose \mathcal{R} is such that \mathcal{R} -algebras are precisely rings. Then there should be “addition” and “multiplication” elements $A, M \in \mathcal{R}(2)$ such that \mathcal{R} ’s compositions realize the identity $a(b + c) = ab + ac$. The issue is that the 3-ary function $ab + ac$ calls a in more than one input spot. **This is NOT expressible using an operad’s notion of composition.** The closest we can get is $\gamma(A, M, M) \in \mathcal{R}(4)$, but this is $ab + cd$, not really what we wanted. Thus, we cannot impose distributivity.

Here’s a formal proof. First note: there’s an operad Set^\times whose i -th object is a set of size i and whose composition arises from the Cartesian product. For an operad \mathcal{C} , a \mathcal{C} -algebra is precisely a morphism $\mathcal{C} \rightarrow \text{Set}^\times$. The category Operad has a terminal object whose algebras are monoids,⁴ so there’s a functor $S : \text{Monoid} \rightarrow \text{Alg}_{\mathcal{C}}$. It takes each map $* \rightarrow \text{Set}^\times$ and postcomposes it with the unique map $\mathcal{C} \rightarrow *$. **I THINK** the functor $S : \text{Monoid} \rightarrow \text{Alg}_{\mathcal{C}}$ just takes a monoid, forgets its structure, and endows it with the structure of a \mathcal{C} -algebra; **THUS**, this functor fits into a commutative triangle with the forgetful functor. **BUT**, the following proposition says that there are NO functors $\text{Monoid} \rightarrow \text{Ring}$ commuting with the forgetful functor, so there must not exist an operad \mathcal{C} in Set such that $\text{Alg}_{\mathcal{C}} \cong \text{Ring}$.

Proposition 0.6.19. *There are no functors $\text{Monoid} \rightarrow \text{Ring}$ commuting with the forgetful functors to Set .*

Zhen Lin proves this in his MSE answer [here](#).

So there are no operads in Set whose algebras are rings. Here’s a workaround: we can replace Set with a category of objects which have addition already built in, that would be Ab . Then we can finish the job with an operad in Ab and its algebras will recover Ring .

Let Comm denote the commutative monoid operad in Ab . Its objects are trivial groups and its actions are trivial also.

⁴If we’re thinking of symmetric operads, the terminal operad is Comm . If we’re thinking of plain operads, it is Assoc .

Suppose G is a $\mathcal{C}\text{omm}$ -algebra, i.e. an abelian group with a map $f : \mathcal{C}\text{omm} \rightarrow \mathcal{E}\text{nd}_G$. This data distinguishes two elements $1_G := f_0(*) \in G$ and $\times := f_2(*) \in \text{Hom}(G^2, G)$. Since $\mathcal{C}(1) = \{*\}$, we have that

$$\gamma(\times, \text{id}, 1_G) = \gamma(\times, 1_G, \text{id}) \in \mathcal{C}(1).$$

Since f must preserve composition, we get that $g \times 1_G = 1_G \times g = g$ for every $g \in G$. (How to finish???)

OK, in the above I started with $\mathcal{C}\text{omm}$ and tried to show its algebras in \mathbf{Ab} are commutative rings, I should've started with $\mathcal{A}\text{ssoc}$ and tried to show its algebras in \mathbf{Ab} are rings, but I'm getting tired. (I couldn't figure out how to get the distributive property anyway...)

Anyway, I'll remark that we do something similar in homotopy theory. We want a good notion of "spectra that are ringlike up-to-homotopy," and we must play the same game: find a category of objects with addition built-in (that would be \mathbf{Sp}) and then take algebras over an operad for commutativity "up to homotopy." (I think that's little disks?)

We are foreshadowing!

0.6.7 The rest of May

There are some notes I have not texed, and I also spent some time working on a condensed math seminar I'll be organizing at the University of Chicago in June. (I will also be there, probably studying more operad stuff with Peter May.) Maybe I will upload the rest of my May notes eventually?

0.7 June 2023

0.7.1 (6/13) June activities, monoidal categories

There are a few things going on.

1. Peter is giving lectures on operads and algebraic K -theory.
2. I'll probably be reading some of [these lecture notes](#) about algebraic K -theory.
3. I'm organizing a seminar on elementary condensed math. (And teaching quite a bit of basic category for that, as well as for other REU participants who just want to learn basic category theory.)

All this will be taking up most of my time. And it all somehow relates somehow to my goal of understanding higher category theory, especially (2). So I'll be sporadically writing here my inner monologue as I learn/do this stuff.

Today Peter spoke about A_∞ -spaces. I already wrote about those. But Peter also mentioned monoidal categories, and this led me to a little question we were not sure about.

Let me get to explaining my thought.

Definition 0.7.1. A *monoidal category* is the data of a category \mathcal{C} together with

- A functor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ called the *tensor product*;
- A distinguished object $\mathbf{1} \in \mathcal{C}$ called the *unit*;
- Isomorphisms $\lambda : \mathbf{1} \otimes - \rightarrow -$ and $\rho : - \otimes \mathbf{1} \rightarrow -$, we call the *left/right unitor* and whose components we denote λ_X, ρ_X ; and
- An isomorphism

$$A : (- \otimes -) \otimes - \cong - \otimes (- \otimes -)$$

which we call the *associator* and whose components we write $A_{X,Y,Z}$.

And these data must have the following properties.

- The *triangle identity* holds:

$$\begin{array}{ccc} (X \otimes \mathbf{1}) \otimes Y & \xrightarrow{A_{X,\mathbf{1},Y}} & X \otimes (\mathbf{1} \otimes Y) \\ \searrow \rho_X \otimes \mathbf{1} & & \swarrow \mathbf{1} \otimes \lambda_Y \\ & X \otimes Y & \end{array}$$

- The *pentagon identity* holds:

Put it in?

0.7.2 (6/15) Initial, final objects

I'm thinking about initial/final objects so that I can define a stable ∞ -category.

Topological categories are one model for ∞ -categories. In a topological category, one potential definition of a final object is obvious: it is a final object in the underlying ordinary category. Call this a "strict final object." Consider (say) the topological category $CGHaus$. There, the point $*$ is a strict final object. However, there are homotopy-equivalent spaces (i.e., contractible spaces) which are not isomorphic to $*$ in $CGHaus$ —that's not good since an ∞ -categorical definition (i.e. of final objects) should be homotopy-invariant.

But this kind of immediately indicates the (first) correct definition.

Definition 0.7.2. An object of a topological category \mathcal{C} is called *final* if it is final in $\mathrm{ho}\mathcal{C}$ (regarded as enriched over hoCW). Thus, an object X is final iff $\mathrm{Hom}_{\mathrm{ho}\mathcal{C}}(Y, X) \in \mathrm{hoCW}$ is contractible for every Y .

Follow-up after discussion

Why weakly contractible in hoCW ? Why not final as an object in the category? Is there a different

How to port this idea to quasicategories (which we must do in a way that results in the same notion upon passage to the homotopy category)? A geometric definition: if $x \in C$ is terminal, then everything “has an arrow to x ,” so the “collection of arrows to x ” should be “like a deformation retract of C .” Maybe that made sense, here’s the definition.

Definition 0.7.3. Let x be a vertex in a quasicategory C . We say x is *initial* if $C_{x/} \rightarrow C$ is a trivial fibration. Likewise, we say x is *terminal* if $C_{/x} \rightarrow C$ is a trivial fibration.

Remark 0.7.4. By definition, a vertex $x \in C$ is terminal \iff every map $f : \partial\Delta^n \rightarrow C$ such that $f(n) = x$ extends to a map $f : \Delta^n \rightarrow C$. Dually for terminal objects.

Remark 0.7.5. If C is a nerve, restriction $\text{Hom}(\Delta^n, C) \rightarrow \text{Hom}(\partial\Delta^n, C)$ is an equivalence for $n \geq 3$. Therefore the terminal objects of nerves are the terminal objects of their underlying categories.

Proposition 0.7.6. Let $\text{Hom}_C^R(x, y)$ denote the ∞ -category of right-fibrations $x \rightarrow y$ in an ∞ -category C . It is a Kan complex. Furthermore, an object y is terminal $\iff \text{Hom}_C^R(x, y)$ is contractible for every object x . (Dually for initial objects.)

Proposition 0.7.7. An object y is terminal $\iff \text{Hom}_C(x, y)$ is contractible for every object x . (Dually for initial objects.)

Proposition 0.7.8. Let C denote a quasicategory and C' the full subcategory of final objects. Then C' is a contractible Kan complex. (Likewise for the full subcategory of initial objects.)

Eventually define over and under-categories.

0.7.3 (6/18) Stable ∞ -categories

Definition 0.7.9. Let C denote an ∞ -category. We say an object in C is a *zero object* if it is initial and terminal. If C has a zero object, we call C *pointed*.

If C is pointed, its full subcategory of zero objects is a contractible Kan complex, so zero objects are unique up to equivalence.

Remark 0.7.10. As in the ordinary case, an ∞ -category C is pointed if and only if it has an initial object \emptyset , a terminal object 1 , and a morphism $1 \rightarrow \emptyset$.

Remark 0.7.11. If C is pointed, then $\text{Hom}_C(X, 0) \times \text{Hom}_C(0, Y)$ is a contractible Kan complex. The natural map $\text{Hom}_C(X, 0) \times \text{Hom}_C(0, Y) \rightarrow \text{Hom}_C(X, Y)$ then locates a unique element $0 \in \text{Hom}_{\text{ho}C}(X, Y)$. We also call this the *zero morphism*.

Definition 0.7.12. An ∞ -category C is called *stable* if it has the following properties.

- (1) C is pointed.
- (2) Every morphism in C admits a fiber and cofiber.
- (3) “Every morphism is the cokernel of its kernel and the kernel of its cokernel.”

Condition (3) is imprecise, but that is to make obvious the analogy with abelian categories. OK now let me make it precise. This will also mean defining the things in (2).

Definition 0.7.13. Let C be a pointed ∞ -category and $f : X \rightarrow Y$ a morphism. A *kernel* or *fiber* of f is a homotopy pullback of f with $0 \rightarrow Y$. We define a *cokernel* or *cofiber* of f dually. As in the following diagram.

$$\begin{array}{ccc}
 \ker(f) & \dashrightarrow & 0 \\
 \downarrow & & \downarrow \\
 X & \xrightarrow{f} & Y
 \end{array}
 \qquad
 \begin{array}{ccc}
 X & \xrightarrow{f} & Y \\
 \downarrow & & \downarrow \\
 0 & \dashrightarrow & \text{coker}(f)
 \end{array}$$

Actually, triangulated categories – understand this?

Definition 0.7.14. Let \mathcal{C} be a pointed ∞ -category. A *triangle* (g, f) in \mathcal{C} is a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow g \\ 0 & \longrightarrow & Z \end{array}$$

We write it (g, f) but a triangle consists of more data – write that in

We say a triangle (g, f) is a *fiber sequence* (resp. *cofiber sequence*) if it is a pullback (resp. pushout).

Remark 0.7.15. If $f : X \rightarrow Y$ is a morphism in \mathcal{C} , then a fiber of f is precisely a fiber sequence (f, g) for some g . Likewise, a cofiber of f is any cofiber sequence (g, f) .

Definition 0.7.16. A category \mathcal{C} is called *stable* if it has the following properties.

- (1) \mathcal{C} is pointed.
- (2) \mathcal{C} admits all fibers and cofibers.
- (3) A triangle in \mathcal{C} is a fiber sequence \iff it is a cofiber sequence.

Remark 0.7.17. Condition (3) is like “cokernels of kernels are isomorphic to kernels of cokernels.” Is there a precise restatement of (3) in this vein that isn’t a mouthful?

Remark 0.7.18. Stability is a property, not a structure.

Remark 0.7.19. At least for me, “stability” here is used with an eye toward “stabilizing the loop and suspension functors on Top^* .” And indeed, spectra will appear as precisely the stabilization of Top^* with respect to those functors.

Definition 0.7.20. Let \mathcal{C} denote a pointed ∞ -category.

- If \mathcal{C} admits cofibers, there is determined a *suspension functor* $\Sigma : \mathcal{C} \rightarrow \mathcal{C}$ which informally associates to X its homotopy pushout against two zero maps

$$\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ 0' & \longrightarrow & \Sigma X \end{array}$$

- If \mathcal{C} admits fibers, there is determined a *loop-space functor* $\Omega : \mathcal{C} \rightarrow \mathcal{C}$ which informally associates to X its homotopy pullback along two basepoint-inclusions

$$\begin{array}{ccc} \Omega X & \longrightarrow & 0 \\ \downarrow & & \downarrow \\ 0' & \longrightarrow & X \end{array}$$

Remark 0.7.21. The formal definition of Σ, Ω in general takes some work, simply because it takes work to give any concrete description of functors between ∞ -categories. Lurie’s brief description of Σ, Ω begins in HA p. 23 at the bottom. A very nice unpacking of this construction is given by Alberto García-Raboso in [this article](#).

The bottom line is that we get a “functorial construction of cofibers,” which means a functor $\text{Fun}(\Delta^1, \mathcal{C}) \rightarrow \text{Fun}(\Delta^0, \mathcal{C}) \cong \mathcal{C}$ which on objects assigns a cofiber to each morphism in \mathcal{C} . This is quite nice—there is really no good nice way to do this for triangulated categories. Grothendieck wrote a 1,976-page manuscript on “derivators” which are (a kind of?) “tool” to handle this problem that is just too hard and more concisely dealt with using ∞ -categories. (Do people use derivators?) Grothendieck’s manuscript affirms the belief that this is *definitely a problem worth solving*, in any case.

Learn triangulated categories and discuss the relation to stable ∞ -categories; essentially, HA p. 24–

0.7.4 (6/20) Idempotents

Trying to read some algebraic K -theory papers, I come across *idempotent completions* (for ∞ -categories). I've heard of these before but never really thought about them. I'll think about them today.

I'm looking at Kerodon [Section 03Y9](#) and HTT 4.4.5 for this.

First we review some ordinary category theory.

Retracts should be maps fixing sub-sets/spaces/objects. Formally, denoting by \mathcal{C} an ordinary category, we say Y is a *retract of X* if there exists some $r : X \rightarrow Y$ factoring $\text{id}_Y : Y \rightarrow Y$, i.e. some diagram

$$\begin{array}{ccc} & X & \\ i \nearrow & & \searrow r \\ Y & \xrightarrow{\text{id}_Y} & Y \end{array}$$

Remark 0.7.22. Let Ret or Idem^+ denote the category consisting of an “abstract retract diagram.” (See [Construction 03YB](#).) There is a tautological bijection $\{\text{functors } \text{Ret} \rightarrow \mathcal{C}\} \cong \{\text{retract diagrams in } \mathcal{C}\}$.

Remark 0.7.23. See that $i \circ r : X \rightarrow X$ is an idempotent. In fact, this idempotent canonically determines X up to isomorphism: X is the equalizer of id_X and $i \circ r$. One may ask about the converse: if $\phi : X \rightarrow X$ is idempotent, does X have a retract Y such that $\phi = i \circ r$? (This Y is uniquely determined if it exists.) If this is the case, i.e. if the injection

$$\{\text{retracts of } X\} / \cong \hookrightarrow \{\text{idempotent morphisms } X \rightarrow X\}$$

is a bijection for all X , we say that \mathcal{C} is *idempotent complete*.

Definition 0.7.24. In general, we say an idempotent $\phi : X \rightarrow X$ is *split* if it arises from a retraction of X , i.e. if $\phi = i \circ r$ for some $r : X \rightarrow Y$ and $i : Y \rightarrow X$ satisfying $r \circ i = \text{id}_Y$.

Remark 0.7.25. Thus, an ordinary category is idempotent complete \iff every idempotent splits.

Proposition 0.7.26. *If an ordinary category \mathcal{C} has equalizers or coequalizers, then it is idempotent complete.*

What's ado about retracts and idempotents in ∞ -categories, then? Whatever they are, they should become ordinary retracts/idempotents upon passage to the homotopy category. Lurie explains two reasons that this is insufficient, though.

The ordinary story suggests that there should be a relationship (correspondence!) between ∞ -categorical retracts and idempotents. Let's start with retracts.

Let \mathcal{C} denote an ∞ -category and X an object of \mathcal{C} .

Definition 0.7.27. We say an object $Y \in \mathcal{C}$ is a *retract of X* if there exists $r : X \rightarrow Y$ and $i : Y \rightarrow X$ such that $r \circ i = \text{id}_Y$, i.e. there exists a 2-cell witnessing that composition.

Proposition 0.7.28. *An object $Y \in \mathcal{C}$ is a retract of $X \iff$ there exists a functor $F : N\text{Ret} \rightarrow \mathcal{C}$ taking the “abstract retract object” to Y and the other object to X . (Lurie would say the F “exhibits Y as a retract of X .”)*

As in the ordinary case, we can classify retracts using (split)idempotents.

Definition 0.7.29. Define the category Idem to have one object \tilde{X} and one non-identity morphism $e : \tilde{X} \rightarrow \tilde{X}$ with composition law $e \circ e = e$.

Definition 0.7.30. Let \mathcal{C} be an ∞ -category. An *idempotent* in \mathcal{C} is a functor $F : N_\bullet \text{Idem} \rightarrow \mathcal{C}$.

Remark 0.7.31 (We recover the ordinary case). Suppose that \mathcal{C} is an ordinary category. Since the nerve functor is fully faithful, functors $N_\bullet \text{Idem} \rightarrow N_\bullet \mathcal{C}$ are in bijection with functors $\text{Idem} \rightarrow \mathcal{C}$. The former are precisely the idempotents in an ∞ -category by definition, and the latter are idempotents in \mathcal{C} by inspection. Thus, ∞ -categorical idempotents recover ordinary ones.

Definition 0.7.32. Let \mathcal{C} be an ∞ -category. An idempotent $I : N_\bullet \text{Idem} \rightarrow \mathcal{C}$ is called *split* if there exists a functor $R : N_\bullet \text{Ret} \rightarrow \mathcal{C}$ (a “retract diagram in \mathcal{C} ”) extending I along the obvious functor $N_\bullet \text{Idem} \hookrightarrow N_\bullet \text{Ret}$.

Explain the reasons? Not super important I think.

Remark 0.7.33. These, too, recover ordinary idempotents.

Proposition 0.7.34. *Splittings of an idempotent are essentially unique. Precisely, we mean that restriction*

$$T : \text{Fun}(N_\bullet \text{Ret}, C) \rightarrow \text{Fun}(N_\bullet \text{Idem}, C)$$

is fully faithful. (By definition, its essential image is the full subcategory of split idempotents in C .)

Proof. Idem sits naturally inside Ret. That's kind of the whole point. Clearly, Ret is generated by Idem under retracts; this implies that any $F : \bullet \text{Ret} \rightarrow C$ is left and right Kan extended from its restriction $T(F) : N_\bullet \text{Idem} \rightarrow C$ ([Proposition 03YQ](#)). This lets us apply [Corollary 030S](#) (which I haven't really unpacked) to conclude that T is a trivial Kan fibration. Trivial fibrations of Kan complexes are fully faithful. (Consult Rune's notes.) \square

Remark 0.7.35. Now let me rehash and give some ideas:

- If C is an ordinary category, we may speak of *idempotents* and *retracts*. By definition, given an object X , retracts of X (up to isomorphism) biject with split idempotents $X \rightarrow X$. But not every idempotent splits. Varying X , we get a fully-faithful functor $\text{Fun}(\text{Ret}, C) \hookrightarrow \text{Fun}(\text{Idem}, C)$. The inverse problem amounts to taking a equalizer or coequalizer; if $\phi : X \rightarrow X$ is idempotent, then the (co)equalizer of ϕ with id_X realizes a retract of X (and retracts in general are uniquely determined).
- If C is an ∞ -category, we define *retracts of X* as objects Y with a certain property analogous to ordinary retracts. If Y is a retract of X in this sense, there is determined a functor $N_\bullet \text{Ret} \rightarrow C$, unique up to isomorphism. We do a bit more work for idempotents: the most obvious "property" of idempotency actually is undesirably ambiguous, so we *define* idempotents as functors $N_\bullet \text{Idem} \rightarrow C$. As in the ordinary case, we get a fully faithful (i.e. a trivial Kan fibration) functor

$$\text{Fun}(N_\bullet \text{Ret}, C) \rightarrow \text{Fun}(N_\bullet \text{Idem}, C).$$

We can characterize its essential image as those idempotents with the *property* of splitness, defined analogously as in the ordinary setting. Once again, the inverse problem (i.e. determining whether an idempotent splits, i.e. extending some idempotent $N_\bullet \text{Idem} \rightarrow C$ to a retract diagram in a Kan-ny way?) amounts to finding the limit or colimit of $N_\bullet \text{Idem} \rightarrow C$.

- Also see [Section 03Y9](#).

0.7.5 (6/21) I hate idempotents today. K -theory?

I physically cannot think about idempotents after yesterday. Luckily there is a different thing I also need to understand for modern algebraic K -theory, that being (modern?) algebraic K -theory. I already understand a little bit.

Let me collect some references.

- Rune Haugseng's 2010 notes [The \$Q\$ -construction for stable \$\infty\$ -categories](#).
- The MO thread [Motivation/interpretation for Quillen's \$Q\$ -construction?](#) and some of what's referenced therein.

Let's start somewhere classical. In *The Geometry of Iterated Loop Spaces*, May defined the little cubes operads C_k , the prototypical E_k -operad. Given a space X , we can form the *free C_k -algebra* $C_k[X]$. Any k -fold loop space is a C_k -algebra, in particular $\Omega^k \Sigma^k X$. We get a natural C_k -algebra map $e : C_k[X] \rightarrow \Omega^k \Sigma^k X$ from the natural map $X \rightarrow \Omega^k \Sigma^k X$ of spaces.⁵ This e is a weak equivalence $\iff X$ is grouplike. This is the *approximation theorem*, which is used to prove the following.

Theorem 0.7.36 (May's recognition theorem). *X is a grouplike C_k -algebra $\iff X$ is weakly equivalent to $\Omega^k Y$ for some Y .*

⁵Our spaces are based, of course. We may also want a based variant of the free C_k -algebra on a space?

Prove that given isomorphic idempotents $I \cong I'$, one splits iff the other splits. This isn't too hard—
I just
Understand 7.3.6.15 one day?
to think about isofibrations?

Write this out?

What monoid structure?

Let me sketch the hard direction of this (not going into detail about the hard parts). Suppose as given X a \mathcal{C}_k -algebra. There is a map of monads $\alpha_k : C_k X \rightarrow \Omega^k \Sigma^k X$ induced by the natural map $X \rightarrow \Omega^k \Sigma^k X$. The map $X \rightarrow \Omega^k \Sigma^k X$ then induces a map $X \rightarrow \Omega^k B(\Sigma^n, C_k, X)$, where that last space is a two-sided bar construction. We think of it as “like a k -fold delooping of X .” Furthermore, α_k was a group completion,⁶ so $X \rightarrow \Omega^k B(\Sigma^n, C_k, X)$ is a group completion also. (Here, the monoid structure on $\pi_0 X$ is induced by the \mathcal{C}_k -algebra structure.) Therefore, it is a weak equivalence if and only if X is grouplike, in which case we’ve realized X as a k -fold loop space.

What’s the relation to K -theory? Take $k = 1$. The above says that an A_∞ structure on X lets us “deloop” X (that delooping being $B(\Sigma^n, C_k, X)$ above) and construct a group completion $X \rightarrow \Omega B X$. In other words, you get what you might expect: a “homotopy” monoid X has a “homotopy” group completion $X \rightarrow \Omega B X$.

Trying to sort my thoughts out about group completions is actually a bit of a pain.⁷ I’m going to write in my stupidity and stop writing for today. Also see [this MO post](#).

0.7.6 (6/26) Structure of $\mathrm{Hom}_{\mathcal{C}}(-, -)$ for (various adjectives) categories

Recall that a *topological category* is one enriched over CG . In some form, Whitehead’s theorem says that every space X is weakly equivalent to a CW-complex X' , unique up to a unique weak equivalence. Therefore, $X \mapsto [X] := X'$ defines a functor $\theta : \mathrm{CG} \rightarrow \mathrm{hoCW}$. This functor exhibits hoCW as a localization $\mathrm{CG}[w^{-1}]$ at weak equivalences. It happens that θ preserves products. By general procedure, given such a nice functor θ , any CG -enriched category may now be canonically enriched (via θ) over hoCW . Now given a topological category \mathcal{C} , we define $\mathrm{ho}\mathcal{C}$ to have the same objects and we define $\mathrm{Hom}_{\mathrm{ho}\mathcal{C}}(X, Y)$ to be the CW-approximation $[\mathrm{Hom}_{\mathcal{C}}(X, Y)] \in \mathrm{hoCW}$.

Proposition 0.7.37. *The homotopy category of a topological category is canonically enriched (via Whitehead’s theorem) over hoCW , i.e. the homotopy category of spaces.*

Now let \mathcal{C} denote a quasicategory. Recall that a *morphism* $f : X \rightarrow Y$ between $X, Y \in \mathcal{C}$ is a 1-simplex such that $d_1(f) = X$ and $d_0(f) = Y$. Morphisms $f : X \rightarrow Y$ are in bijection with the vertices of the *mapping space* $\mathrm{Map}_{\mathcal{C}}(X, Y)$ defined as the fiber product

$$\begin{array}{ccc} \mathrm{Map}_{\mathcal{C}}(X, Y) & \dashrightarrow & \mathrm{Fun}(\Delta^1, \mathcal{C}) \\ \downarrow & & \downarrow \\ \Delta^0 & \xrightarrow{\quad * \mapsto (X, Y) \quad} & \mathrm{Fun}(\partial\Delta^1, \mathcal{C}) \cong \mathcal{C} \times \mathcal{C} \end{array}$$

(One could also define this as a fiber product over $\mathrm{Fun}(\Delta^1, \mathcal{C})$.) This defines $\mathrm{Map}_{\mathcal{C}}(X, Y)$ as a simplicial set. **In fact, $\mathrm{Map}_{\mathcal{C}}(X, Y)$ is a Kan complex.** The shortest proof of this proceeds as follows.

- The morphism $i : \partial\Delta^1 \rightarrow \Delta^1$ is a monomorphism and is bijective on vertices. This implies that the restriction map $i^* : \mathrm{Fun}(\Delta^1, \mathcal{C}) \rightarrow \mathrm{Fun}(\partial\Delta^1, \mathcal{C})$ is conservative (Charles’ notes, 37.1; this works for map of simplicial sets.) Therefore the (trivial) map $\mathrm{Map}_{\mathcal{C}}(X, Y) \rightarrow \Delta^0$ is conservative. Since every edge in Δ^0 is an isomorphism, it must have been so in $\mathrm{Map}_{\mathcal{C}}(X, Y)$. Thus $\mathrm{Map}_{\mathcal{C}}(X, Y)$ is an ∞ -groupoid, i.e. a Kan complex by Joyal’s theorem.

Proposition 0.7.38. *A quasicategory is “enriched over spaces” in the sense that any mapping space between two objects is a Kan complex.*

This is sort of internal to a particular quasicategory. In the *Joyal model structure* on \mathbf{sSet} , the fibrant objects are precisely the quasicategories.

There’s a recurring theme of “infinity-categorical things have a *space* of morphisms between objects.” Whatever “space” means to you. Now, let \mathcal{C} denote a simplicially-enriched category. Simplicial sets are “space-ish,” but we may ask about those simplicially-enriched categories whose hom-sets are all Kan complexes. That makes them *really* space-like.

⁶Is it just a group completion of spaces? Is there more to this statement?

⁷In the process, I found [this](#) nice article from Sanath Devalapurkar. Also [these](#) notes of Dylan Wilson’s.

Hey, that was kind of cool. Think more about this two-sided bar construction? All this operad-monad stuff fits together nicely.

Write more about group completions?

Proposition 0.7.39. *There is a model structure on the category of simplicial categories, called the Bergner model structure whose fibrant objects are precisely the categories enriched in Kan complexes. Furthermore, there is a Quillen equivalence between this model category and sSet with the Joyal model structure.*

This is in HTT? I have heard it is not easy. I wonder if the proof uses anywhere that (by construction) the most natural definition of “the mapping space between objects of a quasicategory” form Kan complexes.

0.7.7 (6/28) Anima and other examples of ∞ -categories via N^c

First, I just wanted to record a characterization of (ordinary) limits and colimits I learned. Let C denote a small category, J a small diagram category, and $\Delta : C \rightarrow \text{Fun}(J, C)$ the “diagonal” functor $c \mapsto (J \mapsto c)$. Furthermore, assume that C has all J -shaped colimits.

The assignment $(\phi : J \rightarrow C) \mapsto \text{colim}_J \phi$ defines a functor $C^J \rightarrow C$. I want to show that this is left-adjoint to Δ . There are many ways to do this (some of which are kind of interesting to think about...) Maybe the quickest way uses the following:

- A functor $F : A \rightarrow B$ is a left adjoint iff one can specify, for each $b \in B$, an object $G_b \in A$ and a “universal arrow” $\epsilon_b : b \rightarrow F(G_b)$.

That this is fulfilled for $F = \text{colim}_J : C^J \rightarrow C$ is true, more-or-less by definition of colimits. The rest is vaguely “determined,” maybe up to choice. One could also proceed via the following.

- Given functors $F : A \rightarrow B : G$, an adjunction $F \dashv G$ is determined by a universal natural transformation $\eta : \text{id}_A \Rightarrow GF$.

Again, in our case, this is immediate by the definition of colim_J . Namely, given a diagram $\phi : J \rightarrow c$, its colimit $\text{colim}_J \phi$ is a universal cone under ϕ , which is precisely the data of a universal morphism (in c^J) $\phi \rightarrow \Delta_{\text{colim}_J} \phi$. Letting ϕ vary, we assemble η .

Proposition 0.7.40. *If C is small and J is a small diagram category, and furthermore C is J -cocomplete, then $\text{colim}_J : C^J$ is left-adjoint to the diagonal $\Delta : C \rightarrow J$.*

Analogously, the limit functor is right-adjoint to Δ . I think, in fact, that the existence of any left (resp. right) adjoint to Δ is equivalent to cocompleteness (resp. completeness) of C .

Anyway, that is not what I wanted to focus on. An *anima* is an ∞ -category whose homotopy category is a groupoid. (In other words, anima are ∞ -groupoids.)

Example 0.7.1. Joyal’s theorem says that anima are precisely the Kan complexes.

Example 0.7.2 (Cores are anima). The core of ∞ -category, i.e. “the maximal ∞ -subgroupoid,” is an anima. Let’s recall how to construct this. If C is an ordinary category, then C^{core} is the subcategory spanned by isomorphisms in C . For C an ∞ -category, we define its core as the pullback

$$\begin{array}{ccc} C^{\text{core}} & \xrightarrow{\quad \quad \quad} & C \\ \downarrow & \lrcorner & \downarrow \\ N(\pi C^{\text{core}}) & \longrightarrow & N(\pi C) \end{array}$$

Example 0.7.3 (Hom-sets are anima). As discussed previously, if C is a quasicategory, then $\text{Hom}_C(x, y)$ is an anima.

We want an ∞ -category of anima. I think, at some point, that I defined the ∞ -category of ∞ -categories explicitly, based on Charles notes. This was *not* really the nerve of the ordinary category of ∞ -categories. For some reason, the nerve isn’t the “right” way to think about these kinds of constructions. This is intuitive: an ∞ -category should express and organize some “homotopical phenomena,” so if our input is an ordinary category C (which has no higher data) then whatever we extract from it (e.g. $N(C)$) probably won’t have much good homotopical information.

Write about N^c one day?

So there are two problems: we start with an insufficient amount of data (an ordinary category), and if we change that (by specifying more data), we need to “upgrade” the nerve $N_\bullet(-)$ to account for the additional data. The *homotopy-coherent nerve* resolves this; it is some technically-defined functor that sends simplicially-enriched categories to simplicial sets.

Definition 0.7.41. The *coherent/simplicial/homotopy-coherent nerve* is some functor

$$N^c(-) : \text{Cat}^{\text{sSet}} \rightarrow \text{sSet}.$$

Proposition 0.7.42 (Fabian’s notes, I.14). *If \mathcal{C} is a category enriched in Kan complexes, then $N^c(\mathcal{C})$ is an ∞ -category. Moreover, there is a canonical homotopy equivalence of Kan complexes*

$$\text{Hom}_{N^c(\mathcal{C})}(x, y) \simeq \text{Hom}_{\mathcal{C}}(x, y).$$

Example 0.7.4. The ordinary full subcategories $\text{Kan}, q\text{Cat} \subseteq \text{sSet}$ are Kan-enriched via

$$\text{Hom}_{\text{Kan}}(X, Y) = \text{Fun}(X, Y) \quad \text{and} \quad \text{Hom}_{q\text{Cat}}(X, Y) = \text{Hom}_{\text{sSet}}(X, Y)^{\text{core}}.$$

The *∞ -category of anima* is the coherent nerve $N^c(\text{Kan})$. The *∞ -category of quasicategories* is the coherent nerve $N^c(q\text{Cat}')$, where $q\text{Cat}'$ is the Kan-enriched category formed by replacing the hom-quasicategories in $q\text{Cat}$ with their maximal ∞ -groupoids.

Example 0.7.5 (Homotopy hypothesis, Fabian I.20). Let X be a space. The singular complex $S_\bullet X$ is an ∞ -groupoid. This functor S_\bullet is right adjoint to geometric realization. A corollary to inspection of the skeletal filtration on simplicial sets says that a geometric realization has a canonical CW structure; thus $|-| : \text{sSet} \rightarrow \text{Top}$ lands in CW. Top, in particular CW, is Kan-enriched via $\text{Hom}(X, Y) := \text{Sing}_\bullet \text{Hom}(X, Y)$. We can therefore pass to the simplicial nerve, and our adjunction becomes an *equivalence*

$$\text{An} := N^c(\text{Kan}) \begin{array}{c} \xrightarrow{|-|} \\ \xleftarrow{\text{Sing}_\bullet} \end{array} N^c(\text{CW})$$

That this is an equivalence is Grothendieck’s *homotopy hypothesis*.

Read example $N^c(\text{Ch}(R))$, i.e. nerve of category of chain complexes of R -modules. See Fabian, I.15(e).

0.8 July 2023

0.8.1 (7/5) Derived ∞ -categories

There are a few ways to define, think about, characterize, etc. the *derived* ∞ -category of an abelian category. I suppose it's the "correct" setting to do homological algebra with complexes modulo quasi-isomorphisms. In particular, they recover the triangulated structure we use in the ordinary setting for the purposes of homological algebra. Naturally, they are related to stable ∞ -categories. I'll understand this eventually. [Today I'll go over my takeaways from Achim Krause's talk about them.](#)

To an abelian category \mathcal{A} , we want to associate an ∞ -category $\mathcal{D}_\infty(\mathcal{A})$ that encodes the homotopy theory (where quasi's should be the weak equivalences) of $\mathbf{Ch}(\mathcal{A})$.

Definition 0.8.1. If \mathcal{A} is an abelian category, we denote by $\mathbf{Ch}(\mathcal{A})$ the category of unbounded chain complexes in \mathcal{A} . Note that for $A, B \in \mathbf{Ch}(\mathcal{A})$, the differential on B gives rise to a dg structure on $\mathrm{Hom}_{\mathbf{Ch}(\mathcal{A})}(A, B)$, i.e. the structure of a complex of abelian groups. Thus, $\mathbf{Ch}(\mathcal{A})$ has a canonical enrichment over $\mathbf{Ch}(\mathbb{Z})$.

There's latent homotopical data in the structure of the $\mathbf{Ch}(\mathbb{Z})$ -enrichment of $\mathbf{Ch}(\mathcal{A})$. (In particular, the $\mathbf{Ch}(\mathbb{Z})$ -enrichment encodes the chain-homotopy equivalences.) We have a good way for translating this into our common language of ∞ -categories: consider the composite

$$K : \mathbf{Ch}(\mathbb{Z}) \xrightarrow{\tau_{\geq 0}} \mathbf{Ch}_{\geq 0}(\mathbb{Z}) \xrightarrow[\Gamma]{\cong} s\mathbf{Ab} \xrightarrow{\text{forget}} \mathbf{Kan}. \quad (2)$$

The forgetful functor lands in \mathbf{Kan} because the underlying simplicial set of a simplicial abelian group is a Kan complex. The functor Γ is one direction of the *Dold-Kan correspondence*. The functor $\tau_{\geq 0}$ is the *canonical truncation*.⁸

Proposition 0.8.3. *Each category in Equation (2) is monoidal, with monoidal product given by (from left to right): the \otimes of chain complexes, the same, pointwise \otimes of abelian groups, the usual product. (In fact, they are symmetric monoidal.)*

Furthermore, each functor in Equation (2) is lax-monoidal. (Note that Γ is not lax-symmetric monoidal, although its inverse is.)

Thus, hitting hom-objects with K describes a functor

$$\{\mathbf{Ch}(\mathbb{Z})\text{-enriched categories}\} \rightarrow \{\mathbf{Kan}\text{-enriched categories}\}.$$

And we (vaguely, I have not yet really worked this out) know how to take the latter sort of category and reformulate it as an ∞ -category.

Definition 0.8.4. If \mathcal{A} is an abelian category, define its *homotopy category* as the ∞ -category

$$\mathcal{K}_\infty(\mathcal{A}) := N^c(\mathbf{Ch}(\mathcal{A})_\Delta),$$

where $\mathbf{Ch}(\mathcal{A})_\Delta$ is the simplicial category obtained by applying K above to Hom-objects. Note that this is in ∞ -category since K lands in \mathbf{Kan} rather than just $s\mathbf{Set}$.

The homotopy category $\mathcal{K}_\infty(\mathcal{A})$ is nice. For example, it is finitely bicomplete. Moreover, for chain complexes A and B we have by construction

$$\pi_n \mathrm{Hom}_{\mathcal{K}_\infty(\mathcal{A})}(A, B) \cong H_n \mathrm{Hom}_{\mathbf{Ch}(\mathcal{A})}(A, B).$$

In particular, $\pi_0 \mathrm{Hom}_{\mathcal{K}_\infty(\mathcal{A})}(A, B)$ consists of chain maps $A \rightarrow B$ modulo chain-homotopy equivalence.

Remark 0.8.5. **There is a problem.** Although chain-homotopic maps are identified upon passage to $\mathcal{K}_\infty(\mathcal{A})$, quasi-isomorphisms do not become isomorphisms. For example, the following chain map is a

⁸In negative degrees, $\tau_{\geq 0}$ naively truncates a chain complex. In degree 0, it sends A_0 to $\ker(d_1)$. It leaves positive degrees unchanged. This has the effect of preserving non-negative homology. This is contrast to the most obvious *stupid truncation*, which leaves A_0 unchanged. See [?, [Section 0118](#)].

quasi-isomorphism but is not invertible in $\mathcal{K}_\infty(\mathcal{A})$.

$$\begin{array}{ccccccccc} \cdots & \longrightarrow & 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{2} & \mathbb{Z} & \longrightarrow & 0 & \longrightarrow & \cdots \\ & & & & \downarrow & & \downarrow \pi & & \downarrow & & \\ \cdots & \longrightarrow & 0 & \longrightarrow & \mathbb{Z}/2 & \longrightarrow & 0 & \longrightarrow & \cdots \end{array}$$

We do not beat around the bush: to fix the problem, we just DK-localize it away. Let W be the set of quasi-isomorphisms of complexes in $\mathcal{K}_\infty(\mathcal{A})$. We define the *derived ∞ -category* of the abelian category \mathcal{A} to be the localization

$$\mathcal{D}_\infty(\mathcal{A}) := \mathcal{K}_\infty(\mathcal{A})[W^{-1}].$$

Write about DWyer-Kan localization, existence, examples?

Bibliography

- [Fri08] Greg Friedman. An elementary illustrated introduction to simplicial sets. 2008.
- [Lur22] Jacob Lurie. Kerodon. <https://kerodon.net>, 2022.
- [Mat] Akhil Mathew. The dold-kan correspondence. <https://math.uchicago.edu/~amathew/doldkan.pdf>.
- [Rie] Emily Riehl. A lesuirely introduction to simplicial sets. <https://math.jhu.edu/~eriehl/ssets.pdf>.