

# 2024 Notebook

Matthew Niemi

# Contents

I	January	2
I.1	(1/1) Cardinality	2
I.2	(1/4) Colimit completions and filtering classes	4
I.3	(1/15) Reminding myself what presentability, dualizability, and stability are	7
I.4	(1/24) Stabilization, the $\infty$ -category of spectra, and the smash product	9
I.5	(1/29) Localizations I	12
II	February	14
II.1	(2/2) Phony multiplication	14
II.2	(2/7) Localizations II	18
II.3	(2/14) Categorifying Mahler's theorem	22
II.4	(2/21) The pushdiagonal of a ring with $G$ -action	23
II.5	(2/22) Linear Galois theory I	23
II.6	(2/26) Linear Galois theory II	25
II.7	(2/28) Descent I	26

# I January

Prior to the advent of the brain, there was no color and no sound in the universe, nor was there any flavor or aroma...before brains the universe was also free of pain and anxiety.

---

Roger Sperry

New year. And a new notebook. Last one was getting tedious to compile, and it was about time I tweak the  $\text{T}_\text{E}\text{X}$  anyway. I should start hearing back from graduate schools soon. I am pretty excited to start grad school, and now that the anxiety of applications is gone, I can feel and appreciate how much time six years (the duration I plan to be in graduate school) is to do things. I have some math-related new year's resolutions: learn some combinatorics, GAP 4, probability theory and ergodic theory, and number theory. Maybe special mention to Cohen-Lenstra statistics. This list reflects personal interests I feel I have neglected. Bonus points if I can make things weird (i.e., categorical). This notebook remains a documentation of what I'm studying in higher category theory.

## I.1 (1/1) Cardinality

A “set of all sets” is set-theoretically impossible, although there is a proper class  $T$  of such things. A group is a set, hence *the class of groups of order 2* is at least the size of  $T$  since each set  $S \in T$  gives rise to a group of order 2, namely  $\{S, S\}$  with whatever<sup>1</sup> group structure you'd like. In particular, there is a *proper* class of groups of order 2. The lesson: for many purposes, we should count up to isomorphism.

Let  $X$  be a finite set and  $\sim$  an equivalence relation. The quotient  $X / \sim$  is finite, hence its *cardinality* is well-defined and equals the number of equivalence classes, which we can count “over  $X$ ” via the easy formula

$$|X / \sim| = \sum_{x \in X} \frac{1}{|[x]|}.$$

Notice this formula does not use the relation  $\sim$  in any meaningful way (it is just a weird way of writing  $|X / \sim| = \sum_{[x]} 1$ ). In certain situations, we may not like this, and let me explain a better and similar-looking quantity to call “the size of  $X / \sim$ .”

**Observation I.1.** Let  $X$  be a finite set and  $\sim$  an equivalence relation. *A priori*, the natural distribution on a finite set  $X$  is the uniform one. You can also put the uniform distribution on  $X / \sim$ , but this does not agree with another distribution we can put on  $|X / \sim|$ , namely that where  $\mathbb{E}([x])$  is induced by the likelihood of obtaining any  $y \in [x]$ .

**Example I.1.** Let  $G = \mathbb{Z}/2\mathbb{Z}$  act on the set

$$X := \{1, 2, 3, 4, 5\}$$

as the cycle (15)(24). Then  $X/G$  has three points, those being the orbits  $\{1, 5\}$ ,  $\{2, 4\}$ , and  $\{3\}$ . If we take the uniform distribution on  $X$ , then the induced distribution on  $X/G$  is *not* uniform. Rather, it weighs the listed orbits  $2/5$ ,  $2/5$ , and  $1/5$ , respectively, but the uniform distribution would weigh them each  $1/3$ .

**Remark I.1.** Our use of probability here is not essential. It just gives a natural language to talk about the “size” of objects—bigger objects should take up more of the distribution.

**Remark I.2.** Let  $G$  be any finite group acting on a finite set  $X$ . If the action is free, then there is no issue like the above, for freeness is equivalent to trivial stabilizers, which implies that the orbits of  $G$  have uniform size, namely the orbits are of size  $|G|$ .

---

<sup>1</sup>Which ever. Of the two.

**Remark I.3.** The category of finite sets is the “categorification” of the natural numbers. We can divide two natural numbers—how to divide two finite sets  $S$  and  $T$ ? If  $T$  is a finite group acting on  $S$ , we may consider the quotient  $S/T$ . By the previous remark, if this action is free, then  $|S/T| = |S|/|T|$ . Hence, quotienting sets by finite *free* actions is a “categorification” of division. However, this process only produces finite sets, which is not what we want because finite sets are categorified natural numbers—we should obtain *rational*s via division! The solution is to consider all action quotients (possibly non-free!). But this suffers from the same problem, namely that a non-free quotient is still *just* a finite set (e.g.,  $|X/G| = 3$  above). Hm...

The problem is that  $3 \in X$  has more automorphisms (a smaller orbit), so it “appears larger” in  $X/G$  than it “really is” because the set  $X/G$  does not see automorphisms. **This suggests that the “size” of an element should be inversely proportional to its number of automorphisms.** We therefore should replace sets with objects that carry automorphism data (groupoids) and extend the notion of cardinality to account for that data.

**Definition I.1.** Let  $X$  be an ordinary groupoid. Define its *homotopy cardinality* as the sum

$$|X| = \sum_{[x] \in \pi_0 X} \frac{1}{|\text{Aut}(x)|}.$$

**Example I.2.** Let  $G = \mathbb{Z}/2\mathbb{Z}$  act on a five-element  $X$  as the cycle  $\sigma = (15)(24)$  again. Consider the associated *action groupoid*  $X//G$  [nLab]. It has  $\text{Ob}(X//G) = X$  and a morphism  $g : x \rightarrow y$  if and only if  $\exists g : gx = y$ . In this case, we can compute its homotopy cardinality as

$$|X| = \frac{1}{|\text{Aut}(1)|} + \frac{1}{|\text{Aut}(2)|} + \frac{1}{|\text{Aut}(3)|} = \frac{1}{|\{\sigma^2 = \text{id}\}|} + \frac{1}{|\{\sigma^2 = \text{id}\}|} + \frac{1}{|\{\text{id}, \sigma\}|} = 5/2.$$

This works. In the above example, it weights elements in the quotient correctly, namely prescribing 3 a weight of  $1/2$ . **More generally, given any (possibly non-free) finite group action  $G \curvearrowright X$ , we have  $|X//G| = |X|/|G|$  as desired.** You may crank out exotic examples. You may ponder the cardinality of the core of your favorite category. For example,  $|\text{FinSet}^\cong| = e$ .

Homotopy cardinality is homotopy invariant. Furthermore, it has the essential properties of ordinary cardinality: it is additive and multiplicative over products and coproducts, respectively. How to extend this notion to  $\infty$ -groupoids? Let me just give you the definition: for an  $\infty$ -groupoid  $X$ , we define its *homotopy cardinality* as the sum

$$|X| := \sum_{[x] \in \pi_0 X} \prod_{n=1}^{\infty} |\pi_n(X, x)|^{(-1)^k} = \sum_{[x] \in \pi_0 X} \frac{|\pi_2(X, x)| \cdot |\pi_4(X, x)| \cdots}{|\pi_1(X, x)| \cdot |\pi_3(X, x)| \cdots}.$$

This satisfies a more general multiplicativity property: given a fibration  $F \rightarrow E \rightarrow B$  over a connected base  $B$ , the homotopy long exact sequence yields  $|E| = |F||B|$ . (This is more general because fiber bundles are “twisted” cartesian products.) This tells us, for instance, that

$$|BG| = \frac{1}{|G|}.$$

**Remark I.4.** What’s with the alternation in the product? **Crudely, it is a manifestation of an iterated inclusion-exclusion argument.** I think of this informally: the cardinality of the underlying set of  $X$  is the number of its points. But homotopy theory says to replace sets with 0-groupoids, in which case the cardinality is the number of connected components. For simplicity, suppose  $X$  is connected; then its underlying 1-groupoid has one point up to isomorphism, but that point may have automorphisms, accounted for by  $\pi_1 X$ , and we know that cardinality should be inversely proportional to this. But wait—those automorphisms “smaller.” But wait—the automorphisms of automorphisms have automorphisms, accounted for by  $\pi_3 X$ , which by parallel reasoning make  $|X|$  *smaller*. So on and so forth *ad infinitum*.

**Remark I.5.** We are implicitly assuming the defining series for  $|X|$  converges. Call spaces for which this is the case *tame*.

Recall that a space  $X$  is called *n-finite* if  $\pi_{k>n}X = 0$  and  $\pi_{k\leq n}X$  is finite, and that  $X$  is called  *$\pi$ -finite* if it is *n-finite* for some  $n$ . It is clear that  $\pi$ -finite spaces are tame. Thus,  $\pi$ -finite spaces seem like a good category of spaces in which to think about homotopy cardinality.

**Definition I.2.** Write  $\mathcal{S}_{\text{fin}}$  for the  $\infty$ -category generated by a point under finite colimits. We also consider  $\mathcal{S}_{n\text{-fin}}$  and  $\mathcal{S}_{\pi\text{-fin}}$ , the full subcategories of *n*- and  $\pi$ -finite spaces.

Homotopy cardinality defines a functor<sup>2</sup>  $\mathcal{S}_{\pi\text{-fin}} \rightarrow \mathbb{Q}_{\geq 0}$ . It is the unique extension of the cardinality of finite sets that is homotopy invariant, additive w.r.t. disjoint unions, and multiplicative w.r.t. fibrations. See the answer to my MO question.

## I.2 (1/4) Colimit completions and filtering classes

Here's a story I really like. Consider an ordinary category  $\mathcal{C}$ . Its presheaf category  $\text{PShv}(\mathcal{C}) := \text{Fun}(\mathcal{C}^{\text{op}}, \text{Set})$  is the “free completion at colimits” of  $\mathcal{C}$  in the following sense.

**Theorem I.1.** The Yoneda embedding  $\mathcal{Y} : \mathcal{C} \rightarrow \text{PShv}(\mathcal{C})$  is a free cocompletion of  $\mathcal{C}$ . That is,  $\text{PShv}(\mathcal{C})$  has small colimits, and if  $\mathcal{D}$  admits small colimits, then restriction along  $\mathcal{Y}$  defines a natural equivalence

$$\mathcal{Y}^* : \text{Fun}^{\text{ccts}}(\text{PShv}(\mathcal{C}), \mathcal{D}) \xrightarrow{\sim} \text{Fun}(\mathcal{C}, \mathcal{D}).$$

This is standard. I think it also characterizes  $\mathcal{Y}$  (and hence  $\text{PShv}(\mathcal{C})$ ) by the usual “thing satisfying universal property is unique” argument. Here is another characterization.

**Definition I.3.** Consider a presheaf  $F : \mathcal{C}^{\text{op}} \rightarrow \text{Set}$ . Its *category of elements*  $\text{el } F$  has (objects) transformations  $\mathcal{Y}c \rightarrow F$  and (arrows) transformations  $\mathcal{Y}c \rightarrow \mathcal{Y}d$  such that the evident triangle commutes.

**Proposition I.1** (Density). There is a canonical map  $\text{el } F \rightarrow \mathcal{C}$ . Its colimit canonically presents  $F$ , i.e.,

$$\text{colim}(\text{el } F \rightarrow \mathcal{C} \rightarrow \text{PShv}(\mathcal{C})) \cong F.$$

The density theorem canonically associates to each presheaf  $F : \mathcal{C}^{\text{op}} \rightarrow \text{Set}$  a diagram of representables  $\text{el } F \rightarrow \text{PShv}(\mathcal{C})$ , which we equivalently regard as its underlying diagram  $\text{el } F \rightarrow \mathcal{C}$ . I think you can upgrade this to say “the category  $\text{PShv}(\mathcal{C})$  is *equivalent* to the category of diagrams in  $\mathcal{C}$ .”

We have roughly provided three equivalent definitions of  $\text{PShv}(\mathcal{C})$ : it is (I) the functor category  $\text{Fun}(\mathcal{C}^{\text{op}}, \text{Set})$ , (II) the category of diagrams in  $\mathcal{C}$ ,<sup>3</sup> and (III) the free colimit completion of  $\mathcal{C}$ . The equivalence (I)  $\rightarrow$  (III) associates to  $F$  its diagram  $\text{el } F \rightarrow \mathcal{C}$ . The equivalence (III)  $\rightarrow$  (I) takes the colimit of the diagram. Either can be shown to fit the description (II).

**Remark I.6.** For a presheaf  $F$ , its category of elements can also be defined as the pullback below. The map  $\text{el } F \rightarrow \mathcal{C}$  is again evident. Perhaps because of this definition, we sometimes write  $\mathcal{C}/F := \text{el } F$ .

$$\begin{array}{ccc} \text{el } F & \dashrightarrow & \text{PShv}(\mathcal{C})/F \\ \downarrow \pi_F & \lrcorner & \downarrow \\ \mathcal{C} & \xrightarrow{\mathcal{Y}} & \text{PShv}(\mathcal{C}) \end{array}$$

**Remark I.7.** The category of elements  $\mathcal{C}/F$  can also be defined as the *comma category*  $\ast \downarrow F$ . I honestly do not know about comma categories and am not sure I need to know about them right now.

---

Often, we want to think about the completion  $\mathcal{C}$  at a *certain class* of colimits—say, filtered or sifted colimits. We described the colimit completion  $\text{PShv}(\mathcal{C})$  in three ways (I), (II), and (III) above; these suggest three ways to complete at a *chosen class* of colimits. Namely, the completion of  $\mathcal{C}$  at a “nice” colimits should be...

<sup>2</sup>I don't think this is actually a functor. It is a function from the set of equivalence classes of  $\pi$ -finite Kan complexes to  $\mathbb{Q}_{\geq 0}$ .

<sup>3</sup>The description (III) needs more attention than we have provided. First of all, you want *small* diagrams. Then you must define morphisms. This requires some care. Will I get around to typing this out?

- (I) A subcategory of “nice” presheaves in  $\text{Fun}(\mathcal{C}^{\text{op}}, \text{Set})$ ;
- (II) A subcategory of “nice” diagrams in  $\mathcal{C}$ ; and
- (III) The “free nice colimit completion” of  $\mathcal{C}$  (in the sense of a universal property).

All three models are useful. For example, *filtered colimits* are easiest to define as colimits over *filtered diagrams*, and this diagrammatic definition suggests we define the *ind-completion*  $\text{Ind}(\mathcal{C})$ <sup>4</sup> as the category of filtered diagrams in  $\mathcal{C}$ , in the likeness of (II).<sup>5</sup> (They are equivalently the diagrams whose colimits in  $\text{Set}$  commute with finite limits.) Meanwhile, *sifted colimits* are not so easily described by the shape<sup>6</sup> of their diagram; we say a diagram is *sifted* if its colimits commute with finite products (in  $\text{Set}$ ). This suggests a definition of the sifted completion in the likeness of (I): when  $\mathcal{C}$  has finite (co?)products, I think you can define its sifted completion as the full subcategory of presheaves commuting with finite products. (You can also define it as the subcategory of sifted diagrams in  $\mathcal{C}$ , but I’m trying to make the point that (I) and (II) are both natural.)

**Remark I.8** (Further reading). I just said lots of stuff, but maybe left out lots of stuff. A little while ago, I gave a detailed account of the above picture in Lecture 6 of our condensed seminar, notes here.

I have not really thought about colimit completions for  $\infty$ -categories before. But, it seems the ordinary picture persists verbatim. Let me spell out an extended example, the motivating example in Charles’ paper [Rez22], which is what made me start thinking about colimit completions again.

**Example I.3** (Completion at  $\kappa$ -filtered colimits). For a regular cardinal  $\kappa$ , an  $\infty$ -category  $\mathcal{C}$  is called  *$\kappa$ -filtered* if every  $\kappa$ -small simplicial set diagram  $K \rightarrow \mathcal{C}$  admits an extension  $K^{\triangleright} \rightarrow \mathcal{C}$  [Lur08, 5.3.1.7]. Parallel to the ordinary case,  $\kappa$ -filteredness is equivalent to commutation with all small limits in  $\text{Space}$  [Lur08, 5.3.3.3]. It turns out that  $\kappa$ -smallness is a generally well-behaved property (e.g., it is preserved by categorical equivalences), enough so that we may *complete*  $\mathcal{C}$  at  $\kappa$ -small colimits to obtain  $\text{Ind}_{\kappa}(\mathcal{C})$ . Also parallel to the ordinary picture,  $\text{Ind}_{\kappa}(\mathcal{C})$  admits models in the likeness of (I), (II), and (III) above:

- (I) In terms of presheaves,  $\text{Ind}_{\kappa}(\mathcal{C})$  is the full subcategory of  $\text{PShv}(\mathcal{C})$  spanned by filtered colimits of representables [Lur08, 5.3.5.4]. Furthermore, if  $\mathcal{C}$  admits small colimits, then  $\text{Ind}_{\kappa}$  can be more concretely described as the full subcategory spanned by presheaves preserving finite limits.
- (II) In terms of point categories,  $\text{Ind}_{\kappa}(\mathcal{C})$  consists of diagrams  $J \rightarrow \mathcal{C}$  such that  $J$  is a  $\kappa$ -filtered simplicial set(?).
- (III) In terms of its universal property,  $\text{Ind}_{\kappa}(\mathcal{C})$  admits  $\kappa$ -filtered colimits, Yoneda factors as  $\mathcal{Y} : \mathcal{C} \rightarrow \text{Ind}_{\kappa}(\mathcal{C})$ , and if  $\mathcal{D}$  admits  $\kappa$ -filtered colimits, then restriction along  $\mathcal{Y}$  defines an equivalence

$$\text{Fun}_{\kappa}(\text{PShv}(\mathcal{C}), \mathcal{D}) \xrightarrow{\sim} \text{Fun}(\mathcal{C}, \mathcal{D}).$$

In other words, if  $\mathcal{D}$  admits  $\kappa$ -filtered colimits, then each arrow  $\mathcal{C} \rightarrow \mathcal{D}$  admits an essentially unique extension to  $\text{Ind}_{\kappa}(\mathcal{C})$  [Lur08, 5.3.5.10].

**Remark I.9.** Lurie’s verbatim definition [Lur08, 5.3.5.1] is “ $\text{Ind}_{\kappa}(\mathcal{C})$  is the full subcategory of  $\text{PShv}(\mathcal{C})$  spanned by presheaves  $F$  which classify right fibrations  $\mathcal{C}/F \rightarrow \mathcal{C}$  such that  $\mathcal{C}/F$  is  $\kappa$ -filtered.” I *feel* like this is just (I) and (II) above at the same time. Lurie basically says in [Lur08, 5.3] that he exhibits model (II), but he does not seem to exhibit that explicitly. But I actually do not know anything about (un)straightening, so I cannot really navigate.

<sup>4</sup>Filtered colimits used to be called inductive limits, so we call the completion of  $\mathcal{C}$  at filtered colimits its “ind-completion” or “indization.”

<sup>5</sup>Then you can recharacterize it in the likeness of (I) and (III). For (I), it turns out  $\text{Ind}(\mathcal{C})$  is the full subcategory of  $\text{PShv}(\mathcal{C})$  spanned by filtered colimits of representable presheaves. If  $\mathcal{C}$  is finitely cocomplete, so that  $\mathcal{C}^{\text{op}}$  has finite limits, then we can be more explicit:  $\text{Ind}(\mathcal{C})$  is precisely those presheaves commuting with finite colimits. For (III), take the universal property of  $\text{PShv}(\mathcal{C})$  but replace “cocontinuous” with “preserves filtered colimits.”

<sup>6</sup>I think there is such a description, but it is not nice.

**Remark I.10.** In the ordinary case, it is standard to characterize (say) ind-objects by their point category, i.e., to say that an ind-object is a filtered diagram in  $\mathcal{C}$ , equivalently a presheaf  $\mathcal{C}^{\text{op}} \rightarrow \text{Set}$  whose category of elements is filtered (and cofinal) [KS06, 3.3.13, 6.1.5].

---

I regard Lurie’s definition of  $\text{Ind}_{\mathcal{F}}(\mathcal{C})$  as an amalgam of models (I) and (II). These are “concrete” models which one shows have the relevant universal property. Charles’ paper [Rez22] considers the general question: **for which classes of  $\infty$ -categories  $\mathcal{F}$  do these “concrete” models still work (i.e., actually have the relevant universal property)?** More precisely, which classes of small  $\infty$ -categories  $\mathcal{F}$  are such that for every  $\infty$ -category  $\mathcal{C}$ , the category  $\{X \in \text{PShv}(\mathcal{C}) : \mathcal{C}/X \in \mathcal{F}\}$  has the “free  $\mathcal{F}$ -colimit completion” property?

For this purpose, we make two definitions. First, we define  $\text{Ind}_{\mathcal{F}}(\mathcal{C}) := \{X \in \text{PShv}(\mathcal{C}) : \mathcal{C}/X \in \mathcal{F}\}$ . Second, we define  $\text{PShv}_{\mathcal{F}}(\mathcal{C})$  as the *minimal full subcategory of  $\text{PShv}(\mathcal{C})$  generated by representables under  $\mathcal{F}$ -colimits*, to mean the smallest full subcategory containing  $\mathcal{Y}(\mathcal{C}) \subseteq \text{PShv}(\mathcal{C})$  and closed under  $\mathcal{F}$ -colimits. Yoneda factors as  $\mathcal{Y} : \mathcal{C} \rightarrow \text{PShv}_{\mathcal{F}}(\mathcal{C})$ , and this functor is a *free  $\mathcal{F}$ -colimit completion* [Lur08, 5.3.6.2]. **Then we seek to compare  $\text{Ind}_{\mathcal{F}}(\mathcal{C})$  and  $\text{PShv}_{\mathcal{F}}(\mathcal{C})$ .**

This is not complicated. First, note that  $\text{Ind}_{\mathcal{F}}(\mathcal{C}) \subseteq \text{PShv}_{\mathcal{F}}(\mathcal{C})$  since any presheaf  $X \in \text{PShv}(\mathcal{C})$  is a colimit of  $\mathcal{C}/X \rightarrow \mathcal{C} \rightarrow \text{PShv}(\mathcal{C})$ , thus  $X$  is an  $\mathcal{F}$ -colimit of representables as soon as  $\mathcal{C}/X \in \mathcal{F}$ . **Now, if we can show that  $\text{Ind}_{\mathcal{F}}(\mathcal{C}) \subseteq \text{PShv}_{\mathcal{F}}(\mathcal{C})$  contains  $\mathcal{Y}(\mathcal{C})$  and is stable under  $\mathcal{F}$ -colimits, then we get the reverse inclusion and conclude**

$$\text{Ind}_{\mathcal{F}}(\mathcal{C}) = \text{PShv}_{\mathcal{F}}(\mathcal{C}).$$

We consider sufficient conditions for  $\text{Ind}_{\mathcal{F}}(\mathcal{C})$  to obtain these properties.

- For which  $\mathcal{F}$  does  $\text{Ind}_{\mathcal{F}}(\mathcal{C})$  always contain the representables? For an object  $c$ , we have  $\mathcal{C}/\mathcal{Y}c \cong \mathcal{C}_{/c}$ . Hence, you could ask that  $\mathcal{F}$  contains all  $\infty$ -categories with a terminal object. Alternatively(?): there  $c$  is terminal, so  $\mathcal{Y}c$  is a terminal presheaf. By definition, we in fact have  $\mathcal{Y}c \in \text{PShv}_{\mathcal{F}}(\mathcal{C})$ . **Hence, if  $\mathcal{F}$  contains  $\mathbf{D}$  whenever  $\text{PShv}_{\mathcal{F}}(\mathbf{D})$  contains a terminal presheaf, then we would have  $\mathcal{Y}c \in \text{Ind}_{\mathcal{F}}(\mathcal{C})$ .**
- I have nothing to say about when  $\text{Ind}_{\mathcal{F}}(\mathcal{C})$  is stable under  $\mathcal{F}$ -colimits. See the proof of Prop. 4.2 in Charles paper [Rez22].

We say that  $\mathcal{F}$  is **filtering** if it contains  $\mathbf{D}$  whenever  $\text{PShv}_{\mathcal{F}}(\mathbf{D})$  contains a terminal presheaf. We seek this property (rather than just ask that  $\mathcal{F}$  contain  $\mathbf{D}$  whenever  $\mathbf{D}$  has a terminal object) because filtration turns out to be closely related to our problem:

**Proposition I.2.** Suppose that a family of  $\infty$ -categories  $\mathcal{F}$  is such that for any  $\mathcal{C}$ , the full subcategory inclusion  $\text{Ind}_{\mathcal{F}}(\mathcal{C}) \subseteq \text{PShv}_{\mathcal{F}}(\mathcal{C})$  is an isomorphism. Then  $\mathcal{F}$  is filtering.

*Proof.* If  $1 \in \text{PShv}(\mathbf{D})$  denotes a terminal presheaf, then  $\mathbf{D}/1 \cong \mathbf{D}$ . Therefore<sup>7</sup>  $\mathbf{D}/1 \cong \mathbf{D}$  lies in  $\text{PShv}_{\mathcal{F}}(\mathbf{D})$  and thus  $\text{Ind}_{\mathcal{F}}(\mathbf{D})$  by assumption. This exactly says that  $\mathcal{F}$  is filtering.  $\square$

To summarize, we explained that if  $\mathcal{F}$  is filtering, then  $\text{Ind}_{\mathcal{F}}(\mathcal{C}) \subseteq \text{PShv}_{\mathcal{F}}(\mathcal{C})$  is an equivalence, and we proved the converse. **Hence, given a class of  $\infty$ -categories  $\mathcal{F}$ , the “diagrammatic” model for the  $\mathcal{F}$ -colimit completion  $\text{Ind}_{\mathcal{F}}(\mathcal{C})$  (presheaves whose point category belong to  $\mathcal{F}$ ) is an actual  $\mathcal{F}$ -colimit completion (in the sense of possessing the relevant universal property, equivalently  $\text{Ind}_{\mathcal{F}}(\mathcal{C}) \hookrightarrow \text{PShv}_{\mathcal{F}}(\mathcal{C})$  being an equality) if and only if  $\mathcal{F}$  is filtering.** I wonder what the diagrammatic model is good for in general. Charles also studies the notion of filtering classes, which I have not read yet. May also be interesting to look at [Du23].

**Remark I.11.** In light of this writing, my mind has changed to consider there as being only *two* truly distinct models: a *diagrammatic* model and a *universal property* model. Maybe it’s not right to even call the latter a model, since it’s really what we’re trying to *model*, but that seems to be the language I’ve found most efficient.

**Remark I.12.** Someone should rename *filtering classes*. But I am not sure what a good name would be. I do not like *filtering* because we already have *filtered/filtering/filtrant* colimits, which are a very specific example of filtering classes. Something to reflect that  $\mathcal{F}$ -diagrams are special. Diagrammatic? Graphical? Figurative? “Graphical classes” has a nice ring to it. Or just call them “graphics.” Or “complete graphics.”

<sup>7</sup>Hmm. Why does  $\mathbf{D}/1$  lie in  $\text{PShv}_{\mathcal{F}}(\mathbf{D})$ ? Does

### I.3 (1/15) Reminding myself what presentability, dualizability, and stability are

Classes just started at UIUC, but I am in Chicago while Ishan and Efimov are giving lectures. Today I want to define a category  $\mathbf{Pr}_{\text{st}}^{\text{dual}}$  which contains the category  $\mathbf{Cat}_{\infty}^{\text{perf}} = \{\text{small, stable, idempotent-complete } \infty\text{-categories}\}$  and out of which we can define *continuous* (“Efimov”) *K-theory*  $\mathbf{Pr}_{\text{st}}^{\text{dual}} \rightarrow \mathbf{Sp}$ , extending the algebraic *K*-theory. I have not actually thought about presentability or dualizability before, nor how they interact with stability. I’m going to review these quickly. Here are some things I am looking at:

- Peter Haine, *Descent for sheaves on compact Hausdorff spaces* here.
- Hoyois’ continuous *K*-theory notes here.
- Parts of Mortiz Groth’s notes here, particularly the bits about presentability.
- Dustin Clausen’s lectures about Efimov *K*-theory (on Youtube).
- He Li’s Efimov *K*-theory notes here.
- Chapter 5 of HTT [Lur08].
- Alberto García-Raboso’s notes on stable  $\infty$ -categories.
- Yonatan’s notes on stable  $\infty$ -categories.

---

First I want to remember what presentability is. In nice cases, colimits are like unions. For instance, a group is the colimit of its directed poset of subgroups. In fact, you can just take that of its *finitely generated* subgroups. We can actually make a categorical equivalence

$$\text{Ind}(\text{Ab}_{\text{fg}}) \cong \text{Ab},$$

the point being that although  $\text{Ab}$  is large, it admits a small subcategory of ind-generators. This should be considered an essential detail of the structure of  $\text{Ab}$ , and it is both practically and philosophically important.

Toward pinning down this “big thing is secretly small and this helps us to work with it” idea, consider the *adjoint functor theorem*: a functor  $F : \mathbf{A} \rightarrow \mathbf{B}$  between cocomplete categories admits a right adjoint if and only if  $F$  preserves colimits and satisfies the *solution set condition*. Without going into details, that condition is a certain smallness condition—then is reasonable to think that it is automatically fulfilled whenever  $\mathbf{A}, \mathbf{B}$  are themselves “small enough.”

**Definition I.4.** Say that an  $\infty$ -category  $\mathbf{C}$  is *presentable* if it is cocomplete and *accessible*: there exists a regular cardinal  $\kappa$  such that  $\mathbf{C}$  has  $\kappa$ -filtered colimits and admits a small subcategory  $\mathbf{C}_0$  such that  $\text{Ind}_{\kappa}(\mathbf{C}_0) \cong \mathbf{C}$ .

**Theorem I.2** (Generalized adjoint functor theorem). A functor between presentable, cocomplete  $\infty$ -categories admits a right adjoint if and only if it preserves colimits.

**Remark I.13.** Since  $\mathcal{Y} : \mathbf{D} \rightarrow \text{Ind}(\mathbf{D})$  is fully faithful, accessibility amounts to the admittance of a small subcategory of ind-generators. Also, presentable categories admit *bilimits*.

**Remark I.14.** See [Lur08, §5.4] for a detailed review of accessibility.

Presentability is not a rare property, and it is generally well-behaved. It can be characterized in terms of presheaves:

**Theorem I.3.** An  $\infty$ -category  $\mathbf{C}$  is presentable if and only if  $\mathbf{C}$  is an *accessible localization* of  $\mathbf{PShv}(\mathbf{D})$  for some small  $\mathbf{D}$ : there exists a functor  $F : \mathbf{PShv}(\mathbf{D}) \rightarrow \mathbf{C}$  such that (loc.)  $F$  admits a fully faithful right adjoint and (acs.)  $F$  commutes with all  $\kappa$ -filtered colimits for some regular  $\kappa$ .

I feel as if I should have something to say about this, but I do not. An idea: let  $\mathbf{D}$  be small and regard the free colimit completion  $\mathbf{PShv}(\mathbf{D})$  as a “free, small collection of generators.” Next, think of an accessible localization  $\mathbf{PShv}(\mathbf{D}) \rightarrow \mathbf{C}$  as “imposing a small number of relations.” Then our characterization seems natural: *presentable categories are those with a “small” presentation by generators and relations*. It is unclear to me how far one can expound this idea, what the role of accessibility is for the localization insofar as it makes this idea work, ... I do not want to waste time on these details right now.

Return to  
this.



**Definition I.5.** We organize presentable  $\infty$ -categories into an  $\infty$ -category. Its objects are (not necessarily small) presentable  $\infty$ -categories and its morphisms are the cocontinuous functors [Lur08, 5.5.3.1]. By the adjoint functor theorem, we can also say this category has left adjoints as morphisms. We write  $\mathbf{Pr}^L$  for this category.

Misc properties.

**Proposition I.3** ([Lur08, 5.5.3.8]). If  $\mathcal{C}, \mathcal{D}$  are presentable, then the full subcategory  $\mathrm{Fun}^L(\mathcal{C}, \mathcal{D}) \subset \mathrm{Fun}(\mathcal{C}, \mathcal{D})$  spanned by left adjoints is presentable. (In fact,  $\mathcal{C}$  can be any simplicial set.)

**Proposition I.4.** If  $\mathcal{C}, \mathcal{D}$  are presentable, then there exists a presentable category  $\mathcal{C} \otimes \mathcal{D}$  which is the universal recipient of a functor from  $\mathcal{C} \times \mathcal{D}$  that is colimit-preserving in both variables separately. One presentation is  $\mathcal{C} \otimes \mathcal{D} \cong \mathrm{Fun}^{cts}(\mathcal{C}^{\mathrm{op}}, \mathcal{D})$ . It inherits this (symmetric monoidal) tensor product as a full subcategory (on the presentables) of  $\hat{\mathrm{Cat}}_\infty(K)$ , the  $\infty$ -category of  $\infty$ -categories with small colimits and colimit-preserving functors. (Which has the Lurie tensor product?) Maybe c.f. [Lur17, 4.8.1.5]. See also nLab.

Ok, now dualizability. In a monoidal category  $\mathcal{C}$ , an object  $x \in \mathcal{C}$  is called *dualizable* if (existence of “dual” with co/evaluation maps). We consider this in the case of vector spaces.

**Example I.4.** Consider a vector space  $V \in \mathrm{Vect}_k$ . A candidate for its dual is  $V^* = \mathrm{Hom}(V, k)$  and there is an obvious map  $\mathrm{ev} : V \otimes V^* \rightarrow k$ . We would like a map  $\mathrm{coev} : k \rightarrow V \otimes V^*$ , which amounts to the choice of an element  $v \in V \otimes V^*$ . If  $(e_i)_I$  is a basis for  $V$ , we have coordinates  $v = \sum c_{i,j} e_i \otimes e_j^*$  such that only finitely many  $c_{i,j}$  are nonzero. The axioms for co/evaluation imply that for each  $i \in I$ , some  $c_{i,j}$  is nonzero, hence  $I$  must be finite, i.e.,  $V$  must be finite-dimensional. Conversely, if  $V$  is finite-dimensional, then it is dualizable with dual  $V^*$ , in fact we may identify  $V \otimes V^* \cong \mathrm{End}(V)$  (which requires a choice of basis) and define  $\mathrm{coev}$  by  $1_k \mapsto \mathrm{id}_V$ .

As this suggests, dualizability is a sort of finiteness condition. But, apparently different from that of presentability. Dualizable objects inherit some of the theory of finite-dimensional vector spaces, for instance a notion of *traces of endomorphisms* and *dimension* (the trace of the identity).

**Remark I.15.** I think that if  $\mathcal{C}$  is monoidal with internal homs, then an object  $X \in \mathcal{C}$  is dualizable if and only if the canonical pairing  $X \otimes \mathrm{Hom}(X, 1) \rightarrow \mathrm{End}(X)$  is an isomorphism. The category  $\mathrm{Vect}_k$  has internal homs, and we saw that the map is an isomorphism therein, so this checks out. Also in  $\mathrm{Vect}_k$ , we know that for non-dualizable (equivalently, infinite dimensional)  $V$ , that  $V \otimes V^*$  is nicer than  $\mathrm{End}(V)$  in general. In other monoidal categories *without* internal homs, I wonder if you can ever treat  $X \otimes X^*$  like a well-behaved substitute for  $\mathrm{End}(X)$ . I had this thought during coffee with Anthony and Sam and they asked, “have you heard of a star-autonomous category?”

We want to think about dualizability in the full subcategory  $\mathbf{Pr}_{\mathrm{st}} \subset \mathbf{Pr}^L$  spanned by the presentable, stable  $\infty$ -categories. A good question is, why are we making the stability hypothesis?

Recall that an  $\infty$ -category is *stable* if it has a zero object, has fibers and cofibers, and fiber sequences coincide with cofiber sequences. I can weakly explain where this comes from: even in classical homotopy theory, we care deeply about pushouts, pullbacks, and (co)fibrations. These are tricky notions when you work with spaces (not intractable), partly due to the rebellious nature of homotopy (co)limits in the classical stable homotopy category. This gives some impetus for *stable* homotopy theory: stable phenomena somehow simplify the story. I wish I could give a concrete, classical example of this, but all I can think of is “fibrations and cofibrations of spectra coincide.” Maybe some helpful discussion here and here.<sup>8</sup>

**Remark I.16.** A functor between stable categories is called *reduced* if it preserves zero, and *exact* if in addition it preserves (co)fiber sequences. Since the zero object and (co)fibration sequences are (co)limits, it is preserved by left adjoints, hence these properties are superfluous in  $\mathbf{Pr}^L$  and  $\mathbf{Pr}^R$ .

Given a pointed category  $\mathcal{C}$ , it has a *suspension functor*  $\Sigma_{\mathcal{C}}$  once it has cofibers, and it has a *loop functor*  $\Omega_{\mathcal{C}}$  once it has fibers. Stability is characterized by either suspension or looping being an equivalence. This presents an idea: does formally inverting  $\Sigma_{\mathcal{C}}$  or  $\Omega_{\mathcal{C}}$  present a “stabilization”? Note that  $\text{Cat}^{\text{ex}}$  occurs as a full subcategory of  $\text{Cat}_*^{\text{fincolim}}$  (resp.  $\text{Cat}_*^{\text{finlim}}$ ), spanned by those categories whose suspension (resp. loop) functor is an equivalence. “Stabilization” should mean an adjoint to these inclusions.

These inclusions in fact have left and right adjoints, and we get the left adjoints in the manner described.

**Proposition I.5.** Let  $\mathcal{C}$  be pointed. If  $\mathcal{C}$  has finite colimits (cofibers and suspension in particular), then the colimit  $\text{Sp}^{\Sigma}(\mathcal{C}) := \text{colim}(\mathcal{C} \xrightarrow{\Sigma} \mathcal{C} \rightarrow \dots) \in \text{Cat}_*^{\text{fincolim}}$  is stable, and this extends to a left adjoint to the inclusion  $\text{Cat}^{\text{ex}} \hookrightarrow \text{Cat}_*^{\text{fincolim}}$ . Dually, if  $\mathcal{C}$  has finite limits, we get a left adjoint to  $\text{Cat}^{\text{ex}} \hookrightarrow \text{Cat}_*^{\text{finlim}}$  given by  $\mathcal{C} \mapsto \text{Sp}^{\Omega}(\mathcal{C})$ .

Rather strangely(?), even if  $\mathcal{C}$  is (finitely) complete and cocomplete, their suspension-spectra and loop-spectra are not generally equivalent. That is, we do not always have  $\text{Sp}^{\Sigma}(\mathcal{C}) \cong \text{Sp}^{\Omega}(\mathcal{C})$ . For example, finite spectra and  $\Omega$ -spectra do not coincide. But...

**Observation I.2.** Presentability puts us somewhere nice: we get *all* limits and colimits, for instance. Also, the natural notion of morphisms of presentable categories (limit or colimit preserving, you choose) already fits that for categories we usually consider stabilizing (finite limit or colimit preserving, you choose). And if your presentable categories were already stable, either notion is automatically an exact functor.

Hence, at least formally, it seems easy and convenient to consider *presentable* stable  $\infty$ -categories. Since presentable demands (co)limits, the characterization of stability simplifies: *if  $\mathcal{C}$  is a pointed, presentable  $\infty$ -category, then  $\Sigma_{\mathcal{C}}, \Omega_{\mathcal{C}}$  are defined, and  $\mathcal{C}$  is stable iff either of these is an equivalence.*

(Stuff; universal “stabilization” property; the realization of  $\Omega$ -spectra as the finite shifts of finite suspension spectra; the definition of the  $\infty$ -category of spectra; the definition of its symmetric monoidal *smash product*!)

## I.4 (1/24) Stabilization, the $\infty$ -category of spectra, and the smash product

My notes previously got me thinking about stability, and then Charles told me some things about presentability and the smash product on the  $\infty$ -category of spectra, so now I am going to think a bit about all that. I have already defined *stable  $\infty$ -categories* at least twice, which behave “like spectra,” or maybe “like chain complexes of abelian groups.” Last time, I also thought a bit about how stability simplifies in the presence of presentability (although it seems I didn’t actually write about that). This makes sense, since presentability forces existence of (co)limits, and stability cares about certain (co)limits (and needs them for the essential  $\Sigma$  and  $\Omega$ ). But that does not do justice to the fun which presentability brings to the party. I’m going to be primarily reading Groth’s notes.

For a pointed category  $\mathcal{C}$ , recall the definition of *triangles*: they are the composable pairs of morphisms  $(g, f)$  together with 2-cells realizing  $gf \simeq h$  and  $h \simeq 0_{X,Y}$ . These form a category, namely the full subcategory of  $\text{Fun}(\Delta^1 \times \Delta^1, \mathcal{C})$  spanned by functors mapping the “bottom-left” vertex to 0. Consider that  $\Delta^1 \times \Delta^1$  is both a left and right cone [Lur24, Tag 0165]:

$$(\Lambda_0^2)^{\triangleleft} \cong \Delta^1 \times \Delta^1 \cong (\Lambda_2^2)^{\triangleright}.$$

We say a triangle is *exact* if it is a limit as a left cone, and *coexact* if it is a colimit as a right cone. If  $\mathcal{C}$  admits all finite (co)limits, we denote by

$$\mathcal{C}^{\Sigma} \subseteq \text{Fun}(\Delta^1 \times \Delta^1, \mathcal{C}) \quad \text{and} \quad \mathcal{C}^{\Omega} \subseteq \text{Fun}(\Delta^1 \times \Delta^1, \mathcal{C})$$

the full subcategories spanned by coexact and exact triangles *with the bottom-left and top-right corners zero*, respectively. To form a suspension  $\Sigma X$ , no data is needed beyond specifying  $X$ ; diagrammatically, a functor  $F \in \mathcal{C}^{\Sigma}$  should be determined by its top-left corner. Dually for  $\Omega X$ . Abstract nonsense says that indeed,  $\text{ev}_{(0,0)} : \mathcal{C}^{\Sigma} \rightarrow \mathcal{C}$  and  $\text{ev}_{(1,1)} : \mathcal{C}^{\Omega} \rightarrow \mathcal{C}$  are acyclic Kan fibrations.

<sup>8</sup>Think about these. Maybe think about connectivity results.

Yonatan’s notes have a bit about suspension and the triangulated structure for stable categories. Maybe read this later.

ponder

Probably give this its own day.

**Definition I.6.** Suppose that  $\mathcal{C}$  is pointed and admits finite (co)limits. Since  $\text{ev}_{0,0} : \mathcal{C}^\Sigma \rightarrow \mathcal{C}$  is acyclic, it admits a section  $s_\Sigma : \mathcal{C} \rightarrow \mathcal{C}^\Sigma$ , well-defined up to a contractible choice. We can thus define (up to a contractible choice) the *suspension functor*  $\Sigma_{\mathcal{C}} = s_\Sigma \circ \text{ev}_{(0,0)}$ . We define the *loops functor*  $\Omega_{\mathcal{C}}$  identically.

The functors  $\Sigma_{\mathcal{C}}, \Omega_{\mathcal{C}}$  are adjoint. Furthermore, if  $\mathcal{C}$  is stable (i.e., exact triangles = coexact triangles), then they are inverse equivalences. The converse is also true.

We are interested in stabilizing categories. As stability is characterized by  $\Omega_{\mathcal{C}}$  being an equivalence, this means inverting  $\Omega_{\mathcal{C}}$ , i.e. by taking the colimit of  $\cdots \rightarrow \mathcal{C} \rightarrow \mathcal{C}$ . That is categorical. But there's an analogy here with algebraic topology, and we wonder about forming “spectrum objects” in  $\mathcal{C}$ . In fact, we can do this, giving a more explicit model for the stabilization. We can apply this in the case of spaces—the categorical properties of the stabilization will give us the *symmetric monoidal smash product* of spectra, and the description as “spectrum objects” will make clear that we are talking *actual* spectra, the kind we care about from classical algebraic topology.

Let  $\mathcal{C}$  be pointed and finitely (co)complete. A *prespectrum* in  $\mathcal{C}$  is a functor  $X : \mathbb{N} \times \mathbb{N} \rightarrow \mathcal{C}$  such that  $X(i, j) = 0$  whenever  $i \neq j$ . A prespectrum  $X$  determines a sequence of triangles in  $\mathcal{C}$ , and thus maps

$$\Sigma X_n \rightarrow X_{n+1} \quad \text{and} \quad X_n \rightarrow \Omega X_{n+1}.$$

We say that  $X$  is a *spectrum* if all the maps  $X_m \rightarrow \Omega X_{m+1}$  are equivalences. We say that  $X$  is a *spectrum below  $n$*  if that is true for  $m < n$ . The following theorem says that (with mild hypotheses) spectrum objects model the stabilization.

**Theorem I.4** (DAG I, 8.14). For an arbitrary  $\infty$ -category  $\mathcal{C}$ , we define its *stabilization*  $\text{Stab}(\mathcal{C}) := \text{Sp}(\mathcal{C}_*)$ . Note that if  $\mathcal{C}$  is pointed, then  $\mathcal{C}_* \rightarrow \mathcal{C}$  is a trivial Kan fibration, and so is the induced  $\text{Stab}(\mathcal{C}) \rightarrow \text{Sp}(\mathcal{C})$ . If  $\mathcal{C}$  is pointed with finite limits, then

$$\text{Sp}(\mathcal{C}) \cong \text{colim}(\cdots \xrightarrow{\Omega} \mathcal{C} \xrightarrow{\Omega} \mathcal{C}).$$

**Remark I.17.** This presentation follows Moritz’s notes and Lurie’s DAG. However, Lurie’s presentation in HA is different. There, he wants to faithfully reenact the story of *infinite loop spaces* (in particular, Brown representability and excision for cohomology) in the setting of  $\infty$ -categories. For  $\mathcal{C}$  with finite limits, he defines  $\text{Sp}(\mathcal{C}) := \text{Fun}(\mathbb{S}_*^{\text{fin}}, \mathcal{C}) \subseteq \text{Fun}(\cdots)$  the full subcategory of reduced, excisive functors [Lur17, 1.4.2.8]. These approaches are equivalent. Given such a functor  $X$ , one may consider  $X_n := X(\mathbb{S}^n)$ , and we get equivalences  $X_n \xrightarrow{\sim} \Omega X_{n+1}$  by considering excision for the following pushout diagram.

$$\begin{array}{ccc} S^n & \xrightarrow{\quad} & * \\ \downarrow & & \downarrow \\ * & \xrightarrow{\quad} & S^{n+1} \end{array}$$

There are forgetful functors  $\text{Sp}(\mathcal{C}) \rightarrow \text{Sp}_{\leq n}(\mathcal{C}) \rightarrow \text{PSp}(\mathcal{C})$ . We can ask about adjoints. Let’s first ask about finding  $\text{PSp}(\mathcal{C}) \rightarrow \text{Sp}_{\leq n}(\mathcal{C})$ . Given a prespectrum  $X$ , one may imagine that “ $L_n(X)$  := the free  $\Omega$ -spectrum below  $n$  on  $X$ ” should be  $X[k]$  in degrees  $k \geq n$  and should be  $\Omega^k X[n]$  in degrees  $k < n$ . Heuristically,  $L_n(X)$  should be determined by the part of  $X$  in degrees  $\geq n$ , obtained by just chopping off lower degrees and refilling them by looping downward.

Lurie spells all this out in [Lur09, §8] (and these approximations  $L_n X$  are needed to prove 8.14 above). The basic idea is as follows. For  $-\infty \leq a \leq b \leq \infty$ , we define

$$Q(a, b) := \{(i, j) : i \neq j \text{ or } a \leq i = j \leq b\} \subseteq \mathbb{N} \times \mathbb{N}.$$

We want to define  $L_n(X) := \text{Ran}_{Q(n, \infty) \hookrightarrow \mathbb{N} \times \mathbb{N}}(X|_{Q(n, \infty)})$ . Since that inclusion  $Q(n, \infty) \hookrightarrow \mathbb{N} \times \mathbb{N}$  is quite “finite,” this is possible as soon as  $\mathcal{C}$  has finite limits.

**Proposition I.6** ([Lur09, 8.12]).  $\mathcal{C}$  pointed with finite limits,  $X_0 \in \mathrm{PSp}_a^\infty(\mathcal{C})$ . Then

- (1) There exists  $X \in \mathrm{PSp}_{a-1}^\infty(\mathcal{C})$  which is a right Kan extension of  $X_0$ .
- (2) There exists  $X \in \mathrm{PSp}_{-\infty}^\infty(\mathcal{C})$  which is a right Kan extension of  $X_0$ .
- (3) An object  $X \in \mathrm{PSp}_{-\infty}^\infty(\mathcal{C})$  is a right Kan extension of  $X_0$  if and only if  $X$  is a spectrum below  $a$ .

**Remark I.18.** Lurie also states a characterization of  $X \in \mathrm{PSp}_{a-1}^\infty(\mathcal{C})$  right Kan-extending  $X_0$ , except there seems to be a typo that renders it unclear.

Hence, for  $\mathcal{C}$  with finite limits, we have described a sequence of functors  $\mathrm{id} \rightarrow L_0 \rightarrow L_1 \rightarrow \dots$  such that

- (1)  $L_n X$  is a spectrum below  $n$ ,
- (2) For  $m \geq n$ , the map  $X[m] \rightarrow L_n X[m]$  is an equivalence,
- (3) If  $X$  is already a spectrum below  $n$ , then the map  $X \rightarrow L_n X$  is an equivalence, and
- (4) As a functor  $\mathrm{PSp}(\mathcal{C}) \rightarrow \mathrm{PSp}_{\leq n}(\mathcal{C})$ , each  $L_n$  is left-adjoint to the inclusion  $\mathrm{PSp}_{\leq n}(\mathcal{C}) \hookrightarrow \mathrm{PSp}(\mathcal{C})$ .

Properties (1) and (2) are immediate if you unwind everything. Property (3) follows from (1) and (2). Property (4) is not hard either. Next, toward an adjoint  $\mathrm{PSp}(\mathcal{C}) \rightarrow \mathrm{Sp}(n)$ , it is natural to ask about the colimit of this tower of approximations  $L_n$ . This works under some mild hypotheses.

**Proposition I.7.** If  $\mathcal{C}$  is pointed and admits finite limits and countable colimits, and  $\Omega_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$  preserves sequential colimits, then  $L := \mathrm{colim} L_n : \mathrm{PSp}(\mathcal{C}) \rightarrow \mathrm{PSp}(\mathcal{C})$  is a localization with essential image  $\mathrm{Sp}(\mathcal{C})$ . Under these conditions, we call  $L$  a *spectrafication functor*.

Finish this

## I.5 (1/29) Localizations I

I secretly never learned anything, ever. This includes most of algebraic topology—somehow, it was my third(?) undergrad course, so I spent that semester learning mathematical maturity, not actual algebraic topology. Rather than patiently fill the holes in my knowledge, I moved forward to homotopy theory, at which point I was mature enough to actually learn homotopy theory, although that was (and still is) complicated since my algebraic topology is lacking. The water settles and some holes get filled idly, but maybe not all of them. This is most apparent when I try to do chromatic things. All that is to say, today I want to review localizations, with an eye toward localizations at spectra (homology theories), and I am going to start from basics. Some references are:

- Tyler Lawson’s expository article about Bousfield localization [Law20].
- nLab.
- Ishan’s notes.
- Paul VanKoughnett’s thesis here.

Fix a category  $\mathcal{C}$  and a class  $S \subseteq \text{Mor}(\mathcal{C})$  of morphisms. We say that an object  $Z$  is  *$S$ -local* if for each  $s \in S$ , the pullback  $s^* : \text{Hom}(Y, Z) \rightarrow \text{Hom}(X, Z)$  is an equivalence. We say that a morphism  $f : X \rightarrow Y$  is an  *$S$ -equivalence* if for each  $S$ -local  $Z$ , the pullback  $f^* : \text{Hom}(Y, Z) \rightarrow \text{Hom}(X, Z)$  is an equivalence. (Hence, each  $s \in S$  is an  $S$ -equivalence.) We call a morphism  $L : X \rightarrow Z$  an  *$S$ -localization* if it is an  $S$ -equivalence and  $Z$  is  $S$ -local.

**Proposition I.8.** If two  $S$ -localizations of an object  $X$  exist, then they are isomorphic under  $X$ .

*Proof.* Given two  $S$ -localizations  $Z \xleftarrow{L} X \xrightarrow{K} Z'$ . Since  $Z, Z'$  are  $S$ -local, their represented functors invert  $S$ -equivalences. In particular, we have bijections  $K^* : \text{Hom}(Z', Z) \cong \text{Hom}(X, Z)$  and  $L^* : \text{Hom}(Z, Z') \cong \text{Hom}(X, Z')$ . So we may consider  $(L^*)^{-1}K : Z \leftrightarrow Z' : (K^*)^{-1}L$ . Pullback is just precomposition, whence the relevant triangles commute, so that these are *morphisms under  $X$* . Furthermore, uniqueness implies that these two morphisms are inverse equivalences.  $\square$

Hence we speak of *the*  $S$ -localization of  $X$ , written  $L^S X$  or just  $LX$  (the issue of existence notwithstanding). Assuming existence, the localization morphisms  $X \rightarrow LX$  are functorial. Here’s that formally:

**Proposition I.9.** Consider  $\text{Loc}^S(\mathcal{C})$  the category of  $S$ -localizing morphisms and commutative squares between them. The forgetful functor  $\text{Loc}^S(\mathcal{C}) \rightarrow \mathcal{C}$  given by  $(X \rightarrow LX) \mapsto X$  is fully faithful.

In particular, if  $\mathcal{C}$  has  $S$ -localizations, then  $\text{Loc}^S(\mathcal{C}) \rightarrow \mathcal{C}$  is an equivalence. In that case we may choose an inverse  $\mathcal{C} \rightarrow \text{Loc}^S(\mathcal{C})$  written as  $X \mapsto (X \rightarrow LX)$ , and forgetting gives us a *functorial localization functor*

$$\mathcal{C} \rightarrow L^S \mathcal{C}.$$

**Proposition I.10.** If  $\mathcal{C}$  has  $S$ -localizations, then we have described a functorial choice of localization  $X \mapsto LX$  which is left adjoint to the forgetful functor  $LX \mapsto X$ .

Some examples.

**Example I.5.** Consider  $\text{Mon}$  and the inclusion  $f : \mathbb{N} \hookrightarrow \mathbb{Z}$ . For a monoid  $M$  to be  $f$ -local, every map  $\mathbb{N} \rightarrow M$  must extend (uniquely) to a map  $\mathbb{Z} \rightarrow M$ . In other words, each element of  $M$  must admit a (unique) inverse. (I parenthesize uniqueness of inverses here because that property is superfluous in monoids.) Hence,

$$L^{\mathbb{N} \hookrightarrow \mathbb{Z}} \text{Mon} = \text{Gp}.$$

Furthermore, group completion  $M \mapsto M^{gc}$  presents the localization functor.

**Example I.6.** Consider  $\mathbf{Gp}$  and the abelianization map  $f : F_2 \rightarrow \mathbb{Z}^2$  given by  $(1, 1) \mapsto (1, 1)$ . A group  $G$  is  $f$ -local if every  $F_2 \rightarrow G$  factors (uniquely) through  $\mathbb{Z}^2$ . This means that for each  $x, y \in G$ , the commutator  $[x, y]$  vanishes, which occurs precisely when  $G$  is abelian. Therefore,

$$L^{F_2 \rightarrow \mathbb{Z}^2} \mathbf{Gp} = \mathbf{Ab},$$

and abelianization  $G \mapsto G^{ab} = G/[G, G]$  presents the localization functor.

**Example I.7.** I think that  $L^{\mathbb{N}^2 \hookrightarrow \mathbb{Z}^2} \mathbf{Mon} = \mathbf{CMon}$ .

**Example I.8.** Consider  $\mathbf{Ab}^{\text{fg}} \subset \mathbf{Ab}$  and let  $S = \{\mathbb{Z} \xrightarrow{p} \mathbb{Z} : p \text{ is prime}\}$ . An abelian group  $G$  is  $S$ -local if every  $p$  is invertible in  $G$ , i.e., when  $G$  is a rational vector space. (Note that this description is valid in both  $\mathbf{Ab}$  and  $\mathbf{Ab}^{\text{fg}}$  because  $\mathbf{Ab}^{\text{fg}}$  is a full subcategory.) Then within  $\mathbf{Ab}^{\text{fg}}$ , the only  $S$ -local object is  $G = 0$ , whence  $S$ -localization is the zero map and all homomorphisms are  $S$ -equivalences. However, the  $S$ -localizations in  $\mathbf{Ab}$  are given by rationalization.

Finish. At least get to Lawson's stuff on unstable and stable settings.

## II February

### II.1 (2/2) Phony multiplication

Some references (in order of discovery as I wrote) are

- Yigal Kamel’s talk for the homotopy theory seminar at UIUC. Good talk!
- Thomason’s article.
- Segal’s paper “Categories and cohomology theories.”
- Dmitri Pavlov’s MO question about a higher  $S^{-1}S$  construction.
- Gurski-Johnson-Osorno’s paper “2-CATEGORICAL OPFIBRATIONS, QUILLEN’S THEOREM  $B$ , AND  $S^{-1}S$ .”
- Dan Grayson’s expository article “Quillen’s work on algebraic  $K$ -theory.”
- Zbigniew Fiedorowicz’s proceedings article “The Quillen-Grothendieck Construction and Extensions of Pairings” c.f. [Fie78].
- Clayton Sherman’s chapter “Group representations and algebraic  $K$ -theory,” c.f. [She82], which has some random details worked out I found helpful.
- Daniel Harrer’s thesis “Comparison of the Categories of Motives defined by Voevodsky and Nori” available here. This is mostly unrelated, but Harrer mentions that somewhere in motivic cohomology, one needs the swap map in a tensor category to be an equality, otherwise you get a problem analogous to what Thomason finds for algebraic  $K$ -theory. Nice
- Baas-Dundas-Rognes “Two-vector bundles and forms of elliptic cohomology” c.f. [BDR03].

---

Given a monoid  $M$ , we may “formally add inverses” to obtain its [group completion](#)  $M^{\text{gp}}$ . If  $M$  is commutative, then  $M^{\text{gp}}$  has a universal property: there exists a monoidal map  $i : M \rightarrow M^{\text{gp}}$  such that for every abelian group  $A$ , monoidal maps  $M \rightarrow A$  extend surjectively and faithfully to homomorphisms  $M^{\text{gp}} \rightarrow A$ . By virtue of universality, this map  $i$  is determined up to isomorphism(?). Alternatively, you could characterize  $(-)^{\text{gp}}$  as left adjoint to  $U : \mathbf{Ab} \rightarrow \mathbf{CMon}$ , and then a choice of maps  $\{M \rightarrow M^{\text{gp}}\}_M$  is a presentation of the left adjoint(?).

That’s categorical. But we can do this with our hands. Two ways! Let’s just take  $M = \mathbb{N}$ .

**Construction II.1.** We define  $\mathbb{N}^{\text{gp}}$  as the set of symbols  $\{m - n : m, n \in \mathbb{N}\} = \mathbb{N} \times \mathbb{N}$  modulo the equivalence relation *generated by*<sup>9</sup> identifying  $a - b \sim c - d$  when there exists  $k \in \mathbb{N}$  such that  $a + k = c$  and  $b + k = d$ . You can check that  $\mathbb{N}^{\text{gp}} \cong \mathbb{Z}$  and that both the maps  $\mathbb{N} \rightarrow \mathbb{N}^{\text{gp}}$  given by  $m \mapsto m - 0$  and  $m \mapsto 0 - m$  are group completions.

**Construction II.2.** We define  $\mathbb{N}^{\text{gp}}$  as  $\text{FAb}(\mathbb{N}) / \langle (m + n) -_F (m +_F n) \rangle$ , where by  $\text{FAb}(\mathbb{N})$  we mean the free abelian group on the set  $\mathbb{N}$ , and by  $+_F, -_F$  we mean the group operations in  $\text{FAb}(\mathbb{N})$ .

Both these constructions readily extend to an arbitrary commutative monoid  $M$ . Note that Construction II.2 works for noncommutative  $M$ . And there’s more: if  $M$  is a semiring, to mean a monoid with a multiplication, then  $(M, +, \times)^{\text{gp}}$  has a canonical and well-defined multiplication  $\times_{M^{\text{gp}}}$  given as

$$((a - b), (c - d)) \mapsto (ac + bd, ad + bc).$$

We have described a functor  $K_0 : \mathbf{CMon} \rightarrow \mathbf{Ab}$ . (The second construction works for  $\mathbf{Mon}$ .) With an eye toward (higher)  $K$ -theory, we would like to consider more general input data. A generation of mathematical work (in geometry, topology, number theory, ...) evidences that *many things have  $K$ -theory*: (homotopy commutative) rings, spaces, schemes, (various adjectives) categories, ... Moreover, we want highly structured output data, perhaps a *sequence* of abelian groups, or better yet a (nice)  $K$ -theory spectrum.

---

<sup>9</sup>“Generated by” here is necessary since  $\sim$  as defined is not reflexive. There are very easy fixes to this, see e.g. the equivalent definitions of the relation on nLab, but this equivalent definition generalizes verbatim.

Often, such things arise by constructing a (category, or higher category/space)  $C$  encoding the important information, in such a way that the data unwinds into an infinite loop space structure on the geometric realization  $BG = |NC|$ .

Let  $M$  be a monoid. Following Grayson, we can easily prototype topological models for  $M^{\text{gp}}$ . For this, we will use simplicial sets, for with simplicial sets we can simultaneously encode the algebra in  $M$  and realize it spatially.

**Construction II.3.** Suppose that  $M$  is commutative. We may form the simplicial set  $X_0(M)$  wherein vertices are formal differences  $m - n$ , edges connect  $(m + k) - (n + k) \rightarrow m - n$  for every  $k \in M$ , faces fill every composable pair of the form shown below, and so on.

$$\begin{array}{ccc}
 m - n & \xrightarrow{k+k'} & m'' - n'' \\
 & \searrow k & \nearrow k' \\
 & m' - n' &
 \end{array}$$

**Construction II.4.** Do not assume that  $M$  is commutative. We may form the simplicial set  $X_1(M)$  wherein there is one vertex, an edge for each  $m \in M$ , a face with edge  $m, m'$ , and  $m + m'$ , and so on.

The constructions of  $X_0(M)$  and  $X_1(M)$  are analogous to Construction II.1 and Construction II.2, respectively. You can check that  $\pi_0 X_0(M) \cong M^{\text{gp}}$  and  $\pi_1 X_1(M) \cong M^{\text{gp}}$ . If  $M$  is already a group (resp. abelian group), then  $X_1(M)$  (resp.  $X_0(M)$ ) has trivial higher homotopy and models the classifying space  $BM$  (resp.  $\Omega BM$ ?). So, we have described two constructions  $X_0, X_1$  which (topologically) model group completion in the sense that  $\pi_i X_i(M) \cong M^{\text{gp}}$ . The  $X_1$  construction works for noncommutative  $M$ , and if  $M$  is already group, then  $X_0(M) = \Omega BM$  and  $X_1(M) = BM$  (in some sense...).

**Remark II.1.** Group completion also changes homology, see e.g. MO. This persists for categories!

We would like to replace  $M$  by a symmetric monoidal category and ask about group completions. We produced  $X_0(M)$  by Construction II.1, and this straightforwardly generalizes.

**Definition II.1.** Suppose that  $(C, \oplus, 0)$  is a symmetric monoidal category. We define its *Quillen completion*  $S^{-1}S(C)$  to be the category whose...

- Objects are pairs  $(A, B)$ .
- Morphisms  $(A, B) \rightarrow (C, D)$  are triples  $(K, A \oplus K \rightarrow C, B \oplus K \rightarrow D)$ , modulo the equivalence relation identifying morphisms when there exists  $K \cong K'$  making the obvious four triangles commute.

**Remark II.2** (Structure of  $S^{-1}S(C)$ ). Here is some basic structure on the Quillen completion.

- The identity morphism  $\text{id}_{(A,B)}$  is  $(0, 0 \rightarrow A, 0 \rightarrow B)$ .
- The monoidal structure on  $C$  induces one on  $S^{-1}S(C)$ . It is computed “coordinate-wise,” i.e.  $(A, B) \oplus (C, D) = (A \oplus C, B \oplus D)$  and likewise for sums of morphisms. If  $s_{A \oplus B} : A \oplus B \rightarrow B \oplus A$  denotes the swap maps in  $C$ , then  $(0, s_{AC}, s_{BD}) : (A, B) \oplus (C, D) \rightarrow (C, D) \oplus (A, B)$  provide swap maps for  $S^{-1}S(C)$ .
- There is a *transposition* functor  $t : S^{-1}S(C) \rightarrow S^{-1}S(C)$  given by

$$(A, B) \mapsto (B, A) \quad \text{and} \quad (K, \alpha, \beta) \mapsto (K, \beta, \alpha).$$



- There are inclusion functors  $i, -i : \mathcal{C} \rightarrow S^{-1}S(\mathcal{C})$  given on objects by  $i(C) = (0, C)$  and  $-i(C) = (C, 0)$ . We want to think of the object  $(A, B) \in S^{-1}S(\mathcal{C})$  as a formal difference  $B - A$ ,<sup>10</sup> whence this notation makes sense.

We would like to understand  $S^{-1}S(\mathcal{C})$  as a “categorical group completion.” For this, consider the following three properties that  $(\mathcal{C}, \oplus, 0)$  might possess.

- (I)  $\mathcal{C}$  is a groupoid.
- (II)  $\mathcal{C}$  is “cancellative:” For every  $A \in \mathcal{C}$ , the functor  $A \oplus - : \mathcal{C} \rightarrow \mathcal{C}$  is faithful.
- (III)  $\mathcal{C}$  has “object-level inverses:” there exists a natural transformation  $0 \rightarrow \text{id} \oplus t$ .

**Proposition II.1.** If  $\mathcal{C}$  satisfies (I) and (II) above, then the inclusion  $i : \mathcal{C} \rightarrow S^{-1}S(\mathcal{C})$  given by  $A \mapsto (0, A)$  induces a group completion on classifying spaces. That is to say

$$\pi_0(BS^{-1}S(\mathcal{C})) = \pi_0(BC)^{\text{gp}} \quad \text{and} \quad H_0(BS^{-1}S(\mathcal{C})) = H_0(BC)[\pi_0 BC^{-1}].$$

Suppose that (I) and (II) hold. Since  $S^{-1}S(\mathcal{C})$  is symmetric monoidal, its classifying space is an  $H$ -space, and by the proposition it is an  $H$ -group.<sup>11</sup> We regard  $(A, B) \in S^{-1}S(\mathcal{C})$  as the formal difference  $B - A$ , and  $(A, B)$  represents this difference in  $\pi_0 BS^{-1}S(\mathcal{C})$ . Also on  $\pi_0$ , the transposition functor  $t$  induces the inverse map. The symmetric monoidal structure gives  $BS^{-1}S(\mathcal{C})$  its monoidal structure, and we would like to say that  $t$  induces its homotopy inverse  $g \mapsto g^{-1}$ .

**Proposition II.2.** If  $\mathcal{C}$  satisfies (I), (II), and (III) above, then the transposition  $t$  for the Quillen completion  $S^{-1}S(\mathcal{C})$  induces a homotopy inverse for the  $H$ -group  $BS^{-1}S(\mathcal{C})$ .

*Proof.* Write  $Z = BS^{-1}S(\mathcal{C})$ . By (III), we may choose a transformation  $\eta : 0 \rightarrow \text{id} \oplus t$ . We get a map  $B\eta : B0 \rightarrow B(\text{id} \oplus t)$ , and this represents a homotopy inducing (in  $[Z, Z]$ )  $0 = [B0] = [B\text{id}] + [Bt] \in [Z, Z]$ , which begets  $[B\text{id}] = -[Bt]$ . Hence,  $Bt$  is a homotopy inverse for the  $H$ -space  $Z$ .  $\square$

Thomason’s essential observation is that (III) is secretly a very strong condition, unfulfilled in even the most standard cases. For example, neither  $\mathcal{C} = \text{Mod}_R^{\text{fg}, \text{proj}}$  nor its maximal subgroupoid have this property unless  $R = 0$ . Fortunately, Proposition II.2 still holds when you do not assume (III)! But results implementing the homotopy inverse for  $BS^{-1}S(\mathcal{C})$  via  $t$  and utilizing the naturality of  $\eta$  in an essential way still fail (since  $\eta$  does not even exist).

The functor  $t \oplus \text{id}$  acts on objects as  $(A, B) \mapsto (B \oplus A, A \oplus B)$ . Hence, a transformation  $\eta : 0 \rightarrow t \oplus \text{id}$  amounts to a natural system of morphisms  $\eta_{AB} = \{K, 0 \oplus K \rightarrow B \oplus A, 0 \oplus K \rightarrow A \oplus B\}$ . One candidate is

$$\eta_{AB} := \{A \oplus B, s_{A \oplus B}, \text{id}_{A \oplus B}\}.$$

**Proposition II.3.** The system of morphisms  $\{\eta_{AB} : (0, 0) \rightarrow [t \oplus \text{id}](A, B)\}_{(A, B) \in \mathcal{C}}$  is natural in  $\mathcal{C}$  if and only if the swap isomorphisms  $s_{S \oplus S}$  are equalities for all  $S \in \mathcal{C}$ .

*Proof.* Choose arbitrary  $A, B, C, D \in S^{-1}S(\mathcal{C})$ . If we denote by  $f$  an arbitrary morphism  $(S, \alpha, \beta) : (A, B) \rightarrow (C, D)$ , then  $(t \oplus \text{id})f$  is the morphism  $(B \oplus A, A \oplus B) \rightarrow (D \oplus C, C \oplus D)$  consisting of the following two arrows.

$$\begin{aligned} B \oplus A \oplus S \oplus S &\xrightarrow{\text{id}_B \oplus s_{AS} \oplus \text{id}_S} B \oplus S \oplus A \oplus S \xrightarrow{\beta \oplus \alpha} D \oplus C \\ A \oplus B \oplus S \oplus S &\xrightarrow{\text{id}_A \oplus s_{BS} \oplus \text{id}_S} A \oplus S \oplus B \oplus S \xrightarrow{\alpha \oplus \beta} C \oplus D \end{aligned}$$

<sup>10</sup>This signage is decided by the direction of edges in  $S^{-1}S(\mathcal{C})$ .

<sup>11</sup>If  $\mathcal{C}$  is an exact category, you could also argue that the  $H$ -space is an  $H$ -group by proving that the map  $K_0(\mathcal{C}) \rightarrow \pi_0(BS^{-1}S(\mathcal{C}))$  is an isomorphism and that the former is a group. I read this in ??.

Naturality amounts to the commutativity of the following diagram, for all  $A, B, C, D$ , and  $f$ .

$$\begin{array}{ccc} (0, 0) & \xrightarrow{\eta_{AB}} & (B \oplus A, A \oplus B) \\ \downarrow 0(f)=0 & & \downarrow (i \oplus \text{id})f \\ (0, 0) & \xrightarrow{\eta_{CD}} & (D \oplus C, C \oplus D) \end{array}$$

Writing out  $\eta_{CD}$  and the composite  $(i \oplus \text{id})f \circ \eta_{AB}$ , Thomason finds commutativity to be equivalent to that of

$$\begin{array}{ccc} BASS & \xrightarrow{s_{AS}} & BSAS \\ \uparrow s_{AB} & & \uparrow s_{(AS)(BS)} \\ ABS S & \xrightarrow{s_{BS}} & ASBS \end{array}$$

Note that this diagram is in  $\mathbf{C}$ . Observe that up-right does not reverse the order of the two  $S$ 's. However, right-up does (because  $s_{(AS)(BS)}$  does). Hence, this diagram commutes if and only if the swap map  $s_{S \oplus S}$  is a *strict* equality. As  $S$  was arbitrary, the result follows.  $\square$

The swap isomorphisms are rarely equalities, and although it is possible to “strictify” a symmetric monoidal category into a permutative one (in which associativity and unitality are strict), commutativity cannot be strictified except in trivial cases. Hence, the result outs (III) a tricky condition to satisfy, for the obvious natural transformation is *not* a natural transformation! A tragedy.

Let me quote some of Peter’s commentary [May80]:

Thomason [...] has given an amusing illustration of the sort of mistake that can arise from a too cavalier attitude towards this kind of categorical distinction when studying pairings of categories, and one of my concerns is to correct a similar mistake of my own.

In [...], I developed a coherence theory of higher homotopies for ring spaces up to homotopy and for pairings of  $H$ -spaces. That theory is entirely correct. I also discussed the analogous categorical coherence, proving some results and asserting others. That theory too is entirely correct, my unproven assertions having been carefully proven by Laplaza [unpublished]. However, my translations from the categorical to the homotopical theories in [...], that of course being the part I thought to be obvious, are quite wrong.

The moral is that to treat the transition from categorical coherence to homotopical coherence smoothly and rigorously, one should take advantage of the definitional framework established by the category theorists.

There’s more to say. But I have been writing some time now, and have generated me thoughts I need to lay bare before the crystal ball (ask Charles about). Let me just say something about the first great victim of Thomason’s observation.

Recall that if we put a semiring structure on a monoid  $M$  (i.e., a unital operation  $\times$  distributing over  $+$ ), for example that already present on the underlying monoid of any unital associative ring, *then this straightforwardly extends to a ring structure on the group  $K_0(M) = M^{\text{gp}}$* . In fact, the full  $K$ -theory  $K_*(A)$  (of rings, schemes, whatever) has a ring structure. Back in the day, Quillen invented higher algebraic  $K$ -theory

Finish this. Talk about ring structure on  $K(R)$ , relation of Quillen completion

## II.2 (2/7) Localizations II

Johnson and Jeremiah spoke for the (anti)telescope seminar today. Johnson wrapped up his previous talk (introducing monochromaticity in  $\text{Cat}_{\text{perf}}$ ) with a proof that the category of  $n$ -monochromatic categories is  $\infty$ -semiadditive for all  $n$ . Jeremiah gave an overview of algebraic  $K$ -theory. We did all this with an eye toward [Ben+23] (wherein spawns the monochromatic language for categories). One takeaway is that I need to learn more about localizations (and the chromatic phenomena motivating our modern theory of localizations, c.f. my monologue last time).

References are:

- Moritz Groth's notes, again
- HTT and maybe HA, again
- Lawson's notes, again
- I also found Martin Gallauer's lecture notes here, although I have not looked at these yet.

---

Let's start with ordinary categories. We did some of this last time. There, we wanted to invert a class of morphisms  $S \subseteq \text{Mor}(\mathcal{C})$  in a universal way. We did not study existence, but we assumed existence and (i) showed uniqueness of a given object's  $S$ -localizations, (ii) exhibited a functorial choice of  $S$ -localizations  $X \mapsto LX$ , and (iii) thought about some basic examples. The resultant basic structure is an adjunction

$$\mathcal{C} \begin{array}{c} \xrightarrow{L^S} \\ \xleftarrow{\perp} \end{array} L^S(\mathcal{C})$$

Which full subcategories arise in this manner? Define a *reflective subcategory* to be a full subcategory  $\mathcal{D} \hookrightarrow \mathcal{C}$  whose inclusion admits a left adjoint. These are essentially our localizations. *It will be a common theme that localizations can be understood as adjoint systems (existence, properties, etc.).*

**Proposition II.4.** Let  $(L \dashv i)$  denote an adjoint pair. TFAE.

- (1) The right adjoint  $i : \mathcal{D} \rightarrow \mathcal{C}$  is fully-faithful ( $\iff \mathcal{D}$  is a reflective subcategory).
- (2) The counit  $Li \rightarrow \text{id}_{\mathcal{D}}$  is a natural isomorphism.
- (3) The left adjoint  $L : \mathcal{C} \rightarrow \mathcal{D}$  is a localization at  $S = L^{-1}(\text{isos})$ .

**Proposition II.5.** Dually, define a *coreflective subcategory* to be a full subcategory whose inclusion admits a right adjoint. Let  $(i \dashv R)$  denotes an adjoint pair. TFAE.

- (1) The left adjoint  $i : \mathcal{D} \rightarrow \mathcal{C}$  is fully faithful ( $\iff \mathcal{D}$  is a coreflective subcategory).
- (2) The unit  $\text{id}_{\mathcal{C}} \rightarrow iR$  is a natural isomorphism.
- (3) The right adjoint  $R : \mathcal{C} \rightarrow \mathcal{D}$  is a colocalization at  $S = R^{-1}(\text{isos})$ .

To my eye, the ordinary theory of localizations is a bit messy. Maybe I feel this way because I do not like how we handle comparisons/notions of equivalence. Maybe that's no discredit to the theory, just nLab's (lack of) presentation, and the fact that most literature I can find works in situations I am trying to not focus on. I'll just move on.

---

Toward real math, we want to think about localizations when  $\mathcal{C}$  is *topologically enriched*. One asks whether this reduces to the localization theory of the homotopy category. The following example suggests this is an awkward approach to the theory.

**Example II.1.** Consider  $\text{Top}$  the topological category of spaces. Pass to its homotopy category and consider the map

$$i_n : S^n \hookrightarrow D^{n+1}.$$

We can ask about inverting  $i_n$ . This requires exhibiting  $i_n$ -localizations. Given a space  $X$ , form  $X'$  by gluing copies of  $D^{n+1}$  onto  $X$  to kill  $\pi_n X$ . Pullback along  $i_n$  induces an equivalence  $[D^{n+1}, X'] \xrightarrow{\sim} [S^n, X']$  since both sides are trivial (in their path components) by construction, hence  $X'$  is  $i_n$ -local. Is the map  $l : X \rightarrow X'$  an  $i_n$ -equivalence? This would require the pullback

$$l^* : [X', Y] \rightarrow [X, Y]$$

to be an isomorphism for every  $i_n$ -local space  $Y$ . This map is surjective:  $Y$  is  $i_n$ -local if and only if  $\pi_n Y$  is zero, in which case an extension of any  $f : X \rightarrow Y$  over any  $n$ -cell  $\partial D^{n+1} \rightarrow X$  exists, exhibiting an inverse in  $l^{-1}(f)$ . **The problem is uniqueness:** if  $\partial D^{n+1} \rightarrow X$  denotes a cell killed to form  $X'$ , then given  $f : X \rightarrow Y$ , two extensions  $g, h$  of  $f$  to  $D^{n+1}$  determine a map  $S^{n+1} \rightarrow Y$ , and  $g \simeq h$  if and only if this map is nullhomotopic. In this way,  $\pi_{n+1} Y$  obstructs the injectivity of  $l^*$ . Since  $l^*$  is not injective, another candidate localization  $X''$  may not be homotopy equivalent to  $X'$ .

**Remark II.3.** The example shows that we cannot solely invert  $i_n : S^n \hookrightarrow D^{n+1}$  in  $h\text{Top}$ , at least not nicely or obviously. The example also showed that inverting  $i_n$  requires at least inverting  $i_m$  for all  $m \geq n$ . One imagines that inverting an arbitrary map  $f$  comes with similar responsibilities as soon as  $f$  is not a homotopy equivalence (i.e., is not already inverted).

Thus, given a topological category  $\mathbf{C}$  and a class  $S \subseteq \text{Mor}(\mathbf{C})$ , rather than work with  $h\mathbf{C}$ , we define  *$S$ -local objects* and  *$S$ -equivalences* in  $\mathbf{C}$  just as before except with “isomorphism” replaced with “weak equivalence.” With these properties defined, we can define  *$S$ -localizations* just as before.

**Proposition II.6.** If  $Y \in \mathbf{C}$  is  $S$ -local, then  $hY \in h\mathbf{C}$  is  $hS$ -local. However, since  $hS$  is generally “smaller” than  $S$ , there are generally more  $hS$ -local objects. Thus, if  $f$  is an  $S$ -equivalence, it is not necessarily true that  $[f]$  is an  $hS$ -equivalence.

We would lastly like a localization *functor*. Unfortunately, even if our topological category  $\mathbf{C}$  has localizations, choosing them *naturally* is a coherence problem. When and how we can do this is nontrivial.

Agony and confusion. How model category theory Bousfield did all this for spaces and spectra. His theory extends to model categories, and resultant localizations get a model structure too. Localization and completions of spectra are examples of Bousfield localizations. **How/in what language (stable  $\infty$ -categories?) can we think about Bousfield localizations as “functors with certain kinds of adjoints,” as we did using (co)reflective subcategories in the case of categorical localization?** Maybe this is not the right way to think about it. Also, triangulated categories and Verdier localization—what’s going on there? I think this is where the “thick subcategory” ideas originate. Maybe H. Krause’s notes [Kra09] will clarify. How do things converge when you use stable  $\infty$ -categories?

(Trigger warning: yapping.) We came to view ordinary localization as the search for left adjoints with fully faithful right adjoints. When considering a topological category  $\mathbf{C}$ , we found it ineffective to restrict to the homotopy category  $h\mathbf{C}$ , so instead we carried out the story of ordinary localizations but with *weak equivalences*. Bousfield did this for spaces and spectra. But things are not so simple now—even if localizations exist, choosing them *functorially* is a nontrivial *coherence problem*. So that complicates things. More generally, this problem occurs for *model categories*. In nice situations, we can view this as a lifting problem, and somehow a functorial choice is possible when “the small object argument works.” Localization becomes a process of changing model structures on a fixed category. This is somewhat different from how you might naturally present and study localizations in ordinary category theory via universal properties. I think you can frame this as an adjoint systems phenomenon *in the language of Quillen adjunctions*, but extracting a plain categorical statement from this is hard? This presents other issues too—e.g., the extension of monoidal structures is very nontrivial.

In contrast, it is quite easy to express and organize  $\infty$ -categorical localizations via universal properties and adjoint systems. This is maybe because the semantics of  $\infty$ -categories, functors, etc. are designed to “see” and naturally handle the associated lifting problems. We can e.g. stay faithful to the story of “reflective subcategories.”

Spiel about small object argument, Lawson §5,6,7

Think about localizations of (based, unbased,  $G$ -) spaces. Fiberwise localization? Also, whatever Lawson talks about at the end of §7.

**Definition II.2.** Let  $\mathcal{C}, \mathcal{D}$  denote  $\infty$ -categories. A functor  $\mathcal{C} \rightarrow \mathcal{D}$  is called a *localization* if it admits a fully faithful right adjoint. We may call  $\mathcal{D}$  a *localization of  $\mathcal{C}$*  and we may call the resultant functor  $\mathcal{C} \rightarrow \mathcal{C}$  a *localization functor*.

**Proposition II.7** (Basic form of localizations, [Lur08, 5.2.7.4]). Let  $\mathcal{C}$  be an  $\infty$ -category and  $L : \mathcal{C} \rightarrow \mathcal{C}$  a functor with essential image  $LC$ . TFAE.

- (1) There exists a localization  $f : \mathcal{C} \rightarrow \mathcal{D}$  with fully faithful right adjoint  $g$  and an equivalence  $g \circ f \simeq L$ .
- (2) As a functor  $L : \mathcal{C} \rightarrow LC$ ,  $L$  is left adjoint to the inclusion  $LC \hookrightarrow \mathcal{C}$ .
- (3) There exists a natural transformation  $\alpha : \mathcal{C} \times \Delta^1 \rightarrow \mathcal{C}$  from  $\text{id}_{\mathcal{C}}$  to  $L$  such that the morphisms  $\alpha(LC), L(\alpha C) : LC \rightarrow LLC$  are equivalences.

**Proposition II.8** ([Lur08, 5.2.7.8]). Consider  $\mathcal{C}_0 \hookrightarrow \mathcal{C}$  an  $\infty$ -category and a full subcategory. TFAE.

- (1) The inclusion  $\mathcal{C}_0 \hookrightarrow \mathcal{C}$  admits a left adjoint.
- (2) Each object  $C \in \mathcal{C}$  admits a morphism  $f : C \rightarrow D$  which *exhibits  $D$  as a  $\mathcal{C}_0$ -localization of  $C$* . By this, we mean that  $D \in \mathcal{C}_0$  and for every  $E \in \mathcal{C}_0$ , the map  $f^* : \text{Map}_{\mathcal{C}_0}(D, E) \rightarrow \text{Map}_{\mathcal{C}}(C, E)$  is an equivalence in the homotopy category.

**Remark II.4.** Note that  $f : C \rightarrow D$  exhibits  $D$  as a  $\mathcal{C}_0$ -localization if and only if  $f \in \mathcal{C}_C / \times_{\mathcal{C}} \mathcal{C}_0$  is an initial object. This implies e.g. uniqueness and invariance under equivalence of categories.

If a full subcategory  $\mathcal{C}_0 \subseteq \mathcal{C}$  satisfies either of the equivalent conditions of the above proposition, we call  $\mathcal{C}_0$  a *reflective subcategory*. It is clear that reflective subcategories are an equivalent formulation of  $\infty$ -categorical localizations.

**Proposition II.9** ([Lur08, 5.2.7.12]). Let  $L : \mathcal{C} \rightarrow \mathcal{C}$  be a localization with essential image  $LC$ . Then for any  $f : \mathcal{C} \rightarrow \mathcal{D}$ , composition induces an equivalence

Recall that an  $\infty$ -category is called *presentable* if it is accessible and admits small colimits. Presentable  $\infty$ -categories have a nice theory of adjunctions. For this and other reasons, they have a nice theory of localizations. I'll quote [Lur08, §5.5]:

Wait, what is this proposition saying?

In view of Theorem 5.5.1.1, the theory of localizations plays a central role in the study of presentable  $\infty$ -categories. In §5.5.4, we will show that the collection of all (accessible) localizations of a presentable  $\infty$ -category  $\mathcal{C}$  can be parametrized in a very simple way. Moreover, there is a good supply of localizations of  $\mathcal{C}$ : given any (small) collection of morphisms  $S$  of  $\mathcal{C}$ , one can construct a corresponding localization functor

$$\mathcal{C} \xrightarrow{L} S^{-1}\mathcal{C} \subseteq \mathcal{C}$$

where  $S^{-1}\mathcal{C}$  is the full subcategory of  $\mathcal{C}$  spanned by the  $S$ -local objects. These ideas are due to Bousfield, who works in the setting of model categories; we will give an exposition here in the language of  $\infty$ -categories. In §5.5.5, we will employ the same techniques to produce examples of factorization systems on the  $\infty$ -category  $\mathcal{C}$ .

I need more time to read §5 of HTT, but here's an important classification result. We say that a localization  $L$  is *accessible* if its fully faithful right adjoint is *accessible*, i.e. if it preserves all  $\kappa$ -filtered colimits for some regular cardinal  $\kappa$ .

**Theorem II.1.** An  $\infty$ -category  $\mathcal{C}$  is presentable if and only if it is an accessible localization of  $\text{PShv}(\mathcal{D})$  for some small  $\infty$ -category  $\mathcal{D}$ .

**Remark II.5.** Recall that presentable  $\infty$ -categories enjoy a (left) adjoint functor theorem: if  $\mathcal{C}, \mathcal{D}$  are presentable, then a functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is a left adjoint if and only if it preserves small colimits. However, the converse needs an additional hypothesis:  $F$  is a right adjoint if and only if it preserves small limits *and is accessible*. In particular, if  $LC \rightarrow \mathcal{D}$  is a localization, then its right adjoint is automatically accessible. This explains the appearance of accessibility in the classification result.

Given a localization  $F : \mathcal{C} \rightarrow \mathcal{D}$ , we denote by  $L : \mathcal{C} \rightarrow \mathcal{C}$  the composition, by which we can study this localization. As in ordinary category theory, the “local” objects play an essential role. Denote by  $S_L$  the collection of morphisms inverted by  $L$ . Say an object  $c \in \mathcal{C}$  is  $S_L$ -local if for all  $f \in S_L$ , the induced  $f^* : \text{Map}(B, c) \rightarrow \text{Map}(A, c)$  is weak equivalence.

**Proposition II.10** ([Lur08, 5.5.4.2]). Given  $\mathcal{C}$  and localization functor  $L : \mathcal{C} \rightarrow \mathcal{C}$ , the essential image  $LC$  is spanned by the  $S_L$ -local objects.

As a localization,  $L$  is left adjoint to the inclusion  $LC \hookrightarrow \mathcal{C}$ . The proposition identifies  $LC = S_L$ , thus  $L$  is determined up to equivalence by  $S_L$ . Hence, **localizations correspond to certain classes of morphisms**. These classes are not arbitrary, but rather possess nice closure properties. An arbitrary class of morphisms having such closure properties (e.g., closure under colimits, pushdowns, 2-out-of-3) is called **strongly saturated** [Lur08, 5.5.4.5]. If  $\mathcal{C}$  admits small colimits, then  $\text{Mor}(\mathcal{C})$  is strongly saturated, and moreover the intersection of strongly saturated classes is strongly saturated, whence any collection  $S_0$  has a minimal containing strongly saturated class  $\overline{S_0}$ . We may say  $\overline{S_0}$  is the strong class of morphisms **generated by  $S_0$**  or call it the **strong saturation** of  $S_0$ . If  $S_0$  is small, we say  $\overline{S_0}$  is **of small generation**.

This language established, we can systematically describe the reverse process: specifying a set of morphisms and exhibiting a localization.

**Proposition II.11** ([Lur08, 5.5.4.15]). Let  $\mathcal{C}$  be a presentable  $\infty$ -category and  $S$  a small class of morphisms. Let  $L^S \mathcal{C}$  denote the full subcategory of  $S$ -local objects. Then:

- (1)  $L^S \mathcal{C}$  is presentable.
- (2) The inclusion  $L^S \mathcal{C} \hookrightarrow \mathcal{C}$  has a left adjoint  $L$ . (Which, by the right adjoint functor theorem for presentable categories, implies that the inclusion is accessible.)
- (3) Each  $c \in \mathcal{C}$  admits an  $S$ -equivalence  $c \rightarrow c'$  to some  $c' \in L^S \mathcal{C}$ .
- (4) TFAE.
  - (a)  $f$  is an  $S$ -equivalence.
  - (b)  $f$  is in  $\overline{S}$ .
  - (c)  $Lf$  is an equivalence.

We knew that localizations  $L : \mathcal{C} \rightarrow \mathcal{C}$  of an arbitrary  $\infty$ -category  $\mathcal{C}$  correspond to certain classes of morphisms (which occur as  $L^{-1}(\text{isos})$ ). The proposition says that if  $\mathcal{C}$  is presentable, then this correspondence can be made simple and precise. In one direction, given *any* set of morphisms  $S$ , there exists an associated accessible localization  $\mathcal{C} \xrightarrow{L} L^S \mathcal{C} \hookrightarrow \mathcal{C}$  inverting  $\overline{S}$ , and all localizations arise this way. In the other direction, given a localization  $L$ , we extract  $S = L^{-1}(\text{isos})$ , which is a strongly saturated set of small generation. This is not a bijective process, but noting that  $\overline{S} = \overline{T} \implies L^S \mathcal{C} = L^T \mathcal{C}$ , we get the following.

**Proposition II.12.** The accessible localizations of presentable  $\infty$ -category  $\mathcal{C}$  correspond with its strongly saturated classes of small generation.

**Example II.2.** Recall that if  $\mathcal{C}$  is pointed with finite limits and finite colimits, we can define its category of *prespectrum objects* and *spectrum objects*. These are modeled concretely and intuitively, c.f. Section I.4. Using this concrete model, it is easy to take a prespectrum  $X$  and freely construct a “spectrum below level  $n$ ”  $L_n X$ . We would like to consider  $\text{colim}_n L_n X$  as the *spectrafication* of  $X$ . With some mild hypotheses on  $\mathcal{C}$  (c.f. Proposition I.7), this works:  $L := \text{colim}_n L_n$  is a localization  $\text{PSp}(\mathcal{C}) \rightarrow \text{PSp}(\mathcal{C})$  with essential image  $\text{Sp}(\mathcal{C})$ . In particular, taking  $\mathcal{C} = \text{Spaces}_*$ , we find  $\text{Sp}$  as a localization of  $\text{PSp}$ . Since  $\text{Spaces}_*$  is presentable, this exhibits  $\text{Sp}$  as an (accessible) localization of a presentable category, whence  $\text{Sp}$  is presentable.

What’s the right way to see that  $\text{Sp}$  is pre-

### II.3 (2/14) Categorifying Mahler's theorem

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Some references for today are:

- Lior Yanovski's thesis on homotopy cardinality and Euler characteristics, which motivated this post.
- Noam Elkies's short note about his generating function proof of Mahler's theorem.
- Bibby-Badish's article using generating functions to study the homology of orbit configuration spaces.
- Mahler's 1958 paper [Mah58] giving criteria for continuously extending  $p$ -adic functions.

---

Given a sequence of numbers  $A = \{a_i\}$ , we define its *binomial transform* as the sequence  $TA$  given by

$$(TA)_n := \sum_{k=0}^n (-1)^k \binom{n}{k} a_k.$$

This defines an involution  $T$  of  $\text{Fun}(\mathbb{N}, M)$  for any  $\mathbb{Z}$ -module  $M$ .<sup>12</sup> The binomial transform is closely related to the *forward difference operator*  $\Delta$  given by  $(\Delta A)_n = a_{n+1} - a_n$ .

**Example II.3.** Let me write out the first few terms of some iterates of  $\Delta^k A$ .

$$\begin{aligned} \Delta A &= a_1 - a_0, & a_1 - a_2, & \dots \\ \Delta^2 A &= a_0 - 2a_1 + a_2, & a_3 - 2a_2 + a_1, & \dots \\ \Delta^3 A &= a_3 - 3a_2 + 3a_1 - a_0, & \dots \end{aligned}$$

The relation between  $\Delta$  and  $T$  is clear:  $(\Delta^n A)_0 = (TA)_n$ .

With that notation established, a classical problem: how to estimate a function? If  $f : \mathbb{R} \rightarrow \mathbb{R}$  is smooth, we may consider its Taylor power series  $\sum f^{(k)}(x)x^k/k!$  about zero. This power series does not always converge, nor does it always converge to  $f$ . We may also consider the *Newton series* of  $f$ , defined as a discrete Taylor series:

$$Nf(x) := \sum_{k=0}^{\infty} \frac{(\Delta^k f)(0)}{k!} x(x-1)(x-2) \dots (x-k+1).$$

This does not always converge either. What's interesting is that  $p$ -adically, *the obstruction to coincidence of  $Nf$  with  $f$  is continuity*. Compare to the real case, where not even smoothness guarantees convergence, let alone coincidence.

**Proposition II.13** (Mahler's theorem). Let  $f : \mathbb{Z}_p \rightarrow \mathbb{Q}_p$  be a  $p$ -adic function. TFAE.

- (1)  $f$  is continuous.
- (2) The Newton series for  $f$  converges to  $f$  everywhere, i.e.  $f(x) = \sum_{k=0}^{\infty} \binom{x}{k} (\Delta^k f)(0)$ .
- (3) The sequence of coefficients  $(Tf)_k = (\Delta^k f)(0)$  converges to  $0 \in \mathbb{Q}_p$ .

---

<sup>12</sup> $\text{Fun}(\mathbb{N}, M)$  is an abelian group under pointwise addition and  $T$  is a group homomorphism. I have several questions...



Given a sequence  $A = \{a_i \in \mathbb{Q}_p\}_{i=0}^\infty$ , we may consider its binomial transformation  $B = TA$ , and if  $b_k \rightarrow 0$  as  $k \rightarrow \infty$  then the  $p$ -adic power series  $A(x) := \sum \binom{x}{k} b_k$  is a uniform limit of polynomials. It therefore converges to a continuous function  $\mathbb{Z}_p \rightarrow \mathbb{Q}_p$  satisfying  $A(n) = a_n$ . This describes a process for extending a “nice” function  $\mathbb{N} \rightarrow \mathbb{Q}_p$  to a continuous function  $\mathbb{Z}_p \rightarrow \mathbb{Q}_p$ . This is how you go from (3) to (1). Here, “nice” is the (above) stated convergence condition on the binomial transform  $BA$ .

Finish this

## II.4 (2/21) The pushdiagonal of a ring with $G$ -action

Let  $\mathcal{C}$  be a nice  $\infty$ -category (I will be more specific later). I want to think about functors  $BG \rightarrow A := \mathcal{C}\text{Alg}(\mathcal{C})$ . For now, just take  $\mathcal{C} = \text{Mod}_k$  for some field  $k$ , so that such a functor  $R$  is precisely a commutative  $k$ -algebra with  $G$ -action (by  $k$ -linear automorphisms). We can ask about the pushforward

$$BG \times BG \xrightarrow{\Delta_* R} A$$

where  $\Delta : BG \rightarrow BG \times BG$  is the diagonal functor. The functor  $\Delta_*$  is right adjoint to the pullback  $\Delta^*$  and hence is characterized as the right Kan extension of  $R$  along  $\Delta$ , as in the following diagram.

$$\begin{array}{ccc} BG & \xrightarrow{R} & A \\ \Delta \downarrow & \dashrightarrow_{\text{Ran}_\Delta R = \Delta_* R} & \\ BG \times BG & & \end{array}$$

The right Kan extension exists because  $A$  has limits, and it is computed by fiberwise limits, i.e.

$$[\Delta_* R](*) = \lim_{\Delta/_*} R\pi.$$

Here,  $\Delta/_*$  is the slice of  $\Delta$  over  $* \in BG \times BG$  and  $\pi : \Delta/_* \rightarrow BG$  is the projection. This is the general formula for a right Kan extension. Let’s acknowledge  $BG$ ’s elementary structure and simplify the formula: the slice category  $\Delta/_*$  has objects  $(g_1, g_2) \in G \times G$  and has a morphism  $h : (g_1, g_2) \rightarrow (g'_1, g'_2)$  for each  $h$  such that  $g_1 = g'_1 h$  and  $g_2 = g'_2 h$ . Each object  $(g_1, g_2) \in \Delta/_*$  is uniquely isomorphic via  $g_1$  to  $(e, g_2 g_1^{-1})$ , hence  $\Delta/_*$  is canonically equivalent to a discrete subcategory isomorphic to  $G$  (embedded on objects by  $g \mapsto (e, g)$ ). We get a much more familiar expression for the underlying object of  $\Delta_* R : BG \times BG \rightarrow A$ :

$$[\Delta_*(R)](*) = \lim_G R\pi = \prod_G R(*).$$

Just as the underlying object of  $R : BG \rightarrow A$  carries a  $G$ -action, that of  $\Delta_* R$  carries a  $G \times G$  action. That action is given by  $(x, y)r_g = x r_{g y^{-1}}$ .

## II.5 (2/22) Linear Galois theory I

An absolutely natural impulse in virtually all of algebra is to do for commutative rings what has already been done for fields.

Daniel Zelinsky

Some references for today are

- (1) Rognes’ seminal *Galois extensions of structured ring spectra* [Rog05],
- (2) Gow-Quinlan’s article “Galois theory and linear algebra” [GQ09],
- (3) Michael Francis’s notes on “Linear Galois theory” found on his site here,
- (4) Sharon Zhou’s great Chicago REU paper found here,
- (5) Farb-Dennis *Noncommutative Algebra* [FD12], and



(6) Dress’s article “One more shortcut to Galois theory” [Dre95].

Recall Galois theory. A field extension  $L/K$  is called a *Galois extension* if it is algebraic, normal, and separable. There are a lot of ways to characterize and study Galois extensions, especially finite ones. For instance, finite Galois extensions of  $K$  are precisely the splitting fields of separable polynomials  $f \in K[x]$ . Alternatively, the *fundamental theorem of Galois theory* says that a finite extension  $L/K$  is Galois if and only if the following correspondence (I won’t write it out) is bijective.

$$\{\text{intermediate field extensions } L/E/K\} \longleftrightarrow \{\text{subgroups of } \text{Aut}(L/K)\}$$

If  $L/K$  is not finite, we can still define the maps both ways, but the analogous statement is more nuanced. The main detail is that  $\text{Aut}(L/K)$  is naturally profinite, hence carries a *profinite topology*, and taking Galois groups of intermediary extensions must produce *closed* subgroups in this topology. Luckily, we can refine the fundamental theorem: a (possibly infinite) field extension  $L/K$  is Galois if and only if the following restricted correspondence is bijective.

$$\{\text{intermediate field extensions } L/E/K\} \longleftrightarrow \{\text{closed subgroups of } \text{Aut}(L/K)\}$$

Another formulation of finite extensions emphasizes the Galois group. Given a finite, *faithful* group action  $G \rightarrow \text{Aut}(L)$ , the extension  $L/L^G$  is Galois, in fact finite with Galois group  $G = \text{Aut}(L/L^G)$ ! That’s a bit funny—faithfulness is equivalent to the *injectivity* of  $G \rightarrow \text{Aut}(L)$ , but it is not in my nature to suspect this restricts to an *equivalence*  $G \cong \text{Aut}(L/L^G)$  (whence  $L/L^G$  is Galois, in fact finite). This is proven in [Sta24, Lemma 09I3]. Conversely, given an extension  $L/K$ , its Galois group acts faithfully on  $L$ . Therefore, for a fixed  $L$ , we have

$$\text{finite Galois extensions } L/K \cong \text{finite, faithful group actions } G \hookrightarrow \text{Aut}(L).$$

I think the picture holds for arbitrary Galois extensions if you consider faithful actions by profinite groups. (Story about Brauer groups, extensions, and Galois extensions?)

We have many approaches to the Galois theory of fields. As to that of *commutative rings*, we are fat with food for thought. What is the correct generalization of Galois theory to rings? Moreover, what would conveniently extend to ring *spectra*? For this, we will bring to the fore the *linear algebra secretly powering the classical Galois theory of fields*. You have already seen footprints of linear algebra in Galois theory, e.g. in dimension counting arguments. Following Francis, we will set up Galois theory using only the linear independence of characters, Dedekind’s theorem (which is an immediate consequence of the independence of characters), and Artin’s lemma.

**Lemma II.1** (Linear independence of characters). Let  $G$  be a group and  $L$  a field. Then the set of characters  $\text{Hom}_{\text{Grp}}(G, L^\times)$  is a linearly independent subset of the  $L$ -vector space  $\text{Fun}_{\text{Set}}(G, L)$ .

Every nontrivial element of a field is invertible, i.e.  $K^\times = K - \{0\}$ , and ring maps must preserve zero, therefore restriction  $r^* : \text{Hom}(K, L) \rightarrow \text{Hom}_{\text{Grp}}(K^\times, L)$  loses no information, i.e.  $r^*$  is injective. Moreover, these are both  $L$ -vector spaces and  $r^*$  is linear, whence the linear independence of characters extends to  $\text{Hom}(K, L)$ .

**Corollary II.1** (Dedekind’s lemma). Let  $\{\sigma_i : K \rightarrow L\}$  denote a finite set of distinct field homomorphisms. Then  $\{\sigma_i\}$  is  $L$ -linearly independent. In other words,  $\text{Hom}(K, L)$  is  $L$ -linearly independent as a subset of  $\text{Fun}_{\text{Set}}(K, L)$ . In particular,  $\text{Aut}(K)$  is  $K$ -linearly independent.

Now consider a subextension  $F \hookrightarrow K \hookrightarrow L$ . Field morphisms  $K \rightarrow L$  fixing  $F$  are  $F$ -linear, that is to say  $\text{Hom}(K, L; /F) \subseteq \text{Vect}_F(K, L)$ . The latter space’s  $L$ -dimension is  $[K : F]$ . Dedekind’s lemma says that  $\text{Hom}(K, L; /F)$  is  $L$ -linearly independent, hence this set is no bigger than that dimension.

**Proposition II.14.** Let  $F \hookrightarrow K \hookrightarrow L$  denote a subextension of fields. Then the number of field maps  $K \rightarrow L$  fixing  $F$  is bounded by the  $L$ -dimension of  $\text{Vect}_F(K, L)$ . In particular, taking  $K = L$ , we find that the number of automorphisms of  $K/F$  is so bounded. That is to say, in equations:

$$|\text{Hom}(K, L; /F)| \leq [K : F] \quad \text{and} \\ |\text{Aut}(L/F)| \leq [L : F].$$

One characterization of finite Galois extensions recognizes their achievement of equality in Proposition II.14. For the purpose of “inventing Galois theory linearly,” we can make this our definition: say  $L/F$  is a *finite Galois extension* if  $|\text{Aut}(L/F)| = [L : F]$ , which (by Dedekind’s lemma) is equivalent to saying that  $\text{Aut}(L/F)$  constitutes a basis for the  $L$ -vector space  $\text{Vect}_F(L, L)$ .

Thus, the basic language takes only linear algebra. What about the basic structure? Let’s see what the orbit-stabilizer theorem has to say. Let me state the version I will use: if  $G$  acts on  $X$  and  $S \subseteq X$  is a subset, then  $|G| = |\text{Stab}(S)| \cdot |\text{Orb}(S)|$  where  $\text{Orb}(S)$  denotes the embeddings  $S \hookrightarrow X$  of the form  $s \mapsto gs$  for some fixed  $g$ . We want to take this with  $G = \text{Aut}(L/F)$  acting on  $L$  with distinguished subset  $K$ .

**Proposition II.15** (Orbit-stabilizer implies subextension structure). Let  $F \hookrightarrow K \hookrightarrow L$  denote a subextension of fields. Then we have

$$|\text{Aut}(L/F)| \leq [L : K][K : F].$$

Furthermore, if  $L/F$  is Galois, then so is  $L/K$ , and the above inequality is an equality.

*Proof.* Take  $G = \text{Aut}(L/F)$ ,  $X = L$ , and  $S = K$ . Then the orbit-stabilizer theorem says that

$$|\text{Aut}(L/F)| = \underbrace{|\text{Stab}(K)|}_{=\text{Aut}(L/K)} \cdot \underbrace{|\text{Orb}(K)|}_{=\{ \phi|_K : \phi \in \text{Aut}(L/F) \} \subseteq \text{Hom}(K, L; F)}.$$

Noting what I’ve indicated with underbraces, we find  $|G| \leq [L : K][K : F]$  by Proposition II.14. This proves the first part of the proposition. Now if  $L/F$  is Galois, then by definition  $|\text{Aut}(L/F)| = [L : F]$ . Since  $[L : F] = [L : K][K : F]$ , the desired equalities follow.  $\square$

**Corollary II.2.** Given a subextension  $L/K/F$ , we know that  $\text{Orb}_{\text{Aut}(L/F)}(K)$  sits within  $\text{Hom}(K, L; F)$ . (Recall  $\text{Hom}(K, L; F)$  is  $L$ -linearly independent by Dedekind’s lemma, thus  $|\text{Hom}(K, L; F)| \leq [K : F]$ .) Our proof argues that if  $L/F$  is Galois, then among other things we have  $|\text{Orb}(K)| = [K : F]$ , necessitating  $\text{Orb}(K) = \text{Hom}(K, L; F)$ .

In other words, we showed that if  $L/F$  is Galois, then every field morphism  $K \rightarrow L$  fixing  $F$  extends to an automorphism of  $L$ , i.e. restriction  $\text{Aut}(L/F) \rightarrow \text{Hom}(K, L; F)$  is surjective.

**Remark II.6.** This setup explains why composing Galois extensions is tricky. Suppose  $L/K$  and  $K/F$  are finite Galois extensions. Orbit-stabilizer says that  $|\text{Aut}(L, F)| = |\text{Aut}(L/K)| |\text{Orb}(K)|$  which equals  $[L : K] \cdot |\text{Orb}(K)|$  since  $L/K$  is Galois. Thus  $L/F$  is Galois if and only if  $|\text{Orb}(K)| = [K : F]$ .

By virtue of subextensions, we have an inclusion  $\text{Orb}(K) \hookrightarrow \text{Hom}(K, L; F)$ , which establishes  $|\text{Orb}(K)| \leq [K : F]$ . Then  $L/F$  is Galois if and only if this is surjective, i.e. iff every  $K \rightarrow L$  fixing  $F$  extends to an automorphism  $L \rightarrow L$ . This is a nontrivial extension problem.

This so far has not acknowledged our assumption that  $K/F$  is Galois. That implies  $[K : F] = |\text{Aut}(K/F)|$ , so  $K/F$  being Galois means  $L/F$  is Galois iff  $|\text{Orb}(K)| = |\text{Aut}(K/F)|$ . Note that  $\text{Aut}(K/F)$  is also a subset of  $\text{Hom}(K, L; F)$  via postcomposing  $K \hookrightarrow L$ , so  $L/F$  is Galois iff  $\text{Orb}(K) \hookrightarrow \text{Hom}(K, L; F)$  has image  $\text{Aut}(K/F)$ . For this to happen, everything must coincide anyway, i.e.  $\text{Orb}(K) = \text{Aut}(K/F) = \text{Hom}(K, L; F)$ , so we’ve not managed to make our problem any easier by assuming  $K/F$  is Galois.

**Theorem II.2** (Fundamental theorem of Galois theory). Let  $L/F$  be a finite Galois extension. Then (blah blah blah)

*Proof.* Prove with the above + Artin’s lemma. Not hard. Essential detail is that for  $L/K/F$ , have equality  $K = L^{\text{Aut}(L/K)}$ , and that for finite  $H \leq \text{Aut}(L)$ , have  $H = \text{Aut}(L/H)$ .  $\square$

## II.6 (2/26) Linear Galois theory II

Last time, we worked up to the fundamental theorem of Galois theory using basic linear algebra and some group theory, following Michael (c.f. the references from last time). This was both a refresher of my Galois theory and an introduction to the linear algebra in Galois theory. The point is to work up to the Galois theory of rings and ring spectra, which “emphasizes (takes as definition) the linear algebraic point of view.” (Although, as I continue to learn things, I’m less convinced this is the correct slogan.) But what exactly

Maybe finish writing the above. Ring-theoretic correspondence?

does that mean? Unfortunately, the picture is not so simple as to somehow seamlessly extend the previous post’s approach. I knew going in that  $\text{End}_K(L)$  played some crucial role, and I wanted the previous post to elucidate this. We *did* get some  $\text{End}_K(L)$  action last time (c.f. the role of  $\text{Vect}_K(L, L) = \text{End}_K(L)$ ), but maybe not enough. (That is not to say the last post was not productive, just maybe did not progress my point.) I’ve since found some more good references:

- Greither’s book *Cyclic Galois extensions of commutative rings* [Gre06] which I accessed here,
- Keith Conrad’s notes “Linear independence of characters” found here, and
- Keith Conrad’s notes “Galois descent” found here, and
- Baez’s nLab post about group cohomology, homotopy fixed points, and Galois descent, and
- This MO question and the links therein.

Let’s keep working on this. That is, let me keep trying to *find*  $\text{End}_K(L)$  and ultimately the definition of *Galois extensions of rings*, rather than serve it up instantly and mysteriously.

Let me return to my earlier comment that one can take an approach to Galois theory “emphasizing the Galois group.” That is, we can characterize finite Galois extensions as faithful finite group actions  $G \rightarrow \text{Aut}(L)$ . Such a thing encodes an extension  $L/L^G$  with the (funny) property that  $\text{Aut}(L/L^G) = G$ . (That property is precisely the magic in classical the Galois correspondence.)

This approach is a little too slick. Recall that Galois extensions are characterized as the extensions which are algebraic, normal, and separable. These are essential properties—both for how they interact to effect Galois extensions and for their independent importance—but it is not obvious how these properties manifest in a faithful action  $G \rightarrow \text{Aut}(L)$ , let alone how to isolate and study them.

**Proposition II.16.**

## II.7 (2/28) Descent I

I have been trying to understand Galois extensions of ring spectra. The definition uses an interpretation of Galois extensions that was, upon my initial reading, unfamiliar to me. I am still trying to unpack this interpretation. (It is easy to read, I am just pedantically trying to make context for it.) This is giving me a rather hard time. Somehow, my impression is that this interpretation is closely related to “descent philosophy.” E.g., my hunch is that a baby case of descent explains the funny fact that if a finite group  $G$  acts by automorphisms on  $L$ , then  $L/L^G$  is  $G$ -Galois. Unclear to me is the precise relation between descent properties and Galois extensions—an example question is, *are the Galois extensions (using this definition) somehow precisely those satisfying (some form of) descent?* Or is the relation just that *when we take this as the definition of Galois extensions, (Galois) descent is more naturally “available”?*

Descent is something I have wanted to understand for a while but which I have avoided. I think I will try to get into it now. But even though Galois theory most recently brought me to think about descent, I will approach it from a more elementary point than is necessary for my earlier purposes, at least to start. I do this in part because I think I will learn it better, more generally, and hopefully in a way that sets up for categorical machinery.

Here are some references I dug up this morning, certainly enough to keep me busy for a while:

- Whatever is in HTT [Lur08], haven’t checked but I’m sure it’s in there,
- Charles’ 2019 Leeds lectures on higher topos theory, the notes for which I find on his website,
- Hess’s “A general framework for homotopic descent and codescent” [Hes10],
- Caenepeel’s “Galois corings from the descent theory point of view” [Cae03],
- Hohl’s “An introduction to field extensions and Galois descent for sheaves of vector spaces” [Hoh23],
- SheafifiedSarah’s blog post about descent in graph theory here,
- Keith Conrad’s notes “Galois descent” found here,

- Baez's nLab post about group cohomology, homotopy fixed points, and Galois descent, and all the links therein; and
- Vistoli's notes [Vis07].

# Bibliography

- [Mah58] Kurt Mahler. “An Interpolation Series for Continuous Functions of a p-adic Variable.” In: (1958).
- [Fie78] Zbigniew Fiedorowicz. “The Quillen-Grothendieck construction and extensions of pairings”. In: *Geometric Applications of Homotopy Theory I: Proceedings, Evanston, March 21–26, 1977*. Ed. by M. G. Barratt and M. E. Mahowald. Berlin, Heidelberg: Springer Berlin Heidelberg, 1978, pp. 163–169. ISBN: 978-3-540-35809-1. DOI: 10.1007/BFb0069233. URL: <https://doi.org/10.1007/BFb0069233>.
- [May80] J.P. May. “Pairings of categories and spectra”. In: *Journal of Pure and Applied Algebra* 19 (1980), pp. 299–346. ISSN: 0022-4049. DOI: [https://doi.org/10.1016/0022-4049\(80\)90105-X](https://doi.org/10.1016/0022-4049(80)90105-X). URL: <https://www.sciencedirect.com/science/article/pii/002240498090105X>.
- [She82] Clayton Sherman. “Group representations and algebraic K-theory”. In: *Algebraic K-Theory*. Ed. by R. Keith Dennis. Berlin, Heidelberg: Springer Berlin Heidelberg, 1982, pp. 208–243. ISBN: 978-3-540-39553-9.
- [Dre95] Andreas WM Dress. “One more shortcut to Galois theory”. In: *Advances in Mathematics* 110.1 (1995), pp. 129–140.
- [BDR03] Nils A. Baas, Bjørn Ian Dundas, and John Rognes. “Two-vector bundles and forms of elliptic cohomology”. In: (2003). arXiv: math/0306027 [math.AT].
- [Cae03] S. Caenepeel. *Galois corings from the descent theory point of view*. 2003. arXiv: math/0311377 [math.RA].
- [Rog05] John Rognes. *Galois extensions of structured ring spectra*. 2005. arXiv: math / 0502183 [math.AT].
- [Gre06] Cornelius Greither. *Cyclic Galois extensions of commutative rings*. Springer, 2006.
- [KS06] Masaki Kashiwara and Pierre Schapira. *Categories and Sheaves*. Springer, 2006.
- [Vis07] Angelo Vistoli. *Notes on Grothendieck topologies, fibered categories and descent theory*. 2007. arXiv: math/0412512 [math.AG].
- [Lur08] Jacob Lurie. *Higher Topos Theory*. 2008. arXiv: math/0608040 [math.CT].
- [GQ09] Rod Gow and Rachel Quinlan. “Galois theory and linear algebra”. In: *Linear Algebra and its Applications* 430.7 (2009), pp. 1778–1789.
- [Kra09] Henning Krause. *Localization theory for triangulated categories*. 2009. arXiv: 0806.1324 [math.CT].
- [Lur09] Jacob Lurie. “Derived Algebraic Geometry I: Stable  $\infty$ -categories”. In: (2009).
- [Hes10] Kathryn Hess. *A general framework for homotopic descent and codescent*. 2010. arXiv: 1001.1556 [math.AT].
- [FD12] Benson Farb and R Keith Dennis. *Noncommutative algebra*. Vol. 144. Springer Science & Business Media, 2012.
- [Lur17] Jacob Lurie. *Higher Algebra*. 2017.

- [Law20] Tyler Lawson. “An introduction to Bousfield localization”. In: (2020). arXiv: 2002.03888 [math.AT].
- [Rez22] Charles Rezk. “Free colimit completion in  $\infty$ -categories”. In: (2022). arXiv: 2210.08582 [math.CT].
- [Ben+23] Shay Ben-Moshe et al. *Descent and Cyclotomic Redshift for Chromatically Localized Algebraic K-theory*. 2023. arXiv: 2309.07123 [math.KT].
- [Du23] Peng Du. *Reshaping limit diagrams and cofinality in higher category theory*. 2023. arXiv: 2308.01798 [math.CT].
- [Hoh23] Andreas Hohl. *An introduction to field extensions and Galois descent for sheaves of vector spaces*. 2023. arXiv: 2302.14837 [math.AG].
- [Lur24] Jacob Lurie. *Kerodon*. <https://kerodon.net>. 2024.
- [Sta24] The Stacks project authors. *The Stacks project*. <https://stacks.math.columbia.edu>. 2024.