

$\mathbb{U}_p$ ,  $\mathbb{U}_{p^k}$ , and  $\mathbb{U}_{2p^k}$  are cyclic

Matthew Li and Zack Yu

July 2023

# Table of Contents

1 Introduction

2  $\mathbb{U}_p$

3  $\mathbb{U}_{p^k}$

4  $\mathbb{U}_{2p^k}$

# Definitions

## Set of Generators

A set of generators  $(g_1, \dots, g_n)$  is a set of elements of a group  $G$  such that performing the group operation on themselves and on each other is capable of producing all the elements in the group.

# Definitions

## Set of Generators

A set of generators  $(g_1, \dots, g_n)$  is a set of elements of a group  $G$  such that performing the group operation on themselves and on each other is capable of producing all the elements in the group.

## Cyclic Group

A cyclic group is a group that is generated by a single generator.

# Motivating Example

Let us observe an example of  $\mathbb{U}_p$ . We will try to see if it is cyclic.

# Motivating Example

Let us observe an example of  $\mathbb{U}_p$ . We will try to see if it is cyclic.

## Example

Consider  $\mathbb{U}_5$ . The elements of  $\mathbb{U}_5$  are 1, 2, 3, and 4. One can see that 2 is a generator of  $\mathbb{U}_5$  because

$$2^1 = 2$$

$$2^2 = 4$$

$$2^3 = 3$$

$$2^4 = 1$$

Thus,  $\mathbb{U}_5$  is cyclic.

## Another Example

Not convinced? Let's consider a bigger  $p$ .

# Another Example

Not convinced? Let's consider a bigger  $p$ .

## Example

Consider  $\mathbb{U}_{13}$ . We claim that 2 is a generator of  $\mathbb{U}_{13}$ .

$$\begin{array}{cccc}
 2^1 = 2 & 2^4 = 3 & 2^7 = 11 & 2^{10} = 10 \\
 2^2 = 4 & 2^5 = 6 & 2^8 = 9 & 2^{11} = 7 \\
 2^3 = 8 & 2^6 = 12 & 2^9 = 5 & 2^{12} = 1
 \end{array}$$



# Proof Sketch of $\mathbb{U}_p$ Cyclic

We need a couple of important results before starting.

# Proof Sketch of $\mathbb{U}_p$ Cyclic

We need a couple of important results before starting.

## Lagrange's Theorem

For a degree  $n$  polynomial mod  $p$ , it has at most  $n$  roots.

# Proof Sketch of $\mathbb{U}_p$ Cyclic

We need a couple of important results before starting.

## Lagrange's Theorem

For a degree  $n$  polynomial mod  $p$ , it has at most  $n$  roots.

## Lemma 1

If  $u \in \mathbb{U}_p$  has order  $d$ , then there exists  $x \in \mathbb{U}_p$  such that  $\text{ord}_p(x) = k$ , where  $k$  is a divisor of  $d$ .

# Proof Sketch of $\mathbb{U}_p$ Cyclic

We need a couple of important results before starting.

## Lagrange's Theorem

For a degree  $n$  polynomial mod  $p$ , it has at most  $n$  roots.

## Lemma 1

If  $u \in \mathbb{U}_p$  has order  $d$ , then there exists  $x \in \mathbb{U}_p$  such that  $\text{ord}_p(x) = k$ , where  $k$  is a divisor of  $d$ .

## Lemma 2

If  $\text{ord}_p(a) = r$ ,  $\text{ord}_p(b) = s$ , and  $\gcd(r, s) = 1$ , then  $\text{ord}_p(ab) = rs$ .

# Proof Sketch of $\mathbb{U}_p$ Cyclic

- We wish to show there is  $u \in \mathbb{U}_p$  of order  $p - 1$ .

# Proof Sketch of $\mathbb{U}_p$ Cyclic

- We wish to show there is  $u \in \mathbb{U}_p$  of order  $p - 1$ .
- $p - 1 = q_1^{e_1} \cdots q_k^{e_k}$

# Proof Sketch of $\mathbb{U}_p$ Cyclic

- We wish to show there is  $u \in \mathbb{U}_p$  of order  $p - 1$ .
- $p - 1 = q_1^{e_1} \cdots q_k^{e_k}$
- There is  $w \in \mathbb{U}_p$  such that  $w^{\frac{p-1}{q_i}} \neq 1$  in  $\mathbb{U}_p$ .

# Proof Sketch of $\mathbb{U}_p$ Cyclic

- We wish to show there is  $u \in \mathbb{U}_p$  of order  $p - 1$ .
- $p - 1 = q_1^{e_1} \cdots q_k^{e_k}$
- There is  $w \in \mathbb{U}_p$  such that  $w^{\frac{p-1}{q_i}} \neq 1$  in  $\mathbb{U}_p$ .
- For sake of contradiction, assume all  $w \in \mathbb{U}_p$  satisfy above. Then, we get contradiction from Lagrange.



# Proof Sketch of $\mathbb{U}_p$ Cyclic

- We wish to show there is  $u \in \mathbb{U}_p$  of order  $p - 1$ .
- $p - 1 = q_1^{e_1} \cdots q_k^{e_k}$
- There is  $w \in \mathbb{U}_p$  such that  $w^{\frac{p-1}{q_i}} \neq 1$  in  $\mathbb{U}_p$ .
- For sake of contradiction, assume all  $w \in \mathbb{U}_p$  satisfy above. Then, we get contradiction from Lagrange.
- So order of  $w$  should not divide  $\frac{p-1}{q_i}$ . Thus, all orders must be divisible by  $q_i^{e_i}$ .

# Proof Sketch of $\mathbb{U}_p$ Cyclic

- We wish to show there is  $u \in \mathbb{U}_p$  of order  $p - 1$ .
- $p - 1 = q_1^{e_1} \cdots q_k^{e_k}$
- There is  $w \in \mathbb{U}_p$  such that  $w^{\frac{p-1}{q_i}} \neq 1$  in  $\mathbb{U}_p$ .
- For sake of contradiction, assume all  $w \in \mathbb{U}_p$  satisfy above. Then, we get contradiction from Lagrange.
- So order of  $w$  should not divide  $\frac{p-1}{q_i}$ . Thus, all orders must be divisible by  $q_i^{e_i}$ .
- Thus, there is  $x \in \mathbb{U}_p$  of order  $q_i^{e_i}$  by Lemma 1.

# Proof Sketch of $\mathbb{U}_p$ Cyclic

- We wish to show there is  $u \in \mathbb{U}_p$  of order  $p - 1$ .
- $p - 1 = q_1^{e_1} \cdots q_k^{e_k}$
- There is  $w \in \mathbb{U}_p$  such that  $w^{\frac{p-1}{q_i}} \neq 1$  in  $\mathbb{U}_p$ .
- For sake of contradiction, assume all  $w \in \mathbb{U}_p$  satisfy above. Then, we get contradiction from Lagrange.
- So order of  $w$  should not divide  $\frac{p-1}{q_i}$ . Thus, all orders must be divisible by  $q_i^{e_i}$ .
- Thus, there is  $x \in \mathbb{U}_p$  of order  $q_i^{e_i}$  by Lemma 1.
- We repeat this process for all  $i$  so  $\text{ord}_p(w_i) = q_i^{e_i}$  for all  $i$ .

# Proof Sketch of $\mathbb{U}_p$ Cyclic

- We wish to show there is  $u \in \mathbb{U}_p$  of order  $p - 1$ .
- $p - 1 = q_1^{e_1} \cdots q_k^{e_k}$
- There is  $w \in \mathbb{U}_p$  such that  $w^{\frac{p-1}{q_i}} \neq 1$  in  $\mathbb{U}_p$ .
- For sake of contradiction, assume all  $w \in \mathbb{U}_p$  satisfy above. Then, we get contradiction from Lagrange.
- So order of  $w$  should not divide  $\frac{p-1}{q_i}$ . Thus, all orders must be divisible by  $q_i^{e_i}$ .
- Thus, there is  $x \in \mathbb{U}_p$  of order  $q_i^{e_i}$  by Lemma 1.
- We repeat this process for all  $i$  so  $\text{ord}_p(w_i) = q_i^{e_i}$  for all  $i$ .
- Then, by Lemma 2,  $\text{ord}_p(w_1 \cdots w_i) = p - 1$ , as desired.

# 1 Introduction

## 2 $\mathbb{U}_p$

## 3 $\mathbb{U}_{p^k}$

## 4 $\mathbb{U}_{2p^k}$

# Examples

It is not immediately obvious that  $\mathbb{U}_{p^k}$  is cyclic for all odd primes  $p$ .

# Examples

It is not immediately obvious that  $\mathbb{U}_{p^k}$  is cyclic for all odd primes  $p$ .

## Example

Consider  $\mathbb{U}_{3^2}$ . The elements of  $\mathbb{U}_9$  are 1, 2, 4, 5, 7, and 8. We claim that 5 is a generator.

$$\begin{array}{ll} 5^1 = 5 & 5^4 = 4 \\ 5^2 = 7 & 5^5 = 2 \\ 5^3 = 8 & 5^6 = 1 \end{array}$$

# Examples

Consider  $\mathbb{U}_{3^3}$ . Through some tedious computation we can show 5 is also a generator of  $\mathbb{U}_{3^3}$ .



# Examples

Consider  $\mathbb{U}_{3^3}$ . Through some tedious computation we can show 5 is also a generator of  $\mathbb{U}_{3^3}$ .

## Example

|            |               |
|------------|---------------|
| $5^1 = 5$  | $5^{10} = 22$ |
| $5^2 = 25$ | $5^{11} = 2$  |
| $5^3 = 17$ | $5^{12} = 10$ |
| $5^4 = 4$  | $5^{13} = 23$ |
| $5^5 = 20$ | $5^{14} = 7$  |
| $5^6 = 19$ | $5^{15} = 8$  |
| $5^7 = 14$ | $5^{16} = 13$ |
| $5^8 = 16$ | $5^{17} = 11$ |
| $5^9 = 26$ | $5^{18} = 1$  |

# Proof Attempt

- One idea is to try the same approach we used in  $\mathbb{U}_p$ . What's wrong with this?

# Proof Attempt

- One idea is to try the same approach we used in  $\mathbb{U}_p$ . What's wrong with this?
- Notice that in proving Lagrange's Theorem, we required the property in  $\mathbb{Z}_p$  that if  $ab = 0$ , then  $a = 0$  or  $b = 0$ .

# Motivating the Proof

- We know that  $|\mathbb{U}_{p^k}| = \varphi(p^k) = p^{k-1}(p-1)$

# Motivating the Proof

- We know that  $|\mathbb{U}_{p^k}| = \varphi(p^k) = p^{k-1}(p-1)$
- We want to find a generator or an element of order  $p^{k-1}(p-1)$

# Motivating the Proof

- We know that  $|\mathbb{U}_{p^k}| = \varphi(p^k) = p^{k-1}(p-1)$
- We want to find a generator or an element of order  $p^{k-1}(p-1)$
- Since  $\gcd(p^{k-1}, p-1) = 1$ , if we can find elements of order  $p^{k-1}$  and order  $p-1$ , their product should have order  $p^{k-1}(p-1)$

# Motivating Example

## Example

Consider 4 and 2  $\in \mathbb{U}_9$ .

$$4^1 = 4 \quad 2^1 = 2$$

$$4^2 = 7 \quad 2^2 = 4$$

$$4^3 = 1 \quad 2^3 = 8$$

$$2^4 = 7$$

$$2^5 = 5$$

$$2^6 = 1$$

Therefore  $\text{ord}_9(4) = 3 = 3^{2-1}$  and  $\text{ord}_9(8) = 2 = (3 - 1)$ .

Then, the product 5 should have order  $3^{2-1}(3 - 1) = 6$  and should be a generator of  $\mathbb{U}_{p^2}$ . We have already showed this.

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We will first find an element of order  $p - 1$



# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We will first find an element of order  $p - 1$
- Consider a generator  $g$  of  $\mathbb{U}_p$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We will first find an element of order  $p - 1$
- Consider a generator  $g$  of  $\mathbb{U}_p$
- Let us say  $\text{ord}_{p^k}(g) = d$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We will first find an element of order  $p - 1$
- Consider a generator  $g$  of  $\mathbb{U}_p$
- Let us say  $\text{ord}_{p^k}(g) = d$
- Since we know  $g^d \equiv 1 \pmod{p^k}$ , we know  $g^d \equiv 1 \pmod{p}$ , so  $p - 1 \mid d$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We will first find an element of order  $p - 1$
- Consider a generator  $g$  of  $\mathbb{U}_p$
- Let us say  $\text{ord}_{p^k}(g) = d$
- Since we know  $g^d \equiv 1 \pmod{p^k}$ , we know  $g^d \equiv 1 \pmod{p}$ , so  $p - 1 \mid d$
- Then, by one of our earlier lemma's, we know there must exist an element with order  $p - 1$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- Now, let us find an element of order  $p^{k-1}$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- Now, let us find an element of order  $p^{k-1}$
- For some motivation, first consider  $\mathbb{U}_{p^2}$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- Now, let us find an element of order  $p^{k-1}$
- For some motivation, first consider  $\mathbb{U}_{p^2}$
- Assume  $x$  is the order of  $p+1$  in  $\mathbb{U}_{p^2}$ , then  $(p+1)^x \equiv 1 \pmod{p^2}$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- Now, let us find an element of order  $p^{k-1}$
- For some motivation, first consider  $\mathbb{U}_{p^2}$
- Assume  $x$  is the order of  $p+1$  in  $\mathbb{U}_{p^2}$ , then  $(p+1)^x \equiv 1 \pmod{p^2}$
- By expanding  $(p+1)^x$  with binomial theorem, we get

$$(p+1)^x = \binom{x}{0}p^x + \binom{x}{1}p^{x-1} + \cdots + \binom{x}{x-1}p + \binom{x}{x}1$$



# Proving $\mathbb{U}_{p^k}$ is Cyclic

- Now, let us find an element of order  $p^{k-1}$
- For some motivation, first consider  $\mathbb{U}_{p^2}$
- Assume  $x$  is the order of  $p+1$  in  $\mathbb{U}_{p^2}$ , then  $(p+1)^x \equiv 1 \pmod{p^2}$
- By expanding  $(p+1)^x$  with binomial theorem, we get

$$(p+1)^x = \binom{x}{0}p^x + \binom{x}{1}p^{x-1} + \cdots + \binom{x}{x-1}p + \binom{x}{x}1$$

- Thus, we have  $(p+1)^x \equiv px + 1 \pmod{p^2}$  since all terms with  $p^2$  disappear

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- Now, let us find an element of order  $p^{k-1}$
- For some motivation, first consider  $\mathbb{U}_{p^2}$
- Assume  $x$  is the order of  $p+1$  in  $\mathbb{U}_{p^2}$ , then  $(p+1)^x \equiv 1 \pmod{p^2}$
- By expanding  $(p+1)^x$  with binomial theorem, we get

$$(p+1)^x = \binom{x}{0}p^x + \binom{x}{1}p^{x-1} + \cdots + \binom{x}{x-1}p + \binom{x}{x}1$$

- Thus, we have  $(p+1)^x \equiv px + 1 \pmod{p^2}$  since all terms with  $p^2$  disappear
- Then, we get  $x \equiv 0 \pmod{p}$  or  $p \mid x$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- Now, let us find an element of order  $p^{k-1}$
- For some motivation, first consider  $\mathbb{U}_{p^2}$
- Assume  $x$  is the order of  $p + 1$  in  $\mathbb{U}_{p^2}$ , then  $(p + 1)^x \equiv 1 \pmod{p^2}$
- By expanding  $(p + 1)^x$  with binomial theorem, we get

$$(p + 1)^x = \binom{x}{0} p^x + \binom{x}{1} p^{x-1} + \cdots + \binom{x}{x-1} p + \binom{x}{x} 1$$

- Thus, we have  $(p + 1)^x \equiv px + 1 \pmod{p^2}$  since all terms with  $p^2$  disappear
- Then, we get  $x \equiv 0 \pmod{p}$  or  $p \mid x$
- Since  $x$  is the order of  $p + 1$  and  $p \mid x$ , there is another element  $h$  with order  $p$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We construct a set  $A = \{k \in \mathbb{Z}^+ \mid \text{the order of } p+1 \text{ in } \mathbb{U}_{p^k} \text{ is not } np^{k-1} \text{ for an integer } n\}$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We construct a set  $A = \{k \in \mathbb{Z}^+ \mid \text{the order of } p+1 \text{ in } \mathbb{U}_{p^k} \text{ is not } np^{k-1} \text{ for an integer } n\}$
- By WOP, there is a minimum element  $l$  of this set

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We construct a set  $A = \{k \in \mathbb{Z}^+ \mid \text{the order of } p+1 \text{ in } \mathbb{U}_{p^k} \text{ is not } np^{k-1} \text{ for an integer } n\}$
- By WOP, there is a minimum element  $l$  of this set
- Therefore, the order of  $p+1$  in  $\mathbb{U}_{p^{l-1}}$  is  $ap^{l-2}$ , for some integer  $a$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We construct a set  $A = \{k \in \mathbb{Z}^+ \mid \text{the order of } p+1 \text{ in } \mathbb{U}_{p^k} \text{ is not } np^{k-1} \text{ for an integer } n\}$
- By WOP, there is a minimum element  $l$  of this set
- Therefore, the order of  $p+1$  in  $\mathbb{U}_{p^{l-1}}$  is  $ap^{l-2}$ , for some integer  $a$
- We know if  $(p+1)^x \equiv 1 \pmod{p^l}$  then  $(p+1)^x \equiv 1 \pmod{p^{l-1}}$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We construct a set  $A = \{k \in \mathbb{Z}^+ \mid \text{the order of } p+1 \text{ in } \mathbb{U}_{p^k} \text{ is not } np^{k-1} \text{ for an integer } n\}$
- By WOP, there is a minimum element  $l$  of this set
- Therefore, the order of  $p+1$  in  $\mathbb{U}_{p^{l-1}}$  is  $ap^{l-2}$ , for some integer  $a$
- We know if  $(p+1)^x \equiv 1 \pmod{p^l}$  then  $(p+1)^x \equiv 1 \pmod{p^{l-1}}$
- Then  $ap^{l-2} \mid x$  or  $x = bap^{l-2}$  for some integer  $b$



# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We construct a set  $A = \{k \in \mathbb{Z}^+ \mid \text{the order of } p+1 \text{ in } \mathbb{U}_{p^k} \text{ is not } np^{k-1} \text{ for an integer } n\}$
- By WOP, there is a minimum element  $l$  of this set
- Therefore, the order of  $p+1$  in  $\mathbb{U}_{p^{l-1}}$  is  $ap^{l-2}$ , for some integer  $a$
- We know if  $(p+1)^x \equiv 1 \pmod{p^l}$  then  $(p+1)^x \equiv 1 \pmod{p^{l-1}}$
- Then  $ap^{l-2} \mid x$  or  $x = bap^{l-2}$  for some integer  $b$
- Now, we can expand  $(p+1)^x$  and obtain  $xp + 1 \equiv 1 \pmod{p^l}$  from binomial theorem

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We construct a set  $A = \{k \in \mathbb{Z}^+ \mid \text{the order of } p+1 \text{ in } \mathbb{U}_{p^k} \text{ is not } np^{k-1} \text{ for an integer } n\}$
- By WOP, there is a minimum element  $l$  of this set
- Therefore, the order of  $p+1$  in  $\mathbb{U}_{p^{l-1}}$  is  $ap^{l-2}$ , for some integer  $a$
- We know if  $(p+1)^x \equiv 1 \pmod{p^l}$  then  $(p+1)^x \equiv 1 \pmod{p^{l-1}}$
- Then  $ap^{l-2} \mid x$  or  $x = bap^{l-2}$  for some integer  $b$
- Now, we can expand  $(p+1)^x$  and obtain  $xp + 1 \equiv 1 \pmod{p^l}$  from binomial theorem
- Thus  $x = np^{l-1}$ , which contradicts  $l \in A$

# Proving $\mathbb{U}_{p^k}$ is Cyclic

- We construct a set  $A = \{k \in \mathbb{Z}^+ \mid \text{the order of } p+1 \text{ in } \mathbb{U}_{p^k} \text{ is not } np^{k-1} \text{ for an integer } n\}$
- By WOP, there is a minimum element  $l$  of this set
- Therefore, the order of  $p+1$  in  $\mathbb{U}_{p^{l-1}}$  is  $ap^{l-2}$ , for some integer  $a$
- We know if  $(p+1)^x \equiv 1 \pmod{p^l}$  then  $(p+1)^x \equiv 1 \pmod{p^{l-1}}$
- Then  $ap^{l-2} \mid x$  or  $x = bap^{l-2}$  for some integer  $b$
- Now, we can expand  $(p+1)^x$  and obtain  $xp + 1 \equiv 1 \pmod{p^l}$  from binomial theorem
- Thus  $x = np^{l-1}$ , which contradicts  $l \in A$
- Then, there is an element with order  $p^{k-1}$

# Finishing the Proof

By multiplying the elements of order  $p^{k-1}$  and  $p-1$ , we obtain an element of order  $p^{k-1}(p-1)$ , which generates  $\mathbb{U}_{p^k}$ , yay!

# 1 Introduction

## 2 $\mathbb{U}_p$

## 3 $\mathbb{U}_{p^k}$

## 4 $\mathbb{U}_{2p^k}$

# Examples

Let us try to convince ourselves  $\mathbb{U}_{2p^k}$  is cyclic for all odd primes  $p$ .

## Example

Consider  $\mathbb{U}_{2 \cdot 3}$ . We know the elements of  $\mathbb{U}_6$  are 1 and 5. Clearly, the element 5 generates  $\mathbb{U}_6$ .

Consider  $\mathbb{U}_{2 \cdot 3^2}$ . Then, we claim 5 is a generator.

$$\begin{array}{ll} 5^1 = 5 & 5^4 = 13 \\ 5^2 = 7 & 5^5 = 11 \\ 5^3 = 17 & 5^6 = 1 \end{array}$$

# Motivating the Proof

- $\mathbb{U}_{2p^k}$  is not something we are very familiar with, let's see if we can find an isomorphism to something “simpler” that we can work with

# Motivating the Proof

- $\mathbb{U}_{2p^k}$  is not something we are very familiar with, let's see if we can find an isomorphism to something “simpler” that we can work with
- Recall that  $\mathbb{Z}_m \times \mathbb{Z}_n \cong \mathbb{Z}_{mn}$



# Motivating the Proof

- $\mathbb{U}_{2p^k}$  is not something we are very familiar with, let's see if we can find an isomorphism to something “simpler” that we can work with
- Recall that  $\mathbb{Z}_m \times \mathbb{Z}_n \cong \mathbb{Z}_{mn}$
- Since  $\gcd(2, p^k) = 1$ , we are motivated to see if there is an isomorphism between  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  and  $\mathbb{U}_{2p^k}$

# Motivating the Proof

- $\mathbb{U}_{2p^k}$  is not something we are very familiar with, let's see if we can find an isomorphism to something “simpler” that we can work with
- Recall that  $\mathbb{Z}_m \times \mathbb{Z}_n \cong \mathbb{Z}_{mn}$
- Since  $\gcd(2, p^k) = 1$ , we are motivated to see if there is an isomorphism between  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  and  $\mathbb{U}_{2p^k}$
- Notice that  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  is very nice to work with since the elements are just  $(1, x)$  where  $x$  is an element of  $\mathbb{U}_{p^k}$

# Motivating the Proof

- $\mathbb{U}_{2p^k}$  is not something we are very familiar with, let's see if we can find an isomorphism to something “simpler” that we can work with
- Recall that  $\mathbb{Z}_m \times \mathbb{Z}_n \cong \mathbb{Z}_{mn}$
- Since  $\gcd(2, p^k) = 1$ , we are motivated to see if there is an isomorphism between  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  and  $\mathbb{U}_{2p^k}$
- Notice that  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  is very nice to work with since the elements are just  $(1, x)$  where  $x$  is an element of  $\mathbb{U}_{p^k}$
- Thus,  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  is basically identical to  $\mathbb{U}_{p^k}$ .

# Motivating the Proof

- $\mathbb{U}_{2p^k}$  is not something we are very familiar with, let's see if we can find an isomorphism to something "simpler" that we can work with
- Recall that  $\mathbb{Z}_m \times \mathbb{Z}_n \cong \mathbb{Z}_{mn}$
- Since  $\gcd(2, p^k) = 1$ , we are motivated to see if there is an isomorphism between  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  and  $\mathbb{U}_{2p^k}$
- Notice that  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  is very nice to work with since the elements are just  $(1, x)$  where  $x$  is an element of  $\mathbb{U}_{p^k}$
- Thus,  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  is basically identical to  $\mathbb{U}_{p^k}$ .
- So, we know  $\mathbb{U}_2 \times \mathbb{U}_{p^k}$  is cyclic and if  $\mathbb{U}_2 \times \mathbb{U}_{p^k} \cong \mathbb{U}_{2p^k}$ , we know  $\mathbb{U}_{2p^k}$  is cyclic

# Establishing the Isomorphisms

- First, we show that  $\mathbb{U}_{2p^k} \cong \mathbb{U}_2 \times \mathbb{U}_{p^k}$

# Establishing the Isomorphisms

- First, we show that  $\mathbb{U}_{2p^k} \cong \mathbb{U}_2 \times \mathbb{U}_{p^k}$
- We claim the mapping  $f : \mathbb{U}_{2p^k} \rightarrow \mathbb{U}_2 \times \mathbb{U}_{p^k}$  defined by  $[a]_{2p^k} \mapsto ([a]_2, [a]_{p^k})$  is bijective and satisfies  $f(ab) = f(a)f(b)$ .

# Establishing the Isomorphisms

- First, we show that  $\mathbb{U}_{2p^k} \cong \mathbb{U}_2 \times \mathbb{U}_{p^k}$
- We claim the mapping  $f : \mathbb{U}_{2p^k} \rightarrow \mathbb{U}_2 \times \mathbb{U}_{p^k}$  defined by  $[a]_{2p^k} \mapsto ([a]_2, [a]_{p^k})$  is bijective and satisfies  $f(ab) = f(a)f(b)$ .
- Note, to prove bijectivity, we only need to prove injectivity because  $|\mathbb{U}_{2p^k}| = |\mathbb{U}_2 \times \mathbb{U}_{p^k}|$ .

# Establishing the Isomorphisms

- First, we show that  $\mathbb{U}_{2p^k} \cong \mathbb{U}_2 \times \mathbb{U}_{p^k}$
- We claim the mapping  $f : \mathbb{U}_{2p^k} \rightarrow \mathbb{U}_2 \times \mathbb{U}_{p^k}$  defined by  $[a]_{2p^k} \mapsto ([a]_2, [a]_{p^k})$  is bijective and satisfies  $f(ab) = f(a)f(b)$ .
- Note, to prove bijectivity, we only need to prove injectivity because  $|\mathbb{U}_{2p^k}| = |\mathbb{U}_2 \times \mathbb{U}_{p^k}|$ .
- Now, we show  $\mathbb{U}_2 \times \mathbb{U}_{p^k} \cong \mathbb{U}_{p^k}$ .



# Establishing the Isomorphisms

- First, we show that  $\mathbb{U}_{2p^k} \cong \mathbb{U}_2 \times \mathbb{U}_{p^k}$
- We claim the mapping  $f : \mathbb{U}_{2p^k} \rightarrow \mathbb{U}_2 \times \mathbb{U}_{p^k}$  defined by  $[a]_{2p^k} \mapsto ([a]_2, [a]_{p^k})$  is bijective and satisfies  $f(ab) = f(a)f(b)$ .
- Note, to prove bijectivity, we only need to prove injectivity because  $|\mathbb{U}_{2p^k}| = |\mathbb{U}_2 \times \mathbb{U}_{p^k}|$ .
- Now, we show  $\mathbb{U}_2 \times \mathbb{U}_{p^k} \cong \mathbb{U}_{p^k}$ .
- Simply consider the map  $f : \mathbb{U}_{p^k} \rightarrow \mathbb{U}_2 \times \mathbb{U}_{p^k}$  defined by  $x \mapsto (1, x)$

# Finishing the Proof

Because  $\mathbb{U}_{p^k} \cong \mathbb{U}_2 \times \mathbb{U}_{p^k} \cong \mathbb{U}_{2p^k}$ , we know  $\mathbb{U}_{2p^k}$  is cyclic, yay!