# Spatial Data is Special Data

HES 505 Fall 2022: Session 6

Matt Williamson

### Space Exploration

### Objectives

- Articulate why we care about space
- Describe elements of spatial data
- Define a coordinate reference system and its importance
- Describe several ways to load spatial data into R
- Identify projections in R

## Locations, Relations, and Understanding

- **Geography** uses *location* to understand how social and physical processes give rise to the environment we experience
- Geographic Information Systems provide a structure for storing, visualizing, and describing *location* data.
- **GeoComputation** and **GIScience** integrate math, stats, and high-performance computing to move beyond description.

### Location lets us ask:

- Questions about geographic distribution
- Questions about geographic interaction
- Questions about geographic change
- Questions about geographic association
- Questions about causation?

#### Location vs. Place


- Place: an area having unique physical and human characteristics interconnected with other places
- Location: the actual position on the earth's surface
- Sense of Place: the emotions someone attaches to an area based on experiences
- Place is *location plus meaning*

### **Describing Location**

- nominal: (potentially contested) place names
- absolute: the physical location on the earth's surface

### Describing Absolute Locations


- **Coordinates:** 2 or more measurements that specify location relative to a *reference system*
- Cartesian coordinate system
- *origin* (*O*) = the point at which both measurement systems intersect
- Adaptable to multiple dimensions (e.g. *z* for altitude)



Cartesian Coordinate System

#### Locations on a Globe

• The earth is not flat...



Latitude and Longitude

- Global Reference Systems (GRS)
- *Graticule*: the grid formed by the intersection of longitude and latitude
- The graticule is based on an ellipsoid model of earth's surface and contained in the *datum*


### Global Reference Systems

The *datum* describes which ellipsoid to use and the precise relations between locations on earth's surface and Cartesian coordinates

- Geodetic datums (e.g., WGS84): distance from earth's center of gravity
- Local data (e.g., NAD83): better models for local variation in earth's surface

### The Earth is Not Flat

- But maps, screens, and publications are...
- **Projections** describe *how* the data should be translated to a flat surface
- Rely on 'developable surfaces'



Developable Surfaces

Projection necessarily induces some form of distortion (tearing, compression, or shearing)

### **Choosing Projections**



Neil Kaye/@neilrkaye

The Mercator Map Projection with the true size and shape of the country overlaid.

- Some projections minimize distortion of angle, area, or distance
- Others attempt to avoid extreme distortion of any kind
- Particularly challenging for raster data


### **Choosing Projections**

- Equal-area for thematic maps
- Conformal for presentations
- Mercator or equidistant for navigation and distance

### Mapping Location in R

## Data Types and R Packages Data Types

- Vector Data
  - Point features
  - Line features
  - Area features (polygons)
- Raster Data
  - Spatially continuous field
  - Based on pixels (not points)





## Mapping loaction: Coordinate Reference Systems

- Includes: Datum, ellipsoid, units, and other information (e.g., False Easting, Central Meridian) to further map the projection to the GCS
- Not all projections have/require all of the parameters
- R stores these data in several formats (EPSG, Proj, and WKT)
- Lots of projection info available at spatialreference.org

## Mapping loaction: Coordinate Reference Systems


Primarily accessed using sf::st\_crs() or terra::crs()


```
1 f <- rast(system.file("ex/meuse.tif", package="terra"))
2 nc <- st_read(system.file("shape/nc.shp", package="sf"))
3 st_crs(nc)
4 crs(nc)</pre>
```

### Mapping location: Extent

• Extent: The amount of the Earth's surface represented by the mapped features

R has a very specific definition of extent: the rectangular region encompassed by the data





### Using R to access the extent

Using st\_bbox() from the sf package

178400 329400 181600 334000

```
1 nc.shp <- st read(system.file("shape/nc.shp", package="sf"))</pre>
Reading layer `nc' from data source
\Library/Frameworks/R.framework/Versions/4.2/Resources/library/sf/shape/nc.shp
  using driver `ESRI Shapefile'
Simple feature collection with 100 features and 14 fields
Geometry type: MULTIPOLYGON
Dimension:
              XY
Bounding box: xmin: -84.32385 ymin: 33.88199 xmax: -75.45698 ymax: 36.58965
Geodetic CRS: NAD27
 1 meuse.rst <- rast(system.file("ex/meuse.tif", package="terra"))</pre>
 2 st bbox(nc.shp)
     xmin
              vmin xmax
                                  ymax
-84.32385 33.88199 -75.45698 36.58965
 1 st bbox(meuse.rst)
  xmin ymin xmax
                       ymax
```

### Using R to access the extent

• Using ext() from the terra package


```
1 ext(nc.shp)
SpatExtent : -84.3238525390625, -75.4569778442383, 33.8819923400879,
36.5896492004395 (xmin, xmax, ymin, ymax)

1 ext(meuse.rst)
SpatExtent : 178400, 181600, 329400, 334000 (xmin, xmax, ymin, ymax)
```

### Mapping location: Resolution

- **Resolution:** the accuracy that the location and shape of a map's features can be depicted
- Minimum Mapping Unit: The minimum size and dimensions that can be reliably represented at a given *map scale*.
- Map scale vs. scale of analysis

### Mapping location: Resolution



### Using R to access resolution

- Thematically defined for vector datasets (check your metadata!!)
- Using res() in terra

```
1 f <- rast(system.file("ex/meuse.tif", package="terra"))
2 res(f)
[1] 40 40</pre>
```

### Recap

### Today's goals

- Articulate why we care about space
- Describe elements of spatial data
- Define a coordinate reference system and its importance
- Describe several ways to load spatial data into R
- Identify projections in R

