

MATH255: Honours Analysis 2

Matthew He

March 30, 2026

Abstract

1 Metric Spaces

1.1 Openness and closedness

1.2 Convergence, completeness and compactness

Definition 1 (Limit and convergence). Let (X, d) be a metric space, $x \in X$, and $(x_k)_{k \in \mathbb{N}}$ be a sequence of elements in X . We say that $(x_k)_k$ converges to x in (X, d) if and we denote $\lim_{k \rightarrow \infty} x_k = x$ or $x_k \rightarrow x$ as $k \rightarrow \infty$ if

$$\forall \varepsilon > 0, \exists k_\varepsilon \in \mathbb{N} \text{ such that } \forall k \geq k_\varepsilon, d(x_k, x) < \varepsilon \quad (1)$$

Moreover, x is called the limit of the sequence $(x_k)_k$ in (X, d) .

Proposition 1. Let (X, d) be a metric space, $x \in X$, and $A \subseteq X$. Then $x \in \overline{A}$ iff there exists a sequence $(x_k)_k$ in A such that $\lim_{k \rightarrow \infty} x_k = x$.

Theorem 1 (Sequential Criterion for Closedness). Let (X, d) be a metric space and $A \subseteq X$. Then the following statements are equivalent:

- A is closed in (X, d) .
- For every $x \in X$ and $(x_k)_{k \in \mathbb{N}}$ sequence of elements in A , if $\lim_{k \rightarrow \infty} x_k = x$ in (X, d) , then $x \in A$.

Definition 2 (Cauchy sequence). Let (X, d) be a metric space and $(x_k)_{k \in \mathbb{N}}$ be a sequence of elements in X . We say that $(x_k)_k$ is a Cauchy sequence in (X, d) if

$$\forall \varepsilon > 0, \exists k_\varepsilon \in \mathbb{N} \text{ such that } \forall m, n \geq k_\varepsilon, d(x_m, x_n) < \varepsilon \quad (2)$$

Proposition 2. Every converging sequence in (X, d) is a Cauchy sequence in (X, d) .

Definition 3 (Completeness, Banach and Hilbert space). We say that a metric space (X, d) is complete if every Cauchy sequence in (X, d) converges to some limit in (X, d) .

- A complete normed vector space is called a Banach space.
- A complete inner product space is called a Hilbert space.

Proposition 3. Let (X, d) be a metric space and $Y \subseteq X$,

1. If (Y, d) is complete, then Y is closed in (X, d) .
2. If (X, d) is complete and Y is closed in (X, d) , then (Y, d) is complete.

Definition 4 (Compactness). Let (X, d) be a metric space and $A \subseteq X$. We say that A is compact (or sequentially compact) in (X, d) if for every sequence $(x_k)_{k \in \mathbb{N}}$ of elements in A , there exists $x \in A$ and a subsequence $(x_{k_j})_{j \in \mathbb{N}}$ such that $\lim_{j \rightarrow \infty} x_{k_j} = x$ in (X, d) .

If this is true for $A = X$, then we say that (X, d) is compact (or sequentially compact).

Compactness forces convergence and existence of limits.

Proposition 4. Let (X, d) be a compact metric space and $A \subseteq X$. Then A is compact in (X, d) iff A is closed in (X, d) .

Theorem 2 (Heine-Borel Theorem). Let $X, \|\cdot\|$ be a normed vector space of finite dimension and $A \subseteq X$. Then, A is compact in $(X, \|\cdot\|)$ iff A is closed and bounded in $(X, \|\cdot\|)$.

Proposition 5 (Finite subcover of compact sets).

Let (X, d) be a metric space and $A \subseteq X$. If A is compact in (X, d) , then for every collection $\{U_i\}_{i \in I}$ of open subsets of (X, d) such that $A \subseteq \bigcup_{\alpha \in I} U_\alpha$, there exists a finite sub-collection $\{U_{i_j}\}_{1 \leq j \leq N}$ where $N \in \mathbb{N}$ and $i_j \in I$ for all $1 \leq j \leq N$ such that $A \subseteq \bigcup_{j=1}^N U_{i_j}$.

Corollary 1. Let (X, d) be a metric space and $A \subseteq X$. If A is compact in (X, d) , then for every $r > 0$, there exists a finite number $N_r \in \mathbb{N}$ of elements $x_1, \dots, x_{N_r} \in A$ such that $A \subseteq \bigcup_{j=1}^{N_r} B(x_j, r)$.

Corollary 2. Let $(X, \|\cdot\|)$ be a normed vector space of infinite dimension. Then, the closed unit ball $\{x \in X : \|x\| \leq 1\}$ is not compact in $(X, \|\cdot\|)$. (Hence the Heine-Borel theorem does not hold in infinite dimension.)

Proposition 6. Every normed vector space of finite dimension is Banach.

1.3 Mapping between metric spaces

Definition 5 (Cluster point).

Let (X, d) be a metric space, $A \subseteq X$, and $x_0 \in X$.

We say that x_0 is a cluster point of A in (X, d) if $x_0 \in \overline{A \setminus \{x_0\}}$.

Proposition 7.

x_0 is a cluster point of A iff there exists a sequence $(x_k)_{k \in \mathbb{N}}$ in A such that $\lim_{k \rightarrow \infty} x_k = x_0$ and $x_k \neq x_0$ for all $k \in \mathbb{N}$.

Definition 6 (Limit of function value). Let (X, d_X) and (Y, d_Y) be two metric spaces, $A \subseteq X$, x_0 be a cluster point of A and $y_0 \in Y$.

1. For $f : A \rightarrow Y$, we say that $f(x)$ converges to y_0 as $x \rightarrow x_0$ and we denote $\lim_{x \rightarrow x_0} f(x) = y_0$ if

$$\forall \varepsilon > 0, \exists \delta_\varepsilon > 0 : \forall x \in A, 0 < d_X(x, x_0) < \delta_\varepsilon \Rightarrow d_Y(f(x), y_0) < \varepsilon \quad (3)$$

2. When $(Y, d_Y) = (\mathbb{R}, |\cdot|)$, for $f : A \rightarrow \mathbb{R}$, we say that $f(x)$ tends to $\pm\infty$ as $x \rightarrow x_0$ and we denote $\lim_{x \rightarrow x_0} f(x) = \pm\infty$ if

$$\forall M > 0, \exists \delta_M > 0 : \forall x \in A, 0 < d_X(x, x_0) < \delta_M \Rightarrow f(x) > M \text{ or } f(x) < -M \quad (4)$$

1.4 Continuous functions on compact sets

2 Infinite Series

Definition 7 (Partial sum). Let $(X, \|\cdot\|)$ be a normed vector space over $\mathbb{R}$ or $\mathbb{C}$. $k_0 \in \mathbb{Z}$ and $(a_k)_{k \geq k_0}$ by a sequence in X . For every $n \geq k_0$, $S_n = \sum_{k=k_0}^n a_k$ is called the partial sum of order n of $(a_k)_k$.

If $\lim_{n \rightarrow \infty} S_n = S$ in $(X, \|\cdot\|)$, for some $S \in X$, then we write

$$\sum_{k=k_0}^{\infty} a_k = S \quad (5)$$

and we say that the series $\sum_{k=k_0}^{\infty} a_k$ and that S is its sum. If $(S_n)_{n \geq k_0}$ does not converge, then we say that the series $\sum_{k=k_0}^{\infty} a_k$ diverges.

Proposition 8. $\forall k'_0 > k_0$, $\sum_{k=k_0}^{\infty} a_k$ converges iff $\sum_{k=k'_0}^{\infty} a_k$ converges. Therefore, we will sometimes assume without loss of generality that $k_0 = 0$ or 1 .

Proof: $\forall n \geq k'_0$, $\sum_{k=k_0}^n a_k = \sum_{k=k_0}^{k'_0-1} a_k + \sum_{k=k'_0}^n a_k$ where the second term is independent of n . Hence, $\sum_{k=k_0}^{\infty} a_k$ converges iff $\sum_{k=k'_0}^{\infty} a_k$ converges. $\square$

Proposition 9. If $\sum_{k=k_0}^{\infty} a_k$ converges, then $a_k \rightarrow 0$ as $k \rightarrow \infty$.

Proof: If $\sum_{k=k_0}^{\infty} a_k$ converges, then $a_n = S_n - S_{n-1} \rightarrow S - S = 0$ as $n \rightarrow \infty$. $\square$

Proposition 10. If $\sum_{k=k_0}^{\infty} a_k$ and $\sum_{k=k_0}^{\infty} b_k$ converge, then $\forall \lambda, \mu \in \mathbb{R}$ or $\mathbb{C}$, $\sum_{k=k_0}^{\infty} (\lambda a_k + \mu b_k) \rightarrow \lambda \sum_{k=k_0}^{\infty} a_k + \mu \sum_{k=k_0}^{\infty} b_k$.

Proof follows from $\sum_{k=k_0}^n (\lambda a_k + \mu b_k) = \lambda \sum_{k=k_0}^n a_k + \mu \sum_{k=k_0}^n b_k$. $\square$

Remark 1 (Examples).

- **Geometric series** $a_k = r^k$ for some $r \in \mathbb{R}$,

$$S_n = \sum_{k=0}^n r^k = \begin{cases} \frac{1-r^{n+1}}{1-r} & \text{if } r \neq 1 \\ n+1 & \text{if } r = 1 \end{cases} \quad (6)$$

It converges iff $|r| < 1$ with $S = \frac{1}{1-r}$ since $\lim_{n \rightarrow \infty} r^{n+1} = 0$ when $|r| < 1$.

- **Series of matrices** For example, $\sum_{k=0}^{\infty} A^k = (I - A)^{-1}$ for all $n \times n$ real value matrices such that $\|A\|_{2,2} = \max_{\|x\|_2=1} \|Ax\|_2 < 1$.
- **p-series** $\sum_{k=1}^{\infty} \frac{1}{k^\alpha}$ converges iff $\alpha > 1$.

Theorem 3 (Cauchy criterion for convergence). *If $(X, \|\cdot\|)$ is Banach, then $\sum_{k=k_0}^{\infty} a_k$ converges iff $\forall \varepsilon > 0, \exists n_\varepsilon \in \mathbb{N}$ such that*

$$\forall m > n \geq n_\varepsilon, \left\| \sum_{k=n+1}^m a_k \right\| < \varepsilon \quad (7)$$

Proof: If $(X, \|\cdot\|)$ is Banach, then $(S_n)_{n \geq k_0}$, $S_n = \sum_{k=k_0}^n a_k$ converges iff it is Cauchy, i.e. $\forall \varepsilon > 0, \exists n_\varepsilon \in \mathbb{N}$ such that

$$\forall m > n \geq n_\varepsilon, \|S_m - S_n\| = \left\| \sum_{k=n+1}^m a_k \right\| < \varepsilon \quad (8)$$

□

Definition 8 (Absolute and conditional convergence). $\sum_{k=k_0}^{\infty} a_k$ is said to be absolutely convergent if $\sum_{k=k_0}^{\infty} \|a_k\|$ converges in $(\mathbb{R}, \|\cdot\|)$. It is said to be conditionally convergent if $\sum_{k=k_0}^{\infty} a_k$ converges but $\sum_{k=k_0}^{\infty} \|a_k\|$ diverges.

Proposition 11. *If $(X, \|\cdot\|)$ is Banach, then for every series $\sum_{k=k_0}^{\infty} a_k$ in $(X, \|\cdot\|)$, if $\sum_{k=k_0}^{\infty} \|a_k\|$ converges absolutely, then $\sum_{k=k_0}^{\infty} a_k$ converges and*

$$\left\| \sum_{k=n}^{\infty} a_k \right\| \leq \sum_{k=n}^{\infty} \|a_k\|, \quad \forall n \geq k_0 \quad (9)$$

Proposition 12. *Let $\sum_{k=k_0}^{\infty} a_k$ be a series in a Banach space $(X, \|\cdot\|)$. If $\sum_{k=k_0}^{\infty} a_k$ converges absolutely, then for every bijective function $f : \{k \geq k_0\} \rightarrow \{k \geq k_0\}$ (permutation), the series $\sum_{k=k_0}^{\infty} a_{f(k)}$ converges and*

$$\sum_{k=k_0}^{\infty} a_{f(k)} = \sum_{k=k_0}^{\infty} a_k \quad (10)$$

We call the series $\sum_{k=k_0}^{\infty} a_{f(k)}$ a rearrangement of $\sum_{k=k_0}^{\infty} a_k$.

Remark 2.

$\tilde{S}_n = \sum_{k=k_0}^n \|a_k\|$ is nondecreasing: $\tilde{S}_{n+1} \geq \tilde{S}_n$ for all $n \in \mathbb{N}$.

Hence, $\tilde{S}_n = \sum_{k=k_0}^n \|a_k\|$ converges iff $(\tilde{S}_n)_{n \geq k_0}$ is bounded. Moreover, $\sum_{k=k_0}^{\infty} \|a_k\| = \sup_{n \geq k_0} \tilde{S}_n$.

Remark 3.

The converse is true, if $\forall \sum a_k$ that converges absolutely, $\sum a_k$ converges, then $(X, \|\cdot\|)$ is Banach.

2.1 Series with nonnegative term

Theorem 4 (Direct Comparison Test). *Let $\sum_{k=k_0}^{\infty} a_k$ and $\sum_{k=k_0}^{\infty} b_k$ be two series in $(\mathbb{R}, \|\cdot\|)$ such that $0 \leq a_k \leq b_k$ for all $k \geq k_0$. Then*

1. *If $\sum_{k=k_0}^{\infty} b_k$ converges, then $\sum_{k=k_0}^{\infty} a_k$ converges and $\sum_{k=k_0}^{\infty} a_k \leq \sum_{k=k_0}^{\infty} b_k$.*
2. *If $\sum_{k=k_0}^{\infty} a_k$ diverges, then $\sum_{k=k_0}^{\infty} b_k$ diverges.*

Corollary 3 (Limit Comparison Test). *Let $\sum_{k=k_0}^{\infty} a_k$ and $\sum_{k=k_0}^{\infty} b_k$ be two series in $(\mathbb{R}, \|\cdot\|)$ such that $a_k > 0$ and $b_k > 0$ for all $k \geq k_0$, and $\lim_{k \rightarrow \infty} \frac{a_k}{b_k} = \ell \in (0, \infty)$. Then $\sum_{k=k_0}^{\infty} a_k$ converges if and only if $\sum_{k=k_0}^{\infty} b_k$ converges.*

Theorem 5 (Root Test). Let $\sum_{k=k_0}^{\infty} a_k$ be a series in $(\mathbb{R}, \|\cdot\|)$ such that $a_k \geq 0$ for all $k \geq k_0$.

1. If $\limsup_{k \rightarrow \infty} \sqrt[k]{a_k} < 1$, then $\sum_{k=k_0}^{\infty} a_k$ converges.

2. If $\limsup_{k \rightarrow \infty} \sqrt[k]{a_k} > 1$, then $\sum_{k=k_0}^{\infty} a_k$ diverges.

If $\limsup_{k \rightarrow \infty} \sqrt[k]{a_k} = 1$, then the test is inconclusive.

Proposition 13 (Ratio Test). Let $\sum_{k=k_0}^{\infty} a_k$ be a series in $(\mathbb{R}, \|\cdot\|)$ such that $a_k > 0$ for all $k \geq k_0$.

1. If $\limsup_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} < 1$, then $\sum_{k=k_0}^{\infty} a_k$ converges.

2. If $\liminf_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} > 1$, then $\sum_{k=k_0}^{\infty} a_k$ diverges.

Proof: (1) Assume that $R = \limsup_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} < 1$. Then $\forall \varepsilon \in (0, 1 - R)$, $\exists n_\varepsilon \in \mathbb{N}$ such that $\forall k \geq n_\varepsilon, \sup_{k \geq n} \frac{a_{k+1}}{a_k} < R + \varepsilon < 1$, which gives for all $k \geq n_\varepsilon, a_{k+1} < (R + \varepsilon)a_k < (R + \varepsilon)^{k-n_\varepsilon+1}a_{n_\varepsilon}$. When $\varepsilon \in (0, 1 - R)$, $0 < R + \varepsilon < 1$, so $\sum_{k=n_\varepsilon}^{\infty} (R + \varepsilon)^{k-n_\varepsilon+1}a_{n_\varepsilon}$ converges, which by direct comparison implies that $\sum_{k=k_0}^{\infty} a_k$ converges.

(2) Assume that $R = \liminf_{k \rightarrow \infty} \frac{a_{k+1}}{a_k} > 1$. Then $\exists k_1 \geq k_0$ such that $\forall k \geq k_1, \inf_{k \geq n} \frac{a_{k+1}}{a_k} > R - \varepsilon > 1$, i.e. $a_{k+1} > a_k$, which implies $a_k \not\rightarrow 0$ as $k \rightarrow \infty$, thus $\sum_{k=k_0}^{\infty} a_k$ diverges. $\square$

2.2 Power series

Now we focus on series in $(\mathbb{R}, |\cdot|)$ with terms of different sign.

Theorem 6 (Alternating Series Test). Let $\sum_{k=k_0}^{\infty} a_k$ be a series in $(\mathbb{R}, |\cdot|)$ such that $0 \leq a_{k+1} \leq a_k$ for all $k \geq k_0$ and $\lim_{k \rightarrow \infty} a_k = 0$. Then $\sum_{k=k_0}^{\infty} a_k$ converges and

$$0 \leq \sum_{k=k_0}^{\infty} (-1)^{k-k_0} a_k \leq a_{k_0} \quad (11)$$

Example 1

$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k}$ converges since $0 \leq \frac{1}{k+1} \leq \frac{1}{k}$ for all $k \geq 1$ and $\lim_{k \rightarrow \infty} \frac{1}{k} = 0$.

Proof: By observing that

$$\sum_{k=k_0}^{\infty} (-1)^k a_k = \sum_{k=0}^{\infty} (-1)^{k+k_0} a_{k+k_0} \quad (12)$$

$$= (-1)^{k_0} \sum_{k=0}^{\infty} (-1)^k a_{k+k_0} \quad (13)$$

we can assume that $k_0 = 0$.

For every $n \in \mathbb{N}$,

$$S_{2n} = a_0 - a_1 + a_2 - \cdots + a_{2n-2} - a_{2n-1} + a_{2n} \quad (14)$$

$$= (a_0 - a_1) + (a_2 - a_3) + \cdots + (a_{2n-2} - a_{2n-1}) + a_{2n} \quad (15)$$

$$\geq 0 \quad (16)$$

$$S_{2n+1} = a_0 - a_1 + a_2 - \cdots + a_{2n-2} - a_{2n-1} + a_{2n} - a_{2n+1} \quad (17)$$

$$= (a_0 - a_1) + (a_2 - a_3) + \cdots + (a_{2n-2} - a_{2n-1}) + (a_{2n} - a_{2n+1}) \quad (18)$$

$$\leq 0 \quad (19)$$

Definition 9 (Cauchy product). Let $\sum_{k=0}^{\infty} a_k$ and $\sum_{k=0}^{\infty} b_k$ be series in $(\mathbb{R}, |\cdot|)$ or $(\mathbb{C}, |\cdot|)$.

For each $n \in \mathbb{N}$, define

$$c_n = \sum_{k=0}^n a_k b_{n-k} = a_0 b_n + a_1 b_{n-1} + \cdots + a_n b_0 \quad (20)$$

Then the series $\sum_{n=0}^{\infty} c_n$ is called the Cauchy product of $\sum_{k=0}^{\infty} a_k$ and $\sum_{k=0}^{\infty} b_k$.

Proposition 14. If $\sum_{k=0}^{\infty} a_k$ and $\sum_{k=0}^{\infty} b_k$ are series in $(\mathbb{R}, |\cdot|)$ or $(\mathbb{C}, |\cdot|)$ such that $\sum_{k=0}^{\infty} a_k$ converges and $\sum_{k=0}^{\infty} b_k$ converges (not necessarily absolutely). Then the Cauchy product $\sum_{n=0}^{\infty} c_n$ converges and $\sum_{n=0}^{\infty} c_n = (\sum_{k=0}^{\infty} a_k)(\sum_{k=0}^{\infty} b_k)$.

Example 2 (Counter example of Cauchy product of two conditionally converging series)

Consider $a_k = b_k = \frac{(-1)^k}{\sqrt{k+1}}$. Then $\sum_{k=0}^{\infty} a_k$ and $\sum_{k=0}^{\infty} b_k$ converge by the alternating series test and p -series convergence.

In this case:

$$|c_n| = \left| \sum_{k=0}^n a_k b_{n-k} \right| \quad (21)$$

$$(22)$$

Definition 10 (Power series). Let $(a_k)_k$ be a sequence in $\mathbb{R}$ or $\mathbb{C}$ and $x_0 \in \mathbb{R}$ or $\mathbb{C}$. The function $x \mapsto \sum_{k=0}^{\infty} a_k (x - x_0)^k$ is defined at points where it converges, is called a power series.

Proposition 15 (Radius of convergence). Let $\sum_{k=0}^{\infty} a_k(x - x_0)^k$ be a power series in $\mathbb{R}$ or $\mathbb{C}$. Define $R = \frac{1}{\limsup_{k \rightarrow \infty} \sqrt[k]{|a_k|}}$ with the convention that $\frac{1}{0} = \infty$ and $\frac{1}{\infty} = 0$. Then R is called the radius of convergence of the power series, and

$$\text{if } |x - x_0| < R, \text{ then } \sum_{k=0}^{\infty} a_k(x - x_0)^k \text{ converges,} \quad (23)$$

$$\text{if } |x - x_0| > R, \text{ then } \sum_{k=0}^{\infty} a_k(x - x_0)^k \text{ diverges.} \quad (24)$$

Proposition 16. Let $f(x) = \sum_{k=0}^{\infty} a_k(x - x_0)^k$ be a power series in $(\mathbb{R}, |\cdot|)$ with the radius of convergence $R > 0$. Then f is infinitely differentiable in $(x_0 - R, x_0 + R)$

Example 3

For every $x \in (0, 2)$, we can write $\ln x = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} (x - 1)^k$.

Indeed, let $\tilde{\ln} x = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} (x - 1)^k$. We want to show that $\tilde{\ln} x = \ln x$, i.e., $\tilde{\ln} e^y = y$ where $y = \ln x$. Observe that

$$\tilde{\ln}'(x) = \sum_{k=1}^{\infty} (-1)^{k-1} (x - 1)^{k-1} = \sum_{k=1}^{\infty} (1 - x)^{k-1} \quad (25)$$

$$= \sum_{k=0}^{\infty} (1 - x)^k \quad (26)$$

$$= \frac{1}{1 - (1 - x)} = \frac{1}{x} \quad (27)$$

More, this is true for all $x \in (0, 2)$ since the radius of convergence of $\sum_{k=0}^{\infty} (1 - x)^k$ is 1. It follows that

$$\frac{d}{dy} [\tilde{\ln} e^y - y] = e^y \tilde{\ln}' e^y - 1 = e^y \frac{1}{e^y} - 1 = 0 \quad (28)$$

hence $\tilde{\ln} e^y - y = \tilde{\ln} e^0 - 0 = 0$, i.e. $\tilde{\ln} x = \ln x$. $\square$

Definition 11 (Taylor series). Let I be an interval in $\mathbb{R}$, $x_0 \in I$ and $f : I \rightarrow \mathbb{R}$ be infinitely differentiable at x_0 . Then the series

$$\sum_{k=0}^{\infty} \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k \quad (29)$$

is called the Taylor series of f at x_0 .

We've proven that every power series $f(x) = \sum_{k=0}^{\infty} a_k(x - x_0)^k$ is equal to its Taylor series at $x_0 \in (x_0 - R, x_0 + R)$.

However, this is not true in general where f is only assumed to be infinitely differentiable at x_0 . For example,

Example 4

The function $x \mapsto e^{-\frac{1}{x^2}}$ for $x \neq 0$ and 0 for $x = 0$ is infinitely differentiable in $\mathbb{R}$ and such that $f^{(k)}(0) = 0$ for all $k \in \mathbb{N}$. That means that its Taylor series at 0 is the zero series.

Thus the function itself is not equal to its Taylor series at 0.

Theorem 7 (Taylor's theorem with mean value reminder).

Let $x, x_0 \in \mathbb{R}, x \neq x_0, I = \begin{cases} [x_0, x] & \text{if } x > x_0 \\ [x, x_0] & \text{if } x < x_0 \end{cases}, \text{int}(I) = \begin{cases} (x_0, x) & \text{if } x > x_0 \\ (x, x_0) & \text{if } x < x_0 \end{cases}$ and $f : I \rightarrow \mathbb{R}$ be n times continuously differentiable in I (i.e. $f^{(n)}$ exists and is continuous in I). and $n + 1$ times differentiable in $\text{int}(I)$.

Then, $\exists x' \in \text{int}(I)$ such that

$$f(x) = \underbrace{\sum_{k=0}^n \frac{f^{(k)}(x_0)}{k!} (x - x_0)^k}_{\text{partial sum of Taylor series at } x_0} + \underbrace{\frac{f^{(n+1)}(x')}{(n+1)!} (x - x_0)^{n+1}}_{\text{mean value reminder}} \quad (30)$$

Define $g(t) = f(x) - \sum_{k=0}^n \frac{f^{(k)}(t)}{k!} (t - t)^k$. We want to show that $g(x_0) = f(x) - \frac{1}{(n+1)!} f^{(n+1)}(x') (x - x_0)^{n+1}$ for some $x' \in \text{int}(I)$.

In particular, $g(x) = f(x) - f(x) = 0$.

g is continuous on I and differentiable on $\text{int}(I)$ (by assumption on f) and

$$g'(t) = \sum_{k=0}^n \frac{f^{(k+1)}(t)}{k!} (t - t)^k \quad (31)$$

$$= \frac{f^{(n+1)}(t)}{n!} (t - t)^n \quad (32)$$

$$= \frac{f^{(n+1)}(t)}{(n+1)!} (t - x_0)^{n+1} - \frac{f^{(n+1)}(t)}{(n+1)!} (t - x)^{n+1} \quad (33)$$

Now define

$$h(t) = g(t)(x - x_0)^{n+1} - g(x_0)(x - t)^{n+1} \quad (34)$$

so that

$$h(x) = 0 \quad (35)$$

$$h(x_0) = 0 \quad (36)$$

Since h is continuous on I and differentiable on $\text{int}(I)$ (by assumption on g), and

$$h'(t) = g'(t)(x - x_0)^{n+1} + (n+1)g(x_0)(x - t)^n \quad (37)$$

$$(38)$$

The point here is to construct a function h such that it vanishes at both x and x_0 so that we can apply Rolle's theorem to it.

By Rolle's Theorem, it follows that

$$\exists x' \in \text{int}(I) \text{ such that } h'(x') = 0 \quad (39)$$

3 Riemann-Stieltjes integral

Proposition 17. $f \in \mathcal{R}([a, b], \alpha)$ iff $\forall \varepsilon > 0, \exists P_\varepsilon$ (partition of $[a, b]$) such that $U(P_\varepsilon, f, \alpha) - L(P_\varepsilon, f, \alpha) < \varepsilon$.

Proof: ($\Rightarrow$) Assume that $f \in \mathcal{R}([a, b], \alpha)$. Then by properties of *inf* and *sup*, $\forall \varepsilon > 0, \exists P_\varepsilon, P'_\varepsilon$ partition of $[a, b]$ such that

$$\int_a^b f(x) d\alpha - \varepsilon < L(P_\varepsilon, f, \alpha) = \int_a^b f(x) d\alpha \leq \overline{\int_a^b f(x) d\alpha} \leq U(P_\varepsilon, f, \alpha) < \overline{\int_a^b f(x) d\alpha} + \varepsilon \quad (40)$$

Let $P_\varepsilon^* = P_\varepsilon \cup P'_\varepsilon$. Then since $P_\varepsilon^* \supset P_\varepsilon$ and $P_\varepsilon^* \supset P'_\varepsilon$,

$$U(P_\varepsilon^*, f, \alpha) - L(P_\varepsilon^*, f, \alpha) \leq U(P'_\varepsilon, f, \alpha) - L(P_\varepsilon, f, \alpha) \quad (41)$$

$$< (\overline{\int_a^b f(x) d\alpha} + \frac{1}{2}\varepsilon) - (\int_a^b f(x) d\alpha - \frac{1}{2}\varepsilon) \quad (42)$$

$$= \overline{\int_a^b f(x) d\alpha} - \int_a^b f(x) d\alpha + \varepsilon \quad (43)$$

$$= \varepsilon \quad (44)$$

where $\overline{\int_a^b f(x) d\alpha} - \int_a^b f(x) d\alpha = 0$ since $f \in \mathcal{R}([a, b], \alpha)$.

($\Leftarrow$) Assume that $\forall \varepsilon > 0, \exists P_\varepsilon$ partition of $[a, b]$ such that $U(P_\varepsilon, f, \alpha) - L(P_\varepsilon, f, \alpha) < \varepsilon$. We have

$$L(P_\varepsilon, f, \alpha) \leq \int_a^b f(x) d\alpha \leq \overline{\int_a^b f(x) d\alpha} \leq U(P_\varepsilon, f, \alpha) \quad (45)$$

It follows that

$$0 \leq \overline{\int_a^b f(x) d\alpha} - \int_a^b f(x) d\alpha \leq U(P_\varepsilon, f, \alpha) - L(P_\varepsilon, f, \alpha) \leq \varepsilon \quad (46)$$

Letting $\varepsilon \rightarrow 0$, we have

$$\overline{\int_a^b f(x) d\alpha} - \int_a^b f(x) d\alpha = 0 \quad (47)$$

therefore $f \in \mathcal{R}([a, b], \alpha)$. □

Look at Question 2 in Assignment 6.

Proposition 18. Let $f, \alpha: [a, b] \rightarrow \mathbb{R}$ be such that f is bounded in $[a, b]$ and α increasing on $[a, b]$.

Let D be the set of points in $[a, b]$ where f is not continuous. Assume that D is countable and that α is continuous at every point in D . Then $f \in \mathcal{R}([a, b], \alpha)$.

Let $\varepsilon > 0$. For every $x \in [a, b] \setminus D$, since f is continuous at x , there exists $\delta_x > 0$ such that $|f(y) - f(x)| < c\varepsilon$ for all $y \in (x - \delta_x, x + \delta_x) \cap [a, b]$, where the constant $c > 0$ will be determined later.

Since D is countable, $\exists (x_k)_k$ in $[a, b]$ such that $D = \{x_k : k \in \mathbb{N}\}$. Since α is continuous at x_k , there exists $\delta_k > 0$ such that $|\alpha(y) - \alpha(x_k)| < c_k \varepsilon$ for all $y \in (x_k - \delta_k, x_k + \delta_k) \cap [a, b]$, where the constant $c_k > 0$ will be determined later.

Since $[a, b]$ is compact, $[a, b] \subset \bigcup_{x \in [a, b]} (x - \delta_x, x + \delta_x)$, there exists a finite subcover, i.e. $\exists y_1 < \dots < y_{N_\varepsilon} \in [a, b]$ such that

$$[a, b] \subset \bigcup_{i=1}^{N_\varepsilon} (y_i - \delta_{y_i}, y_i + \delta_{y_i}) \quad (48)$$

δ_x is defined for every $x \in [a, b]$, since either it's defined by the continuity of f at x (if $x \in [a, b] \setminus D$) or by the continuity of α at x (if $x \in D$).

By ordering the points $a, b, y_i - \delta_{y_i}$ (if ≥ 0), and $y_i + \delta_{y_i}$ (if $\leq b$), we then obtain a partition $P_\varepsilon = \{x_0, \dots, x_{m_\varepsilon}\}$ such that $\forall j \in \{0, \dots, m_\varepsilon\}$, $\exists i \in \{1, \dots, N_\varepsilon\}$ such that $(x_{j-1}, x_j) \subset (y_i - \delta_{y_i}, y_i + \delta_{y_i}) \cap [a, b]$.

We again separate this into two cases (depending on whether $y_i \in D$ or not):

- If $y_i \notin D$, then

$$M_j - m_j = \sup_{x \in [x_{j-1}, x_j]} f(x) - \inf_{x \in [x_{j-1}, x_j]} f(x) \quad (49)$$

$$\leq \left(\sup_{x \in [x_{j-1}, x_j]} f(x) - f(y_i) \right) + \left(f(y_i) - \inf_{x \in [x_{j-1}, x_j]} f(x) \right) \quad (50)$$

$$< 2c\varepsilon \quad (51)$$

- If $y_i \in D$, then we have

$$M_j - m_j \leq M - m = \sup_{x \in [x_{j-1}, x_j]} f(x) - \inf_{x \in [x_{j-1}, x_j]} f(x) \quad (52)$$

$$\Delta\alpha_j = \alpha(x_j) - \alpha(x_{j-1}) \leq |\alpha(x_j) - \alpha(y_i)| + |\alpha(y_i) - \alpha(x_{j-1})| \quad (53)$$

$$< 2c_k \varepsilon \quad (54)$$

Therefore,

$$U(P_\varepsilon, f, \alpha) - L(P_\varepsilon, f, \alpha) = \sum_{j=1}^{m_\varepsilon} (M_j - m_j) \Delta\alpha_j \quad (55)$$

$$= \sum_{j=1, y_{i_j} \in D}^{m_\varepsilon} (M_j - m_j) \Delta\alpha_j + \sum_{j=1, y_{i_j} \notin D}^{m_\varepsilon} (M_j - m_j) \Delta\alpha_j \quad (56)$$

$$\leq 2c\varepsilon \underbrace{\sum_{j=1, y_{i_j} \notin D}^{m_\varepsilon} \Delta\alpha_j}_{ss} + (M - m) \sum_{j=1, y_{i_j} \in D}^{m_\varepsilon} \Delta\alpha_j \quad (57)$$

$$(58)$$

 References