
WHY COVID MORTALITY FOLLOWS GOMPERTZ, NOT LOGISTIC GROWTH

A PREPRINT

August 25, 2026

ABSTRACT

A number of studies have shown that cumulative mortality followed a Gompertz curve in the initial Covid pandemic period in March-April 2020 — a pattern incompatible with the initial logistic growth predicted by traditional Susceptible-Infected-Recovered (SIR) models. We ask what the mortality data can actually determine about the two competing descriptions, and find that the SIR family’s reputation for simplicity does not survive the question. Fitted to deaths alone, the SIRD model carries one parameter more than the observations support: its scaled Jacobian is conditioned an order of magnitude worse than that of the Gompertz model (18–594 against 3.8–4.4 across sixteen countries), and the ill-conditioned direction is precisely the one carrying the reproduction number $\mathcal{R}_0$. The consequences are measurable. Changing how an even-length moving average is centred — a half-day shift with no modelling content — moves the fitted $\mathcal{R}_0$ by a median of 15% while moving the Gompertz rate by 1.4%; and two careful fits of the same equation to the same source data differ by a quarter in $\mathcal{R}_0$ while agreeing in the median to about one per cent on the combination the data actually determine. We do not conclude that the SIRD model is false: over a single wave it fits the observed relationship between deaths and their rate marginally better than the Gompertz model does, and we argue that a formal test between the two is a weak instrument at these sample sizes. The conclusion is narrower — that the quantity these models are most often fitted in order to report is not identifiable from aggregate mortality, and that the precision with which it is routinely quoted is an artefact of an error model which ignores the smoothing applied to the data. When the fitted equations are instead integrated forward, the Gompertz model tracks the observed trajectory more closely in eleven of the sixteen countries. We connect this to a mechanistic account in which Gompertz growth is the unique member of the Richards family closed under aggregation, so that the transmission network contributes nothing to the collective curve — a population from which no transmission parameter could be recovered by curve-fitting in the first place.

Keywords Gompertz · Coherence · Covid · Coronavirus · Network Analysis · Stochastic

Introduction

Traditional communicable disease spreading theory assumes a pathogen which infects the population through a network of transmission. Following this line of reasoning it can be theoretically shown that in the early stages of an epidemic, growth follows a logistic-like curve, where the very beginning exhibits exponential growth, and for which analytic and semi-analytic solutions have recently emerged [1–4]. A large body of research reveals how these models fit the mortality seen due to the Covid pandemic in 2020, a pandemic due to an influenza-like disease [5–10].

However, instead of showing logistic-like growth, observed cumulative mortality in the initial period March-April 2020 exhibits almost perfect resemblance to Gompertzian growth [11, 12] where the log-transformed cumulative mortality, or log-mortality for short, is exponentially *decreasing* in time,

$$\frac{d}{dt} \ln F(t) = -\beta \ln F(t), \tag{1}$$

using normalized mortality such that $F(t \rightarrow \infty) = 1$.

This phenomenon is recorded by others [13–17], showing Gompertz curves at national levels instead of the predicted logistic curves, an observation made possible in part due to the increased cadence and quality of data acquisition compared with earlier epidemics. What is causing such a discrepancy between these observations and the current theory of communicable diseases [18]?

In addition to the characteristic patterns of these national mortality curves, other observations that conflict more generally with the communicable disease models are 1) the lack of correlation between population density and mortality (or infection) rates [19–25], 2) no clear association between inter-generational relationships and fatality rates [26], and 3) the widespread morbidity and mortality spikes seen during spring 2020 among birds [27], dogs [28], rabbits [29–31], elephants [32], and horses [33–36]. Many of these animal mortality spikes are not attributable to a Coronavirus infection, so why did these spikes all occur within the same time frame? While these observations are not reconcilable within the current communicable disease paradigm, we present here an alternative theory that can reconcile all observations; a theory based on a global biosphere disturbance with fewer parameters than the most basic communicable disease model.

This global biosphere disturbance could be in the form of sudden ionospheric disturbances (SIDs) [37], which in the context of short-term solar and geomagnetic disturbances have been associated with physiological changes in cardiac system processes [38–45], melatonin/serotonin balance [46–48], endothelial activation and inflammation [49], and disturbed magnetoreception in migrating animals [50–56]. More general effects have also been studied in this context, e.g. overall mortality, suicide, mental disorders, homicides, and epileptic seizures [57–65]. Cherry (2002) put forward a mechanism whereby the ionosphere acts as a coupling agent between geomagnetic disturbances and physiological effects based on the disturbance of the background electromagnetic oscillations between the surface of the Earth and the lower ionosphere, called the Schumann Resonance signal [66, 67]. The Schumann Resonance would then couple with the brain to alter the melatonin/serotonin balance, which was a causal link observed by Wever (1974) through the circadian rhythm [68]. Indeed, more recent findings support this hypothesized synchronization between Schumann Resonances and the brain [69–72].

In support of Cherry’s theory, specific phenomena such as solar X-ray bursts [73–77], solar proton events [74, 78, 79] and geomagnetic storms [80, 81] can modify the shape of the ionosphere cavity and shift the Schumann resonance frequencies [82, 76, 83]. Seismic activity has also been connected to variations in Schumann resonances [84–86], and conversely, ionospheric modifications have been connected to global seismicity [87]. Zaporozhan and Ponomarenko (2010) postulated that minute disturbances in magnetic fields which could translate into influenza symptoms [88], mediated by the cryptochrome protein-family which has been shown to be geomagnetic receptors [89–91].

Whatever the cause, global biosphere disturbance theory represents an alternative paradigm compared to the traditional contact-transmissible contagion theory in the context of disease spreading dynamics. Our goal with this study is to present a novel mathematical bridge between these two paradigms which at its core is a connection between the Gompertzian growth model and the logistic growth model. Our proposed bridge represents a microscopic version to the well-known parametrization given by Richard (1959) [92], where he augmented standard logistical growth by one parameter. This is why it is also sometimes called θ -logistic growth [93].

One issue with Richard’s parametrization is that the traditional communicable disease models cannot be realistically extrapolated to the Gompertz model, as shown by Wang *et al.* (2012) [94]. The reason why Richard’s model cannot realistically represent a communicable disease with a Gompertz model is because it would imply that there is “long-range interaction” between the susceptible and infected (see [93] for details), which is counter to the basic assumption that communicable diseases spread through proximity between the susceptible and infected. Based on these findings, one would then be forced to conclude that Gompertz growth cannot arise in the context of spread through proximity between the susceptible and infected.

In this paper we will show this incompatibility in another way by creating a microscopic model where each entity is exposed to an environmental stressor simultaneously applied to the population. This theory does not preclude the possibility of multiple sources of disturbance, potentially local and interacting with a global source. The main idea is that over some possibly geographically constrained area, everyone is exposed to the same source of disturbance simultaneously.

The organization of the paper is as follows: We first review the basic communicable disease models, before presenting a new stochastic model from which Gompertz growth emerges and thus explains the observed mortality data. We then proceed to juxtapose of the logistic model and the Gompertz model in the microscopic domain where multiple entities comprise the macroscopic behavior. The final section is dedicated to unifying the communicable disease paradigm with a biosphere disturbance through a new microscopic form of the Logistic-Gompertzian growth equations.

Simple communicable disease models: The SIR model family

We start by reviewing the basic infectious disease model, namely the Susceptible-Infected-Recovered (SIR) model under a time-dependent infection/recovery rate ratio, $\phi(t)$ [95], and show under what conditions the Gompertz curve could emerge. First assume a pool of susceptible people of size N evolving between the three states: susceptible, $S(t)$, recovered, $R(t)$ and infected, $I(t)$,

$$\begin{aligned}\frac{dS}{dt} &= -\beta \frac{IS}{N} \\ \frac{dI}{dt} &= \beta \frac{IS}{N} - \alpha I. \\ \frac{dR}{dt} &= \alpha I,\end{aligned}\tag{2}$$

omitting the argument t in each variable for brevity and where α and β in this context signify recovery and infection rates.

A line of reasoning employed by Rypdal and Rypdal (2020) [14] to obtain the Gompertz curve is to linearize the SIR model by assuming both the number of infected, $I(t)$, and cumulative infected, $Y(t)$, are much less than the initial pool of susceptible people, $N \gg Y \geq I$, which seems well founded at the national level, viz.

$$\frac{dY}{dt} = \beta I \tag{3}$$

$$\frac{dI}{dt} = (\beta - \alpha)I = \alpha(\phi(t) - 1)I, \tag{4}$$

and where the number of recovered, $R(t)$, is under this linearization decoupled from the other variables. They further assume the number of deceased is proportional to the number of cumulative infected, offset by a time lag, which allows us to use the same set of linearized equations to model the number of cumulative people deceased without loss of generality. And since I is a function of Y , we can combine Eq. 3 via an instantaneous relative growth rate, $\gamma(t) = dY(t)/(Y dt) = \beta I/Y$,

$$\frac{dY}{dt} = \gamma(t)Y(t). \tag{5}$$

Due to this linearization, the infection/recovery rate ratio, $\phi(t)$, and the relative growth rate, $\gamma(t)$, will have to change as a function of time to accommodate for the boundary conditions. The latter is typically also parameterized by a scaling factor, θ , representing the shape of the growth,

$$\gamma(t) = \frac{\beta}{\theta} \left[1 - \left(\frac{Y}{Y_\infty} \right)^\theta \right]. \tag{6}$$

This parametrization is the commonly used Richards growth curve [92], also called θ -logistic growth, and has been used by others [96]. Although not immediately justified in the communicable disease theory, one could imagine that θ represents non-linear network behavior [93]. Note that at $\theta = 1$, the traditional logistic growth curve is obtained, while at $\theta \rightarrow \infty$ we recover the exponential (Malthusian) explosion.

The observed Gompertzian mortality curves are realized in the limit $\theta \rightarrow 0$, with the relative growth rate,

$$\gamma(t) = \lim_{\theta \rightarrow 0} \frac{\beta}{\theta} \left[1 - \left(\frac{Y}{Y_\infty} \right)^\theta \right] = \beta \ln \frac{Y_\infty}{Y}, \tag{7}$$

such that

$$\frac{d \ln Y}{dt} = -\beta \ln \frac{Y}{Y_\infty}. \tag{8}$$

The Gompertzian limit implies an exponentially decreasing relative growth rate from the very first time point, which under the SIR model is unlikely given the large pool of susceptible people in the beginning. One would rather expect a near-constant relative growth rate in the beginning due to a disease propagation stage. Rypdal and Rypdal (2020) [14] suggest that the decreasing relative growth rate is caused by social and political mitigating efforts, but these hardly justify such coherent and consistent mortality characteristics across countries, particularly with an exponentially decreasing growth rate. In fact, Wang et al. (2012) [94] showed that any values of $\theta < 1$ are not physically realistic under a similar SIR model as the one given above.

Perhaps a more likely scenario from which a Gompertz curve would emerge is the selective infection of central nodes in the transmission network resulting in an immediate decrease in relative growth. Herrmann and Schwartz (2020) [97] studied a networked SIR model on a variety of networks, but did not elaborate on a possible fit to a Gompertz curve. Estrada and Bartesaghi (2022) [98] found that the Gompertz curve can emerge in later stages of an SIS (Susceptible-Infected-Susceptible) model, but not in the beginning of an SIR-like model. Although it may be possible to realize Gompertzian growth from networks, firm theoretical work has yet to be done to establish this connection. We will touch upon how the Gompertz curve emerges from such networks below.

In general, the SIR models and their extensions which do not consider network interactions explicitly consider rather averaged macroscopic behavior of an ensemble of microscopic units justified through mean-field theory [99]. In such models, all entities are implicitly assumed connected and communicating through long-range interactions as shown by Mombach et al. (2002) [100]. However, even with such strong assumptions, it is odd that observed mortality never exhibits an early stage of near-exponential growth as this macroscopic SIR model family predicts, but rather a constant negative slope in the double-log-domain, which we will now show.

The extended SIR model family is almost Gompertzian

Without the linearization and the θ -parametrization of the Richards model, one can still almost obtain Gompertzian growth from the communicable disease SIR-like models. Under a double-log-transform of Gompertzian growth, the growth curve is a simple straight line which is one of its characteristic traits [17],

$$\ln(\ln(Y_\infty/Y)) = -\beta t + k, \quad (9)$$

with $k = \ln \ln \frac{Y_\infty}{Y_0}$. To show this discrepancy between the SIR-like models and the Gompertz model, we follow Carletti et al. (2020) [5] and consider the extended version of the SIR model by including a group of deceased, $D(t)$, such that $N = S(t) + I(t) + R(t) + D(t)$, also called the SIRD model. This extension requires an addition to our original set of equations 2, with an equation for the deceased group's growth at some rate δ relative to the current infected group,

$$\frac{dD}{dt} = \delta I. \quad (10)$$

Algebraic manipulations reveal that the deceased group is described by a single equation, viz.

$$\frac{dD}{dt} = \eta(1 - e^{-\xi D}) - \kappa D, \quad (11)$$

where $\eta = \delta N$, $\xi = \beta/(\delta N)$, $\kappa = \delta + \alpha$, and where N is again synonymous to the initial susceptible pool of people. This pool of people cannot be obtained from deaths alone, but can be inferred by assuming a known ratio between mortality and recovery rates, δ and α . However, when only modeling the deceased as we do here, knowledge of N is irrelevant and the three parameters in Eq. 11 are sufficient.

In the beginning of an epidemic one can assume small values of D and thus the SIRD model follows logistic growth with

$$\frac{dD}{dt} = cD(1 - D/K), \quad (12)$$

obtained with a second order expansion of the exponential term in Eq. 11 and with $c = \eta\xi - \kappa$ and $K = 2c/\eta\xi^2$. Both the SIR and SEIR model exhibit similar properties and their discrepancy with initial observations related to the Covid pandemic has been noted by others [101].

Without this approximation, the full 3-parameters in Eq. 11 of the deceased group can show a curve quite close to a straight line in the double-log-domain even in the initial observations, although there will always be a non-zero concavity (Fig. 1 and the SI Appendix). This concavity is a property of the model rather than a discriminating measurement: over a single wave it is small enough that both models track the observations closely, and as we show below the two are separated far more sharply by how well their parameters can be determined than by how closely they fit.

Meanwhile, the Gompertz model is estimated through linear regression of log-mortality [17] with only two parameters,

$$\frac{d}{dt} \ln Y(t) = -\beta [\ln Y(t) - \ln Y_\infty], \quad (13)$$

a rate β at which the relative growth rate decays and an asymptote Y_∞ . Its solution is closed-form,

$$Y(t) = Y_\infty \left(\frac{Y_0}{Y_\infty} \right)^{e^{-\beta t}}, \quad (14)$$

where $Y_0 = Y(t = 0)$.

With two parameters the Gompertz model is parsimonious, and thus avoids the non-identifiability that plagues more complex models [102].

In summary, we see that the SIR model family both lacks the observed immediate decrease in relative growth rate, and makes strong assumptions about the disease spreading network structure that imply non-local interactions. We are thus prompted to look for another model which can explain the observed mortality patterns. For this challenge, we shall first present a stochastic approach without a network and then present an approach with a network which unifies the SIR family models with the Gompertz curve.




Figure 1: Cumulative number of deceased from the Covid pandemic, transformed by $g(Y) = \ln \ln(Y_\infty/Y(t))$ plotted against the number of days elapsed since the WHO declared a pandemic on 11 March 2020, marked on each panel by the dotted rule, comparing an SIRD model (green) with a Gompertz model (yellow) for a variety of countries in the period Jan-May 2020. Fitting begins where $Y(t)/Y_{max} > 0.005$, which for most countries falls after the declaration and for Italy, Iran and China before it; the common origin is what makes the panels comparable, since dating each country from its own take-off silently aligns sixteen different calendar dates at zero. Both models are fit using non-linear least squares according to equations 13 and 11 in the text. Although the Gompertz curve can also be obtained by linear regression of log-mortality, it is here obtained using non-linear least squares to put both models on equal footing. The temporal evolution of the SIRD model is obtained from the Runge-Kutta algorithm, while the Gompertz has a closed form for its temporal evolution. Each plot is annotated with the basic reproduction number defined as $R_0 = \gamma\xi/\kappa$ [5]. The 95th percent confidence interval is shown in brackets and computed with the parametric bootstrap assuming a multivariate normal distribution on the parameters and 10000 generated bootstrap samples [103]. Fitting is done with Python-Scipy and a 6-day moving average of deaths. Observations are taken from the Github repository compiled by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, Baltimore, USA [104].

The third SIRD parameter is not supported by the mortality data

Both models describe the observed $(\dot{D}, D)$ relationship well, and judged by the criterion on which they were fitted the SIRD is marginally the better of the two: across the sixteen countries of Fig. 2 the median coefficient of determination is 0.968 for Eq. 11 against 0.957 for Eq. 13, and the Akaike criterion prefers the SIRD in eleven of the sixteen countries even after charging it for its extra parameter. We therefore do not claim that these data reject the SIRD model, and we would caution against resting such a claim on a formal test of one model against the other. Both curves lie close to the observations over a single wave; a test that must distinguish them is being asked to resolve a difference of the same order as the noise, and the verdicts it returns are correspondingly unstable.

The more informative question is not which curve passes closer to the data but which parameters the data are able to determine at all. A convenient diagnostic is the condition number of the fitted Jacobian with each column scaled by its own parameter,

$$\text{cond} = \frac{\sigma_{\max}(J\Theta)}{\sigma_{\min}(J\Theta)}, \quad J_{ij} = \frac{\partial \dot{D}_i}{\partial \theta_j}, \quad \Theta = \text{diag}(\theta), \quad (15)$$

which measures how sharply *relative* changes in the different parameters are distinguished by the data, and which is invariant both to the units of the individual parameters and to any rescaling of the death counts. Table 1 reports it for both models. For Eq. 13 it lies between 3.8 and 4.4 for every country except China, and barely varies from one epidemic to the next: the rate and the asymptote are close to orthogonal directions, whatever the data. For Eq. 11 it runs from 18 to 594, an order of magnitude larger and strongly dependent on which country is supplied. The SIRD model is not unidentifiable — its Jacobian has full rank — but the third direction is weakly determined, and it is precisely the direction that carries $\mathcal{R}_0$.

The structure of that weakness is transparent in the rescaled coordinates of Carletti et al. [5]. Writing $x = \kappa D / \eta$ and measuring time in units of $1/\kappa$, Eq. 11 collapses onto the one-parameter family $dx/d(\kappa t) = 1 - e^{-\mathcal{R}_0 x} - x$: of the three parameters, η sets the vertical scale, η/κ the horizontal scale — the implied final death toll — and $\mathcal{R}_0$ alone the shape. The two scales can be read almost directly off the observed peak height and the observed total, and are determined accordingly. What the conditioning analysis reports is that the shape is not.

The practical consequence is a sensitivity to choices that ought to be immaterial. The smoothing window applied to $\dot{D}$ is even in length, so it cannot be centred exactly; placing it half a day to one side rather than the other changes no modelling assumption whatsoever. Yet it moves the fitted $\mathcal{R}_0$ by a median of 15% and by as much as 18.6% (Table 1), while the Gompertz rate β moves by a median of 1.4% and the asymptote Y_∞ by 0.8%. A quantity that shifts by a sixth under a half-day realignment of a moving average is not being measured to the three significant figures at which it is commonly reported.

The same instability is visible between independent implementations. Re-fitting Eq. 11 to the same source data as Carletti et al. [5], we recover their published γ/β — the ratio that fixes the implied final death toll — with a median ratio of 0.99, within 5% for eleven of sixteen countries, and their γ and β individually to within about 20%, the two moving together. But our ξ is systematically some 25% smaller, and since $\mathcal{R}_0 = (\gamma/\beta)\xi$ the discrepancy passes undiluted into the reproduction number, which comes out a median 24% below the published values. Half of that gap closes on adopting the odd-length window above. The point is not that either set of numbers is wrong, but that two careful fits of one model to one data set can differ by a third in the quantity the model is used to report, while the quantity the data actually constrain agrees to one per cent.

Finally, the models are fitted in the $(\dot{D}, D)$ plane but read in time. These are not the same test. Integrating each fitted equation forward and comparing with the observed trajectory under the transform $g(Y) = \ln \ln(Y_\infty/Y)$ of Fig. 1, the ordering reverses: the Gompertz residual is smaller in eleven of sixteen countries, by a median factor of 2.1 and by as much as a factor of 10 (Germany) and 8 (Austria). The SIRD's slender advantage in the phase plane is systematic rather than random — it is the concavity noted above — and systematic error accumulates on integration, whereas the Gompertz residual, though larger, does not. A model may fit the instantaneous relationship better and still track the epidemic worse.

Taken together these observations suggest a different reading of the SIR family's reputation for simplicity. Eq. 11 is simple in form, and it is derived from a transparent mechanism; but as an estimation problem posed against mortality data alone it carries one parameter more than the observations can support, and the resulting looseness is inherited by every quantity computed from the fit. Gompertzian growth, with one parameter fewer, gives up a little of the fit and returns estimates that are stable under the arbitrary choices of the analysis pipeline.

Table 1: Estimated parameters of the Gompertz model (Eq. 13), with standard deviations in parentheses corresponding to the last reported digit, alongside the scaled condition number of Eq. 15 for the Gompertz (cond_G) and SIRD (cond_S) fits. The final two columns give the relative change in the SIRD reproduction number and in the Gompertz rate when the even-length smoothing window applied to $\dot{D}$ is centred to the other side, a shift of half a day that alters no modelling assumption. Y_∞ is reported as a multiple of the death toll observed by 16 May 2020, so values near unity indicate a wave that had run its course by the end of the fitting window and larger values one that had not.

	β (days $^{-1}$)	Y_∞/Y_{obs}	cond_G	cond_S	$\Delta\mathcal{R}_0$	$\Delta\beta$
Belgium	0.0808(22)	1.076(11)	4.0	63	+15.3%	+1.9%
Portugal	0.0623(9)	1.140(8)	3.8	34	+17.2%	+1.2%
Austria	0.0762(13)	1.064(7)	4.0	67	+16.0%	+1.4%
Switzerland	0.0813(14)	1.050(6)	4.1	92	+17.5%	+1.3%
Sweden	0.0428(29)	1.474(77)	4.4	56	+1.9%	+3.8%
Italy	0.0588(9)	1.074(7)	3.9	23	+11.3%	+1.0%
France	0.0817(19)	1.052(9)	4.1	48	+15.3%	+1.4%
Spain	0.0750(15)	1.036(7)	4.1	21	+13.9%	+0.8%
Germany	0.0653(11)	1.149(10)	3.8	56	+15.5%	+2.1%
Denmark	0.0656(20)	1.099(14)	3.9	24	+3.8%	+1.4%
U. K.	0.0654(9)	1.136(8)	3.9	28	+15.0%	+1.7%
Netherlands	0.0653(11)	1.100(8)	3.9	26	+13.7%	+1.5%
Turkey	0.0638(11)	1.180(11)	3.8	66	+15.4%	+2.2%
Iran	0.0504(9)	1.103(9)	3.9	18	+12.7%	+0.7%
US	0.0532(10)	1.275(16)	4.1	20	+13.6%	+2.0%
China	0.0945(20)	1.006(4)	6.0	594	+18.6%	+0.6%

A stochastic theory for observed mortality

An alternative line of reasoning that does not rely on the framework of communicable diseases is that the environment was perturbed by an instantaneous external stressor, initiating a stress response to eventually bring mortality rates back to stability.

Inspired by De Lauro et al. (2014) [105], the stressor can be modeled as a multiplicative stochastic dampening term along with a countering force of the immune system. As with all multiplicative processes, it is convenient to log-transform mortality and work in a dimensionless space such that $F(t) = Y(t)/Y_\infty$ and $Z(t) = \ln F(t)$, from which a natural perturbation model emerges,

$$dZ(t) = -\beta Z(t)dt + \sqrt{\sigma}dB(t), \quad (16)$$

where $dB(t)$ is a delta-correlated Wiener process with zero mean, making $Z(t)$ a stochastic process. The second term on the right hand side represents the ongoing perturbation due to the stressor, while the first represents the stress response.

This perturbation model is an Ornstein–Uhlenbeck process [106], where $Z(t)$ is Gaussian. It is also interpreted as Newton’s equation of motion with friction and a random force (Langevin’s equation), and a continuous version of an Auto-Regressive(1) model [107]. As argued by De Lauro et al. (2014) [105], the diffusion coefficient, σ , represents the strength of the perturbation. This is seen if we recast our perturbation model in terms of mortality while introducing a new parameter $C = \exp(-\frac{\sigma}{2\beta})$, viz.

$$dF(t) = \left\{ \frac{\sigma}{2}F(t) - \beta F(t) \ln \left[\frac{F(t)}{C} \right] \right\} dt + \sqrt{\sigma}F(t)dB(t). \quad (17)$$

To obtain the deterministic observable, we first take the average in the log-domain (Eq. 16),

$$d\mathbb{E}[\ln F(t)] = -\beta\mathbb{E}[\ln F(t)]dt, \quad (18)$$

where $\mathbb{E}$ signifies the averaging operation. Then we use the property that for the log-normal quantity $F(t)$, the average of its logarithm is the logarithm of the *median* quantity [108], where we denote the median of $F(t)$ as $M(t)$,

$$d\ln M(t) = -\beta\ln M(t)dt, \quad (19)$$

which is the deterministic Gompertz equation of Eq. 13 in the normalised variable, with $Y(t) = Y_\infty M(t)$ and asymptote $M_\infty = 1$. The two terms of Eq. 17 separate the roles cleanly: the positive drift $\sigma F/2$ is generated by the stressor, so its

strength σ is the source of growth, while β is the growth-limiting factor. It is the constant $C = \exp(-\sigma/2\beta)$ that places the asymptote exactly at $M_\infty = 1$, so the two are not independent, and neither can be read off a fitted mortality curve, on which only β and Y_∞ appear. This comparison is clear after taking the median of Eq. 17 wherein the cross-term of $F(t)dB(t)$ goes to zero due to the independence of $F(t)$ and $dB(t)$.

Thus, a more parsimonious interpretation of the observations not reliant on a transmission network is that mortality was caused by a perturbation of the biosphere, modeled as a random process, to which organisms gradually develop resistance at a geometric rate in the log-transformed domain [109, 110], which is the natural transformation for many processes in nature [111]. Under this model, the distribution of the abundance of $F(t)$ is log-normal, a result that can be obtained by directly solving the stochastic equation 17 [112, 108], or from thermodynamic principles [113–115]. Intuitively, this is seen by noting that the solution to the perturbation model in the log-domain (Eq. 16) is Gaussian in the variable $Z(t)$, thus suggesting a log-normal distribution of $F(t)$.

Coherent growth as a mechanism for the observed Gompertz curve

The stochastic account above is a single-entity model. It is natural to ask what distinguishes logistic from Gompertzian dynamics once a population of interacting entities is admitted, and the answer bears directly on the estimation problem of the previous section. We summarise the relevant results here; the derivations, the interpolating family and the supporting simulations are given in a companion paper [116].

The distinguishing property is closure under aggregation. Splitting the system into N microscopic abundances x_i and writing $z_i = \ln x_i$, Gompertz dynamics are linear in z and therefore closed under the geometric mean: the microscopic and macroscopic domains obey the same equation, exactly and at every stage of the epidemic. Logistic dynamics with transmission, $\dot{x}_i = \beta(1 - x_i) \sum_j w_{ij} x_j$, are not. Averaging leaves a term $\frac{\beta}{N} \sum_i \sum_j w_{ij} x_i x_j$ which equals $\beta \bar{x}^2$ only if neighbouring states are uncorrelated or all nodes are perfectly synchronised — and neither is compatible with a communicable disease, since the first denies the transmission mechanism and the second denies any heterogeneity in health status. Macroscopic logistic growth is thus an approximation conditioned on a property its own microscopic model violates, whereas macroscopic Gompertz growth is exact.

Embedding both in the Richards family through an interaction order θ , which recovers Gompertz as $\theta \rightarrow 0$ and logistic-type dynamics at $\theta = 1$, the Gompertz limit is the unique member for which the coupling term contributes exactly zero to the aggregate growth [116]. The transmission network becomes dynamically invisible in the aggregate: it still governs each individual trajectory, but it leaves no trace in the population curve. This is what connects the mechanism to Table 1. If the aggregate mortality of a coherent population is blind to the network, then no property of that network — $\mathcal{R}_0$ among them — can be recovered by fitting a curve to aggregate deaths, and an attempt to do so will produce exactly what we observe: a parameter that is formally estimable, but so weakly determined that a half-day shift in a moving average moves it by a sixth.

We stress that the empirical results of the previous section do not depend on this mechanism. That $\mathcal{R}_0$ is poorly conditioned in Eq. 11, and that the Gompertz rate is not, are statements about the fit that hold whatever one believes about the underlying dynamics. The coherent-growth account is offered as an explanation consistent with those facts, not as a premise on which they rest.

Conclusion

We have argued that the SIR family’s reputation for simplicity does not survive contact with mortality data. Eq. 11 is simple in form and transparent in derivation, but as an estimation problem posed against deaths alone it carries one parameter more than the observations support. Its fitted Jacobian is an order of magnitude worse conditioned than that of the Gompertz model, and the ill-conditioned direction is precisely the one carrying $\mathcal{R}_0$. The consequences are measurable: a half-day change in how an even-length moving average is centred — a choice with no modelling content whatsoever — moves the fitted reproduction number by a median of 15%, while the Gompertz rate moves by 1.4%; and two careful fits of the same equation to the same source data differ by a quarter in $\mathcal{R}_0$ while agreeing in the median to about one per cent on the combination the data actually determine. Reproduction numbers obtained this way are routinely reported to three significant figures with uncertainties an order of magnitude smaller than these shifts.

We do not conclude from this that the SIRD model is false. Over a single wave it fits the observed $(\dot{D}, D)$ relationship marginally better than the Gompertz model does, and we have been explicit that a formal test between the two is a weak instrument at these sample sizes. The conclusion is narrower and, we think, more useful: the quantity these models are most often fitted in order to report is not identifiable from aggregate mortality, and the apparent precision with which it is quoted is an artefact of an error model that ignores the smoothing applied to the data. The Gompertz model, with one

parameter fewer, gives up a little of the fit and returns a rate and an asymptote that are stable under the arbitrary choices of the analysis pipeline — and, when the fitted equations are integrated forward, tracks the observed trajectory more closely in eleven of the sixteen countries examined.

The Gompertz curve also admits a mechanistic reading. It is the unique member of the Richards family that is closed under aggregation, and in that limit the transmission network contributes nothing to the aggregate growth [116]. A population whose collective mortality is blind to its own contact network is one from which no transmission parameter can be recovered by curve-fitting, which is what the conditioning analysis measures from the opposite direction. Read together with the observation that relative growth rates decrease from the first reported death, without the propagation phase a finite-speed transmission process requires, this favours an account in which the population is perturbed simultaneously by a common external stressor rather than sequentially by a communicable agent. Whether that account is correct is a separate question from the estimation result, which stands on its own.

Methods

Estimating model parameters for Covid data

The time-evolution of the SIRD and the Gompertz model are obtained with a Runge-Kutta algorithm, after first fitting parameters using non-linear least squares according to equations 13 and 11 in the text (see SI Appendix for details). Fitting is done using Python-Scipy’s non-linear least squares algorithm with specified Jacobians and a 6-day windowed average of observed deaths, similar to the technique by Carletti et al. [5]. Death counts are normalised by the toll observed at the end of the fitting window before fitting, which leaves β unchanged and expresses Y_∞ as a multiple of the observed total.

Identifiability is assessed with the scaled condition number of Eq. 15. The column scaling is what makes the diagnostic comparable between models with parameters of different physical dimension, and it renders the number invariant to the normalisation above; as a check, the SIRD fit for Italy returns 18.1 whether deaths are counted in units of one or of one thousand. Sensitivity to the smoothing convention is measured by repeating every fit with a 7-day window, which unlike the 6-day window admits an exact centre, and recording the relative change in each estimated quantity.

We caution against reading the reported standard errors as a full account of the uncertainty. They are obtained from the least-squares covariance under the assumption of independent errors, whereas $\dot{D}$ is a centred moving average, so residuals a day apart share five of their six constituent observations and are strongly correlated by construction. The intervals so obtained are consequently far narrower than the spread between equally defensible analyses of the same data: the $\mathcal{R}_0$ values of Table 1 and those of ref. [5] differ by amounts that are tens of times the uncertainty either analysis quotes. The comparisons in Table 1 are reported for this reason — a perturbation study of the analysis pipeline bounds the reproducibility of an estimate more honestly than its nominal standard error does. Observations are taken from the Github repository compiled by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, Baltimore, USA [104].

The basic reproduction number is modeled as the product of $\eta\xi/\kappa$ in Eq. 11, and its associated variance is obtained through a parametric bootstrap assuming a multivariate normal distribution with mean and covariance equal to the estimate from the non-linear least squares technique. The 95th percent confidence interval is obtained after 10000 parameter samples drawn from this distribution, each computing a resampled estimate of $\mathcal{R}_0$ [103].

Code Availability

Scripts used to produce the plots are downloadable in the form of a Jupyter notebook using Python from https://nbviewer.org/urls/emf-research.fra1.digitaloceanspaces.com/gompertz/gompertz_vs_sird.ipynb and <https://nbviewer.org/urls/emf-research.fra1.digitaloceanspaces.com/gompertz/hypergeometric.ipynb>

Data Availability

Observations are taken from the Github repository compiled by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, Baltimore, USA [104], and are available at <https://github.com/CSSEGISandData/COVID-19>, specifically the table `csse_covid_19_data/csse_covid_19_time_series/time_series_covid19_deaths_global.csv`, downloaded Aug. 5th 2022.

Competing interests

The authors declare no competing interests.

Supplement




Figure 2: Cumulative diseased plotted against number of days after $Y(t)/Y_{max} > 0.005$, comparing an SIRD with a Gompertz model. Both models are fit using non-linear least squares according to equations (13) and (11) in the text. Fitting is done with Python-Scipy and a 6-day windowed average of deaths. The basic reproduction number, R_0 is computed as specified in the Methods. Observations are taken from the Github repository compiled by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, Baltimore, USA [104].

References

- [1] Harko, T., Lobo, F. S., and Mak, M. Exact analytical solutions of the susceptible-infected-recovered (sir) epidemic model and of the sir model with equal death and birth rates. *Applied Mathematics and Computation*, 236:184–194, 2014.
- [2] Kröger, M. and Schlickeiser, R. Analytical solution of the SIR-model for the temporal evolution of epidemics. part a: time-independent reproduction factor. *Journal of Physics A: Mathematical and Theoretical*, 53(50): 505601, 2020.
- [3] Schlickeiser, R. and Kröger, M. Analytical solution of the SIR-model for the temporal evolution of epidemics: part b. semi-time case. *Journal of Physics A: Mathematical and Theoretical*, 54(17):175601, 2021.
- [4] Heng, K. and Althaus, C. L. The approximately universal shapes of epidemic curves in the susceptible-exposed-infectious-recovered (SEIR) model. *Scientific Reports*, 10(1):1–6, 2020.

Table 2: Estimated values for the SIRD model fitted to various countries' observations as shown in Fig. 2 using the notation of Eq. 11. Standard deviation estimates is given in the parenthesis corresponding to the last significant digit of the reported number.

	$\eta \times 10^{-1}$	$\xi \times 10^4$	$\kappa \times 10^2$
Belgium	94(8)	3.2(2)	9.3(7)
Portugal	6.6(4)	35(2)	4.5(3)
Austria	6.9(5)	40(3)	8.7(6)
Switzerland	22(1)	13.6(7)	11.0(5)
Sweden	19(5)	8(2)	3(1)
Italy	137(3)	1.77(5)	3.71(1)
France	240(10)	1.24(7)	8.1(4)
Spain	145(2)	2.27(5)	4.75(8)
Germany	61(4)	3.9(3)	6.5(5)
Denmark	2.7(2)	96(8)	4.3(3)
United Kingdom	181(5)	1.39(5)	4.3(1)
Netherlands	29(1)	8.9(5)	4.3(2)
Turkey	34(3)	6.8(5)	6.9(6)
Iran	22.3(4)	10.4(4)	2.51(7)
US	312(5)	0.76(2)	1.83(7)
China	130(30)	3.5(5)	26(4)

- [5] Carletti, T., Fanelli, D., and Piazza, F. COVID-19: The unreasonable effectiveness of simple models. *Chaos, Solitons & Fractals*, X, 5:100034, 2020.
- [6] Cooper, I., Mondal, A., and Antonopoulos, C. G. A SIR model assumption for the spread of COVID-19 in different communities. *Chaos, Solitons & Fractals*, 139:110057, 2020.
- [7] Postnikov, E. B. Estimation of COVID-19 dynamics “on a back-of-envelope”: Does the simplest SIR model provide quantitative parameters and predictions? *Chaos, Solitons & Fractals*, 135:109841, 2020.
- [8] Muñoz-Fernández, G. A., Seoane, J. M., and Seoane-Sepúlveda, J. B. A SIR-type model describing the successive waves of COVID-19. *Chaos, Solitons & Fractals*, 144:110682, 2021.
- [9] Cooper, I., Mondal, A., Antonopoulos, C. G., and Mishra, A. Dynamical analysis of the infection status in diverse communities due to COVID-19 using a modified SIR model. *Nonlinear Dynamics*, pages 1–14, 2022.
- [10] Saikia, D., Bora, K., and Bora, M. P. COVID-19 outbreak in india: an SEIR model-based analysis. *Nonlinear Dynamics*, 104(4):4727–4751, 2021.
- [11] Gompertz, B. XXIV. on the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. in a letter to francis baily, esq. f. r. s. &c. *Philosophical Transactions of the Royal Society of London*, 115:513–583, dec 1825. doi:10.1098/rstl.1825.0026.
- [12] Bajzer, Ž., Vuk-Pavlović, S., and Huzak, M. Mathematical modeling of tumor growth kinetics. In *A survey of models for tumor-immune system dynamics*, pages 89–133. Springer, 1997.
- [13] Ohnishi, A., Namekawa, Y., and Fukui, T. Universality in COVID-19 spread in view of the gompertz function. *Progress of Theoretical and Experimental Physics*, 2020(12), oct 2020. doi:10.1093/ptep/ptaa148.
- [14] Rypdal, K. and Rypdal, M. A parsimonious description and cross-country analysis of COVID-19 epidemic curves. *International Journal of Environmental Research and Public Health*, 17(18):6487, sep 2020. doi:10.3390/ijerph17186487.
- [15] Català, M., Alonso, S., Alvarez-Lacalle, E., López, D., Cardona, P.-J., and Prats, C. Empirical model for short-time prediction of COVID-19 spreading. *PLOS Computational Biology*, 16(12):e1008431, dec 2020. doi:10.1371/journal.pcbi.1008431.
- [16] Rodrigues, T. and Helene, O. Monte carlo approach to model COVID-19 deaths and infections using gompertz functions. *Physical Review Research*, 2(4):043381, 2020.
- [17] Levitt, M., Scaiewicz, A., and Zonta, F. Predicting the trajectory of any COVID19 epidemic from the best straight line. *medRxiv*, jun 2020. doi:10.1101/2020.06.26.20140814.
- [18] Castro, M., Ares, S., Cuesta, J. A., and Manrubia, S. The turning point and end of an expanding epidemic cannot be precisely forecast. *Proceedings of the National Academy of Sciences*, 117(42):26190–26196, 2020.

- [19] Hamidi, S., Sabouri, S., and Ewing, R. Does density aggravate the COVID-19 pandemic? *Journal of the American Planning Association*, 86(4):495–509, June 2020. doi:10.1080/01944363.2020.1777891.
- [20] Hamidi, S., Ewing, R., and Sabouri, S. Longitudinal analyses of the relationship between development density and the COVID-19 morbidity and mortality rates: Early evidence from 1,165 metropolitan counties in the united states. *Health & Place*, 64:102378, July 2020. doi:10.1016/j.healthplace.2020.102378.
- [21] Carozzi, F. Urban density and covid-19. *SSRN Electronic Journal*, 2020. doi:10.2139/ssrn.3643204.
- [22] Liu, L. Emerging study on the transmission of the novel coronavirus (covid-19) from urban perspective: Evidence from china. *Cities*, 103:102759, 2020.
- [23] A.G., G., S., C., M.R., M., J.C., V., G., S., G.W., C., and J.B., S. Covid-19 mortality rates in the european union, switzerland, and the uk: effect of timeliness, lockdown rigidity, and population density. *Minerva Med.*, 111(4), 2020.
- [24] Khavarian-Garmsir, A. R., Sharifi, A., and Moradpour, N. Are high-density districts more vulnerable to the COVID-19 pandemic? *Sustainable Cities and Society*, 70:102911, 2021.
- [25] Barak, N., Sommer, U., and Mualam, N. Urban attributes and the spread of COVID-19: The effects of density, compliance and socio-political factors in israel. *Science of the Total Environment*, 793:148626, 2021.
- [26] Arpino, B., Bordone, V., and Pasqualini, M. No clear association emerges between intergenerational relationships and COVID-19 fatality rates from macro-level analyses. *Proceedings of the National Academy of Sciences*, 117(32):19116–19121, jul 2020. doi:10.1073/pnas.2008581117.
- [27] Fischer, L., Peters, M., Merbach, S., Eydner, M., Kuczka, A., Lambertz, J., Kummerfeld, M., Kahnt, K., Weiss, A., and Petersen, H. Increased mortality in wild tits in north rhine-westphalia (germany) in 2020 with a special focus on suttonella ornithocola and other infectious pathogens. *European Journal of Wildlife Research*, 67(3), may 2021. doi:10.1007/s10344-021-01500-7.
- [28] Perisé-Barrios, A. J., Tomeo-Martín, B. D., Gómez-Ochoa, P., Delgado-Bonet, P., Plaza, P., Palau-Concejo, P., González, J., Ortiz-Díez, G., Meléndez-Lazo, A., Gentil, M., García-Castro, J., and Barbero-Fernández, A. Humoral responses to SARS-CoV-2 by healthy and sick dogs during the COVID-19 pandemic in spain. *Veterinary Research*, 52(1), feb 2021. doi:10.1186/s13567-021-00897-y.
- [29] Duff, P., Fenimore, C., Holmes, P., Hopkins, B., Jones, J., He, M., Everest, D., and Rocchi, M. Disease surveillance in england and wales, december 2019. *Veterinary Record*, 186(1):15–16, Jan. 2020. doi:10.1136/vr.m68.
- [30] Hu, B., Wei, H., Fan, Z., Song, Y., Chen, M., Qiu, R., Zhu, W., Xu, W., Xue, J., and Wang, F. Emergence of rabbit haemorrhagic disease virus 2 in china in 2020. *Veterinary Medicine and Science*, 7(1):236–239, Aug. 2020. doi:10.1002/vms3.332.
- [31] Fukui, H., Shimoda, H., Kadekaru, S., Henmi, C., and Une, Y. Rabbit hemorrhagic disease virus type 2 epidemic in a rabbit colony in japan. *Journal of Veterinary Medical Science*, 83(5):841–845, 2021. doi:10.1292/jvms.21-0007.
- [32] van Aarde, R. J., Pimm, S. L., Guldemon, R., Huang, R., and Maré, C. The 2020 elephant die-off in botswana. *PeerJ*, 9:e10686, jan 2021. doi:10.7717/peerj.10686.
- [33] Kambayashi, Y., Bannai, H., Tsujimura, K., Hirama, A., Ohta, M., and Nemoto, M. Outbreak of equine coronavirus infection among riding horses in tokyo, japan. *Comparative Immunology, Microbiology and Infectious Diseases*, 77:101668, Aug. 2021. doi:10.1016/j.cimid.2021.101668.
- [34] King, S., Rajko-Nenow, P., Ashby, M., Frost, L., Carpenter, S., and Batten, C. Outbreak of african horse sickness in thailand, 2020. *Transboundary and Emerging Diseases*, jul 2020. doi:10.1111/tbed.13701.
- [35] Lu, G., Pan, J., Ou, J., Shao, R., Hu, X., Wang, C., and Li, S. African horse sickness: Its emergence in thailand and potential threat to other asian countries. *Transboundary and Emerging Diseases*, jun 2020. doi:10.1111/tbed.13625.
- [36] Castillo-Olivares, J. African horse sickness in thailand: Challenges of controlling an outbreak by vaccination. *Equine Veterinary Journal*, 53(1):9–14, oct 2020. doi:10.1111/evj.13353.
- [37] Cannon, P. and Rycroft, M. J. Schumann resonance frequency variations during sudden ionospheric disturbances. *Journal of Atmospheric and Terrestrial Physics*, 44(2):201–206, 1982.
- [38] Vieira, C. L. Z., Chen, K., Garshick, E., Liu, M., Vokonas, P., Ljungman, P., Schwartz, J., and Koutrakis, P. Geomagnetic disturbances reduce heart rate variability in the normative aging study. *Science of The Total Environment*, page 156235, 2022.

- [39] Alabdulgader, A., McCraty, R., Atkinson, M., Dobyns, Y., Vainoras, A., Ragulskis, M., and Stolc, V. Long-term study of heart rate variability responses to changes in the solar and geomagnetic environment. *Scientific Reports*, 8(1), feb 2018. doi:10.1038/s41598-018-20932-x.
- [40] Ghione, S., Mezzasalma, L., Seppia, C., and Papi, F. Do geomagnetic disturbances of solar origin affect arterial blood pressure? *Journal of human hypertension*, 12(11):749–754, 1998.
- [41] Chernouss, S., Vinogradov, A., and Vlassova, E. Geophysical hazard for human health in the circumpolar auroral belt: evidence of a relationship between heart rate variation and electromagnetic disturbances. *Natural hazards*, 23(2):121–135, 2001.
- [42] Stoupel, E. Ambulatory blood pressure monitoring in patients with hypertension on days of high and low geomagnetic activity. *J Hum Hypertens*, 9:293–294, 1995.
- [43] Stoupel, E., Kalediene, R., Petrauskiene, J., Starkuviene, S., Abramson, E., Israelevich, P., Sulkes, J., et al. Twenty years study of solar, geomagnetic, cosmic ray activity links with monthly deaths number (n-850304). *Journal of Biomedical Science and Engineering*, 4(06):426, 2011.
- [44] Malin, S. and Srivastava, B. Correlation between heart attacks and magnetic activity. *Nature*, 277(5698):646–648, 1979.
- [45] Stoupel, E. Sudden cardiac deaths and ventricular extrasystoles on days with four levels of geomagnetic activity. *Journal of basic and clinical physiology and pharmacology*, 4(4):357–366, 1993.
- [46] Burch, J., Reif, J., and Yost, M. Geomagnetic disturbances are associated with reduced nocturnal excretion of a melatonin metabolite in humans. *Neuroscience Letters*, 266(3):209–212, 1999.
- [47] Rapoport, S., Malinovskaia, N., Oraevskii, V., Komarov, F., Nosovskii, A., and Vetterberg, L. Effects of disturbances of natural magnetic field of the earth on melatonin production in patients with coronary heart disease. *Klinicheskaiia Meditsina*, 75(6):24–26, 1997.
- [48] Bergiannaki, J.-D., Paparrigopoulos, T., and Stefanis, C. N. Seasonal pattern of melatonin excretion in humans: relationship to daylength variation rate and geomagnetic field fluctuations. *Experientia*, 52(3):253–258, 1996.
- [49] Schiff, J. E., Vieira, C. L., Garshick, E., Wang, V., Blomberg, A., Gold, D. R., Schwartz, J., Tracy, S. M., Vokonas, P., and Koutrakis, P. The role of solar and geomagnetic activity in endothelial activation and inflammation in the nas cohort. *PloS one*, 17(7):e0268700, 2022.
- [50] Keeton, W. T., Larkin, T. S., and Windsor, D. M. Normal fluctuations in the earth’s magnetic field influence pigeon orientation. *Journal of comparative physiology*, 95(2):95–103, 1974.
- [51] Schreiber, B. and Rossi, O. Correlation between race arrivals of homing pigeons and solar activity. *Italian Journal of Zoology*, 43(3):317–320, 1976.
- [52] Kowalski, U., Wiltchko, R., and Füller, E. Normal fluctuations of the geomagnetic field may affect initial orientation in pigeons. *Journal of Comparative Physiology A*, 163(5):593–600, 1988.
- [53] Bianco, G., Ilieva, M., and Åkesson, S. Magnetic storms disrupt nocturnal migratory activity in songbirds. *Biology letters*, 15(3):20180918, 2019.
- [54] Ferrari, T. E. Cetacean beachings correlate with geomagnetic disturbances in earth’s magnetosphere: an example of how astronomical changes impact the future of life. *International Journal of Astrobiology*, 16(2):163–175, 2017.
- [55] Vanselow, K. H., Jacobsen, S., Hall, C., and Garthe, S. Solar storms may trigger sperm whale strandings: explanation approaches for multiple strandings in the north sea in 2016. *International Journal of Astrobiology*, 17(4):336–344, 2018.
- [56] Granger, J., Walkowicz, L., Fitak, R., and Johnsen, S. Gray whales strand more often on days with increased levels of atmospheric radio-frequency noise. *Current Biology*, 30(4):R155–R156, 2020.
- [57] Zenchenko, T. A. and Breus, T. K. The possible effect of space weather factors on various physiological systems of the human organism. *Atmosphere*, 12(3):346, 2021.
- [58] Zilli Vieira, C. L., Alvares, D., Blomberg, A., Schwartz, J., Coull, B., Huang, S., and Koutrakis, P. Geomagnetic disturbances driven by solar activity enhance total and cardiovascular mortality risk in 263 us cities. *Environmental Health*, 18(1):1–10, 2019.
- [59] Gordon, C. and Berk, M. The effect of geomagnetic storms on suicide. *South African Psychiatry Review*, 6(3): 24–27, 2003.
- [60] Kay, R. W. Geomagnetic storms: association with incidence of depression as measured by hospital admission. *The British Journal of Psychiatry*, 164(3):403–409, 1994.

- [61] Kay, R. W. Schizophrenia and season of birth: relationship to geomagnetic storms. *Schizophrenia Research*, 66 (1):7–20, 2004.
- [62] Halberg, F., Cornélissen, G., Panksepp, J., Otsuka, K., and Johnson, D. Chronomics of autism and suicide. *Biomedicine & pharmacotherapy*, 59:S100–S108, 2005.
- [63] Berk, M., Dodd, S., and Henry, M. Do ambient electromagnetic fields affect behaviour? a demonstration of the relationship between geomagnetic storm activity and suicide. *Bioelectromagnetics: Journal of the Bioelectromagnetics Society, The Society for Physical Regulation in Biology and Medicine, The European Bioelectromagnetics Association*, 27(2):151–155, 2006.
- [64] Doronin, V., Parfentëv, V., SZh, T., Namvar, R., Somsikov, V., Drobzhev, V., and Chemeris, A. Effect of variations of the geomagnetic field and solar activity on human physiological indicators. *Biofizika*, 43(4):647–653, 1998.
- [65] Persinger, M. and Psych, C. Sudden unexpected death in epileptics following sudden, intense, increases in geomagnetic activity: prevalence of effect and potential mechanisms. *International Journal of Biometeorology*, 38(4):180–187, 1995.
- [66] Cherry, N. Schumann resonances, a plausible biophysical mechanism for the human health effects of solar. *Natural hazards*, 26(3):279–331, 2002.
- [67] Schumann, W. O. Über die strahlungslosen eigenschwingungen einer leitenden kugel, die von einer luftschicht und einer ionosphärenhülle umgeben ist. *Zeitschrift für Naturforschung A*, 7(2):149–154, 1952.
- [68] Wever, R. ELF-effects on human circadian rhythms. In *ELF and VLF electromagnetic field effects*, pages 101–144. Springer, 1974.
- [69] Pobachenko, S., Kolesnik, A., Borodin, A., and Kalyuzhin, V. The contingency of parameters of human encephalograms and schumann resonance electromagnetic fields revealed in monitoring studies. *Biophysics*, 51 (3):480–483, 2006.
- [70] Saroka, K. S. and Persinger, M. A. Quantitative evidence for direct effects between earth-ionosphere schumann resonances and human cerebral cortical activity. *International Letters of Chemistry, Physics and Astronomy*, 20, 2014.
- [71] Persinger, M. A., Saroka, K. S., et al. Human quantitative electroencephalographic and schumann resonance exhibit real-time coherence of spectral power densities: implications for interactive information processing. *Journal of Signal and Information Processing*, 6(02):153, 2015.
- [72] König, H. L., Krueger, A. P., Lang, S., and Sönning, W. *Biologic effects of environmental electromagnetism*. Springer Science & Business Media, 2012.
- [73] Dyrda, M., Kulak, A., Mlynarczyk, J., and Ostrowski, M. Novel analysis of a sudden ionospheric disturbance using schumann resonance measurements. *Journal of Geophysical Research: Space Physics*, 120(3):2255–2262, 2015.
- [74] Roldugin, V., Maltsev, Y. P., Vasiljev, A., Schokotov, A., and Belyajev, G. Schumann resonance frequency increase during solar x-ray bursts. *Journal of Geophysical Research: Space Physics*, 109(A1), 2004.
- [75] Sători, G., Williams, E., and Mushtak, V. Response of the earth–ionosphere cavity resonator to the 11-year solar cycle in x-radiation. *Journal of Atmospheric and Solar-Terrestrial Physics*, 67(6):553–562, 2005.
- [76] Sători, G., Williams, E., Price, C., Boldi, R., Koloskov, A., Yampolski, Y., Guha, A., and Barta, V. Effects of energetic solar emissions on the earth–ionosphere cavity of schumann resonances. *Surveys in Geophysics*, 37(4): 757–789, 2016.
- [77] Shvets, A., Nickolaenko, A., and Chebrov, V. Effect of solar flares on the schumann-resonance frequencies. *Radiophysics and Quantum Electronics*, 60(3):186–199, 2017.
- [78] Schlegel, K. and Füllekrug, M. Schumann resonance parameter changes during high-energy particle precipitation. *Journal of Geophysical Research: Space Physics*, 104(A5):10111–10118, 1999.
- [79] Singh, B., Tyagi, R., Hobara, Y., and Hayakawa, M. X-rays and solar proton event induced changes in the first mode schumann resonance frequency observed at a low latitude station agra, india. *Journal of Atmospheric and Solar-Terrestrial Physics*, 113:1–9, 2014.
- [80] Pazos, M., Mendoza, B., Sierra, P., Andrade, E., Rodríguez, D., Mendoza, V., and Garduño, R. Analysis of the effects of geomagnetic storms in the schumann resonance station data in mexico. *Journal of Atmospheric and Solar-Terrestrial Physics*, 193:105091, 2019.
- [81] Salinas, A., Toledo-Redondo, S., Navarro, E., Fornieles-Callejón, J., and Portí, J. Solar storm effects during saint patrick’s days in 2013 and 2015 on the schumann resonances measured by the elf station at sierra nevada (spain). *Journal of Geophysical Research: Space Physics*, 121(12):12–234, 2016.

- [82] Kudintseva, I., Galuk, Y. P., Nickolaenko, A., and Hayakawa, M. Modifications of middle atmosphere conductivity during sudden ionospheric disturbances deduced from changes of schumann resonance peak frequencies. *Radio Science*, 53(5):670–682, 2018.
- [83] Zhou, H. and Qiao, X. Studies of the variations of the first schumann resonance frequency during the solar flare on 7 march 2012. *Journal of Geophysical Research: Atmospheres*, 120(10):4600–4612, 2015.
- [84] Christofilakis, V., Tatsis, G., Votis, C., Contopoulos, I., Repapis, C., and Tritakis, V. Significant elf perturbations in the schumann resonance band before and during a shallow mid-magnitude seismic activity in the greek area (kalpaki). *Journal of Atmospheric and Solar-Terrestrial Physics*, 182:138–146, 2019.
- [85] Galuk, Y. P., Kudintseva, I. G., Nickolaenko, A. P., and Hayakawa, M. Scattering of elf radio waves by a localized non-uniformity in the lower ionosphere. *Journal of Atmospheric and Solar-Terrestrial Physics*, 194:105093, 2019.
- [86] Hayakawa, M., Nickolaenko, A., Galuk, Y. P., and Kudintseva, I. Scattering of extremely low frequency electromagnetic waves by a localized seismogenic ionospheric perturbation: Observation and interpretation. *Radio Science*, 55(12):1–26, 2020.
- [87] Marchitelli, V., Harabaglia, P., Troise, C., and Natale, G. D. On the correlation between solar activity and large earthquakes worldwide. *Scientific Reports*, 10(1), jul 2020. doi:10.1038/s41598-020-67860-3.
- [88] Zaporozhan, V. and Ponomarenko, A. Mechanisms of geomagnetic field influence on gene expression using influenza as a model system: basics of physical epidemiology. *International journal of environmental research and public health*, 7(3):938–965, 2010.
- [89] Rodgers, C. T. and Hore, P. J. Chemical magnetoreception in birds: the radical pair mechanism. *Proceedings of the National Academy of Sciences*, 106(2):353–360, 2009.
- [90] Engels, S., Schneider, N.-L., Lefeldt, N., Hein, C. M., Zapka, M., Michalik, A., Elbers, D., Kittel, A., Hore, P., and Mouritsen, H. Anthropogenic electromagnetic noise disrupts magnetic compass orientation in a migratory bird. *Nature*, 509(7500):353–356, 2014.
- [91] Hiscock, H. G., Worster, S., Kattnig, D. R., Steers, C., Jin, Y., Manolopoulos, D. E., Mouritsen, H., and Hore, P. The quantum needle of the avian magnetic compass. *Proceedings of the National Academy of Sciences*, 113(17):4634–4639, 2016.
- [92] Richards, F. A flexible growth function for empirical use. *Journal of experimental Botany*, 10(2):290–301, 1959.
- [93] Petroni, N. C., De Martino, S., and De Siena, S. Logistic and θ -logistic models in population dynamics: General analysis and exact results. *Journal of Physics A: Mathematical and Theoretical*, 53(44):445005, 2020.
- [94] Wang, X.-S., Wu, J., and Yang, Y. Richards model revisited: Validation by and application to infection dynamics. *Journal of theoretical biology*, 313:12–19, 2012.
- [95] Kermack, W. O. and McKendrick, A. G. A contribution to the mathematical theory of epidemics. *Proceedings of the royal society of london. Series A, Containing papers of a mathematical and physical character*, 115(772):700–721, 1927.
- [96] Wu, K., Darcet, D., Wang, Q., and Sornette, D. Generalized logistic growth modeling of the covid-19 outbreak: comparing the dynamics in the 29 provinces in china and in the rest of the world. *Nonlinear dynamics*, 101(3):1561–1581, 2020.
- [97] Herrmann, H. A. and Schwartz, J.-M. Why COVID-19 models should incorporate the network of social interactions. *Physical Biology*, 17(6):065008, 2020.
- [98] Estrada, E. and Bartesaghi, P. From networked SIS model to the gompertz function. *Applied Mathematics and Computation*, 419:126882, 2022.
- [99] Smilkov, D., Hidalgo, C. A., and Kocarev, L. Beyond network structure: How heterogeneous susceptibility modulates the spread of epidemics. *Scientific reports*, 4(1):1–7, 2014.
- [100] Mombach, J. C., Lemke, N., Bodmann, B. E., and Idiart, M. A. P. A mean-field theory of cellular growth. *EPL (Europhysics Letters)*, 59(6):923, 2002.
- [101] Vattay, G. Forecasting the outcome and estimating the epidemic model parameters from the fatality time series in COVID-19 outbreaks. *Physical Biology*, 17(6):065002, 2020.
- [102] Roda, W. C., Varughese, M. B., Han, D., and Li, M. Y. Why is it difficult to accurately predict the COVID-19 epidemic? *Infectious disease modelling*, 5:271–281, 2020.
- [103] Efron, B. Bayesian inference and the parametric bootstrap. *The annals of applied statistics*, 6(4):1971, 2012.

- [104] Dong, E., Du, H., and Gardner, L. An interactive web-based dashboard to track COVID-19 in real time. *The Lancet infectious diseases*, 20(5):533–534, 2020.
- [105] De Lauro, E., De Martino, S., De Siena, S., and Giorno, V. Stochastic roots of growth phenomena. *Physica A: Statistical Mechanics and its Applications*, 401:207–213, 2014.
- [106] Risken, H. *The Fokker-Planck Equation*. Springer, 1996.
- [107] Akaike, H. Statistical predictor identification. *Annals of the institute of Statistical Mathematics*, 22(1):203–217, 1970.
- [108] Petroni, N. C., De Martino, S., and De Siena, S. Gompertz and logistic stochastic dynamics: Advances in an ongoing quest. *arXiv preprint arXiv:2002.06409*, 2020.
- [109] Boxenbaum, H. Hypotheses on mammalian aging, toxicity, and longevity hormesis: Explication by a generalized gompertz function. In *Biological Effects of Low Level Exposures to Chemicals and Radiation*, pages 1–39. CRC Press, 2017.
- [110] Neafsey, P. J., Boxenbaum, H., Ciraulo, D. A., and Fournier, D. J. A gompertz age-specific mortality rate model of aging, hormesis, and toxicity: Fixed-dose studies. *Drug metabolism reviews*, 19(3-4):369–401, 1988.
- [111] Zhang, C.-L. and Popp, F.-A. Log-normal distribution of physiological parameters and the coherence of biological systems. *Medical Hypotheses*, 43(1):11–16, 1994.
- [112] Skiadas, C. H. Exact solutions of stochastic differential equations: Gompertz, generalized logistic and revised exponential. *Methodology and Computing in Applied Probability*, 12(2):261–270, 2010.
- [113] Sitaram, B. and Varma, V. Statistical mechanics of the gompertz model of interacting species. *Journal of theoretical biology*, 110(2):253–256, 1984.
- [114] Gunasekaran, N. and Pande, L. Log normal distribution for the intrinsic abundance of species from the gompertz model. *Journal of Theoretical Biology*, 98(2):301–305, 1982.
- [115] Chakrabarti, C. and Bhadra, S. Non equilibrium thermodynamics and stochastics of gompertzian growth. *J. Biol. Systems*, 4(2):151–157, 1996. doi:doi.org/10.1142/S0218339096000119.
- [116] Haugen, M. A. and Gilbert, D. From logistic to gompertz: A microscopic theory of coherent growth in biological systems. *arXiv preprint arXiv:2211.05653*, 2026.