
GOMPERTZ GROWTH AS THE COHERENT LIMIT OF EPIDEMIC DYNAMICS

A PREPRINT

May 17, 2026

ABSTRACT

A number of studies have shown that cumulative mortality followed a Gompertz curve in the initial Covid pandemic period in March-April 2020 – a pattern incompatible with expected initial logistic growth curves as predicted by traditional Susceptible-Infected-Recovered (SIR) models. We show that this incompatibility is not merely empirical but structural: logistic growth cannot close exactly under macroscopic aggregation unless nodes are either synchronized or statistically independent of the network — both conditions contradicting the premise of communicable disease transmission. We propose an alternative theory based on a simultaneous environmental stressor, and introduce a novel microscopic model that unifies logistic and Gompertzian growth through an order- θ power-mean coupling between network nodes. This model reveals Gompertz growth ($\theta \rightarrow 0$) as the unique coherent limit: the macroscopic dynamics decouple exactly from microscopic fluctuations, the system is unconditionally stable, and each entity contributes equally to the collective growth rate regardless of network topology. In contrast, Richards ($\theta > 0$) and logistic ($\theta = 1$) growth require synchronization as a precondition for macroscopic validity — a condition that Gompertz growth instead imposes. The coherence parameter θ thus plays the role of a symmetry-breaking parameter, with the Gompertz limit exhibiting infinite-order correlations in the natural domain, structural independence from microscopic noise, and a spectral gap protecting the macroscopic state — properties reminiscent of coherent states in quantum systems.

Keywords Gompertz · Coherence · Covid · Coronavirus · Network Analysis · Stochastic

Introduction

Traditional communicable disease spreading theory assumes a pathogen which infects the population through a network of transmission. Following this line of reasoning it can be theoretically shown that in the early stages of an epidemic, growth follows a logistic-like curve, where the very beginning exhibits exponential growth, and for which analytic and semi-analytic solutions have recently emerged [1–4]. A large body of research reveals how these models fit the mortality seen due to the Covid pandemic in 2020, a pandemic due to an influenza-like disease [5–10].

However, instead of showing logistic-like growth, observed cumulative mortality in the initial period March-April 2020 exhibits almost perfect resemblance to Gompertzian growth [11, 12] where the log-transformed cumulative mortality, or log-mortality for short, is exponentially *decreasing* in time,

$$\frac{d}{dt} \ln F(t) = -\beta \ln F(t), \quad (1)$$

using normalized mortality such that $F(t \rightarrow \infty) = 1$.

This phenomenon is recorded by others [13–17], showing Gompertz curves at national levels instead of the predicted logistic curves, an observation made possible in part due to the increased cadence and quality of data acquisition compared with earlier epidemics. What is causing such a discrepancy between these observations and the current theory of communicable diseases [18]?

In addition to the characteristic patterns of these national mortality curves, other observations that conflict more generally with the communicable disease models are 1) the lack of correlation between population density and mortality (or infection) rates [19–25], 2) no clear association between inter-generational relationships and fatality rates [26], and 3) the widespread morbidity and mortality spikes seen during spring 2020 among birds [27], dogs [28], rabbits [29–31], elephants [32], and horses [33–36]. Many of these animal mortality spikes are not attributable to a Coronavirus infection, so why did these spikes all occur within the same time frame? While these observations are not reconcilable within the current communicable disease paradigm, we present here an alternative theory that can reconcile all observations; a theory based on a global biosphere disturbance with fewer parameters than the most basic communicable disease model.

This global biosphere disturbance could be in the form of sudden ionospheric disturbances (SIDs) [37], which in the context of short-term solar and geomagnetic disturbances have been associated with physiological changes in cardiac system processes [38–45], melatonin/serotonin balance [46–48], endothelial activation and inflammation [49], and disturbed magnetoreception in migrating animals [50–56]. More general effects have also been studied in this context, e.g. overall mortality, suicide, mental disorders, homicides, and epileptic seizures [57–65]. Cherry (2002) put forward a mechanism whereby the ionosphere acts as a coupling agent between geomagnetic disturbances and physiological effects based on the disturbance of the background electromagnetic oscillations between the surface of the Earth and the lower ionosphere, called the Schumann Resonance signal [66, 67]. The Schumann Resonance would then couple with the brain to alter the melatonin/serotonin balance, which was a causal link observed by Wever (1974) through the circadian rhythm [68]. Indeed, more recent findings support this hypothesized synchronization between Schumann Resonances and the brain [69–72].

In support of Cherry’s theory, specific phenomena such as solar X-ray bursts [73–77], solar proton events [74, 78, 79] and geomagnetic storms [80, 81] can modify the shape of the ionosphere cavity and shift the Schumann resonance frequencies [82, 76, 83]. Seismic activity has also been connected to variations in Schumann resonances [84–86], and conversely, ionospheric modifications have been connected to global seismicity [87]. Zaporozhan and Ponomarenko (2010) postulated that minute disturbances in magnetic fields which could translate into influenza symptoms [88], mediated by the cryptochrome protein-family which has been shown to be geomagnetic receptors [89–91].

Whatever the cause, global biosphere disturbance theory represents an alternative paradigm compared to the traditional contact-transmissible contagion theory in the context of disease spreading dynamics. Our goal with this study is to present a novel mathematical bridge between these two paradigms which at its core is a connection between the Gompertzian growth model and the logistic growth model. Our proposed bridge represents a microscopic version to the well-known parametrization given by Richard (1959) [92], where he augmented standard logistical growth by one parameter. This is why it is also sometimes called θ -logistic growth [93].

One issue with Richard’s parametrization is that the traditional communicable disease models cannot be realistically extrapolated to the Gompertz model, as shown by Wang *et al.* (2012) [94]. The reason why Richard’s model cannot realistically represent a communicable disease with a Gompertz model is because it would imply that there is “long-range interaction” between the susceptible and infected (see [93] for details), which is counter to the basic assumption that communicable diseases spread through proximity between the susceptible and infected. Based on these findings, one would then be forced to conclude that Gompertz growth cannot arise in the context of spread through proximity between the susceptible and infected.

In this paper we will show this incompatibility in another way by creating a microscopic model where each entity is exposed to an environmental stressor simultaneously applied to the population. This theory does not preclude the possibility of multiple sources of disturbance, potentially local and interacting with a global source. The main idea is that over some possibly geographically constrained area, everyone is exposed to the same source of disturbance simultaneously.

The organization of the paper is as follows: We first review the basic communicable disease models, before presenting a new stochastic model from which Gompertz growth emerges and thus explains the observed mortality data. We then proceed to juxtapose of the logistic model and the Gompertz model in the microscopic domain where multiple entities comprise the macroscopic behavior. The final section is dedicated to unifying the communicable disease paradigm with a biosphere disturbance through a new microscopic form of the Logistic-Gompertzian growth equations.

Simple communicable disease models: The SIR model family

We start by reviewing the basic infectious disease model, namely the Susceptible-Infected-Recovered (SIR) model under a time-dependent infection/recovery rate ratio, $\phi(t)$ [95], and show under what conditions the Gompertz curve could emerge. First assume a pool of susceptible people of size N evolving between the three states: susceptible, $S(t)$,

recovered, $R(t)$ and infected, $I(t)$,

$$\begin{aligned}\frac{dS}{dt} &= -\beta \frac{IS}{N} \\ \frac{dI}{dt} &= \beta \frac{IS}{N} - \alpha I. \\ \frac{dR}{dt} &= \alpha I,\end{aligned}\tag{2}$$

omitting the argument t in each variable for brevity and where α and β in this context signify recovery and infection rates.

A line of reasoning employed by Rypdal and Rypdal (2020) [14] to obtain the Gompertz curve is to linearize the SIR model by assuming both the number of infected, $I(t)$, and cumulative infected, $Y(t)$, are much less than the initial pool of susceptible people, $N \gg Y \geq I$, which seems well founded at the national level, viz.

$$\frac{dY}{dt} = \beta I \tag{3}$$

$$\frac{dI}{dt} = (\beta - \alpha)I = \alpha(\phi(t) - 1)I, \tag{4}$$

and where the number of recovered, $R(t)$, is under this linearization decoupled from the other variables. They further assume the number of deceased is proportional to the number of cumulative infected, offset by a time lag, which allows us to use the same set of linearized equations to model the number of cumulative people deceased without loss of generality. And since I is a function of Y , we can combine Eq. 3 via an instantaneous relative growth rate, $\gamma(t) = dY(t)/(Y dt) = \beta I/Y$,

$$\frac{dY}{dt} = \gamma(t)Y(t). \tag{5}$$

Due to this linearization, the infection/recovery rate ratio, $\phi(t)$, and the relative growth rate, $\gamma(t)$, will have to change as a function of time to accommodate for the boundary conditions. The latter is typically also parameterized by a scaling factor, θ , representing the shape of the growth,

$$\gamma(t) = \frac{\beta}{\theta} \left[1 - \left(\frac{Y}{Y_\infty} \right)^\theta \right]. \tag{6}$$

This parametrization is the commonly used Richards growth curve [92], also called θ -logistic growth, and has been used by others [96]. Although not immediately justified in the communicable disease theory, one could imagine that θ represents non-linear network behavior [93]. Note that at $\theta = 1$, the traditional logistic growth curve is obtained, while at $\theta \rightarrow \infty$ we recover the exponential (Malthusian) explosion.

The observed Gompertzian mortality curves are realized in the limit $\theta \rightarrow 0$, with the relative growth rate,

$$\gamma(t) = \lim_{\theta \rightarrow 0} \frac{\beta}{\theta} \left[1 - \left(\frac{Y}{Y_\infty} \right)^\theta \right] = \beta \ln \frac{Y_\infty}{Y}, \tag{7}$$

such that

$$\frac{d \ln Y}{dt} = -\beta \ln \frac{Y}{Y_\infty}. \tag{8}$$

The Gompertzian limit implies an exponentially decreasing relative growth rate from the very first time point, which under the SIR model is unlikely given the large pool of susceptible people in the beginning. One would rather expect a near-constant relative growth rate in the beginning due to a disease propagation stage. Rypdal and Rypdal (2020) [14] suggest that the decreasing relative growth rate is caused by social and political mitigating efforts, but these hardly justify such coherent and consistent mortality characteristics across countries, particularly with an exponentially decreasing growth rate. In fact, Wang et al. (2012) [94] showed that any values of $\theta < 1$ are not physically realistic under a similar SIR model as the one given above.

Perhaps a more likely scenario from which a Gompertz curve would emerge is the selective infection of central nodes in the transmission network resulting in an immediate decrease in relative growth. Herrmann and Schwartz (2020) [97] studied a networked SIR model on a variety of networks, but did not elaborate on a possible fit to a Gompertz curve. Estrada and Bartesaghi (2022) [98] found that the Gompertz curve can emerge in later stages of an SIS (Susceptible-Infected-Susceptible) model, but not in the beginning of an SIR-like model. Although it may be possible to realize

Gompertzian growth from networks, firm theoretical work has yet to be done to establish this connection. We will touch upon how the Gompertz curve emerges from such networks below.

In general, the SIR models and their extensions which do not consider network interactions explicitly consider rather averaged macroscopic behavior of an ensemble of microscopic units justified through mean-field theory [99]. In such models, all entities are implicitly assumed connected and communicating through long-range interactions as shown by Mombach et al. (2002) [100]. However, even with such strong assumptions, it is odd that observed mortality never exhibits an early stage of near-exponential growth as this macroscopic SIR model family predicts, but rather a constant negative slope in the double-log-domain, which we will now show.

The extended SIR model family is almost Gompertzian

Without the linearization and the θ -parametrization of the Richards model, one can still almost obtain Gompertzian growth from the communicable disease SIR-like models. Under a double-log-transform of Gompertzian growth, the growth curve is a simple straight line which is one of its characteristic traits [17],

$$\ln(\ln(Y_\infty/Y)) = -\beta t + k, \quad (9)$$

with $k = \ln \ln \frac{Y_\infty}{Y_0}$. To show this discrepancy between the SIR-like models and the Gompertz model, we follow Carletti et al. (2020) [5] and consider the extended version of the SIR model by including a group of deceased, $D(t)$, such that $N = S(t) + I(t) + R(t) + D(t)$, also called the SIRD model. This extension requires an addition to our original set of equations 2, with an equation for the deceased group's growth at some rate δ relative to the current infected group,

$$\frac{dD}{dt} = \delta I. \quad (10)$$

Algebraic manipulations reveal that the deceased group is described by a single equation, viz.

$$\frac{dD}{dt} = \eta(1 - e^{-\xi D}) - \kappa D, \quad (11)$$

where $\eta = \delta N$, $\xi = \beta/(\delta N)$, $\kappa = \delta + \alpha$, and where N is again synonymous to the initial susceptible pool of people. This pool of people cannot be obtained from deaths alone, but can be inferred by assuming a known ratio between mortality and recovery rates, δ and α . However, when only modeling the deceased as we do here, knowledge of N is irrelevant and the three parameters in Eq. 11 are sufficient.

In the beginning of an epidemic one can assume small values of D and thus the SIRD model follows logistic growth with

$$\frac{dD}{dt} = cD(1 - D/K), \quad (12)$$

obtained with a second order expansion of the exponential term in Eq. 11 and with $c = \eta\xi - \kappa$ and $K = 2c/\eta\xi^2$. Both the SIR and SEIR model exhibit similar properties and their discrepancy with initial observations related to the Covid pandemic has been noted by others [101].

Without this approximation, the full 3-parameters in Eq. 11 of the deceased group can show a curve quite close to a straight line in the double-log-domain even in the initial observations, although there will always be a non-zero concavity (Fig. 1 and the SI Appendix).

Meanwhile, even though the Gompertz model can be fit with a 3-parameter model, it can also be simplified to a 2-parameter model estimated through linear regression of log-mortality [17],

$$\frac{d}{dt} \ln Y(t) = -\beta [\ln Y(t) - \ln Y_\infty]. \quad (13)$$

The 3-parameter model has one extra degree of freedom which can stabilize the asymptotic abundance estimate,

$$\frac{d}{dt} \ln Y(t) = -\beta \ln \frac{Y(t)}{\tilde{Y}} + \nu, \quad (14)$$

with constants $\tilde{Y}$, β , and ν . The Gompertz growth solution is given as

$$Y(t) = Y_\infty \left(\frac{Y_0}{Y_\infty} \right)^{e^{-\beta t}}, \quad (15)$$

where $Y_0 = Y(t=0)$ and $Y_\infty = \tilde{Y}e^{\nu/\beta}$.

Whether though the 3-parameter model or the 2-parameter linear model, the Gompertz model is relatively parsimonious in its parameters and thus mitigates the possibility of non-identifiability issues, which plagues more complex models [102].

In summary, we see that the SIR model family both lacks the observed immediate decrease in relative growth rate, and makes strong assumptions about the disease spreading network structure that imply non-local interactions. We are thus prompted to look for another model which can explain the observed mortality patterns. For this challenge, we shall first present a stochastic approach without a network and then present an approach with a network which unifies the SIR family models with the Gompertz curve.




Figure 1: Cumulative number of deceased from the Covid pandemic, transformed by $g(Y) = \ln \ln(Y_\infty/Y(t))$ plotted against number of days elapsed after $Y(t)/Y_{max} > 0.005$, comparing an SIRD model (green) with a Gompertz model (yellow) for a variety of countries in the period Jan-May 2020. Both models are fit using non-linear least squares according to equations 14 and 11 in the text. Although the Gompertz curve can also be obtained with a simple linear fit using Eq. 13, it is here obtained using non-linear least squares to put both models on equal footing. The temporal evolution of the SIRD model is obtained from the Runge-Kutta algorithm, while the Gompertz has a closed form for its temporal evolution. Each plot is annotated with the basic reproduction number defined as $\mathcal{R}_0 = \gamma\xi/\kappa$ [5]. The 95th percent confidence interval is shown in brackets and computed with the parametric bootstrap assuming a multivariate normal distribution on the parameters and 10000 generated bootstrap samples [103]. Fitting is done with Python-Scipy and a 6-day moving average of deaths. Observations are taken from the Github repository compiled by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, Baltimore, USA [104].

A stochastic theory for observed mortality

An alternative line of reasoning that does not rely on the framework of communicable diseases is that the environment was perturbed by an instantaneous external stressor, initiating a stress response to eventually bring mortality rates back to stability.

Inspired by De Lauro et al. (2014) [105], the stressor can be modeled as a multiplicative stochastic dampening term along with a countering force of the immune system. As with all multiplicative processes, it is convenient to log-transform mortality and work in a dimensionless space such that $F(t) = Y(t)/Y_\infty$ and $Z(t) = \ln F(t)$, from which a natural perturbation model emerges,

$$dZ(t) = -\beta Z(t)dt + \sqrt{\sigma}dB(t), \quad (16)$$

where $dB(t)$ is a delta-correlated Wiener process with zero mean, making $Z(t)$ a stochastic process. The second term on the right hand side represents the ongoing perturbation due to the stressor, while the first represents the stress response.

This perturbation model is an Ornstein–Uhlenbeck process [106], where $Z(t)$ is Gaussian. It is also interpreted as Newton’s equation of motion with friction and a random force (Langevin’s equation), and a continuous version of an Auto-Regressive(1) model [107]. As argued by De Lauro et al. (2014) [105], the diffusion coefficient, σ , represents the strength of the perturbation. This is seen if we recast our perturbation model in terms of mortality while introducing a new parameter $C = \exp(-\frac{\sigma}{2\beta})$, viz.

$$dF(t) = \left\{ \frac{\sigma}{2}F(t) - \beta F(t) \ln \left[\frac{F(t)}{C} \right] \right\} dt + \sqrt{\sigma}F(t)dB(t). \quad (17)$$

To obtain the deterministic observable, we first take the average in the log-domain (Eq. 16),

$$d \mathbb{E}[\ln F(t)] = -\beta \mathbb{E}[\ln F(t)]dt, \quad (18)$$

where $\mathbb{E}$ signifies the averaging operation. Then we use the property that for the log-normal quantity $F(t)$, the average of its logarithm is the logarithm of the *median* quantity [108], where we denote the median of $F(t)$ as $M(t)$,

$$d \ln M(t) = -\beta \ln M(t)dt, \quad (19)$$

which corresponds with the familiar deterministic Gompertz differential equation in Eq. 14 with $Y(t) = Y_\infty M(t)$. Comparing the stochastic stressor σ in Eq. 17 with the deterministic growth equation in Eq. 14 gives $\nu = \sigma/2$, suggesting that the stressor is indeed the source of growth, while β is the growth-limiting factor. This comparison is clear after taking the median of Eq. 17 wherein the cross-term of $F(t)dB(t)$ goes to zero due to the independence of $F(t)$ and $dB(t)$.

Thus, a more parsimonious interpretation of the observations not reliant on a transmission network is that mortality was caused by a perturbation of the biosphere, modeled as a random process, to which organisms gradually develop resistance at a geometric rate in the log-transformed domain [109, 110], which is the natural transformation for many processes in nature [111]. Under this model, the distribution of the abundance of $F(t)$ is log-normal, a result that can be obtained by directly solving the stochastic equation 17 [112, 108], or from thermodynamic principles [113–115]. Intuitively, this is seen by noting that the solution to the perturbation model in the log-domain (Eq. 16) is Gaussian in the variable $Z(t)$, thus suggesting a log-normal distribution of $F(t)$.

Comparing logistic and Gompertzian growth in the microscopic domain

Given that both logistic and Gompertz models describe a sigmoid saturation at the macroscopic level, what distinguishes them at the microscopic level? The answer will turn out to be the domain in which dynamics are linear, which in turn determines the correlation structure.

First, juxtapose the logistic model with the Gompertz model,

$$\dot{M} = \frac{d}{dt}M(t) = \beta M(t)(1 - M(t)) \quad \text{Logistic} \quad (20a)$$

$$\frac{d}{dt} \ln(M(t)) = -\beta \ln(M(t)) \quad \text{Gompertz}, \quad (20b)$$

where $M(t)$ is deterministic.

Note that the Gompertz model is naturally expressed as a polynomial in the log-domain $\ln M(t)$, while the logistic function is naturally expressed as a polynomial in the original domain. In the logistic model, the right-hand side is the transmission term in an SIR model, but also as an interaction term between the two macroscopic states, $M(t)$ and $1 - M(t)$.

A key point is that only Gompertz growth is closed under the geometric mean of microscopic equations, i.e. both micro- and macroscopic domains obey the same dynamics. To see this, split the system into N uncoupled microscopic abundance variables, $x_1(t), x_2(t), \dots, x_N(t)$, where the lower case x_i emphasizes the microscopic quality of the variables, and is here interpreted as probability of infection. We will later drop the assumption regarding uncoupled variables, but either case preserves closure as we shall see. Macroscopic Gompertzian growth is retained when we compute the arithmetic mean in the log-domain, with $z_i = \ln x_i$,

$$\ln M(t) = \frac{1}{N} \sum_i^N z_i(t) \quad (21)$$

$$\frac{d}{dt} z_i(t) = -\beta z_i(t) \quad (22)$$

$$\therefore \frac{d}{dt} \ln M(t) = \frac{1}{N} \sum_i \frac{d}{dt} z_i(t) = -\beta \frac{1}{N} \sum_i z_i(t) \quad (23)$$

$$\frac{d}{dt} \frac{1}{N} \sum_i z_i(t) = -\beta \ln M(t). \quad (24)$$

The same closure does not hold for logistic dynamics. To see this, we must apply the microscopic logistic model with transmission where x_j transmits a pathogen to x_i via a coupling constant w_{ij} that is assumed column-normalized such that $\sum_i w_{ij} = 1$,

$$\dot{x}_i = \beta(1 - x_i) \sum_j w_{ij} x_j, \quad (25)$$

where we dropped t for brevity.

The coupling assumption is absolutely necessary in a transmission model, but not in an environmental stressor model. So while we must include coupling with a network in a logistic model involving microscopic dynamics it is not necessary in Gompertzian dynamics.

Now, we proceed with the logistic model to average over the microscopic entities, with $\bar{x} = \frac{1}{N} \sum_i x_i$,

$$\frac{1}{N} \sum_i \dot{x}_i = \frac{\beta}{N} \sum_i (1 - x_i) \sum_j w_{ij} x_j = \frac{\beta}{N} \sum_i \sum_j w_{ij} x_j - \frac{\beta}{N} \sum_i \sum_j w_{ij} x_i x_j. \quad (26)$$

The first term is $\frac{\beta}{N} \sum_j x_j = \beta \bar{x}$. Now, remember that for closure to hold, we need the right hand side to equal $\beta \bar{x}(1 - \bar{x})$. The second term is the weighted average of pairwise products, $\frac{\beta}{N} \sum_i \sum_j w_{ij} x_i x_j$, which is $\beta \bar{x}^2$ only if either x_i and x_j are uncorrelated across the network, $\text{Cov}(x_i, x_j) = 0$, or all nodes are completely synchronized i.e. $x_i = \bar{x}$ for all i . Neither condition is compatible with a communicable disease: synchronization precludes any heterogeneity in health status, while the uncorrelated condition requires that infection status be independent of one's contacts' infection status — contradicting the very transmission mechanism on which communicable disease models are predicated. In a real epidemic, highly connected nodes are infected earlier, creating strong positive network-state correlations that violate both conditions.

As a side note, the aggregation operator is determined by the condition that the dynamics should be closed under aggregation, thus the geometric mean is not a modeler's choice but rather an operation that preserves dynamics. That logistic growth is not exact at the macroscopic level implies some noise is expected in real observations or simulations (see Results section).

Note also that the geometric mean couples at all orders in the original domain, seen by just looking at the first derivative

$$M(t) = \left[\prod_i x_i(t) \right]^{1/N} \quad (27)$$

$$\therefore \frac{d}{dx_i} M(t) = \frac{1}{N} \frac{M}{x_i}, \quad (28)$$

and so any new partial derivative with respect to another $x_j \neq x_i$, will also be a function of $M(x_1, x_2, \dots)$, implying long-range interaction. In contrast, the logistic model is only interacting maximally at the second order in the original domain.

With this phenomenological argument, the emergence of the Gompertz curve at the macroscopic level suggests that the system is correlated, or coherent, presumably as a result of the simultaneous exposure to the same underlying stressor and due to the implied log-normal nature of the microscopic entities, where multiplication replaces addition as the aggregating operator [111]. We thus show *what* distinguishes the logistic from the Gompertz model, namely the closure conditions and the correlation order, and now turn to show *how* these models connect through a continuous parameter.

Unifying microscopic logistic growth with Gompertzian growth

The Richards model (Eq. 6) unifies the macroscopic logistic and Gompertzian growth models, but by proposing a microscopic version of this model we shall see that gompertz growth at the microscopic level is the only type of growth which is coherent in the sense that a) internal fluctuations do not perturb the macroscopic mean, b) the variance of the fluctuations is decoupled from the macroscopic mean, and c) each entity contributes equally to the global growth of the system. Let us start with N nodes in the log-transformed domain, $z_i = \ln x_i$, connected by a network of edges represented by a doubly stochastic connectivity matrix, W , where the w_{ij} element is the positive coupling strength, between node i and j . A doubly stochastic matrix is one where both rows and columns are normalized and sum to 1 [116]. This condition holds exactly for regular networks; for networks with heterogeneous degree distributions, it is effectively satisfied under the mean-field approximation used in our simulations (see Methods). With these assumptions, the microscopic model for growth becomes

$$dz_i = \left[\frac{\beta}{\theta} (1 - e^{\theta z_i}) + \beta(\bar{z}_{\theta,i} - z_i) \right] dt + \sqrt{\sigma} dB_i(t) \quad (29)$$

where $\bar{z}_{\theta,i} = \frac{1}{\theta} \ln(\sum_j w_{ij} e^{\theta z_j})$ is the order- θ power mean of neighbors' states. The parameter θ smoothly interpolates between Gompertz ($\theta \rightarrow 0$) and logistic ($\theta = 1$) dynamics, with intermediate values recovering the Richards growth family. The noise, $dB_i(t)$ is assumed to be independent normally distributed, which is interpreted as the environmental stressor having a unique response on each individual. Note that for perfectly synchronized nodes, with $z_i = \bar{z}$, i , Equation 29 reduces to our single-entity model (Eq. 16).

First, we see that in the unique case of Gompertz, i.e. $\theta \rightarrow 0$, the dynamics are linear in the log-space and microscopic fluctuations are decoupled from the macroscopic state. This is seen by noting the average macroscopic state reduces to $d\bar{z}/dt = -\beta\bar{z}$, where $\bar{z}$ is the arithmetic weighted mean and we have used the doubly stochastic property of the adjacency matrix, $\frac{1}{N} \sum_i \sum_j w_{ij} z_j = \bar{z}$. Both the second and third terms in Eq. 29 go to zero. Due to the linearity of the Gompertz case, microscopic fluctuations are decoupled from the macroscopic state.

Secondly, any microscopic fluctuations revert to the mean, producing a stable evolution independent on the fluctuation profile. We see this by first defining a set of zero-mean fluctuations to be

$$\delta_i = z_i - \bar{z} \quad (30)$$

then,

$$\frac{d\delta_i}{dt} = \frac{dz_i}{dt} - \frac{d\bar{z}}{dt} \quad (31)$$

$$= -2\beta(\bar{z} + \delta_i) + \beta \sum_j w_{ij}(\bar{z} + \delta_j) + \beta\bar{z} \quad (32)$$

$$= -2\beta\bar{z} - 2\beta\delta_i + \beta\bar{z} \sum_j w_{ij} + \beta \sum_j w_{ij}\delta_j + \beta\bar{z} \quad (33)$$

$$= -2\beta\delta_i + \beta \sum_j w_{ij}\delta_j, \quad (34)$$

where we have used the assumption that columns of w_{ij} sum to 1. The result is a linear system $\dot{\delta} = \beta(-2I + W)\delta$. The eigenvalues of W for a row-normalized matrix lie in $[-1, 1]$, so the eigenvalues of $-2I + W$ lie in $[-3, -1]$. All eigenvalues are strictly negative, meaning every deviation mode decays exponentially with rate at least β . Thus, one could say the system is unconditionally stable, where deviations shrink regardless of their initial magnitude.

Combining the first and the second point we see that the gompertz model does not require synchronicity, but rather imposes it. One could call this a characteristic of a coherent system.

Thirdly, the gompertz model's mean and variance are decoupled, which is crucially not the case for any other model in the Richard spectrum where $\theta > 0$. To see this, first recognize that the variance of z_i is $\sum_j w_{ij} \delta_j^2$, our zero-mean random fluctuations. Then, by Taylor expanding the order- θ mean around $\theta = 0$, and using the property that δ_i has mean of 0,

$$\bar{z}_{\theta,i} = \frac{1}{\theta} \ln \left(\sum_j w_{ij} e^{\theta z_j} \right) = \frac{1}{\theta} \ln \left(e^{\theta \bar{z}} \sum_j w_{ij} e^{\theta \delta_j} \right) = \bar{z} + \frac{1}{\theta} \ln \left(\sum_j w_{ij} (1 + \theta \delta_j + \theta^2 \delta_j^2 / 2 + O(\theta^3)) \right) \quad (35)$$

$$\bar{z}_{\theta,i} = \bar{z} + \frac{\theta}{2} \text{Var}(z_i) + O(\theta^2), \quad (36)$$

and we immediately see that the mean and the variance are uncoupled only when θ goes to 0. Thus, for the general- θ Richard's model, only if all variables are in complete synchronicity, $z_i = \bar{z}$, will the microscopic model reduce to the macroscopic model, and if $\theta = 0$, the single entity Gompertz model in Eq. 17. Overall, the macroscopic Richards model is *conditioned* on synchronicity, while the Gompertz model *imposes* synchronicity in the deterministic case ($\sigma = 0$) and maintains exact macroscopic reduction despite residual fluctuations in the stochastic case, with $\text{Var}(\bar{z}) = \frac{\sigma}{4\beta N}$. This is due to each δ_i having variance $\sigma/4\beta$ and being weakly correlated for large N , so that when the variance of the mean is computed the answer is as shown.

Lastly, as an illustration that each entity contributes equally to the growth in the Gompertz model, we will present a simple example. Consider a two-variable system at a particular point in time where they are at unequal abundance, e.g. $x_1 = 0.01$ and $x_2 = 0.1$. In this state, the geometric mean is $\sqrt{0.01 \cdot 0.1} \approx 0.032$, while $\bar{z} = (\ln 0.01 + \ln 0.1)/2 \approx -3.45$ and we see equidistance in additive terms in the log-domain and in multiplicative terms in the original domain; $z_1 - \bar{z} = \bar{z} - z_2$ and $x_1/e^{\bar{z}} = e^{\bar{z}}/x_2$. This implies that the second term in Eq. 29, the collective forcing term, has the same magnitude for both variables. The same could not be said in the logistic growth model, where the mean would be the arithmetic mean of the two states, 0.055, and closer to the higher value.

To show this generally, we just have to show that this collective growth term, $\beta(\bar{z}_{\theta \rightarrow 0,i} - z_i)$ is independent of $z_i \forall i$ and only a function of the fluctuations, δ_i . To this end, we use the fluctuation from the mean $z_i = \bar{z} + \delta_i$, viz.

$$\begin{aligned} \bar{z}_{\theta \rightarrow 0,i} - z_i &= \lim_{\theta \rightarrow 0} \frac{1}{\theta} \sum_j w_{ij} e^{\theta z_j} - z_i \\ &= \sum_j w_{ij} z_j - z_i \\ &= \sum_j w_{ij} (\bar{z} + \delta_j) - (\bar{z} + \delta_i) \\ &= \sum_j w_{ij} \delta_j - \delta_i \end{aligned}$$

where the first right-hand-side term vanishes in expectation in the large- N limit, thus reducing the collective growth term to

$$\lim_{N \rightarrow \infty, \theta \rightarrow 0} \bar{z}_{\theta \rightarrow 0,i} - z_i = -\delta_i, \quad (37)$$

which completes the argument.

In contrast, logistic growth ($\theta = 1$) has a coupling term $\bar{z}_{1,i} = \ln(\sum_j w_{ij} e^{z_j})$ which depends on all neighbours and any node $z_i \gg \bar{z}$ will dominate the exponential sum.

Thus, in logistic growth the outliers drive the growth of the system, while in Gompertz growth, growth is egalitarian, and the forcing magnitude is the same for nodes equidistant from the mean regardless of which specific neighbors they have.

Nevertheless, the Gompertz model is correlated at all orders in the natural domain. This is seen because the dynamics are linear in the log-domain, producing a jointly Gaussian distribution there, and hence fully categorized with second order correlations, which is one of the key properties of Gaussian distributions. The geometric mean then maps this second-order structure through the exponential transform into an infinitely-correlated original abundance domain, because the log-normal distribution has non-zero cumulants at all orders, as opposed to the Gaussian distribution. Thus, infinitely correlated growth is a natural consequence of macroscopic Gompertzian dynamics, which we here choose to call *coherent growth* in accordance with previous work [111, 108, 117].

By contrast, mean-field approximated logistic growth does not have linear dynamics in the log-transformed domain, but through the arithmetic mean in the natural domain of x_i , has second order correlations only, and would thus be an ideal candidate for a communicable disease model which is completely describable through peer-to-peer connections, with the caveat of its limitations when using this mean-field macroscopic approximation.

Simulations and Results

Simulated growth scenarios for a variety of Richards "coherence" θ parameter all show close correspondence to their theoretical counterparts, except at high values of θ where the mean-variance coupling identified in Eq. 31 produces visible deviations from theory, particularly in the late stages where the double-log transform amplifies residual fluctuations near saturation (Figure 2). The numerical component of this deviation can be mitigated with adaptive sub-stepping, but the physical component — arising from the mean-variance coupling — persists and is left visible to illustrate this structural difference between the Gompertz and logistic regimes. We do not see a large difference when using different networks in the simulations, consistent with the mean-field approximation described in the Methods section, under which the network topology enters only through degree-weighted averages (Figure 3).




Figure 2: Different simulated Richard growth models are compared with their theoretical curves, using a Log-Normal network with 10,000 nodes and mean-field approximation. Note that the higher values of θ makes the growth unstable as discussed in the text.




Figure 3: Comparison of relative growth rates for different networks under a microscopic Gompertz mode with 10,000 nodes using a mean field approximation, showing fidelity to the theoretical limit regardless of the underlying network. Different networks plotted in color, showing minimal deviation from the theoretical gompertz curve. A logistic and Richards $\theta = 0.3$ theoretical curve is plotted for comparison. The abundance variable is normalized to values between 0 and 1.

Conclusion

In conclusion, we have shown that Gompertzian growth is a maximally coherent growth process, though a mathematical parameter θ , with specific mathematical properties: 1. closure under aggregation, 2. stability in spite of internal

fluctuation, 3. mean-variance uncoupling, and 4. egalitarian forcing with equal contributions to the relative growth. All of these properties shed new light on infinitely correlated and coherent dynamics exhibited at the macroscopic level.

We can now further appreciate the Richards parametrization as a transition from non-collaborative logistic growth to maximally collaborative Gompertz growth as $\theta \rightarrow 0$. This feature of θ was also obtained by Petroni et al. (2020) [93] by interpreting the θ -logistic growth rate in Eq. 6 as non-linear resource availability dependent on the overall abundance, with linearity reached at $\theta = 1$, without resorting to microscopic modelling. Coherence was there defined as maximally synergistic or collaborating elementary components. Used in a mortality model, the maximally coherent scenario implies that all units in the network are exposed to the stressor simultaneously. Molski and Konarski (2003) [117] made a similar observation that Gompertzian growth is the coherent state in a quantum mechanical system with a time-dependent potential, an interpretation which sheds further light on the temporal nature of the postulated stress response. This quantum mechanical system has also been used to describe coherent energy states of diatomic molecules in space [118]. There is also an uncanny analogy between the mean-variance decoupling of the Gompertz at $\theta \rightarrow 0$ and symmetry-breaking dynamics in various physical systems. The spectral gap of the deviation dynamics $\dot{\delta} = \beta(-2I + W)\delta$ plays an analogous role to the energy gap in BCS superconductivity — both protect the macroscopic state from microscopic perturbations through a structural property of the dynamics rather than through fine-tuning of parameters. In the field of quantum physics, *coherence* is a well-defined mathematical property first explored by Glauber (1963) [119] in the context of electromagnetic fields. The fact that stable macroscopic dynamics emerge from the microscopic scales only from the Gompertz model compared to other Richards and logistic models is also noteworthy.

Observing Gompertzian growth through the communicable diseases paradigm, the perceived pathogen exposes the population simultaneously, precluding finite-speed propagation, exemplified by the immediate decrease in the relative growth rate of the infected. We are thus forced to reject the concept of a communicable pathogen in the context of Gompertzian growth.

We further show that the observed mortality across countries can be explained by a model where the biosphere is disturbed by a ubiquitous, simultaneous, and non-local stressor eliciting a corresponding stress response through which a new stress tolerance baseline is gradually established. The stressor is modeled as a stochastic perturbation in the log-transformed domain, where each entity experiences its own stress response which is multiplicative and not additive in the natural domain. From this model, we draw parallels between the coherent behavior of the population’s mortality evolution during an epidemic and the spatial coherence of quantum mechanical systems.

Thus, we see growth on a spectrum: In one extreme we find non-collaborative growth models or models with parameterized linear interaction effects, and in the other extreme we see a field of microscopic entities coherently sharing information as if exposed to the same environmental stressor. This observation adds to the list of emerging biological phenomena that exhibit coherent quantum behavior at the macroscopic level [120]. Although a global biosphere disturbance can explain our observations, one might still be surprised to find that temporal evolution of human mortality during epidemics can behave like the spatial energy distribution of quantum coherent systems.

Methods

Estimating model parameters for Covid data

The time-evolution of the SIRD and the Gompertz model are obtained with a Runge-Kutta algorithm, after first fitting parameters using non-linear least squares according to equations 14 and 11 in the text (see SI Appendix for details). Fitting is done using Python-Scipy’s non-linear least squares algorithm with specified Jacobians and a 6-day windowed average of observed deaths, similar to the technique by Carletti et al. [5]. Observations are taken from the Github repository compiled by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, Baltimore, USA [104].

The basic reproduction number is modeled as the product of $\eta\xi/\kappa$ in Eq. 11, and its associated variance is obtained through a parametric bootstrap assuming a multivariate normal distribution with mean and covariance equal to the estimate from the non-linear least squares technique. The 95th percent confidence interval is obtained after 10000 parameter samples drawn from this distribution, each computing a resampled estimate of $\mathcal{R}_0$ [103].

Running networked growth model simulations

The simulations of growth in a theoretical network structure are done using Equation 29 with a mean-field approximation, where each node’s local neighbor-average is replaced by a degree-weighted global average. Specifically, the coupling target $\bar{z}_{\theta,i} = \frac{1}{\theta} \ln(\sum_j w_{ij} e^{\theta z_j})$ is approximated as $\bar{z}_{\theta} = \frac{1}{\theta} \ln(\sum_j w_j e^{\theta z_j})$ with weights $w_j = k_j / \sum_k k_k$, where k_j is

the degree of node j . This approximation is standard for configuration-model random graphs and becomes exact for large N when the correlation between node state z_j and degree k_j is weak.

We simulate $N = 10,000$ nodes with parameters $\beta = 0.25$, $\sigma = 0.001$, and time step $\Delta t = 0.02$. Initial conditions are drawn as $z_i \sim \mathcal{N}(-13.8, 0.03^2)$, corresponding to initial abundance $x_i \approx 10^{-6}$. Integration is performed using the Euler-Maruyama scheme. For θ near 1, mean-variance coupling will create some noise in the later stages of the simulations.

The node degrees are drawn from the following distributions, each representing a distinct network topology:

- **Scale-free:** $P(k) \propto k^{-3}$, $k_{\min} = 2$, generated via inverse CDF sampling, approximating a Barabási-Albert network.
- **Power law:** $P(k) \propto k^{-2.2}$, $k_{\min} = 2$, a heavier-tailed variant generated via inverse CDF sampling.
- **Log-normal:** $\ln k \sim \mathcal{N}(1.5, 0.8^2)$, $k_{\min} = 2$.
- **Weibull:** $k \sim \text{Weibull}(\text{shape} = 1.5, \text{scale} = 5)$, $k_{\min} = 2$.
- **Exponential:** $k \sim \text{Exp}(\text{scale} = 4)$, $k_{\min} = 2$.
- **Erdős-Rényi:** $k \sim \text{Poisson}(\lambda = 6)$, $k_{\min} = 2$.

For the Richards interpolation experiments, the coherence parameter θ is varied in $\{0, 0.1, 0.3, 0.5, 0.7, 1.0\}$ on a single network (log-normal degree distribution). Since logistic-like dynamics ($\theta \rightarrow 1$) exhibit slower growth from low initial abundance, the simulation duration is extended as $T_{\max} = 40 \cdot \max(1, 1 + 4\theta)$ to ensure all trajectories reach saturation. The macroscopic observable is the geometric mean $M(t) = \exp(\frac{1}{N} \sum_i z_i)$, and the relative growth rate is computed numerically as $\gamma(t) = \dot{M}(t)/M(t)$ using a central difference scheme for $\dot{M}$.

Code Availability

Scripts used to produce the plots are downloadable in the form of a Jupyter notebook using Python from https://nbviewer.org/urls/emf-research.fra1.digitaloceanspaces.com/gompertz/gompertz_vs_sird.ipynb and <https://nbviewer.org/urls/emf-research.fra1.digitaloceanspaces.com/gompertz/hypergeometric.ipynb>

Data Availability

Observations are taken from the Github repository compiled by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, Baltimore, USA [104], and are available at <https://github.com/CSSEGISandData/COVID-19>, specifically the table `csse_covid_19_data/csse_covid_19_time_series/time_series_covid19_deaths_global.csv`, downloaded Aug. 5th 2022.

Competing interests

The authors declare no competing interests.

Supplement




Figure 4: Cumulative diseased plotted against number of days after $Y(t)/Y_{max} > 0.005$, comparing an SIRD with a Gompertz model. Both models are fit using non-linear least squares according to equations (14) and (11) in the text. Fitting is done with Python-Scipy and a 6-day windowed average of deaths. The basic reproduction number, R_0 is computed as specified in the Methods. Observations are taken from the Github repository compiled by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University, Baltimore, USA [104].

References

- [1] Harko, T., Lobo, F. S., and Mak, M. Exact analytical solutions of the susceptible-infected-recovered (sir) epidemic model and of the sir model with equal death and birth rates. *Applied Mathematics and Computation*, 236:184–194, 2014.
- [2] Kröger, M. and Schlickeiser, R. Analytical solution of the SIR-model for the temporal evolution of epidemics. part a: time-independent reproduction factor. *Journal of Physics A: Mathematical and Theoretical*, 53(50): 505601, 2020.
- [3] Schlickeiser, R. and Kröger, M. Analytical solution of the SIR-model for the temporal evolution of epidemics: part b. semi-time case. *Journal of Physics A: Mathematical and Theoretical*, 54(17):175601, 2021.
- [4] Heng, K. and Althaus, C. L. The approximately universal shapes of epidemic curves in the susceptible-exposed-infectious-recovered (SEIR) model. *Scientific Reports*, 10(1):1–6, 2020.
- [5] Carletti, T., Fanelli, D., and Piazza, F. COVID-19: The unreasonable effectiveness of simple models. *Chaos, Solitons & Fractals*: X, 5:100034, 2020.
- [6] Cooper, I., Mondal, A., and Antonopoulos, C. G. A SIR model assumption for the spread of COVID-19 in different communities. *Chaos, Solitons & Fractals*, 139:110057, 2020.

Table 1: Estimated values for the SIRD model fitted to various countries' observations as shown in Figure 1 using notation from eq. (12). Standard deviation estimates is given in the parenthesis corresponding to the last significant digit of the reported number.

	$\eta \times 10^{-1}$	$\xi \times 10^4$	$\kappa \times 10^2$
Belgium	94(8)	3.2(2)	9.3(7)
Portugal	6.6(4)	35(2)	4.5(3)
Austria	6.9(5)	40(3)	8.7(6)
Switzerland	22(1)	13.6(7)	11.0(5)
Sweden	19(5)	8(2)	3(1)
Italy	137(3)	1.77(5)	3.71(1)
France	240(10)	1.24(7)	8.1(4)
Spain	145(2)	2.27(5)	4.75(8)
Germany	61(4)	3.9(3)	6.5(5)
Denmark	2.7(2)	96(8)	4.3(3)
United Kingdom	181(5)	1.39(5)	4.3(1)
Netherlands	29(1)	8.9(5)	4.3(2)
Turkey	34(3)	6.8(5)	6.9(6)
Iran	22.3(4)	10.4(4)	2.51(7)
US	312(5)	0.76(2)	1.83(7)
China	130(30)	3.5(5)	26(4)

Table 2: Estimated values for the Gompertz model fitted to various countries' observations, as shown in Figure 1 using notation from eq. (1).

	$\nu \times 10$	$\beta \times 10^2$	$\tilde{Y} \times 10^2$
Belgium	9.3(1)	8.1(2)	10.1(9)
Portugal	6.15(3)	6.22(7)	7.0(3)
Austria	6.98(5)	7.6(1)	7.8(4)
Switzerland	8.08(6)	8.1(1)	8.9(5)
Sweden	4.9(1)	4.3(2)	6(1)
Italy	7.60(6)	5.88(7)	8.4(4)
France	10.2(1)	8.2(1)	11.0(8)
Spain	9.41(9)	7.5(1)	10.2(7)
Germany	7.56(6)	6.53(8)	8.4(4)
Denmark	5.94(7)	6.5(2)	6.8(7)
United Kingdom	8.46(5)	6.54(7)	9.3(4)
Netherlands	7.34(6)	6.52(8)	8.2(4)
Turkey	7.04(5)	6.38(8)	7.9(4)
Iran	5.87(6)	5.04(8)	6.7(4)
US	7.53(7)	5.32(7)	8.4(5)
China	9.8(1)	9.4(2)	10(1)

- [7] Postnikov, E. B. Estimation of COVID-19 dynamics “on a back-of-envelope”: Does the simplest SIR model provide quantitative parameters and predictions? *Chaos, Solitons & Fractals*, 135:109841, 2020.
- [8] Muñoz-Fernández, G. A., Seoane, J. M., and Seoane-Sepúlveda, J. B. A SIR-type model describing the successive waves of COVID-19. *Chaos, Solitons & Fractals*, 144:110682, 2021.
- [9] Cooper, I., Mondal, A., Antonopoulos, C. G., and Mishra, A. Dynamical analysis of the infection status in diverse communities due to COVID-19 using a modified SIR model. *Nonlinear Dynamics*, pages 1–14, 2022.
- [10] Saikia, D., Bora, K., and Bora, M. P. COVID-19 outbreak in india: an SEIR model-based analysis. *Nonlinear Dynamics*, 104(4):4727–4751, 2021.
- [11] Gompertz, B. XXIV. on the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. in a letter to francis baily, esq. f. r. s. &c. *Philosophical Transactions of the Royal Society of London*, 115:513–583, dec 1825. doi:10.1098/rstl.1825.0026.
- [12] Bajzer, Ž., Vuk-Pavlović, S., and Huzak, M. Mathematical modeling of tumor growth kinetics. In *A survey of models for tumor-immune system dynamics*, pages 89–133. Springer, 1997.
- [13] Ohnishi, A., Namekawa, Y., and Fukui, T. Universality in COVID-19 spread in view of the gompertz function. *Progress of Theoretical and Experimental Physics*, 2020(12), oct 2020. doi:10.1093/ptep/ptaa148.
- [14] Rypdal, K. and Rypdal, M. A parsimonious description and cross-country analysis of COVID-19 epidemic curves. *International Journal of Environmental Research and Public Health*, 17(18):6487, sep 2020. doi:10.3390/ijerph17186487.
- [15] Català, M., Alonso, S., Alvarez-Lacalle, E., López, D., Cardona, P.-J., and Prats, C. Empirical model for short-time prediction of COVID-19 spreading. *PLOS Computational Biology*, 16(12):e1008431, dec 2020. doi:10.1371/journal.pcbi.1008431.
- [16] Rodrigues, T. and Helene, O. Monte carlo approach to model COVID-19 deaths and infections using gompertz functions. *Physical Review Research*, 2(4):043381, 2020.
- [17] Levitt, M., Scaiewicz, A., and Zonta, F. Predicting the trajectory of any COVID19 epidemic from the best straight line. *medRxiv*, jun 2020. doi:10.1101/2020.06.26.20140814.
- [18] Castro, M., Ares, S., Cuesta, J. A., and Manrubia, S. The turning point and end of an expanding epidemic cannot be precisely forecast. *Proceedings of the National Academy of Sciences*, 117(42):26190–26196, 2020.
- [19] Hamidi, S., Sabouri, S., and Ewing, R. Does density aggravate the COVID-19 pandemic? *Journal of the American Planning Association*, 86(4):495–509, June 2020. doi:10.1080/01944363.2020.1777891.
- [20] Hamidi, S., Ewing, R., and Sabouri, S. Longitudinal analyses of the relationship between development density and the COVID-19 morbidity and mortality rates: Early evidence from 1,165 metropolitan counties in the united states. *Health & Place*, 64:102378, July 2020. doi:10.1016/j.healthplace.2020.102378.
- [21] Carozzi, F. Urban density and covid-19. *SSRN Electronic Journal*, 2020. doi:10.2139/ssrn.3643204.
- [22] Liu, L. Emerging study on the transmission of the novel coronavirus (covid-19) from urban perspective: Evidence from china. *Cities*, 103:102759, 2020.
- [23] A.G., G., S., C., M.R., M., J.C., V., G., S., G.W., C., and J.B., S. Covid-19 mortality rates in the european union, switzerland, and the uk: effect of timeliness, lockdown rigidity, and population density. *Minerva Med.*, 111(4), 2020.
- [24] Khavarian-Garmsir, A. R., Sharifi, A., and Moradpour, N. Are high-density districts more vulnerable to the COVID-19 pandemic? *Sustainable Cities and Society*, 70:102911, 2021.
- [25] Barak, N., Sommer, U., and Mualam, N. Urban attributes and the spread of COVID-19: The effects of density, compliance and socio-political factors in israel. *Science of the Total Environment*, 793:148626, 2021.
- [26] Arpino, B., Bordone, V., and Pasqualini, M. No clear association emerges between intergenerational relationships and COVID-19 fatality rates from macro-level analyses. *Proceedings of the National Academy of Sciences*, 117(32):19116–19121, jul 2020. doi:10.1073/pnas.2008581117.
- [27] Fischer, L., Peters, M., Merbach, S., Eydner, M., Kuczka, A., Lambertz, J., Kummerfeld, M., Kahnt, K., Weiss, A., and Petersen, H. Increased mortality in wild tits in north rhine-westphalia (germany) in 2020 with a special focus on suttonella ornithocola and other infectious pathogens. *European Journal of Wildlife Research*, 67(3), may 2021. doi:10.1007/s10344-021-01500-7.
- [28] Perisé-Barrios, A. J., Tomeo-Martín, B. D., Gómez-Ochoa, P., Delgado-Bonet, P., Plaza, P., Palau-Concejo, P., González, J., Ortiz-Díez, G., Meléndez-Lazo, A., Gentil, M., García-Castro, J., and Barbero-Fernández, A. Humoral responses to SARS-CoV-2 by healthy and sick dogs during the COVID-19 pandemic in spain. *Veterinary Research*, 52(1), feb 2021. doi:10.1186/s13567-021-00897-y.

- [29] Duff, P., Fenemore, C., Holmes, P., Hopkins, B., Jones, J., He, M., Everest, D., and Rocchi, M. Disease surveillance in england and wales, december 2019. *Veterinary Record*, 186(1):15–16, Jan. 2020. doi:10.1136/vr.m68.
- [30] Hu, B., Wei, H., Fan, Z., Song, Y., Chen, M., Qiu, R., Zhu, W., Xu, W., Xue, J., and Wang, F. Emergence of rabbit haemorrhagic disease virus 2 in china in 2020. *Veterinary Medicine and Science*, 7(1):236–239, Aug. 2020. doi:10.1002/vms3.332.
- [31] Fukui, H., Shimoda, H., Kadekaru, S., Henmi, C., and Une, Y. Rabbit hemorrhagic disease virus type 2 epidemic in a rabbit colony in japan. *Journal of Veterinary Medical Science*, 83(5):841–845, 2021. doi:10.1292/jvms.21-0007.
- [32] van Aarde, R. J., Pimm, S. L., Guldemon, R., Huang, R., and Maré, C. The 2020 elephant die-off in botswana. *PeerJ*, 9:e10686, jan 2021. doi:10.7717/peerj.10686.
- [33] Kambayashi, Y., Bannai, H., Tsujimura, K., Hiram, A., Ohta, M., and Nemoto, M. Outbreak of equine coronavirus infection among riding horses in tokyo, japan. *Comparative Immunology, Microbiology and Infectious Diseases*, 77:101668, Aug. 2021. doi:10.1016/j.cimid.2021.101668.
- [34] King, S., Rajko-Nenow, P., Ashby, M., Frost, L., Carpenter, S., and Batten, C. Outbreak of african horse sickness in thailand, 2020. *Transboundary and Emerging Diseases*, jul 2020. doi:10.1111/tbed.13701.
- [35] Lu, G., Pan, J., Ou, J., Shao, R., Hu, X., Wang, C., and Li, S. African horse sickness: Its emergence in thailand and potential threat to other asian countries. *Transboundary and Emerging Diseases*, jun 2020. doi:10.1111/tbed.13625.
- [36] Castillo-Olivares, J. African horse sickness in thailand: Challenges of controlling an outbreak by vaccination. *Equine Veterinary Journal*, 53(1):9–14, oct 2020. doi:10.1111/evj.13353.
- [37] Cannon, P. and Rycroft, M. J. Schumann resonance frequency variations during sudden ionospheric disturbances. *Journal of Atmospheric and Terrestrial Physics*, 44(2):201–206, 1982.
- [38] Vieira, C. L. Z., Chen, K., Garshick, E., Liu, M., Vokonas, P., Ljungman, P., Schwartz, J., and Koutrakis, P. Geomagnetic disturbances reduce heart rate variability in the normative aging study. *Science of The Total Environment*, page 156235, 2022.
- [39] Alabdulgader, A., McCraty, R., Atkinson, M., Dobyns, Y., Vainoras, A., Ragulskis, M., and Stolc, V. Long-term study of heart rate variability responses to changes in the solar and geomagnetic environment. *Scientific Reports*, 8(1), feb 2018. doi:10.1038/s41598-018-20932-x.
- [40] Ghione, S., Mezzasalma, L., Seppia, C., and Papi, F. Do geomagnetic disturbances of solar origin affect arterial blood pressure? *Journal of human hypertension*, 12(11):749–754, 1998.
- [41] Chernouss, S., Vinogradov, A., and Vlassova, E. Geophysical hazard for human health in the circumpolar auroral belt: evidence of a relationship between heart rate variation and electromagnetic disturbances. *Natural hazards*, 23(2):121–135, 2001.
- [42] Stoupel, E. Ambulatory blood pressure monitoring in patients with hypertension on days of high and low geomagnetic activity. *J Hum Hypertens*, 9:293–294, 1995.
- [43] Stoupel, E., Kalediene, R., Petrauskiene, J., Starkuviene, S., Abramson, E., Israelevich, P., Sulkes, J., et al. Twenty years study of solar, geomagnetic, cosmic ray activity links with monthly deaths number (n-850304). *Journal of Biomedical Science and Engineering*, 4(06):426, 2011.
- [44] Malin, S. and Srivastava, B. Correlation between heart attacks and magnetic activity. *Nature*, 277(5698):646–648, 1979.
- [45] Stoupel, E. Sudden cardiac deaths and ventricular extrasystoles on days with four levels of geomagnetic activity. *Journal of basic and clinical physiology and pharmacology*, 4(4):357–366, 1993.
- [46] Burch, J., Reif, J., and Yost, M. Geomagnetic disturbances are associated with reduced nocturnal excretion of a melatonin metabolite in humans. *Neuroscience Letters*, 266(3):209–212, 1999.
- [47] Rapoport, S., Malinovskaia, N., Oraevskii, V., Komarov, F., Nosovskii, A., and Vetterberg, L. Effects of disturbances of natural magnetic field of the earth on melatonin production in patients with coronary heart disease. *Klinicheskaya Meditsina*, 75(6):24–26, 1997.
- [48] Bergiannaki, J.-D., Paparrigopoulos, T., and Stefanis, C. N. Seasonal pattern of melatonin excretion in humans: relationship to daylength variation rate and geomagnetic field fluctuations. *Experientia*, 52(3):253–258, 1996.
- [49] Schiff, J. E., Vieira, C. L., Garshick, E., Wang, V., Blomberg, A., Gold, D. R., Schwartz, J., Tracy, S. M., Vokonas, P., and Koutrakis, P. The role of solar and geomagnetic activity in endothelial activation and inflammation in the nas cohort. *PloS one*, 17(7):e0268700, 2022.

- [50] Keeton, W. T., Larkin, T. S., and Windsor, D. M. Normal fluctuations in the earth's magnetic field influence pigeon orientation. *Journal of comparative physiology*, 95(2):95–103, 1974.
- [51] Schreiber, B. and Rossi, O. Correlation between race arrivals of homing pigeons and solar activity. *Italian Journal of Zoology*, 43(3):317–320, 1976.
- [52] Kowalski, U., Wiltchko, R., and Füller, E. Normal fluctuations of the geomagnetic field may affect initial orientation in pigeons. *Journal of Comparative Physiology A*, 163(5):593–600, 1988.
- [53] Bianco, G., Ilieva, M., and Åkesson, S. Magnetic storms disrupt nocturnal migratory activity in songbirds. *Biology letters*, 15(3):20180918, 2019.
- [54] Ferrari, T. E. Cetacean beachings correlate with geomagnetic disturbances in earth's magnetosphere: an example of how astronomical changes impact the future of life. *International Journal of Astrobiology*, 16(2):163–175, 2017.
- [55] Vanselow, K. H., Jacobsen, S., Hall, C., and Garthe, S. Solar storms may trigger sperm whale strandings: explanation approaches for multiple strandings in the north sea in 2016. *International Journal of Astrobiology*, 17(4):336–344, 2018.
- [56] Granger, J., Walkowicz, L., Fitak, R., and Johnsen, S. Gray whales strand more often on days with increased levels of atmospheric radio-frequency noise. *Current Biology*, 30(4):R155–R156, 2020.
- [57] Zenchenko, T. A. and Breus, T. K. The possible effect of space weather factors on various physiological systems of the human organism. *Atmosphere*, 12(3):346, 2021.
- [58] Zilli Vieira, C. L., Alvares, D., Blomberg, A., Schwartz, J., Coull, B., Huang, S., and Koutrakis, P. Geomagnetic disturbances driven by solar activity enhance total and cardiovascular mortality risk in 263 us cities. *Environmental Health*, 18(1):1–10, 2019.
- [59] Gordon, C. and Berk, M. The effect of geomagnetic storms on suicide. *South African Psychiatry Review*, 6(3): 24–27, 2003.
- [60] Kay, R. W. Geomagnetic storms: association with incidence of depression as measured by hospital admission. *The British Journal of Psychiatry*, 164(3):403–409, 1994.
- [61] Kay, R. W. Schizophrenia and season of birth: relationship to geomagnetic storms. *Schizophrenia Research*, 66(1):7–20, 2004.
- [62] Halberg, F., Cornélissen, G., Panksepp, J., Otsuka, K., and Johnson, D. Chronomics of autism and suicide. *Biomedicine & pharmacotherapy*, 59:S100–S108, 2005.
- [63] Berk, M., Dodd, S., and Henry, M. Do ambient electromagnetic fields affect behaviour? a demonstration of the relationship between geomagnetic storm activity and suicide. *Bioelectromagnetics: Journal of the Bioelectromagnetics Society, The Society for Physical Regulation in Biology and Medicine, The European Bioelectromagnetics Association*, 27(2):151–155, 2006.
- [64] Doronin, V., Parfentëv, V., SZh, T., Namvar, R., Somsikov, V., Drobzhev, V., and Chemeris, A. Effect of variations of the geomagnetic field and solar activity on human physiological indicators. *Biofizika*, 43(4):647–653, 1998.
- [65] Persinger, M. and Psych, C. Sudden unexpected death in epileptics following sudden, intense, increases in geomagnetic activity: prevalence of effect and potential mechanisms. *International Journal of Biometeorology*, 38(4):180–187, 1995.
- [66] Cherry, N. Schumann resonances, a plausible biophysical mechanism for the human health effects of solar. *Natural hazards*, 26(3):279–331, 2002.
- [67] Schumann, W. O. Über die strahlungslosen eigenschwingungen einer leitenden kugel, die von einer luftschicht und einer ionosphärenhülle umgeben ist. *Zeitschrift für Naturforschung A*, 7(2):149–154, 1952.
- [68] Wever, R. ELF-effects on human circadian rhythms. In *ELF and VLF electromagnetic field effects*, pages 101–144. Springer, 1974.
- [69] Pobachenko, S., Kolesnik, A., Borodin, A., and Kalyuzhin, V. The contingency of parameters of human encephalograms and schumann resonance electromagnetic fields revealed in monitoring studies. *Biophysics*, 51(3):480–483, 2006.
- [70] Saroka, K. S. and Persinger, M. A. Quantitative evidence for direct effects between earth-ionosphere schumann resonances and human cerebral cortical activity. *International Letters of Chemistry, Physics and Astronomy*, 20, 2014.

- [71] Persinger, M. A., Saroka, K. S., et al. Human quantitative electroencephalographic and schumann resonance exhibit real-time coherence of spectral power densities: implications for interactive information processing. *Journal of Signal and Information Processing*, 6(02):153, 2015.
- [72] König, H. L., Krueger, A. P., Lang, S., and Sönning, W. *Biologic effects of environmental electromagnetism*. Springer Science & Business Media, 2012.
- [73] Dyrda, M., Kulak, A., Mlynarczyk, J., and Ostrowski, M. Novel analysis of a sudden ionospheric disturbance using schumann resonance measurements. *Journal of Geophysical Research: Space Physics*, 120(3):2255–2262, 2015.
- [74] Roldugin, V., Maltsev, Y. P., Vasiljev, A., Schokotov, A., and Belyajev, G. Schumann resonance frequency increase during solar x-ray bursts. *Journal of Geophysical Research: Space Physics*, 109(A1), 2004.
- [75] Sători, G., Williams, E., and Mushtak, V. Response of the earth–ionosphere cavity resonator to the 11-year solar cycle in x-radiation. *Journal of Atmospheric and Solar-Terrestrial Physics*, 67(6):553–562, 2005.
- [76] Sători, G., Williams, E., Price, C., Boldi, R., Koloskov, A., Yampolski, Y., Guha, A., and Barta, V. Effects of energetic solar emissions on the earth–ionosphere cavity of schumann resonances. *Surveys in Geophysics*, 37(4): 757–789, 2016.
- [77] Shvets, A., Nickolaenko, A., and Chebrov, V. Effect of solar flares on the schumann-resonance frequencies. *Radiophysics and Quantum Electronics*, 60(3):186–199, 2017.
- [78] Schlegel, K. and Füllekrug, M. Schumann resonance parameter changes during high-energy particle precipitation. *Journal of Geophysical Research: Space Physics*, 104(A5):10111–10118, 1999.
- [79] Singh, B., Tyagi, R., Hobara, Y., and Hayakawa, M. X-rays and solar proton event induced changes in the first mode schumann resonance frequency observed at a low latitude station agra, india. *Journal of Atmospheric and Solar-Terrestrial Physics*, 113:1–9, 2014.
- [80] Pazos, M., Mendoza, B., Sierra, P., Andrade, E., Rodríguez, D., Mendoza, V., and Garduño, R. Analysis of the effects of geomagnetic storms in the schumann resonance station data in mexico. *Journal of Atmospheric and Solar-Terrestrial Physics*, 193:105091, 2019.
- [81] Salinas, A., Toledo-Redondo, S., Navarro, E., Fornieles-Callejón, J., and Portí, J. Solar storm effects during saint patrick’s days in 2013 and 2015 on the schumann resonances measured by the elf station at sierra nevada (spain). *Journal of Geophysical Research: Space Physics*, 121(12):12–234, 2016.
- [82] Kudintseva, I., Galuk, Y. P., Nickolaenko, A., and Hayakawa, M. Modifications of middle atmosphere conductivity during sudden ionospheric disturbances deduced from changes of schumann resonance peak frequencies. *Radio Science*, 53(5):670–682, 2018.
- [83] Zhou, H. and Qiao, X. Studies of the variations of the first schumann resonance frequency during the solar flare on 7 march 2012. *Journal of Geophysical Research: Atmospheres*, 120(10):4600–4612, 2015.
- [84] Christofilakis, V., Tatsis, G., Votis, C., Contopoulos, I., Repapis, C., and Tritakis, V. Significant elf perturbations in the schumann resonance band before and during a shallow mid-magnitude seismic activity in the greek area (kalpaki). *Journal of Atmospheric and Solar-Terrestrial Physics*, 182:138–146, 2019.
- [85] Galuk, Y. P., Kudintseva, I. G., Nickolaenko, A. P., and Hayakawa, M. Scattering of elf radio waves by a localized non-uniformity in the lower ionosphere. *Journal of Atmospheric and Solar-Terrestrial Physics*, 194:105093, 2019.
- [86] Hayakawa, M., Nickolaenko, A., Galuk, Y. P., and Kudintseva, I. Scattering of extremely low frequency electromagnetic waves by a localized seismogenic ionospheric perturbation: Observation and interpretation. *Radio Science*, 55(12):1–26, 2020.
- [87] Marchitelli, V., Harabaglia, P., Troise, C., and Natale, G. D. On the correlation between solar activity and large earthquakes worldwide. *Scientific Reports*, 10(1), jul 2020. doi:10.1038/s41598-020-67860-3.
- [88] Zaporozhan, V. and Ponomarenko, A. Mechanisms of geomagnetic field influence on gene expression using influenza as a model system: basics of physical epidemiology. *International journal of environmental research and public health*, 7(3):938–965, 2010.
- [89] Rodgers, C. T. and Hore, P. J. Chemical magnetoreception in birds: the radical pair mechanism. *Proceedings of the National Academy of Sciences*, 106(2):353–360, 2009.
- [90] Engels, S., Schneider, N.-L., Lefeldt, N., Hein, C. M., Zapka, M., Michalik, A., Elbers, D., Kittel, A., Hore, P., and Mouritsen, H. Anthropogenic electromagnetic noise disrupts magnetic compass orientation in a migratory bird. *Nature*, 509(7500):353–356, 2014.

- [91] Hiscock, H. G., Worster, S., Kattnig, D. R., Steers, C., Jin, Y., Manolopoulos, D. E., Mouritsen, H., and Hore, P. The quantum needle of the avian magnetic compass. *Proceedings of the National Academy of Sciences*, 113(17): 4634–4639, 2016.
- [92] Richards, F. A flexible growth function for empirical use. *Journal of experimental Botany*, 10(2):290–301, 1959.
- [93] Petroni, N. C., De Martino, S., and De Siena, S. Logistic and θ -logistic models in population dynamics: General analysis and exact results. *Journal of Physics A: Mathematical and Theoretical*, 53(44):445005, 2020.
- [94] Wang, X.-S., Wu, J., and Yang, Y. Richards model revisited: Validation by and application to infection dynamics. *Journal of theoretical biology*, 313:12–19, 2012.
- [95] Kermack, W. O. and McKendrick, A. G. A contribution to the mathematical theory of epidemics. *Proceedings of the royal society of london. Series A, Containing papers of a mathematical and physical character*, 115(772): 700–721, 1927.
- [96] Wu, K., Darcet, D., Wang, Q., and Sornette, D. Generalized logistic growth modeling of the covid-19 outbreak: comparing the dynamics in the 29 provinces in china and in the rest of the world. *Nonlinear dynamics*, 101(3): 1561–1581, 2020.
- [97] Herrmann, H. A. and Schwartz, J.-M. Why COVID-19 models should incorporate the network of social interactions. *Physical Biology*, 17(6):065008, 2020.
- [98] Estrada, E. and Bartesaghi, P. From networked SIS model to the gompertz function. *Applied Mathematics and Computation*, 419:126882, 2022.
- [99] Smilkov, D., Hidalgo, C. A., and Kocarev, L. Beyond network structure: How heterogeneous susceptibility modulates the spread of epidemics. *Scientific reports*, 4(1):1–7, 2014.
- [100] Mombach, J. C., Lemke, N., Bodmann, B. E., and Idiart, M. A. P. A mean-field theory of cellular growth. *EPL (Europhysics Letters)*, 59(6):923, 2002.
- [101] Vattay, G. Forecasting the outcome and estimating the epidemic model parameters from the fatality time series in COVID-19 outbreaks. *Physical Biology*, 17(6):065002, 2020.
- [102] Roda, W. C., Varughese, M. B., Han, D., and Li, M. Y. Why is it difficult to accurately predict the COVID-19 epidemic? *Infectious disease modelling*, 5:271–281, 2020.
- [103] Efron, B. Bayesian inference and the parametric bootstrap. *The annals of applied statistics*, 6(4):1971, 2012.
- [104] Dong, E., Du, H., and Gardner, L. An interactive web-based dashboard to track COVID-19 in real time. *The Lancet infectious diseases*, 20(5):533–534, 2020.
- [105] De Lauro, E., De Martino, S., De Siena, S., and Giorno, V. Stochastic roots of growth phenomena. *Physica A: Statistical Mechanics and its Applications*, 401:207–213, 2014.
- [106] Risken, H. *The Fokker-Planck Equation*. Springer, 1996.
- [107] Akaike, H. Statistical predictor identification. *Annals of the institute of Statistical Mathematics*, 22(1):203–217, 1970.
- [108] Petroni, N. C., De Martino, S., and De Siena, S. Gompertz and logistic stochastic dynamics: Advances in an ongoing quest. *arXiv preprint arXiv:2002.06409*, 2020.
- [109] Boxenbaum, H. Hypotheses on mammalian aging, toxicity, and longevity hormesis: Explication by a generalized gompertz function. In *Biological Effects of Low Level Exposures to Chemicals and Radiation*, pages 1–39. CRC Press, 2017.
- [110] Neafsey, P. J., Boxenbaum, H., Ciraulo, D. A., and Fournier, D. J. A gompertz age-specific mortality rate model of aging, hormesis, and toxicity: Fixed-dose studies. *Drug metabolism reviews*, 19(3-4):369–401, 1988.
- [111] Zhang, C.-L. and Popp, F.-A. Log-normal distribution of physiological parameters and the coherence of biological systems. *Medical Hypotheses*, 43(1):11–16, 1994.
- [112] Skiadas, C. H. Exact solutions of stochastic differential equations: Gompertz, generalized logistic and revised exponential. *Methodology and Computing in Applied Probability*, 12(2):261–270, 2010.
- [113] Sitaram, B. and Varma, V. Statistical mechanics of the gompertz model of interacting species. *Journal of theoretical biology*, 110(2):253–256, 1984.
- [114] Gunasekaran, N. and Pande, L. Log normal distribution for the intrinsic abundance of species from the gompertz model. *Journal of Theoretical Biology*, 98(2):301–305, 1982.
- [115] Chakrabarti, C. and Bhadra, S. Non equilibrium thermodynamics and stochastics of gompertzian growth. *J. Biol. Systems*, 4(2):151–157, 1996. doi:doi.org/10.1142/S0218339096000119.

-
- [116] Sinkhorn, R. and Knopp, P. Concerning nonnegative matrices and doubly stochastic matrices. *Pacific Journal of Mathematics*, 21(2):343–348, 1967.
 - [117] Molski, M. and Konarski, J. Coherent states of gompertzian growth. *Physical review E*, 68(2):021916, 2003.
 - [118] Morse, P. M. Diatomic molecules according to the wave mechanics. ii. vibrational levels. *Physical review*, 34(1): 57, 1929.
 - [119] Glauber, R. J. Coherent and incoherent states of the radiation field. *Physical Review*, 131(6):2766, 1963.
 - [120] Lambert, N., Chen, Y.-N., Cheng, Y.-C., Li, C.-M., Chen, G.-Y., and Nori, F. Quantum biology. *Nature Physics*, 9(1):10–18, 2013.