Draft version November 11, 2026
Typeset using LaTeX **RNAAS** style in AASTeX631

HD 170312 is a bright, long-period eclipsing binary

Mauri de Souza Meneguzzo[1]

[1] *Independent researcher*

ABSTRACT

I identify HD 170312 ($V = 8.7$), a star with an evolved F primary, as a long-period eclipsing binary. Its TESS light curves show four eclipses centered on the star, two 8.5 percent deep and two 15.4 percent deep; the fourth lies in cadences flagged for scattered light. Only two orbital periods fit them, 137.90 and 68.95 d. A possible egress at the start of sector 101, an eclipse in a NEOWISE visit of 2014 and ASAS-3 photometry favor 68.95 d. A deeper eclipse at BTJD 4475.8152 (2027 March 11), in TESS sector 114, would settle the period.

Keywords: Eclipsing binary stars (444) — Stellar photometry (1620) — Light curves (918)

I report an eclipsing binary that had not been recognized: HD 170312 (TIC 292292315; F3 IV/V, $V = 8.71$; Stassun et al. 2019). No catalog I searched lists it as variable, including the TESS eclipsing binary catalog (Prša et al. 2022) and the Gaia DR3 variability tables (Gaia Collaboration 2016, 2023). I call the F star A and its companion B.

The star was found by a search of multi-sector TESS 2-minute light curves (Ricker et al. 2015; Jenkins et al. 2016) with transitgpu 0.4.0 (Meneguzzo 2026). Its peak, at 12.9 d = 1792.694/139 d, is a spacing of the eclipses, not the orbital period.

TESS recorded four complete eclipses (Figure 1b). Three lie in unflagged data, in sectors 39, 102 and 103, at BTJD 2377.1543 ± 0.0002, 4131.0663 ± 0.0001 and 4169.8482 ± 0.0005. They are of two kinds: the first and third are 8.5 percent deep, the second 15.4 percent, each uncertain by about 1.5 percentage points because the PDC correction tilts the baseline beside the sector 102 eclipse.

The fourth eclipse lies in data that the pipeline sets aside. Sector 93 continues for a day after its last unflagged cadence, in 715 flagged cadences, 359 of them for scattered light, which PDCSAP omits. Their SAP flux holds a deep eclipse. Fitted with the shape of the sector 102 eclipse, it is centered at BTJD 3855.2675, 20 s from the predicted time, at 1.003 times the depth of that eclipse. Two tests show that it is the star's and not the scattered light's. The fall in each of 121 pixels is proportional to that pixel's light out of eclipse (correlation −0.998). The depth changes by under 0.1 percentage point across five estimates of the background, which rises more than tenfold over the eclipse.

The eclipses come from the HD 170312 system. The dimming lies within 2.2″ of A, and every Gaia DR3 source within 5″ is $G = 19.6$ or fainter, 11 magnitudes too faint to make a 15 percent dip.

The eclipses also show that B is a star, too large for a planet or a brown dwarf. I do not know which eclipse is B passing in front of A, so the shallow eclipse bounds the size of B: $R_B \geq R_A\sqrt{\delta/c}$, where $c = 1.134$ is the ratio of the central to the mean intensity of A for quadratic limb darkening at 6645 K and $\log g = 3.5$ (Claret 2017). This gives $R_B \geq 0.95\,R_\odot$ with the TIC radius $R_A = 3.46\,R_\odot$. Models of two eclipsing stars fitted to the eclipses need A to be 24 to 34 percent smaller than its TIC radius, which I cannot explain. With their smallest radius of A, $2.27\,R_\odot$, the bound is $R_B \geq 0.62\,R_\odot$.

Two periods fit the four eclipses. The shallow eclipses are $\Delta T = 1792.694$ d apart, so each kind recurs with $P = \Delta T/N$ for an integer N. Of 3,200 values of N, only $N = 13$ and 26, $P = 137.90$ and 68.95 d, predict all four eclipses and no eclipse where the data show none. The eclipses that only 68.95 d predicts fall in gaps of the TESS coverage, except at the start of sector 101 (Figure 1a). The two-star models fit the eclipses equally well at both periods, so the eclipse shapes cannot choose between them. The deep eclipse falls at phase 0.44 of the 68.95 d orbit or 0.72 of the 137.90 d one, so the orbit is eccentric, with $e \geq 0.10$ or 0.34.

Three observations favor 68.95 d. First, flagged cadences at the start of sector 101 rise as the egress of a shallow eclipse, at the time that only 68.95 d predicts (BTJD 4100.898) and at 0.997 times the depth of that kind (Figure 1b). Five stars of similar brightness on the same CCD show no such rise; two brighter ones show ramps at that sector start, and the event was examined after the data were seen. Second, ASAS-3 observed the star every two days from 2000 to 2009 (Pojmanski 1997). On the ten nights that only 68.95 d places in eclipse, a test whose rule was fixed before it was run, though after its points had been examined, finds them fainter by 1.04 times the TESS depths, with $p \approx 0.01$ over random shifts of the ephemeris (0.006 to 0.012 by the shifts excluded; Figure 1c). Each deficit is a single point, so ASAS-3 leans toward 68.95 d without deciding. Third, a NEOWISE visit (Mainzer et al. 2014) on 2014 September 24 covers the deep eclipse that only 68.95 d predicts: HD 170312 fades by 0.13 mag in W2, faintest at the predicted time within the 95 minutes between exposures, while none of 11 stars in the same frames, one as bright, fades by more than 0.02 mag. With no visit at an eclipse both periods predict, the test has no control, and the visit was found after the period analysis, so it favors 68.95 d without establishing it. ASAS-SN photometry of 2014 to 2026 (Shappee et al. 2014; Kochanek et al. 2017; Winecki & Kochanek 2024) is also fainter at the eclipses both periods predict, by 2.07 times the TESS depths ($p < 5 \times 10^{-5}$), but at those only 68.95 d predicts by 0.89 ± 0.37 times, so it does not support 68.95 d.

The period can be tested on 2026 December 2, when only 68.95 d predicts a shallow eclipse, and settled by a deeper eclipse at BTJD 4475.8152 ± 0.0002 (2027 March 11), which TESS sector 114 covers and 137.90 d does not predict. The classification rests on photometry: I found no public spectrum of the system, and Gaia DR3 gives it no radial velocity.




Figure 1. (a) Phases of the 137.90 d orbit (0 at the shallow eclipse) covered by each TESS sector and, in the top rows, by ASAS-SN camera-nights, ASAS-3 nights and NEOWISE visits. Solid lines mark the shallow (blue) and deep (red) eclipses that both periods predict; dashed orange lines, those that only 68.95 d predicts. Orange ticks are the ten ASAS-3 nights of panel (c), the orange triangle the NEOWISE visit at such a deep eclipse, and hollow boxes the ASAS-SN nights in those windows, whose test is undecided. The marks show when data exist, not whether the star was fainter. Markers show the eclipses seen, a possible egress in flagged cadences of sector 101 and the predicted eclipse of 2027 March 11. (b) The four eclipses in SAP flux with 1 h medians, aligned on their deepest point; black, the flagged start of sector 101 at the time that 68.95 d predicts. (c) ASAS-3 nightly V means on those ten nights and, in grey, other nights, with the TESS eclipse shapes; error bars are scaled by 2.9 to the scatter of the data. Inset: amplitudes at 20,000 shifted epochs (log scale), 0.6 percent of which reach the fitted 1.04.

This paper uses TESS data from the MAST archive at STScI (doi:10.17909/t9-nmc8-f686), which is operated by AURA under NASA contract NAS5–26555; TESS is funded by the NASA Explorer Program. It uses data products of NEOWISE (doi:10.26131/IRSA144), a project of JPL/Caltech funded by NASA, and the SIMBAD database. This work has made use of data from the European Space Agency (ESA) mission *Gaia* (https://www.cosmos.esa.int/gaia), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, https://www.cosmos.esa.int/web/gaia/dpac/consortium). Funding for the DPAC has been provided by national institutions, in particular the institutions participating in the *Gaia* Multilateral Agreement. An AI language model (Anthropic Claude) assisted with the code and the writing.

REFERENCES

Claret, A. 2017, A&A, 600, A30

Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, A&A, 595, A1

Gaia Collaboration, Vallenari, A., Brown, A. G. A., et al. 2023, A&A, 674, A1

Jenkins, J. M., Twicken, J. D., McCauliff, S., et al. 2016, Proc. SPIE, 9913, 99133E

Kochanek, C. S., Shappee, B. J., Stanek, K. Z., et al. 2017, PASP, 129, 104502

Mainzer, A., Bauer, J., Cutri, R. M., et al. 2014, ApJ, 792, 30

Meneguzzo, M. de S. 2026, Parallel Search for Alien Worlds: GPU-Accelerated BLS and TLS for Exoplanet Transit Detection, preprint, github.com/mauri870/transitgpu

Pojmanski, G. 1997, Acta Astron., 47, 467

Prša, A., Kochoska, A., Conroy, K. E., et al. 2022, ApJS, 258, 16

Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2015, JATIS, 1, 014003

Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, ApJ, 788, 48

Stassun, K. G., Oelkers, R. J., Paegert, M., et al. 2019, AJ, 158, 138

Winecki, D., & Kochanek, C. S. 2024, ApJ, 971, 61