

Subsurface warming of West Antarctica during El Niño

Maurice F. Huguenin^{1,2,3,*}, Ryan M. Holmes^{3,4}, Paul Spence^{2,3,5} and Matthew H. England^{1,2,3}


¹Climate Change Research Centre, University of New South Wales, Sydney, Australia, ²ARC Australian Centre for Excellence in Antarctic Science, University of New South Wales, Sydney, New South Wales, Australia, ³ARC Centre of Excellence in Climate Extremes, University of New South Wales, Sydney, Australia, ⁴School of Geosciences, University of Sydney, Sydney, Australia, ⁵Institute for Marine and Antarctic Studies, University of Tasmania, Australia



UNSW
SYDNEY

climate extremes
ARC centre of excellence

*E-mail: m.huguenin-virchaux@unsw.edu.au



This poster as .PDF

UNSW
Climate Change
Research Centre



ACEAS
Australian Centre for Excellence in Antarctic Science
A Special Research Initiative of the Australian Research Council

- El Niño-Southern Oscillation (ENSO) modulates West Antarctic shelf water temperatures and can cause rapid basal melting of grounded ice shelves, accelerating sea level rise
- However we lack understanding of the oceanic response to ENSO in this region due to **sparse observations**


Idealised El Niño & La Niña simulations

- **ACCESS-OM2-01** Kiss et al. (2020)
 - 1/10° global ocean-sea ice model with 75 z* levels
 - forced by JRA55-do, atmospheric reanalysis Tsujino et al. (2018)
 - investigate warming and cooling on the shelf during ENSO

Idealised simulations

- climatological repeat-year forcing[x,y,t] + ENSO anomalies
(time series[t] × spatial pattern[x,y])

Fig. 1. a, c, Composite time series associated with ENSO sea surface temperature anomalies based on observed events. b, d, Spatial patterns of sea level pressure (hPa) and surface winds ($m s^{-1}$) during the shaded El Niño (pink) and La Niña (blue) periods in a, c.



Response of the West Antarctic shelf to El Niño & La Niña

- El Niño: weaker Amundsen Sea Low → more northward Ekman transport & advection of warm Circumpolar Deep Water onto shelf
- La Niña: response inhibited by stronger Amundsen Sea Low & surface easterlies




Fig. 2. a, b, Peak event 100-1000 m mean temperature response ($^{\circ}C$). c, d, Mean across-shelf temperature responses $150^{\circ}W$ - $60^{\circ}W$ ($^{\circ}C$). e, f, Mean Ekman transport velocities ($m^3 s^{-1}$). g, h, Eulerian heat budget anomalies ($10^6 W m^{-2}$) throughout the simulations.




Fig. 3. a, b, Schematic of anomalous physical processes on the West Antarctic shelf during El Niño and La Niña. ¹ = key findings in Paolo et al. (2018).