

Problem Sheet 7.2: Lambda Calculus Types

Exercise 1: What are the types of the following typed λ -terms? In each case, give a typing derivation to demonstrate that the term really is well typed.

- a) $\lambda x^o.x$
- b) $\lambda x^o.\lambda y^o.\lambda z^{o \rightarrow o}.zx$
- c) $\lambda f^{(o \rightarrow o) \rightarrow o}.\lambda g^{o \rightarrow o \rightarrow o}.f(\lambda x^o.gxx)$

Exercise 2: Can the term $(\lambda x.x)(\lambda y.y)$ be typed? That is, are there types with which one can label the variables x and y such that the term is well typed? If not, why not?

Exercise 3: We have seen that the Church numerals could be typed, with type

$$(o \rightarrow o) \rightarrow o \rightarrow o.$$

(Let us call this type N for short.)

- a) Give a type derivation to show that the successor operator

$$\lambda n^N.\lambda f^{o \rightarrow o}.\lambda x^o.f(n f x)$$

can be given type $N \rightarrow N$.

- b) Write down the type you would expect the addition operator to have.
- c) What does the term for addition look like when its bound variables are labelled appropriately?
- d) Give a type derivation for the addition term.
- e) Now consider the term for exponentiation: can that be similarly typed?