

SSC CGL (COMBINED GRADUATE LEVEL - INSPECTOR, ASO, TAX ASST) — SUBJECT COMPLETE QUESTION BANK

PRACTICE MATERIAL • VERIFIED QUESTION BANK

Total Questions: 585 • Mode: Single Subject Comprehensive Practice

Q.1. In SSC CGL Tier-1 context (Number Systems & Divisibility, Problem #1): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.2. In SSC CGL Tier-1 context (HCF and LCM Applications, Problem #2): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.3. In SSC CGL Tier-1 context (Simplification & Surds & Indices, Problem #3): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.4. In SSC CGL Tier-1 context (Percentages & Successive Change, Problem #4): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.5. In SSC CGL Tier-1 context (Profit, Loss & Discount / Marked Price, Problem #5): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.6. In SSC CGL Tier-1 context (Simple & Compound Interest Differences, Problem #6): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.7. In SSC CGL Tier-1 context (Ratio, Proportion & Mixture / Alligation, Problem #7): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.8. In SSC CGL Tier-1 context (Time, Work, Pipes & Cisterns, Problem #8): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.9. In SSC CGL Tier-1 context (Speed, Time, Distance & Trains / Boats, Problem #9): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) $\frac{1}{2}$ B) $\frac{1}{3}$ C) $\frac{1}{4}$ D) $\frac{1}{5}$

Q.10. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('18'B "Y "%>'ÙG'> @, '©M" 6"Ù("% 'iM'ù> #10): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #10): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i"é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+2B)MABWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.11. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('É@'É 9?'I@'ò 8" M"Y8'é?Y>•ù •ù5•", 'TM'A'©&, '©M" 6"Ù("% 'iM'ù> #11): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #11): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" **é\$ "ù '%oM'ùB"%@'Y0'2 "é0'R &"Ü5"é0"â **Ü0"é**ù\$ '©0B)%(ù\$ 'é\$+2B)MABWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.12. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- (" H'i?R "Y "iM'Y?%>B 8'é@'Y0'2Ä *"Ü0'iM", 8•) "Ü"ä 3 "" •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #12): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'é>%o •i"é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+2B)MABWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.13. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?'ÜA" "Ü"é"ü"0"y "iM" "Y '©M" .")y), '©M" 6"Ù("% 'iM'ù> #13): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #13): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ü%"é*, **Ü0"éG'ò &"Ü5"é0"â 8'iM'ù>©?B 'iM'Y?"I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.14. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ("YC'iM'B *"Ü0"éG'ò %8B"Ü" M"i0"y "é •" "Y 'É@'Y>•ù), '©M" 6"Ù("% 'iM'ù> #14): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #14): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i"é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+2B)MABWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.15. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ("%?" M'iG'>•) "EM'ù>é?I? •ù5•" **Ü0"Y#I>, '©M" 6"Ù("% 'iM'ù> #15): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #15): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" **é\$ "ù '%oM'ùB"%@'Y0'2 "é0'R &"Ü5"é0"â **Ü0"é**ù\$ '©0B)%(ù\$ 'é\$+2B)MABWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.16. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?'YK'9,"ù\$) •Y(" **é\$ "ü5•" 8" M"Y8'é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #16): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #16): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'é>%o •i"é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+2B)MABWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.17. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- (•© >©•, "Y 'iB' @, '©M" 6"Ù("% 'iM'ù> #17): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #17): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ü%"é*, **Ü0"éG'ò &"Ü5"é0"â 8'iM'ù>©?B 'iM'Y?"I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.18. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM'Y?"Y?é@'ò "Ü7"y\$Ü0"é?I?, '©M" 6"Ù("% 'iM'ù> #18): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #18): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i"é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+2B)MABWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.19. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?"Y?"Y?é@'ò "Ü7"y\$Ü0"é?I?, '©M" 6"Ù("% 'iM'ù> #19): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #19): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" **é\$ "ù '%oM'ùB"%@'Y0'2 "é0'R &"Ü5"é0"â **Ü0"é**ù\$ '©0B)%(ù\$ 'é\$+2B)MABWVÇBFA Bed via proportional reduction factor (Case: #Q31)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2"%MÉ0'ÙB B0%0"é.F-Ö"Ä 66QWFCda 802 QS-ér, %8FÜWVÇBFA Bed via proportional reduction factor (Case: #Q31)

Q.30. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('18'B "Y "%>'ÙG'>" @, '©M" 6"Ù("% 'iM'ù> #30): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #30): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+69)MABlææNValuëmRtdlù\$ "ëançadlùBasùBëqunçrret(%)dù\$
(Case:#Q"
C) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é

Q.31. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('É@'É '9?'l@'ò 8" M"Y8'é?Y>•ù •ù5•" , "™A'©&, '©M" 6"Ù("% 'iM'ù> #31): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #31): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ 'é\$+69)MABlææNValuëmRtdlù\$ "ëançadlùBasùBëqunçrret(%)dù\$
(Case:#Q"
C) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é

Q.32. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- (" H'i?R "Y "iM'Y? "%> B 8'éY@'Y0'2A *"Ù0'iM, 8•) "Ù"à 33"" •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #32): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é
(Case:#Q"
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+69)MABlææNValuëmRtdlù\$ "ëançadlùBasùBëqunçrret(%)dù\$

Q.33. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?'ÙA" "Ù"é"ù\$ "0"y "iM" "Y '©M" .")y), '©M" 6"Ù("% 'iM'ù> #33): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #33): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é
(Case:#Q"
C) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é

Q.34. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ("YCiM'B *"Ù0'ég'ò "ù\$ "0"y "iM" "Y 'É@"Y>•ù), '©M" 6"Ù("% 'iM'ù> #34): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #34): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+69)MABlææNValuëmRtdlù\$ "ëançadlùBasùBëqunçrret(%)dù\$
(Case:#Q"
C) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é

Q.35. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ("%"? " M'iG'>•) "EM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é
[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #35): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ 'é\$+69)MABlææNValuëmRtdlù\$ "ëançadlùBasùBëqunçrret(%)dù\$
(Case:#Q"
C) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é

Q.36. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?'YK9" "ù\$ "0"y "iM" "Y "Y8'é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #36): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #36): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é
(Case:#Q"
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+69)MABlææNValuëmRtdlù\$ "ëançadlùBasùBëqunçrret(%)dù\$

Q.37. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- (•© >•, "Y "iB" @, '©M" 6"Ù("% 'iM'ù> #37): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"ù\$ •
[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #37): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é
(Case:#Q"
C) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é

Q.38. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM'Y?"Y?é@'ò "Ù7"y\$ "Ù0'é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é
[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #38): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é\$+69)MABlææNValuëmRtdlù\$ "ëançadlùBasùBëqunçrret(%)dù\$
(Case:#Q"
C) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é

Q.39. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?"Y?"Y?é@'ò "Ù7"y\$ "Ù0'é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é
[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #39): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ 'é\$+69)MABlææNValuëmRtdlù\$ "ëançadlùBasùBëqunçrret(%)dù\$
(Case:#Q"
C) 'ÉM'ù>é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "M(\$ùB" 0"%:30" 50" 66" 6- & Ù\$ 60CWMa-60? B- iæYV24@,ùVQ0W6-é

Q.70. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('18'B "Y "%>'ÙG'>" @, '©M" 6"Ù("% 'iM'ù> #70): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #70): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+14M)DAtSou@Y0'2alté0RaChù5gérantù0'èà\$SineQaàné(é(Ca
(Case:#Q#
C) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q##
theorem (Case :C03S•

Q.71. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('É@'É'9?'I@'ò 8" M"Y8é?Y>•ù •ù5•" , "™A'©&, '©M" 6"Ù("% 'iM'ù> #71): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #71): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$ "ù '%oM'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©OB)##6-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q#
C) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q##
theorem (Case :C03SR•

Q.72. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- (" H'i?'R "Y "iM'Y?'>B 8é@'Y0'2A *"Ù0'iM', 8•) "Ù/"â 3s"" •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #72): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q#
C) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+14M)DAtSou@Y0'2alté0RaChù5gérantù0'èà\$SineQaàné(é(Ca
(Case:#Q#
theorem (Case :C03SR•

Q.73. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?'ÙA" "Ùe"ù\$ "0"y "iM" "Y '©M" .")y), '©M" 6"Ù("% 'iM'ù> #73): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #73): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>'©?B 'iM'Y?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q#
C) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q##
parameter (Case :Ö Cb•

Q.74. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ("YC'iM'B *"Ù0'ég'ò "ù\$ "0"y "iM" "Y 'É@'Y>•ù), '©M" 6"Ù("% 'iM'ù> #74): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #74): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+14M)DAtSou@Y0'2alté0RaChù5gérantù0'èà\$SineQaàné(é(Ca
(Case:#Q#
C) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q##
theorem (Case :C03S•

Q.75. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ("%"? M'ic'>•) "EM'ù>é?'I@'òù •) "ù5•" *"Ù0"Y#I>, '©M" 6"Ù("% 'iM'ù> #75): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #75): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$ "ù '%oM'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©OB)##6-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q#
C) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q##
parameter (Case :Ö S•

Q.76. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?'YK'9,"ù\$ "Y("é\$ "ù5•" 8" M"Y8é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #76): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #76): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q#
C) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+14M)DAtSou@Y0'2alté0RaChù5gérantù0'èà\$SineQaàné(é(Ca
(Case:#Q#
theorem (Case :C03SR•

Q.77. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- (•©'©>•, "Y 'iB' @, '©M" 6"Ù("% 'iM'ù> #77): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"ù\$ •
[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #77): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>'©?B 'iM'Y?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q#
C) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q##
parameter (Case :Ö SB•

Q.78. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM'Y?"Y?é@'ò "Ù7"y\$ "Ù0'é?I?, '©M" 6"Ù("% 'iM'ù> #78): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #78): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+14M)DAtSou@Y0'2alté0RaChù5gérantù0'èà\$SineQaàné(é(Ca
(Case:#Q#
C) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q##
theorem (Case :C03S•

Q.79. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM" ?"Y?"Y?é@'ò "Ù7"y\$ "Ù0'é?I?, '©M" 6"Ù("% 'iM'ù> #79): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #79): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$ "ù '%oM'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©OB)##6-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q#
C) 'ÉM'ù>é?'I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)E2-%MVE0'ÙB B0%006" F-ÖrÄ"66QVBF-àà602"ù\$ éar,%8F"ÙW0\$W6=I
(Case:#Q##
theorem (Case :C03R•

Q.90. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('18'B "Y "%>'ÙG'>" @, '©M" 6"Ù("% 'iM'ù> #90): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #90): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+18M'DatSou@Y0'2alté0Rat'ù5géantù0'basé0Qua#né(%)C#
(Case: #Q#s •
C) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é

Q.91. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('É@'É'9?'ù0'8" M'Y8'é?Y>•ù •ù5•" , "™A'©&, '©M" 6"Ù("% 'iM'ù> #91): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #91): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$"ù '%oM'ùB"%@'Y0'2 "é0'R &"ù5"é0"â *""0"é**"ù\$ '©0B)%(ù\$'é\$+18M'DatSou@Y0'2alté0Rat'ù5géantù0'basé0Qua#né(%)C#
(Case: #Q#s •
C) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é

Q.92. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- (" H'i?R "Y "iM'Y?%> B'8'é@'Y0'2Â *""0'iM', 8•) "ù"à 3"" "ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #92): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é
(Case: #Q#s •
C) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+18M'DatSou@Y0'2alté0Rat'ù5géantù0'basé0Qua#né(%)C#

Q.93. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM' ?'ÙA" "Ù"é"ù\$ "0"y "iM' "Y '©M" .")y), '©M" 6"Ù("% 'iM'ù> #93): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #93): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *""0'ég'ò &"ù5"é0"â 8'iM'ù>?B 'iM'Y?"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é
(Case: #Q#s •
C) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é

Q.94. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ("Yc'iM'B "Ù0'ég'ò "ù\$ "0"y "iM' "Y 'É@'Y>•ù), '©M" 6"Ù("% 'iM'ù> #94): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #94): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+18M'DatSou@Y0'2alté0Rat'ù5géantù0'basé0Qua#né(%)C#
(Case: #Q#s •
C) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é

Q.95. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ("%"? " M'ic'>•) "EM'ù>?é?l@'òù •) "ù\$ "0"y "iM' "Y 'É@'Y>•ù), '©M" 6"Ù("% 'iM'ù> #95): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #95): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$"ù '%oM'ùB"%@'Y0'2 "é0'R &"ù5"é0"â *""0"é**"ù\$ '©0B)%(ù\$'é\$+18M'DatSou@Y0'2alté0Rat'ù5géantù0'basé0Qua#né(%)C#
(Case: #Q#s •
C) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é

Q.96. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM' ?'YK'9,"ù\$ "0"y "iM' "Y "é\$ "ù5•" 8" M'Y8'é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #96): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #96): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é
(Case: #Q#s •
C) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+18M'DatSou@Y0'2alté0Rat'ù5géantù0'basé0Qua#né(%)C#

Q.97. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- (•©'©>•, "Y "iB' @, '©M" 6"Ù("% 'iM'ù> #97): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"ù\$ •
[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #97): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *""0'ég'ò &"ù5"é0"â 8'iM'ù>?B 'iM'Y?"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é
(Case: #Q#s •
C) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é

Q.98. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM'Y?"Y?é@'ò "ù7"y\$ "Ù0'é?l?, '©M" 6"Ù("% 'iM'ù> #98): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #98): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>%o •i""é0'ÙB'B *"%0"é." " "r '%oA" B'¢ ** '?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+18M'DatSou@Y0'2alté0Rat'ù5géantù0'basé0Qua#né(%)C#
(Case: #Q#s •
C) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é

Q.99. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('iM' ?'Y?"Y?é@'ò "ù7"y\$ "Ù0'é?l?, '©M" 6"Ù("% 'iM'ù> #99): •ù •™."Ù." &"Y>" ("ù0"Ù""é0"
[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #99): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$"ù '%oM'ùB"%@'Y0'2 "é0'R &"ù5"é0"â *""0"é**"ù\$ '©0B)%(ù\$'é\$+18M'DatSou@Y0'2alté0Rat'ù5géantù0'basé0Qua#né(%)C#
(Case: #Q#s •
C) 'ÉM'ù>?é?l@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IB0@'Y0'2F"0' R 60QW60a"60?B -ImY74@,òV9QW60é

- Q.100.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (•i 'Y!'É> "%? " M'Y ' , %, "é0 'yM" >'2Â ""é '©>" M'ò'Â ""'Ù0"iM', 8•) "Ù/"â 3 " •ù
- [English]: In SSC CGL Tier-1 context (Data Interpretation (Bar Graphs, Pie Charts, Tables), Problem #100): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'é>"% •i""é0'ÜB'B *"%0"é." " "r" '%0A" B'Ç *"" ?'Y2"ù\$ 'é>', %8B) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.101.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("%" 'iM'ù> '©M" # "é2" "Y" 'Y'Ù>'ÉM'ù\$ "âÂ ""'Ù0"iM', 8•) "Ù/"â 3 " •ù •TM."Ù."
- [English]: In SSC CGL Tier-1 context (Number Systems & Divisibility, Problem #101): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) '©M" \$"ù8"Ù%"é*, ""'Ù0'ÉG'ò &"Ù5"é0"â 8'IM'ù>'©?B 'iM'Y?'I@) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.102.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('âé8.Çâ •ù5" 2.é* "Y(" "Ù0"iM', 8•) "Ù/"â 3 "" •ù •TM."Ù." &"Y>" (
- [English]: In SSC CGL Tier-1 context (HCF and LCM Applications, Problem #102): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) 'é>"% •i""é0'ÜB'B *"%0"é." " "r" '%0A" B'Ç *"" ?'Y2"ù\$ 'é>', %8B) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.103.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("%"0)@ 'Y02 'Y" %>'I>•) "R" " # " Â ""'Ù0"iM', 8•) "Ù/"â 3 2" •ù •TM."Ù." &"Y>" (
- [English]: In SSC CGL Tier-1 context (Simplification & Surds & Indices, Problem #103): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) 'i(" *""é\$ "ù" '%M'ùB"%@ 'Y0'2 "é0'R &"Ù5"é0"â ""'Ù0"é""Ù\$ '©B) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.104.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('©M" \$"ù6"\$â "Y" YM" .ù" '©0"ù5" M'I(, '©M" 6"Ù("% 'iM'ù> #104): •ù •TM."Ù.".
- [English]: In SSC CGL Tier-1 context (Percentages & Successive Change, Problem #104): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'é>"% •i""é0'ÜB'B *"%0"é." " "r" '%0A" B'Ç *"" ?'Y2"ù\$ 'é>', %8B) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.105.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (")>'ÒÂ 9"é(" "Y" 'E" "U"â ö •) "ù\$ 'éB")M'òÂ ""'Ù0"iM', 8•) "Ù/"â 3 R" •ù •TM."Ù.".
- [English]: In SSC CGL Tier-1 context (Profit, Loss & Discount / Marked Price, Problem #105): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) '©M" \$"ù8"Ù%"é*, ""'Ù0'ÉG'ò &"Ù5"é0"â 8'IM'ù>'©?B 'iM'Y?'I@) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.106.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("%">'y>" # "ù5" "YM" 5'9&"U"ò , "Ù/"é •Y 'i0, '©M" 6"Ù("% 'iM'ù> #106): •ù •TM."Ù.".
- [English]: In SSC CGL Tier-1 context (Simple & Compound Interest Differences, Problem #106): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) 'é>"% •i""é0'ÜB'B *"%0"é." " "r" '%0A" B'Ç *"" ?'Y2"ù\$ 'é>', %8B) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.107.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (•Y(" *""é\$, "%."é(" *""é\$ "ù5" . "ù6"Ù0'2Â ""'Ù0"iM', 8•) "Ù/"â 3 r" •ù •TM."Ù." &"Y>" (
- [English]: In SSC CGL Tier-1 context (Ratio, Proportion & Mixture / Alligation, Problem #107): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) 'i(" *""é\$ "ù" '%M'ùB"%@ 'Y0'2 "é0'R &"Ù5"é0"â ""'Ù0"é""Ù\$ '©B) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.108.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('Y>" M'òÂ 8'è "ù5" ("U" •) " Â ""'Ù0"iM', 8•) "Ù/"â 3 , " •ù •TM."Ù." &"Y>" ("ù0"Ù
- [English]: In SSC CGL Tier-1 context (Time, Work, Pipes & Cisterns, Problem #108): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'é>"% •i""é0'ÜB'B *"%0"é." " "r" '%0A" B'Ç *"" ?'Y2"ù\$ 'é>', %8B) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- Q.109.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('©>" "Â 8'è, "iB" @, "G") "é!"É@ •ù5" ("é5"y>" >, '©M" 6"Ù("% 'iM'ù> #109): •ù
- [English]: In SSC CGL Tier-1 context (Speed, Time, Distance & Trains / Boats, Problem #109): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
- A) '©M" \$"ù8"Ù%"é*, ""'Ù0'ÉG'ò &"Ù5"é0"â 8'IM'ù>'©?B 'iM'Y?'I@) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •
- C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y-" 7"ùB) "r2"(éS M'Y% M' (B)C@'Y0'F-"é0'R6QÜV6é0â 6Ü0'G-"ùS"©0F#V0'ÖW'ÖN' (Case : #Q3#B) parameter (Case : COS •

Q.110. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('I8'B "Y "%>'ÙG'i>" @, '©M" 6"Ù("% 'iM'ù> #110): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #110): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+228M'DatSou@Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•

Q.111. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('É@'É '9?'I@'ò 8" M'Y8é?Y>•ù •ù5•" , "™A'©&, '©M" 6"Ù("% 'iM'ù> #111): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #111): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$"ù "%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$'é\$+228M'DatSou@Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•

Q.112. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (" H'i?R "Y "IM'Y?"Y>B 8é?Y0'2A *"Ù0'iM', 8•) "Ù/"â 3 "" •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #112): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+228M'DatSou@Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•

Q.113. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('IM' ?'UA'A "Ùé."ùs"ò "% 'y 'iM' "Y '©M" ."y), '©M" 6"Ù("% 'iM'ù> #113): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #113): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y>B 8é?Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•

Q.114. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('YC'IM'B *"Ù0'ég'ò "%8"Ù" M'io'y "é •" "Y 'É@"Y>•ù), '©M" 6"Ù("% 'iM'ù> #114): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #114): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+228M'DatSou@Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•

Q.115. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('%?" M'ig'> "EM'ù>é?I? "ù5•" *"Ù0'Y#I>, '©M" 6"Ù("% 'iM'ù> #115): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #115): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$"ù "%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$'é\$+228M'DatSou@Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•

Q.116. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('IM' ?'YK'9 "ù\$" / •Y(" *"é\$ •ù5•" 8" M'Y8'é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #116): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #116): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+228M'DatSou@Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•

Q.117. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('©' ©>, Y 'IB' @, '©M" 6"Ù("% 'iM'ù> #117): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #117): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y>B 8é?Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•

Q.118. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM'Y?"Y>é@'ò "U7'y\$U0é?I?, '©M" 6"Ù("% 'iM'ù> #118): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #118): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$'é\$+228M'DatSou@Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•

Q.119. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('IM' ?"Y?é@'ò "U7'y\$U0é?I?, '©M" 6"Ù("% 'iM'ù> #119): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùs

[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #119): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *"é\$"ù "%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$'é\$+228M'DatSou@Y0'2'Valé0'RaChù5géantù0'BasSineQ'aramé(%)C (Case:#Q33•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'BOC@'Y0'2'F'É0'RaQ'WFO'a'60'¢'ImeY'4@'p'V'ù0'W6 theorem (Case ;CÔSS•

Q.130. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('I8'B "Y "%>'ÙG'i>" @, '©M" 6"Ù("% 'iM'ù> #130): •ù •™."Ù." &"Y>" ("ù0"Ù"é
[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #130): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+26M Data@Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"Ù\$ '©OB)##-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;CÔcS" •

Q.131. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('É@'É '9?'I@'ò 8" M"Y8é?Y>•ù •ù5•" , "™A'©&, '©M" 6"Ù("% 'iM'ù> #131): •ù
[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #131): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"Ù\$ '©OB)##-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;CÔcS" •

Q.132. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (" H'i?R "Y "iM'Y?Y?>B 8é@'Y0'2A *"Ù0"iM', 8•) "Ù/"â 3 3"" •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #132): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+26M Data@Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"Ù\$ '©OB)##-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •

Q.133. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'ÙA" "Ù"é."ù\$ 0 % "y 'iM' "Y '©M" ."y), '©M" 6"Ù("% 'iM'ù> #133): •ù
[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #133): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B "iM'Y?"I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •

Q.134. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('YCiM'B *"Ù0éG'ò %8"Ù" M"i0"y "é •" "Y 'É@"Y>•ù), '©M" 6"Ù("% 'iM'ù> #
[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #134): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+26M Data@Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"Ù\$ '©OB)##-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •

Q.135. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("%?" M'iG'> "ÉM'ù>é?I? •ù5•" *"Ù0"Y#I>, '©M" 6"Ù("% 'iM'ù> #135): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #135): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"Ù\$ '©OB)##-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •

Q.136. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'YK'9."ù\$ / •Y(" *é\$ •ù5•" 8" M"Y8'é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #136): •
[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #136): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+26M Data@Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"Ù\$ '©OB)##-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •

Q.137. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (•© '©>•, Y 'iB' @, '©M" 6"Ù("% 'iM'ù> #137): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ù\$
[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #137): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B "iM'Y?"I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •

Q.138. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM'Y?"Y?é@'ò "U7"ys"Ù0é?I?, '©M" 6"Ù("% 'iM'ù> #138): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #138): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+26M Data@Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"Ù\$ '©OB)##-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •

Q.139. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?"Y?é@'ò "U7"ys"Ù0é?I?, '©M" 6"Ù("% 'iM'ù> #139): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #139): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"Ù\$ '©OB)##-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2-%M'è0'ÙB'BO3206&W7VÇB.FAVBéd"vay2pùpùèh&B"ù0"ù"é
(Case:#Q3" parameter (Case:;Ô#c" theorem (Case:;CÔcS" •

Q.140. •ù8•ù8“%@@ “%@‘É@•ù2 ‘ù?’‘ù0-1 “% ‘i0”Ù- (•¡‘Y!‘É> ‘%0?“ M“Y ‘, %,“é0 ‘yM“ >²Â ““é ‘©>“ M‘d‘Â ““Ù0“iM‘, 8•) ”Ù/“â 3 C “ •ù
 [English]: In SSC CGL Tier-1 context (Data Interpretation (Bar Graphs, Pie Charts, Tables), Problem #140): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.141. •ù8•ù8"%@ @ "%@•É@•ù2 u?•ù0-1 "%% 'i0"Ù- ("%" iM"ù> "©M" #e2 "Y "Y?"Ù>ÉM"ù\$ââ *""Ù0"iM", 8•) "Ù"/â 3 C " •ù •™."Ù.
 [English]: In SSC CGL Tier-1 context (Number Systems & Divisibility, Problem #141): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.142. [English]: In SSC CGL Tier-1 context (HCF and LCM Applications, Problem #142): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.143. [English]: In SSC CGL Tier-1 context (Simplification & Surds & Indices, Problem #143): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.144. [English]: In SSC CGL Tier-1 context (Percentages & Successive Change, Problem #144): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.145. [English]: In SSC CGL Tier-1 context (Profit, Loss & Discount / Marked Price, Problem #145): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.146. [English]: In SSC CGL Tier-1 context (Simple & Compound Interest Differences, Problem #146): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.147. [English]: In SSC CGL Tier-1 context (Ratio, Proportion & Mixture / Alligation, Problem #147): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.148. •ù8•ù8“%0@ “%0@É@•ù2 `ù?`ù0-1 “%0 `ì0”Ù- (‘Y>“ M`dÀ 8`è/ •ù5. (“U•) “Á *”Ù0“ìM, 8•) ”Ù/“á 3 C,“ •ù •™.“ù.” &“Y>“ (“ù0”Ù
 [English]: In SSC CGL Tier-1 context (Time, Work, Pipes & Cisterns, Problem #148): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.149. [English]: In SSC CGL Tier-1 context (Speed, Time, Distance & Trains / Boats, Problem #149): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

parameter (Case , $U_{\#}$, •

Q.150. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('I8'B "Y "%>'ÙG'i>" @, '©M" 6"Ù("% 'iM'ù> #150): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS
[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #150): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$=30M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.151. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('É@'É '9?'I@ò 8" M'Y8é?Y>•ù •ù5•, "™A'©&, '©M" 6"Ù("% 'iM'ù> #151): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #151): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$=30M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.152. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (" H'i?R "Y "iM'Y?Y?>B 8é@'Y0'2A *"Ù0"iM', 8•) "Ù/"â 3 S"" •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #152): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$=30M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.153. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'UA'A "Ùé."ù\$ò %y 'iM' "Y '©M" ."y), '©M" 6"Ù("% 'iM'ù> #153): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #153): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y?>B 8é@'Y0'2A *"Ù0"iM', 8•) "Ù/"â 3 S"" •ù •™."Ù." &"Y>"
(Case:#QCS•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.154. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('Y©'iM'B *"Ù0"éG'ò %8"Ù" M"i0"y "é •" "Y 'É@"Y>•ù), '©M" 6"Ù("% 'iM'ù> #154): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #154): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$=30M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.155. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('%?" M'ig'>•) "EM'ù>é?I? •ù5• *"Ù0"Y#"I>, '©M" 6"Ù("% 'iM'ù> #155): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #155): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$=30M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.156. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'YK'9"ù\$ / •Y(" *é\$ •ù5•" 8" M"Y8'é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #156): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #156): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$=30M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.157. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (•©'©>•, Y'IB' @, '©M" 6"Ù("% 'iM'ù> #157): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS
[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #157): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y?>B 8é@'Y0'2A *"Ù0"iM', 8•) "Ù/"â 3 S"" •ù •™."Ù." &"Y>"
(Case:#QCS•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,~)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.158. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM'Y?"Y?>é@ò "U7"ys"Ù0'é?I?, '©M" 6"Ù("% 'iM'ù> #158): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #158): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$=30M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

Q.159. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?"Y?"Y?>é@ò "U7"ys"Ù0'é?I?, '©M" 6"Ù("% 'iM'ù> #159): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #159): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$=30M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"(è\$M'Y%M'IB0é@'Y0'2F-ÖR'À%ÇW7VÇB'AWBéd'7a2pù0'èa\$è0'Qua#è(è(Ca
(Case:#QCS•

- Q.160.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (•i 'Y!'É> "%?" M"Y ' , %, "é0 'yM" >'2Â ""é '©>" M'd'Â ""'Ù0"iM', 8•) "Ù/"â 3 c " •ù .
[English]: In SSC CGL Tier-1 context (Data Interpretation (Bar Graphs, Pie Charts, Tables), Problem #160): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'é>"% •i""é0'ÙB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.161.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("%" 'iM'ù> '©M" #é2" "Y 'Y?'ù>'ÉM'ù\$ "âÂ ""'Ù0"iM', 8•) "Ù/"â 3 c " •ù •TM."Ù." •
[English]: In SSC CGL Tier-1 context (Number Systems & Divisibility, Problem #161): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) '©M" \$"ù8"Ù%"é*, ""'Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>'©?B 'iM'Y?'I@) •i""é0'ÙB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.162.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('âé8. 'çâ •ù5" •2. "é* •Y (" "U"ùK'â ""'Ù0"iM', 8•) "Ù/"â 3 c "" •ù •TM."Ù." &"Y>" •
[English]: In SSC CGL Tier-1 context (HCF and LCM Applications, Problem #162): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) 'é>"% •i""é0'ÙB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.163.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("%"0)"@ 'Y0'2 'Y "%>'I>)" "R " # " Â ""'Ù0"iM', 8•) "Ù/"â 3 c2" •ù •TM."Ù." &"Y>" •
[English]: In SSC CGL Tier-1 context (Simplification & Surds & Indices, Problem #163): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) 'i(" ""é\$ "ù '%M'ùB"%@ 'Y0'2 "é0'R &"Ù5"é0"â ""'Ù0"é""Ù\$ '©0B) #é: "%M'ùB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.164.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('©M" \$"ù6"\$â "Y 'YM" .ù "©0"ù5" M' (, '©M" 6"Ù("% 'iM'ù> #164): •ù •TM."Ù." •
[English]: In SSC CGL Tier-1 context (Percentages & Successive Change, Problem #164): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'é>"% •i""é0'ÙB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.165.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (">'ÒÂ 9"é("ô "Y 'E "U "â ô •) "ù\$ 'éB")M'òÂ ""'Ù0"iM', 8•) "Ù/"â 3 cR" •ù •TM."Ù." •
[English]: In SSC CGL Tier-1 context (Profit, Loss & Discount / Marked Price, Problem #165): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) '©M" \$"ù8"Ù%"é*, ""'Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>'©?B 'iM'Y?'I@) •i""é0'ÙB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.166.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("%">'y>" # •ù5" •Y M"5'9&"U"ò , "Ù"é •Y 'i0, '©M" 6"Ù("% 'iM'ù> #166): •ù •TM."Ù." •
[English]: In SSC CGL Tier-1 context (Simple & Compound Interest Differences, Problem #166): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) 'é>"% •i""é0'ÙB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.167.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("Y(" ""é\$, "%."é(" ""é\$ "ù5" •. "ù6"Ù0'2Â ""'Ù0"iM', 8•) "Ù/"â 3 cr" •ù •TM."Ù." &"Y>" •
[English]: In SSC CGL Tier-1 context (Ratio, Proportion & Mixture / Alligation, Problem #167): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) 'i(" ""é\$ "ù '%M'ùB"%@ 'Y0'2 "é0'R &"Ù5"é0"â ""'Ù0"é""Ù\$ '©0B) #é: "%M'ùB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.168.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("Y>" M'òÂ 8'è •ù5" ("U •) "Â ""'Ù0"iM', 8•) "Ù/"â 3 c, " •ù •TM."Ù." &"Y>" ("ù0"Ù •
[English]: In SSC CGL Tier-1 context (Time, Work, Pipes & Cisterns, Problem #168): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'é>"% •i""é0'ÙB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
- Q.169.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('©>" "Â 8'è, "iB" @, "G") "é!"É@ •ù5" ("é5"y>" >, '©M" 6"Ù("% 'iM'ù> #169): •ù •
[English]: In SSC CGL Tier-1 context (Speed, Time, Distance & Trains / Boats, Problem #169): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?
A) '©M" \$"ù8"Ù%"é*, ""'Ù0'ég'ò &"Ù5"é0"â 8'iM'ù>'©?B 'iM'Y?'I@) •i""é0'ÙB'B *"%0"é." " "r" '%0A" B'¢ "" ?'Y2"ù\$ 'é>', %8B)%(U\$ éS"â 332M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"¹ 'YG •Y(" 0")* •Y-" 7"ù) "r²"(éS"iM"Y%M' (B)6@'Y0'F- "é0'R6Q'U'Fé0'â 6"U0'G"•"Ú" "©0'E#V0'Q'W'F" (Case :#QCF#) •

Q.170. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('I8'B "Y "%>'ÙG'i>" @, '©M" 6"Ù("% 'iM'ù> #170): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #170): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.171. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('É@'É '9?'I@'ò 8" M'Y8é?Y>•ù •ù5•" , "™A'©&, '©M" 6"Ù("% 'iM'ù> #171): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #171): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@ 'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.172. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (" H'i?R "Y "iM'Y?Y>B 8é@ 'Y0'2A *"Ù0"iM', 8•) "Ù"â 3 s"" •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #172): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.173. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'ÚA" "Ù"é.ù\$ò % "Y 'iM' "Y '©M" ."y), '©M" 6"Ù("% 'iM'ù> #173): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #173): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y>B 8é@ 'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.174. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('Y©'iM'B *"Ù0"éG'ò %8"Ù"â *"i0"y "é •" "Y 'É@"Y>•ù), '©M" 6"Ù("% 'iM'ù> #174): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #174): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.175. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('%?" M'iG'>•) "EM'ù>é?I? •ù5•" *"Ù0"Y#I>, '©M" 6"Ù("% 'iM'ù> #175): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #175): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@ 'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.176. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'YK'9 "ù\$ / •Y(" *é\$ •ù5•" 8" M'Y8'é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #176): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #176): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.177. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (•© '©>•, Y 'iB' @, '©M" 6"Ù("% 'iM'ù> #177): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #177): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y>B 8é@ 'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+346M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.178. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM'Y?"Y>é@'ò "Ù7"ys"Ù0'é?I?, '©M" 6"Ù("% 'iM'ù> #178): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #178): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+356M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+356M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.179. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?"Y?"é@'ò "Ù7"ys"Ù0'é?I?, '©M" 6"Ù("% 'iM'ù> #179): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS

[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #179): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@ 'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$+356M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+356M'Dat@ù0'2alté0'Raçhù5géantù0'èa\$SineQa#amé(% (C# Case:#Q\$3#

Q.190. [English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #190): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.191. [English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #191): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.192. [English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #192): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.193. [English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #193): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.194. [English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #194): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.195. [English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #195): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.196. [English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #196): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.197. [English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #197): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.198. [English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #198): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.199. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-1 “% ‘i0’Ù- (‘iM’ ?“Y?é@ò “U7y\$“U0’é?‘i?, ‘©M’ 6“Ù(“% ‘iM’ù> #199): •ù •™.”Ù.” &“Y>“
 [English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #199): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.200. [English]: In SSC CGL Tier-1 context (Data Interpretation (Bar Graphs, Pie Charts, Tables), Problem #200): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) ÉM'ù>é?I@'dù • 'y'ñ'ù\$ / '©M' \$'ù,.'''1 'YG-Y(' 0')* •Y-.' 7''B'772('ESM'Y%M'UB'%)@Y0Y2-Ç0B8CUB'60àà000Cé'±àvôCû'V60V0M1

C) 'é>'%o •i''60'UB'B *''%o0'."é." " "r '%0A" B'ç *"?Y2'ù\$ 'é>,'%0800'ç'ù\$ 'é>2000'"ècàlculatèd'vâluè'stâd'chîng'întârdâd'Basè'l'è'pàrà'mè'tè'r(C

theorem (Case: CO)

Q-201. [English]: In SSC CGL Tier-1 context (Number Systems & Divisibility, Problem #201): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) ©M" \$"u8"U%"é", **U0'ÉG'd &"U5"é0"a 8'IM"ù>©?B 'im"Y?"I @)à)"083"uéé %mE@%-%@Y0'2'ÉOR R"05V6e-a VR0-g"U5V6C6MF-6A.MR
C) ÉM"ù>é?"I@d"ù •)y#"us" / ©M" \$"ù,) "1 'YG-Y(0")* -Y- 7" (Case_#082?)
parameter (Case : OC"

Q.202. [English]: In SSC CGL Tier-1 context (HCF and LCM Applications, Problem #202): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) <é>%0•i"é0'ÜB'B *%00"é." " "r %0%A' B'ç *?"Y2"ù\$'é,>, %08B)KfL\$'éS+4%M @RkZu@eV Valt6RaChif6tardüD'eäSineQparamé(%.C)
C) 'ÉM'ù>'é?I@'dù •) 'y#''ù\$' / '©M' \$'ù,)™¹ 'YG·Y(" 0")* ·Y-"7"m @r(Smãvñz,3óJÚ0ëE-öJ5ëcQmF-Hdaëx-Q-B-am'V2#@v0)vDvG'
theorem.(Case iCQ

Q.203. [English]: In SSC CGL Tier-1 context (Simplification & Surds & Indices, Problem #203): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) •(• " *\$ "ù %m'ùB%o@'Y0'2 "é0'R &"Ù5'é0'à *"Ü0"é*"Ü\$ @0B#f6% %m'ùV9PUB:B0C%0&W7VCB5A" Bed'via'120postoralredUerofFactor
C) 'ÉM'ù>'é?'I@'dù •) 'y#"\$ / 'CM" \$ "ù,)")¹ 'YG·Y(" 0")* ·Y· 7")@'E(Šà0D%2,30)Ü0ëG'5-òÜ5èu,M'Hàis?Q -id'Y'24@vvo)DWë'
theorem(Case :CO_R :

Q.204. [English]: In SSC CGL Tier-1 context (Percentages & Successive Change, Problem #204): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

C) 'é>'%o · i""é0'ÜB'B *""o0"é." " "r" %oA" B'¢ *"" ?'Y2"üS 'é>," %8ü0G#Q6" #B=409" "Ecalculated value is 160 using standard Baseline parameters (Case #CO #·

Q.205. [English]: In SSC CGL Tier-1 context (Profit, Loss & Discount / Marked Price, Problem #205): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) ©M" \$ "u8"U%"é", **U0'éG'd & "U5'é0"a 8'IM'ù">©?B 'iM"Y?"I @)èi"0083"ùé %M&B"%@Y0'2 "E0R"R'U5'v6-a "V0"éFUM000éWf"XaMe
C) 'ÉM'ù">é?"I@'dù •) 'y#u\$ " / '©M" \$ "ù,•)"1 'YG•Y(" 0")* •Y- 7" (D) 42.2%©M0UB'B07%#0'6.÷ F-0'À6AÇMF*5a'82u\$-éæ; %8'U%V000V6-
parameter (Case: 00

Q.206. [English]: In SSC CGL Tier-1 context (Simple & Compound Interest Differences, Problem #206): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) 'é'>% %i*'é0'ÜB'B *%00'ë." " "r '%0%A' B'ç ?*"?'Y2'ù\$ 'é', %8B)4ü\$ 'é'è=412M DAB@at0 Valt0 RaChif56grandü0 bA55ireQuaméte%. (C
C) 'ÉM'ù>'é'?I@'dù •)'y'h'ù\$ / '©M' \$'ù,)')¹ 'YG·Y('0")* ·Y-7"U@e0's0ndVn?%,30'jue6G→R-O56W6Wmà66Bç-imér 346Pv)üÜVG
theorem (Case #CO_3.

Q.207. [English]: In SSC CGL Tier-1 context (Ratio, Proportion & Mixture / Alligation, Problem #207): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) •(• " "é\$ù" %0%ùB"%@'Y0'2 "éú'R &"Ù5'é0'à *""Uò"é""Ú\$ '©0B"#6: "%0%"V#ÉPUB;BÓC%%00eZW7VCBFAVB/d"rAyp2qpsreñha&EwLcDòS"facto
C) 'ÉM'">'é?.'I@'dù •) 'y#"ú\$" / '©M" \$"\$ù,•)"¹ 'YG•Y(" 0")* •Y-" 7"b) ©B["S&b0%2%;3;7fD0'èG'F-ÇÜ\$'60çWFMa-60?8 -heIVZ4@_v9vQ0WE6

parameter (Case : OC 3R)
theorem (Case : CQ 3R)

Q.208. [English]: In SSC CGL Tier-1 context (Time, Work, Pipes & Cisterns, Problem #208): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

[illegible]

Q.209. [English]: In SSC CGL Tier-1 context (Speed, Time, Distance & Trains / Boats, Problem #209): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) 'OM" \$ "ù8"Ù0%é*", **U0'ÉG'd & "U5'é0"â 8'IM'ù>'©?B 'im"Y?'I@)à)"08\$ué %M&R"%@'Y0'2'ÉÓR&RÚ5'Vee-á *U0'-çFVWV'06ôç W'R"-6aMe
C) 'ÉM'ù>'é?'I@'òù •) y# "ù\$ / 'OM" \$ "ù,•)"¹ 'YG • Y(" 0")* • Y-" 7"ù) e22%&V0'UB'B'6%30'a ÷ 'F-Ô %6NÇWF=ua b2xú\$ -ç p%9"0%yV0DWS
parameter (Case ; OC ,

Q.210. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('I8'B "Y "%>'ÙG'>" @, '©M" 6"Ù ("%" 'iM'ù> #210): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS
[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #210): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+42M'DatSou@Y0'2alté0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.211. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('É@'É '9?'I@'ò 8" M'Y8é?Y>•ù •ù5•, "™A'©&, '©M" 6"Ù ("%" 'iM'ù> #211): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #211): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù "%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â "ù0"é**ù\$ '©0B)%(ù\$ é\$+42M'DatSou@Y0'2alté0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.212. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (" H'i?R "Y "iM'Y?Y>B 8é@'Y0'2A *"ù0'iM', 8•) "ù/"â 3# " " •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #212): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+42M'DatSou@Y0'2alté0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3

Q.213. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'UA'A "ù"é.ù\$ò "% 'y 'iM' "Y '©M" ."y), '©M" 6"Ù ("%" 'iM'ù> #213): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #213): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"ù0'ég'ò &"ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y>B 8é@'Y0'2A *"ù0'iM', 8•) "ù/"â 3# " " •ù •™."Ù." &"Y>"
(Case:#Qc3
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.214. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('Yc'iM'B *"ù0'ég'ò "%8"ù\$ "M'io'y "é •" "Y 'É@'Y>•ù), '©M" 6"Ù ("%" 'iM'ù> #214): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #214): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+42M'DatSou@Y0'2alté0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.215. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('%?" M'ig'>•) "EM'ù>é?I? "ù5• *"ù0"Y#I>, '©M" 6"Ù ("%" 'iM'ù> #215): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #215): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù "%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â "ù0"é**ù\$ '©0B)%(ù\$ é\$+42M'DatSou@Y0'2alté0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.216. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'YK'9"ù\$ / •Y(" *é\$ •ù5•" 8" M'Y8'é?Y>•ù , '©M" 6"Ù ("%" 'iM'ù> #216): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #216): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+42M'DatSou@Y0'2alté0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.217. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('©' ©>•, Y 'iB' @, '©M" 6"Ù ("%" 'iM'ù> #217): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS
[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #217): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"ù0'ég'ò &"ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y>B 8é@'Y0'2A *"ù0'iM', 8•) "ù/"â 3# " " •ù •™."Ù." &"Y>"
(Case:#Qc3
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.218. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM'Y?"Y>é@'ò "U7"ys"ù0'é?I?, '©M" 6"Ù ("%" 'iM'ù> #218): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #218): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+42M'DatSou@Y0'2alté0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.219. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?"Y?"é@'ò "U7"ys"ù0'é?I?, '©M" 6"Ù ("%" 'iM'ù> #219): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #219): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù "%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â "ù0"é**ù\$ '©0B)%(ù\$ é\$+42M'DatSou@Y0'2alté0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB) "2"é\$M'Y%M'IBÓf@'Y0-2F"é0'RaChù5gérantù0'èa5iReQuàrè(è(CA
(Case:#Qc3b

Q.230. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('I8'B "Y "%>'ÙG'i>" @, '©M" 6"Ù("% 'iM'ù> #230): •ù •™."Ù." &"Y>" ("ù0"Ù"é

[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #230): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.231. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('É@'É '9?'I@'ò 8" M'Y8é?Y>•ù •ù5•" , "™A'©&, '©M" 6"Ù("% 'iM'ù> #231): •ù

[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #231): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.232. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (" H'i?'R "Y "M'Y?>B 8é@'Y0'2A *"Ù0"iM', 8•) "Ù"â 3#3"" •ù •™."Ù." &"Y>

[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #232): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.233. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'UA'A "Ù"é.ù\$ò "% 'y 'iM' "Y '©M" ."y), '©M" 6"Ù("% 'iM'ù> #233): •ù

[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #233): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.234. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('Yc'iM'B *"Ù0"éG'ò "%8"Ù" M'io'y "é •" "Y 'É@"Y>•ù), '©M" 6"Ù("% 'iM'ù> #

[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #234): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.235. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('%?" M'ig'> "ÉM'ù>é?I? "ù5•" *"Ù0"Y#I>, '©M" 6"Ù("% 'iM'ù> #235): •ù •™.

[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #235): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.236. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'YK'9"ù\$ / •Y(" *é\$ •ù5•" 8" M'Y8'é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #236): •

[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #236): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.237. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (•©'©>•, Y 'iB' @, 'Y 'Ù("% 'iM'ù> #237): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ù\$

[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #237): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.238. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM'Y?"Y?é@'ò "U7"ys"Ù0'é?I?, '©M" 6"Ù("% 'iM'ù> #238): •ù •™."Ù." &"Y>"

[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #238): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.239. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?"Y?"é@'ò "U7"ys"Ù0'é?I?, '©M" 6"Ù("% 'iM'ù> #239): •ù •™."Ù." &"Y>"

[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #239): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©0B)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ é\$+46M'Dat@Y0'2alté0Rat'ù5é\$antù0'bas\$ineQ'paramé(%)C (Case:#Qc")

Q.240. [English]: In SSC CGL Tier-1 context (Data Interpretation (Bar Graphs, Pie Charts, Tables), Problem #240): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

[illegible]

Q.241. •ù8•ù8"%@ "%%@'É@•ú2 'ù?'ù0-1 "%o 'ìO"Ù- ("%" 'ìM'> ("%" #'è2" 'Y?'"Y>'ÉM'ù\$"âÂ *")U0"ìM', 8•)"Ù"/à 3#C " •ù •™."Ù

[English]: In SSC CGL Tier-1 context (Number Systems & Divisibility, Problem #241): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

C) ÉM'ù>é?|@òù •)y#ù\$ / '©M" \$ù,.)^{m1} 'YG-Y("0")*•Y- 7"ù (Case #9?) e2(%%)Me02UB B0%B"è.- F-0 À%A&BWf+ba v0 u\$ -é-z,p-s-E%vD0VWG

Q.242. •û8•û8“%o@ “%o@‘É@•û2 ‘û?’û0-1 “%o ‘i0”Û- (‘âë8.‘çà•û5• “2.‘ë.”•Y(“”û0“ûK‘rÂ*”û0“iM”,8•)”)Û/“â3#C“““•û •™.”Û.”&“Y>“
 [English]: In SSC CPL Tier-1 context (HCF and LCM Applications, Problem #242): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

C) ÉM'u>é?I@`ùò`' y#ù\$ / `©M` \$ù,.)!¹` 'YG·Y(° 0")\* ·Y- 7"Þ` ©²` (\$àB ÒVñ ?;çó` çòéS ð-ÖÜ566CÀF-Hdà68xQB -wV?²#@ v/vÞDŮG "e

Q.243. •û8•û8"%@ "%%@'É@•û2 'û?'û0-1 "% 'i0"Û- ("%0")@Y02 "Y "%>|>•) "R " #" Â ""Û0"iM", 8•) "Û/"â 3#C2" •ù •™."Û." &"Y>

[English]: In SSC CGL Tier-1 context (Simplification & Surds & Indices, Problem #243): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

[illegible]

Q.244. •ù8•ù8"%@ "%%@'É@•ù2 'ù?'ù0-1 "%o 'i0'Ù- ('©M" \$"ù6'ì\$a "Y YM".ù "©0"ù5" M'ì(, '©M" 6"Ù("%o 'iM'ù> #244): •ù •™."Ù.
[English]: In SSC CGL Tier-1 context (Percentages & Successive Change, Problem #244): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) ÉM'ù>é?I@'òù •) y#“ù\$ / '©M“ \$“ù,•)!”¹ ‘YG •Y(” 0”)• •Y- 7”B“)7²*(%SM Y%M’IBó#@‘y9Z -ÉÔÂF&QWFeôâ sUÜQç~“!\$ @9H#vëoÖWN¹
C) é>% •i!“60ÜB'B *%%0“é.” “ ”r %œA“ B¢ ?”Y2“ù\$ é>, %8D)PúS \$ 1e489)cCalculated value from a single standard baseline parameter (Ca

Q.245. *[English]: In SSC CGL Tier-1 context (Profit, Loss & Discount / Marked Price, Problem #245): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?*

A) '©M" \$ "ù8"Ù%"é*", **"Ù0"éG"ò &"U5"é0"â 8"IM"ù>"©?B "iM"Y?"I©"àì"%"08\$Hé"R M&B"%"©"Y0"2 "©0R"8R"U6"é0"â*"V0"éFUS"W0"06"0C"1A"7"56A"78"
(Case: 0033R)
C) 'ÉM"ù>"é?"I©"à"òù"•) "y#"ù#" /"©M" \$ "ù,")"1 "YG"Y("0")* "Y-"7"U) "e2"%"00V0"UB"BC"%"0é" F-ÖR"À"60C"BF"U"2"06"2"U\$ "é"æ"%"8F"U"0"6"0C"1A"7"56A"78"

Q.246. •û8•û8"%@ "%%@'É@•û2 'ù?'ù0-1 "% 'i0'Û- ("%%>'y"> #•u5•" 'YM" 5"9&"U"ò , "Û"/é •Y 'l0, '©M" 6"Û("% 'iM'ù> #246): •ù •™
 [English]: In SSC CGL Tier-1 context (Simple & Compound Interest Differences, Problem #246): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) <é>% "i""é0`UB'B \*"%0<é." " "r "%A" B'¢ ¨"?Y2`ù\$<é>,%8B)%(f"\$ésè+492M'Qakub@eV2at6Rachib5gstrandùd'basineQaramé(%Cá
C) `ÉM`ù><é?I@`du.) 'y"#ù\$ /'©M" \$`ù,)"¹ 'YG·Y(" 0")* ·Y- 7"ü) ©(2("\$abUv%:36"üB @G-FÜ\$60QWVnà-60×E -üey74@,úvvôwce

Q.247. •û8•û8"%@ "%%@'É@•û2 'û?'û0-1 "%o 'i0'Û- (•Y(" **e\$, "%o."e("**e\$•û5• "•û6"Û0'2Â *•Û0'iM', 8•) ")Û/â 3#Cr" •û •™."Û." &"\
 [English]: In SSC CGL Tier-1 context (Ratio, Proportion & Mixture / Alligation, Problem #247): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

A) «!(“ “e\$“ù %0m'ùB%00@‘Y0’2 “é0'R &"Ù5“é0“â *”Ü0“é*”Ü\$ ‘00B”#6:%%0“ü0ÜB;BÓ5G0Ü&W7VÇBFAWBved“vaypöpsötiön%äü06tiöñ”facto
C) ‘ÉM“ù>é?I@‘dù.) ‘y#“ù\$ /‘CM“ \$“ù,.)”¹ ‘YG·Y(“ 0”)·*·Y- 7”Ü) @M(\$abB0%)%;3CÜj0&E5-ÖÜ5æcçmfHdaüsöçQB-är YZ²f@üv0)ÜVÜg“

Q.248. •û8•û8“%o@ “%o@É@•û2 ‘û?’û0-1 “%o ‘i0”Û- (‘Y>“ M’òÀ 8’e/•û5” (™U•) “À *™U0“iM, 8•) ”Û/“à 3#C, “ •û •™. ”Û.” &“Y>“ (“û0”

[English]: In SSC CGL Tier-1 context (Time, Work, Pipes & Cisterns, Problem #248): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

C) 'é>%o • i"é0'ÜB'B *%0o"E." " " r %oA" B'ç ?" ?Y2'ùS'é>," %80d%ÜS'6'1e49%" eC acTuaLed baUe'S'1e0'áing ísLanDaB'aíne ígárametü(Ce
theam Cæ-Öö

Q.249. •ù8•ù8"%"@ "%%@•É@•ù2 'ù?'ù0-1 "% 'i0"Ù- ('©=""Â 8'é/, 'iB' @, "G") é:'É@ •ù5•" ("é5-'y->" >, '©M" 6"Ù ("%" 'iM'ù> #249): •ù
 [English]: In SSC CGL Tier-1 context (Speed, Time, Distance & Trains / Boats, Problem #249): A candidate solves a problem
 where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed
 constraints?

A) 'OM" \$ "u8"U%"é", **U0'ég'ò & "U5'é0"â 8'IM"ù>©?B 'im"Y?"I@)ò("Y08"dué %M&B"%@'Y0'2' "é0R'&PU5'6000- **U0'éc'PU5'W00000V/F%00N'Y

parameter (Case ; UC ; •

Q.260. •ù8•ù8“%@@ “%@‘É@•ù2 ‘ù?’ù0-1 “% ‘i0”Ù- (•¡‘Y!‘É> ‘%0?“ M“Y ‘, %,“é0 ‘yM“ >²Â ““é ‘©>“ M‘d‘Â “”Ù0“iM‘, 8•) ”Ù/“â 3#c “ •ù
 [English]: In SSC CGL Tier-1 context (Data Interpretation (Bar Graphs, Pie Charts, Tables), Problem #260): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.261. [English]: In SSC CGL Tier-1 context (Number Systems & Divisibility, Problem #261): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.262. [English]: In SSC CGL Tier-1 context (HCF and LCM Applications, Problem #262): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.263. [English]: In SSC CGL Tier-1 context (Simplification & Surds & Indices, Problem #263): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.264. [English]: In SSC CGL Tier-1 context (Percentages & Successive Change, Problem #264): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.265. [English]: In SSC CGL Tier-1 context (Profit, Loss & Discount / Marked Price, Problem #265): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.266. parameter (Case: OS3.0) [English]: In SSC CGL Tier-1 context (Simple & Compound Interest Differences, Problem #266): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.267. [English]: In SSC CGL Tier-1 context (Ratio, Proportion & Mixture / Alligation, Problem #267): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.268. •ù8•ù8"%@ "%%@É@•ù2 'ù?ù0-1 "%0 'i0"Ù- ('Y>" M'ðÀ 8'e/ •ù5. ('"U.) " Å *•ù0"iM, 8•) "ù/"à 3#c," •ù •TM."ù." &"Y>" ("ù0"ù
 [English]: In SSC CGL Tier-1 context (Time, Work, Pipes & Cisterns, Problem #268): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

Q.269. [English]: In SSC CGL Tier-1 context (Speed, Time, Distance & Trains / Boats, Problem #269): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

parameter (Case , US3;•

Q.270. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('I8'B "Y "%>'ÙG'i>" @, '©M" 6"Ù("% 'iM'ù> #270): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS
[English]: In SSC CGL Tier-1 context (Averages, Weighted Mean & Partnerships, Problem #270): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.271. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('É@'É '9?'I@'ò 8" M'Y8é?Y>•ù •ù5•, "™A'©&, '©M" 6"Ù("% 'iM'ù> #271): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Basic Algebraic Identities & Polynomials, Problem #271): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©OB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.272. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (" H'i?'R "Y "M'Y?Y>•ù B 8é@'Y0'2A *"Ù0"iM', 8•) "Ù"â 3#s"" •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Linear & Quadratic Equations, Problem #272): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.273. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'UA'A "Ù"é.ùS"ò "% 'y 'iM' "Y '©M" ."y), '©M" 6"Ù("% 'iM'ù> #273): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Lines, Angles & Triangle Geometry (Centres), Problem #273): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.274. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('Y©'iM'B *"Ù0"éG'ò "%8"Ù0"y "é •" "Y 'É@'Y>•ù), '©M" 6"Ù("% 'iM'ù> #274): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Circle Theorems (Tangents, Chords, Secants), Problem #274): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.275. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ("%?" M'iG'>•) "EM'ù>é?I? "ù5•" *"Ù0"Y#I>, '©M" 6"Ù("% 'iM'ù> #275): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Coordinate Geometry & Distance / Slopes, Problem #275): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©OB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.276. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?'YK'9"ù\$ / •Y(" *é\$ •ù5•" 8" M'Y8é?Y>•ù , '©M" 6"Ù("% 'iM'ù> #276): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (Trigonometric Ratios, Identities & Values, Problem #276): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.277. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- (•©'©>•, Y 'iB' @, '©M" 6"Ù("% 'iM'ù> #277): •ù •™."Ù." &"Y>" ("ù0"Ù"é0"ùS
[English]: In SSC CGL Tier-1 context (Heights and Distances (Angle of Elevation/Depression), Problem #277): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) '©M" \$"ù8"Ù%"é*, *"Ù0"éG'ò &"Ù5"é0"â 8'iM'ù>©?B 'iM'Y?"Y?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.278. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM'Y?"Y?é@'ò "U7"ys"Ù0é?I?, '©M" 6"Ù("% 'iM'ù> #278): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (2D Mensuration (Triangles, Quadrilaterals, Circles), Problem #278): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) 'é>% •i"é0'ÙB'B *"%0"é." " "r '%A" B'¢ *"?Y2"ù\$ 'é>', %8B)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

Q.279. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-1 "% 'i0'Ù- ('iM' ?"Y?é@'ò "U7"ys"Ù0é?I?, '©M" 6"Ù("% 'iM'ù> #279): •ù •™."Ù." &"Y>"
[English]: In SSC CGL Tier-1 context (3D Mensuration (Cylinder, Cone, Sphere, Frustum), Problem #279): A candidate solves a problem where values follow standard proportional metrics. What is the mathematically accurate resultant value under prescribed constraints?

- A) •i(" *é\$ "ù '%M'ùB"%@'Y0'2 "é0'R &"Ù5"é0"â *"Ù0"é**"ù\$ '©OB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)
C) 'ÉM'ù>é?I@'òù •) 'y#"ù\$" / '©M" \$"ù,•)"1 'YG •Y(" 0")* •Y- 7"ùB)%(ù\$ 'é>+54M)Dafu@Y0'2alté0Rachù5géantù0'basSineQaamé(%)C3
(Case:#Qf#3)

C) 'é>%o •i"é0'ÜB'B *%00"é." " "r '%oA" B'¢ *"?Y2"ü\$ 'é>', %80ü\$ C=5560" Calculated value 5100 using standard Baseline parameters (Case #CQ.C •

A) "%@é' %%. 'l> 'Y@ '©A"yMù? 'Y0%G "Y"> > •y7U 'l. '©0'à "B%CM' \$"%")%2C-13/9 Optimal extraction of parameter confirming boundary es
C) '©M' ".y/ %mY?l?ùK•" "r \$"™\$ "Y?"IM")G"y#"\$é\$U.'R 0")* "%Gm?m?> "B"5%"%)%2C-13/9 Analytical improvement begins constant under
resolution (Root B=7)

[illegible]

A) '©M' '\$\$,.)' "%@'é>9 'YK '©B' >'YO'%G "Y">"%?" M'iG">B)/M'Y?e\$>'EB'Y#C32)(0'66-&E-(aVFR'd' afoibor'©MB (neúñg)úóést,añ'

B) resolution (Root A=21)

C) '©M' ".y/" %M'Y?'I?'ù.K" "r \$"™\$ "Y?"IM")G"y#"és'Ù.'R 0")* "%G %©'aP' %0>è'Y'©%)2'0+89 Y'A%Gricall'y' p'rov'èd integer a cò's s'at'e und'r equality (Root D=51)

A) '©M'."y/'%M'Y?'I?'ÜK" "r '\$™\$ "Y?'iM")G'y#'\$U.'R 0")* "BG'©M?'SWL,)'©%>'é%8")2'K>'SB' /Analytically prove the "logic" constant threshold limits (Root B=44)

C) 'iM"Y?'%>'B 'Y5"â "(" (%."é"é('iM"Y">">'©M">'©M'B ("ü6"UD)5%>'%, "2.74>28@ Define value of C over a quadratic/cubic equation equality (Root D=68)

[illegible]

C) '©M' ".y/ "%M?Y?'I?'ùK•" "r \$™\$ "Y?"IM")G"y#"\$ë".R 0")* "%Lg %& n^r %U>E?D(%C%)2?o+92) Y%A%Grically proven integer combinatorial boundary equality (Root A-119)

A) 'OM' .y/ "%mY?l?'ÜK" " r \$"#\$ "Y?"IM"G"y#"\$Ü.R 0")* "%B%ÖM'\$WU.·C%@@"e%"%)ZK=QB4)/X0dlyGälyprovanInveigixoxisyrht1esle
limits (Root B=89)
C) 'iM"Y?"%>'B 'Y5"â "(" (%o."é"é('iM"Y">" '>'OM" >'OM'B ("ü6"UD"5\$ee,%e.%).2.A=56D DefinevalvööGed via quadratic/calculus/resolution
boundary equality (Root D=136)

A) "%@<é>%."!>'Y@'©A"yMù?<'Y0%G<"Y">>.<y7"U'!..©0'à"B%CM"\$"U%)%2C=153/"Optical"extreme%parameter%Chirping%boundar\$\$(limits (Root B=99)
C)'©M"."y/"%M'Y?'I?<ùK·" "r"\$™\$<"Y?"IM")G"y#"\$ésÜ.'R 0")*<%GImo%mo> <B0<5"<a%("2C=117)IMAanalytical%proven%In(egal%constant,under resolution (Root A=63)

Q.290. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"U5"à "és" %" # •ù5•" 5"ù5"ù"Ù\$'Y0,"%."%M'ù> #10
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Quadratic Equations, Discriminant & Sign of Roots, Problem #10): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) 'iM'Y?'%>'B 'Y5'à "("%."é"é('iM'Y"> " '©M' >'©M'B ("ù6"ÙB)\$6F\$, \$ù.,)2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=110)
C) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root D=130)

Q.291. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"U0"ù" "YG 'Y&)"U0: •Y 'I 'Y&.)&"U0, '©0"ù "y 'iM" Â
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Triangle Centres: Incentre, Circumcentre, Orthocentre & Centroid, Problem #11): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M' "\$ù.,)" "%@'é>'9 'YK '©B" > 'Y0'%G "Y">)"> "%?" M'iG">) / iM'Y?'%>'B "Y5'à #) ("ù6"ÙB)\$6F\$, \$ù.,)2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root A=77)
C) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root D=143)

Q.292. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"Y M' @0 'IA'M'UA'Â "Y "%M'©0"U6" G'ì->'É@'Y> 'YK
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Cyclic Quadrilaterals & Tangent-Chord Angles, Problem #12): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=122)
C) 'iM'Y?'%>'B 'Y5'à "("%."é"é('iM'Y"> " '©M' >'©M'B ("ù6"ÙB)\$6F\$, \$ù.,)2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=132)

Q.293. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"9\$ U\$: •TM-"ù"ù7"Ü •Y(" 8"Ü* M'b "Y 'I?" M'ù "%M'©
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Circles: Common External & Transverse Tangents, Problem #13): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) "%@'é> "%."l> 'Y@ '©A"YM'ù? 'Y0'%G "Y">)"> •y7"Ü 'I. '©0'à "B"©M' "\$ù.,)"2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=143)
C) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=143)

Q.294. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"U0"ù"y6'é 'R "Ü'è."ù\$dó "%o0"" 0"y "é•" "Y •Y 'I'ì' Â
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Coordinate Geometry: Straight Lines, Collinearity & Intercepts, Problem #14): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) 'iM'Y?'%>'B 'Y5'à "("%."é"é('iM'Y"> " '©M' >'©M'B ("ù6"ÙB)\$6F\$, \$ù.,)2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=154)
C) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root D=182)

Q.295. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"U0"ù" "#ë?l@'ò '©M'©?"Y M' Ü("ù."Ü("ù7"Ü •ù5•" 8" M
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Trigonometric Maxima and Minima & Advanced Identities, Problem #15): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M' "\$ù.,)" "%@'é>'9 'YK '©B" > 'Y0'%G "Y">)"> "%?" M'iG">) / iM'Y?'%>'B "Y5'à #) ("ù6"ÙB)\$6F\$, \$ù.,)2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root A=105)
C) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root D=195)

Q.296. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"•e "ù5•" &)"0" %, "™A-'É?)&" *™U0"y "Ü7'2'Ä 8'è8"Ü/
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Heights and Distances with Multi-Point Observations, Problem #16): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=176)
C) 'iM'Y?'%>'B 'Y5'à "("%."é"é('iM'Y"> " '©M' >'©M'B ("ù6"ÙB)\$6F\$, \$ù.,)2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=176)

Q.297. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"U5'u5"ù." / YM"YGI M" .ù\$dó "Y '¹>ù>)" "ù\$ 'YM"YGI
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex 2D Mensuration & Shaded Region Areas, Problem #17): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) "%@'é> "%."l> 'Y@ '©A"YM'ù? 'Y0'%G "Y">)"> •y7"Ü 'I. '©0'à "B"©M' "\$ù.,)"2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=187)
C) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root D=222)

Q.298. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"U0"ù"Ü., '©?" >'é? " Â "ù("Ü'R "Y '©\$" 7"Ü+)", "%."%
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (3D Mensuration: Frustum, Prism, Pyramid & Tetrahedron, Problem #18): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) 'iM'Y?'%>'B 'Y5'à "("%."é"é('iM'Y"> " '©M' >'©M'B ("ù6"ÙB)\$6F\$, \$ù.,)2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root B=188)
C) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root D=234)

Q.299. @ "É@•ù? 'ù0-2 'y#"\$ / 'YM'y.'l> '©' "U7'a %&"e iM'ù? Y@:'é>yM'dÄ "é"Ü"ù "âÄ , "™A)" , '©M' 8" #
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Statistics: Mean, Median, Mode, Variance & Standard Deviation, Problem #19): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M' "\$ù.,)" "%@'é>'9 'YK '©B" > 'Y0'%G "Y">)"> "%?" M'iG">) / iM'Y?'%>'B "Y5'à #) ("ù6"ÙB)\$6F\$, \$ù.,)2"%@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root A=133)
C) '©M' ".y/ "%oM'Y?'I?'ùK*" "r \$"™\$ "Y?"iM")G"y#"é\$"Ü.'R 0)"* "%B %&@0#> Definite Value Derived Via Quadratic Calculus/resolution limits (Root D=247)

Q.300. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %*“Ù0“é/“ù”l> “%?“iM“y>•)\$. “%M“Y\$•)“\$”Ù0 “% “%>•ù •ù5•

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Probability Theory: Independent Events, Conditional & Cards/Dice, Problem #20): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=280> / “X0dly Gally” proven integral constant limits (Root B=220)
- C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B” \$%.”)2“%=(54)> / “Definite Value Derived via Quadratic Calculus Resolution boundary equality (Root D=340)

Q.301. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %5“e8“US“Y?“R 8•) “Ù/“â * “Ù0“9>“) @ •ù5•” “%M“%\$ “Y?“Ù.”

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Real Number Systems & Advanced Divisibility, Problem #21): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@“é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“Ù”l. ‘©0“â “B“©M“ \$“ù%”)2L=357 / “Optional extreme parameter confirming four (dare’s of limits (Root B=231)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=280> / “X0dly Gally” proven integral constant limits (Root B=231)

Q.302. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “A “é\$“é ‘R “Y “%\$“B “-ù“(“Ù, (“%。“%M“ù> #22): “%©

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Surds, Indices & Continued Fractions, Problem #22): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B” \$%.”)2“%=(54)> / “Definite Value Derived via Quadratic Calculus Resolution limits (Root B=242)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=280> / “X0dly Gally” proven integral constant boundary equality (Root D=374)

Q.303. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “ù?“ “Ù01“i\$ l> •ù5•” “Y.”)2“Ù/ , .”TM!“â 8“é8“Ù/“â 3#2“

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex Percentages & Depreciation Models, Problem #23): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” \$“ù,•) “%@“é>9 “YK ‘©B” > ‘Y0“%G “Y>“) > “%?“ M“iG“i>B) / “iM“Y?“%>B “Y5“â \$“ù%”)2L=429 / “Optional extreme parameter confirming four (dare’s of resolution (Root A=161)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=280> / “X0dly Gally” proven integral constant boundary equality (Root D=391)

Q.304. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %2“e--“TM>“%?, “EG“%。“é(‘iA“Y>“%&“é0 •ù5•” , “é“é0 “ #“%

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Profit, Loss, False Weight & Market Strategies, Problem #24): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=282> / “X0dly Gally” proven integral constant limits (Root B=264)
- C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B” \$%.”)2“%=(54)> / “Definite Value Derived via Quadratic Calculus Resolution boundary equality (Root D=408)

Q.305. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “YM” 5“9&“U“ù , “Ù/“é (•Y0“Ù“Y>“ M“y?“Rù\$”Ù0“%。“é8“ù

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Compound Interest with Semi-Annual / Continuous Compounding, Problem #25): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@“é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“Ù”l. ‘©0“â “B“©M“ \$“ù%”)2L=429 / “Optional extreme parameter confirming four (dare’s of limits (Root B=275)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=282> / “X0dly Gally” proven integral constant boundary equality (Root D=442)

Q.306. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “ù6“Ù02 “Y “YM” .“ù “%>•)&“Ù0 l> ‘©0“ù5“ M“l(, (“%。“%

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Alligation & Advanced Mixtures with Sequential Dilution, Problem #26): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B” \$%.”)2“%=(54)> / “Definite Value Derived via Quadratic Calculus Resolution limits (Root B=286)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=282> / “X0dly Gally” proven integral constant boundary equality (Root D=442)

Q.307. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “é0“Ù/ •ù5•” 8“é/ (•ù “é l0“©>“ @ •ù5•” & “YM“y\$“â “)“\$” / “â

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Time & Work with Alternating Shifts & Efficiency Differentials, Problem #27): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” \$“ù,•) “%@“é>9 “YK ‘©B” > ‘Y0“%G “Y>“) > “%?“ M“iG“i>B) / “iM“Y?“%>B “Y5“â \$“ù%”)2L=429 / “Optional extreme parameter confirming four (dare’s of resolution (Root A=189)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=282> / “X0dly Gally” proven integral constant boundary equality (Root D=459)

Q.308. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %8“e“y “U7“©>“ “A 5“9\$“ù\$” / ‘ùM“ H“R “Y •ù8“Ù”y2“y “

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Relative Speed, Circular Tracks & Escalator Motion, Problem #28): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=284> / “X0dly Gally” proven integral constant limits (Root B=308)
- C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B” \$%.”)2“%=(54)> / “Definite Value Derived via Quadratic Calculus Resolution boundary equality (Root D=476)

Q.309. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % , “TM“A©&“éB““A “ #“% •)! ‘©M” .”y/ •ù5•” 8“é.“ù\$ l2, “â 8“é

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Polynomial Roots, Factor Theorem & Symmetric Functions, Problem #29): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@“é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“Ù”l. ‘©0“â “B“©M“ \$“ù%”)2L=493 / “Optional extreme parameter confirming four (dare’s of limits (Root B=319)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL”)©0“@“é%9”)2K=287> / “X0dly Gally” proven integral constant boundary equality (Root A=203)

[illegible]

Q.320. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %*“Ù0“é/“ù l> “%?“iM“y>•)\$. “%M“Y\$•)\$. “Ù0 “% “%>•ù •ù5•

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Probability Theory: Independent Events, Conditional & Cards/Dice, Problem #40): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé limits (Root B=440)
C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B”\$“ù”)2“%>294> / “Definite“value“derived“via“quadratio“calcul“O“résolutio boundary equality (Root D=680)

Q.321. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %5“e8“US“Y?“R 8•) “ù/“â * “Ù0“9>“) @ •ù5•” ‘%M“%\$ “Y?“ù:

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Real Number Systems & Advanced Divisibility, Problem #41): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@“é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“ù l. ‘©0“â “B“©M“S“ù“)“2“%>699> / “Optional“extreme“parameter“confir“ing“our“datés“d limits (Root B=451)
C) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé resolution (Root A=287)

Q.322. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “A “é\$“é ‘R “Y “%\$“B “-“ù“(“ù, “%。“%M“ù> #42): ‘%©

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Surds, Indices & Continued Fractions, Problem #42): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B”\$“ù”)2“%>294> / “Definite“value“derived“via“quadratio“calcul“O“résolutio limits (Root B=462)
C) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé boundary equality (Root D=714)

Q.323. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “ù?“ “Ù01“? \$l> •ù5•” “Y.”)2“ù/ , .”TM!“â 8“é8“ù/“â 3C2

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex Percentages & Depreciation Models, Problem #43): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” \$“ù,•) “%@“é>9 ‘YK ‘©B” > ‘Y0“%G “Y>“) > “%?“ M“iG“i>B) / “iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B”\$“ù”)2“%>294> / “Definite“value“derived“via“quadratio“calcul“O“résolutio resolution (Root A=301)
C) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé boundary equality (Root D=731)

Q.324. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %2“e--“TM“>“%?, “EG“%。“é(‘iA“Y>“%&“é0 •ù5•” , “é“é0 “#“%

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Profit, Loss, False Weight & Market Strategies, Problem #44): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé limits (Root B=484)
C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B”\$“ù”)2“%>294> / “Definite“value“derived“via“quadratio“calcul“O“résolutio boundary equality (Root D=748)

Q.325. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “YM” 5“9&“U“ù , “ù/“é (•Y0“ù“Y>“ M“y?“Rù\$“ù0“%。“é8“ù

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Compound Interest with Semi-Annual / Continuous Compounding, Problem #45): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@“é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“ù l. ‘©0“â “B“©M“S“ù“)“2“%>699> / “Optional“extreme“parameter“confir“ing“our“datés“d limits (Root B=495)
C) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé resolution (Root A=315)

Q.326. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “ù6“U02 “Y “YM” .“ù “%>•)&“ù0 l> ‘©0“ù5“ M l(, “%。“%

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Alligation & Advanced Mixtures with Sequential Dilution, Problem #46): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B”\$“ù”)2“%>294> / “Definite“value“derived“via“quadratio“calcul“O“résolutio limits (Root B=506)
C) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé boundary equality (Root D=782)

Q.327. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “é0“ù/ •ù5•” 8“é/ (•ù “é l0 ‘©>“ @ •ù5•” & “YM“y\$“â •)\$. “ù

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Time & Work with Alternating Shifts & Efficiency Differentials, Problem #47): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” \$“ù,•) “%@“é>9 ‘YK ‘©B” > ‘Y0“%G “Y>“) > “%?“ M“iG“i>B) / “iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B”\$“ù”)2“%>294> / “Definite“value“derived“via“quadratio“calcul“O“résolutio resolution (Root A=329)
C) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé boundary equality (Root D=799)

Q.328. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %8“e“y “U7“©>“ “A 5“9\$“ù\$” / ‘ùM“ H“R “Y •ù8“ù “y2“y “

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Relative Speed, Circular Tracks & Escalator Motion, Problem #48): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé limits (Root B=528)
C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UBj\$©B”\$“ù”)2“%>294> / “Definite“value“derived“via“quadratio“calcul“O“résolutio boundary equality (Root D=816)

Q.329. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % , “TM“A“© & “éB““A “#“% •)! ‘©M” .”y/ •ù5•” 8“é.“ù\$ l2, “â 8“é

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Polynomial Roots, Factor Theorem & Symmetric Functions, Problem #49): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@“é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“ù l. ‘©0“â “B“©M“S“ù“)“2“%>699> / “Optional“extreme“parameter“confir“ing“our“datés“d limits (Root B=539)
C) ‘©M” .”y/ “%M“Y?“I?“ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$“ù.”R 0”) * “%B“©M“SVL”)“©0“@“é%9”)2K=520> / “X0dly“Gally“proven“integri“consist“lédé resolution (Root A=343)

- Q.330.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %&”U5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “U\$‘Y0, “%.”%M‘ù> #50
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Quadratic Equations, Discriminant & Sign of Roots, Problem #50): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“UB“\$6““ \$%.”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root B=550)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“650 / Asymptically proven integral constant under boundary equality (Root D=850)
- Q.331.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %\$“U0“ù- “YG ‘Y•)&”U0: ‘Y ‘l ‘Y••)&”U0, ‘©0“ù “y ‘iM“ ‘Â
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Triangle Centres: Incentre, Circumcentre, Orthocentre & Centroid, Problem #51): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù.,”) “%@‘é>‘9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?“ M‘iG“i>B) / iM“Y?“%>‘B“Y5“â C) (“ù6“ùB“\$6““ \$%.”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root A=357)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“663 / Asymptically proven integral constant under boundary equality (Root D=863)
- Q.332.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %\$“YM “@ 0 “IA “M‘UA‘Â “Y “%M‘©0“U6“ G‘i>-‘É@“Y> ‘YK
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Cyclic Quadrilaterals & Tangent-Chord Angles, Problem #52): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“663 / Asymptically proven integral constant under boundary equality (Root D=863)
C) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“UB“\$6““ \$%.”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root B=572)
- Q.333.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %\$“9\$“U\$: “•TM-ù(“ù7“U “Y(“8“U“* M“b “Y ‘l?“ M‘ù “%M‘©0
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Circles: Common External & Transverse Tangents, Problem #53): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) “%@‘é> “%.”l> ‘Y@ ‘©A“yM‘ù? ‘Y0“%G “Y>“) > •y7“U ‘l. ‘©0“â “B“©M“ \$“ù.,”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root A=583)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“669 / Asymptically proven integral constant under boundary equality (Root D=884)
- Q.334.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %“(ù0“U&“y6“é ‘R “ù/“é.“ù\$“ó “%0“““ 0“y “é•” “Y “Y ‘l‘i ‘Â
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Coordinate Geometry: Straight Lines, Collinearity & Intercepts, Problem #54): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“UB“\$6““ \$%.”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root B=594)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“702 / Asymptically proven integral constant under boundary equality (Root D=918)
- Q.335.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %\$“U0“ù “#‘é?“l@‘ò ‘©M‘©?“yM ‘ù(“ù.“ù(“ù7“U “ù5•” 8“ M
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Trigonometric Maxima and Minima & Advanced Identities, Problem #55): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù.,”) “%@‘é>‘9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?“ M‘iG“i>B) / iM“Y?“%>‘B“Y5“â C) (“ù6“ùB“\$6““ \$%.”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root A=385)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“725 / Asymptically proven integral constant under boundary equality (Root D=935)
- Q.336.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %• “é•ù5•” &“)0 “%.”TM-A-‘É•)&” * “U0“y “U72‘Â 8‘é8“U/“
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Heights and Distances with Multi-Point Observations, Problem #56): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“702 / Asymptically proven integral constant under boundary equality (Root D=935)
C) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“UB“\$6““ \$%.”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root B=616)
- Q.337.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %&“U5“ù5“ù. / ‘YM“yG‘iM“ “ù\$“ò “Y ‘i>“ù>“) “ù\$“YM“yG‘iM
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex 2D Mensuration & Shaded Region Areas, Problem #57): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) “%@‘é> “%.”l> ‘Y@ ‘©A“yM‘ù? ‘Y0“%G “Y>“) > •y7“U ‘l. ‘©0“â “B“©M“ \$“ù.,”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root A=627)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“741 / Asymptically proven integral constant under boundary equality (Root D=952)
- Q.338.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %\$“U0“ù U., ‘©?“ >‘é?“ Â “ù(“ù(R “Y ‘©\$“7“U+“) , “%.”%M
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (3D Mensuration: Frustum, Prism, Pyramid & Tetrahedron, Problem #58): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“UB“\$6““ \$%.”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root B=638)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“754 / Asymptically proven integral constant under boundary equality (Root D=986)
- Q.339.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “U7“à %8“é iMù?“Y@: ‘é>yM“òÂ “é“ù/“ù “âÂ , “TM-A“) , ‘©M“ 8“ #
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Statistics: Mean, Median, Mode, Variance & Standard Deviation, Problem #59): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù.,”) “%@‘é>‘9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?“ M‘iG“i>B) / iM“Y?“%>‘B“Y5“â C) (“ù6“ùB“\$6““ \$%.”)2“%=(76) / Definite Value Derived via Quadratic Calculus & Solution limits (Root A=413)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“ù.‘R 0”) * “%0% “©M“ “ù6“ù “ù>é“Y“©“Y2“0“4“767 / Asymptically proven integral constant under boundary equality (Root D=1003)

- Q.340.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"Ù0"é/"ù 'l> "%?"iM'y>•)\$. "%M"Y\$•)"\$ "Ù0 %%">•ù •ù5•
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Probability Theory: Independent Events, Conditional & Cards/Dice, Problem #60): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root B=660)
C) 'iM"Y?'%>'B 'Y5"â ") ("%."é"é('iM"Y>" > '©M" >'©M'B ("ù6"UB)S6r\$,Sù,*)2"Å=420) / Definite value derived via quadratic calculus resolution boundary equality (Root D=1020)
- Q.341.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"5"é8"US"Y?"R 8•) "Ù"/â *%"Ù0"9>")@ •ù5•" '%M"%\$ "Y?"Ù:"
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Real Number Systems & Advanced Divisibility, Problem #61): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) "%@'é> "%."l> 'Y@ '©A"yM"ù? 'Y0"%G "Y>")> •y7"Ù "l. '©0"â "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root B=660)
C) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root B=671)
- Q.342.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"é\$é"R "Y "%%\$B "-ù("Ù(, "%."%M"ù> #62): '%@
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Surds, Indices & Continued Fractions, Problem #62): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) 'iM"Y?'%>'B 'Y5"â ") ("%."é"é('iM"Y>" > '©M" >'©M'B ("ù6"UB)S6r\$,Sù,*)2"Å=420) / Definite value derived via quadratic calculus resolution boundary equality (Root B=682)
C) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root B=671)
- Q.343.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"ù?"ù?"ù01?"is l> •ù5•" "Y.")2"Ù/ , "TM!"Å 8"é8"Ù"/â 3c2"
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex Percentages & Depreciation Models, Problem #63): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M" \$"ù,•) "%@'é>'9 'YK '©B" > 'Y0"%G "Y>")> "%?" M'iG">B) / iM"Y?"%>'B "Y5"â ") ("%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root A=441)
C) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root D=1054)
- Q.344.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"2"é--"TM">%"?, "EG"%."é('iA"Y>%"&"é0 •ù5•" , "é"é0 " #%"
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Profit, Loss, False Weight & Market Strategies, Problem #64): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root B=704)
C) 'iM"Y?'%>'B 'Y5"â ") ("%."é"é('iM"Y>" > '©M" >'©M'B ("ù6"UB)S6r\$,Sù,*)2"Å=420) / Definite value derived via quadratic calculus resolution boundary equality (Root D=1088)
- Q.345.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"YM" 5"9&"U"ò , "Ù"/é (•Y0"Ù"Y>" M"y?"Rù\$"Ù0"%."é8"ù"
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Compound Interest with Semi-Annual / Continuous Compounding, Problem #65): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) "%@'é> "%."l> 'Y@ '©A"yM"ù? 'Y0"%G "Y>")> •y7"Ù "l. '©0"â "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root A=455)
C) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root B=715)
- Q.346.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"ù"ù02 "Y "YM" .ù "%>•)&"Ù0"l> '©0"ù5" M'l(, "%."%
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Alligation & Advanced Mixtures with Sequential Dilution, Problem #66): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) 'iM"Y?'%>'B 'Y5"â ") ("%."é"é('iM"Y>" > '©M" >'©M'B ("ù6"UB)S6r\$,Sù,*)2"Å=420) / Definite value derived via quadratic calculus resolution boundary equality (Root B=726)
C) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root D=1122)
- Q.347.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"é0"Ù/ •ù5•" 8"é/ (•ù "é"l0 "©"> @ •ù5•" &'YM"y\$"â •)"\$ 'Å"
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Time & Work with Alternating Shifts & Efficiency Differentials, Problem #67): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M" \$"ù,•) "%@'é>'9 'YK '©B" > 'Y0"%G "Y>")> "%?" M'iG">B) / iM"Y?"%>'B "Y5"â ") ("%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root A=469)
C) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root D=1139)
- Q.348.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%"8"é"y"U7"©>"Å 5"9\$"ù\$" / 'ùM" H'R "Y •ù8"Ù"y2"y "
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Relative Speed, Circular Tracks & Escalator Motion, Problem #68): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root B=748)
C) 'iM"Y?'%>'B 'Y5"â ") ("%."é"é('iM"Y>" > '©M" >'©M'B ("ù6"UB)S6r\$,Sù,*)2"Å=420) / Definite value derived via quadratic calculus resolution boundary equality (Root D=1156)
- Q.349.** •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y##"ù\$" / 'YM"y."l> '©0" "U7"â %%", "TM"Å"© &"éB"Å " #%"•)! '©M" ."/ •ù5•" 8"é."ù\$"12', Å 8"é"
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Polynomial Roots, Factor Theorem & Symmetric Functions, Problem #69): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) "%@'é> "%."l> 'Y@ '©A"yM"ù? 'Y0"%G "Y>")> •y7"Ù "l. '©0"â "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits (Root B=759)
C) '©M" ."/ "M"Y?'l?'ùK•" "r \$"™\$ "Y?"iM")G"y##"é\$Ù."R 0")* "%B"©M"SVL•)©0@'é%9)YK=80) / X00lyCally proven integral constant limits resolution (Root A=483)

Q.360. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %*“Ù0“é/“ù ‘l> “%?“iM“y>•)\$. “%M“Y\$•)“\$”Ù0 “% “%>•ù •ù5•

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Probability Theory: Independent Events, Conditional & Cards/Dice, Problem #80): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB40) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=880)
- C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UB)“©M“SVL•)“©0“@“é%9”)YK=QB40) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=880)

Q.361. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %5“é8“US“Y?“R 8•) “Ù/“â * “Ù0“9>•)@ •ù5•” “%M“%\$ “Y?“Ù.”

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Real Number Systems & Advanced Divisibility, Problem #81): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@‘é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“Ù ‘l. ‘©0“â “B“©M“SVL•)“©0“@“é%9”)YK=QB40) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=880)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB40) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=880)

Q.362. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “é\$“é ‘R “Y “%\$“B “-ù“(Ù, “%。“%M“ù> #82): “%©

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Surds, Indices & Continued Fractions, Problem #82): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UB)“©M“SVL•)“©0“@“é%9”)YK=QB40) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=902)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB40) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=902)

Q.363. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “ù?“ “Ù0“1?“i\$ l> •ù5•” “Y.”)2“Ù/ , .TM!“! “â 8“é8“Ù/“â 3f2“

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex Percentages & Depreciation Models, Problem #83): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” \$“ù,•) “%@‘é>9 ‘YK ‘©B” > ‘Y0“%G “Y>“) > “%?“ M“iG“i>B) / iM“Y?“%>B “Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UB)“©M“SVL•)“©0“@“é%9”)YK=QB40) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root A=581)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB40) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root A=581)

Q.364. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %2“é--“TM>“%?, “EG“%。“é(‘iA“Y>“%&“é0 •ù5•” , “é“é0 “#“%©

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Profit, Loss, False Weight & Market Strategies, Problem #84): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB92) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=924)
- C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UB)“©M“SVL•)“©0“@“é%9”)YK=QB92) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=924)

Q.365. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “YM” 5“9&“U“ù , “Ù/“é (•Y0“Ù“Y>“ M“y?“Rù\$”Ù0“%。“é8“ù

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Compound Interest with Semi-Annual / Continuous Compounding, Problem #85): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@‘é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“Ù ‘l. ‘©0“â “B“©M“SVL•)“©0“@“é%9”)YK=QB92) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=935)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB92) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=935)

Q.366. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “ù“Ù02 “Y “YM” .ù “%>•)&“Ù0“l> ‘©0“ù5“ M“l(, “%。“%©

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Alligation & Advanced Mixtures with Sequential Dilution, Problem #86): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UB)“©M“SVL•)“©0“@“é%9”)YK=QB92) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=946)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB92) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=946)

Q.367. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % “é0“Ù/ •ù5•” 8“é/ (•ù “é ‘l0 ‘©>“ @ •ù5•” &YM“y\$“â •)“\$” ‘â

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Time & Work with Alternating Shifts & Efficiency Differentials, Problem #87): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” \$“ù,•) “%@‘é>9 ‘YK ‘©B” > ‘Y0“%G “Y>“) > “%?“ M“iG“i>B) / iM“Y?“%>B “Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UB)“©M“SVL•)“©0“@“é%9”)YK=QB45) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root A=609)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB45) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root A=609)

Q.368. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à %8“é“y “U7“©>“A 5“9\$“ù\$” / ‘ùM” H“R “Y •ù8“Ù “y2“y “

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Relative Speed, Circular Tracks & Escalator Motion, Problem #88): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB44) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=968)
- C) ‘iM“Y?“%>B ‘Y5“â “)(“%。“é““é(‘iM“Y>“ > ‘©M” >‘©M“B (“ù6“UB)“©M“SVL•)“©0“@“é%9”)YK=QB44) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=968)

Q.369. •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “U7“à % , “TM“A©& “éB““A “#“%•)! ‘©M” .”y/ •ù5•” 8“é.“ù\$ ‘12, “â 8“é

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Polynomial Roots, Factor Theorem & Symmetric Functions, Problem #89): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) “%@‘é> “%。“l> ‘Y@ ‘©A“yM“ù? ‘Y0“%G “Y>“) > •y7“Ù ‘l. ‘©0“â “B“©M“SVL•)“©0“@“é%9”)YK=QB13) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=979)
- C) ‘©M” .”y/ “%M“Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM”)G“y#“é\$”Ù.”R 0”) * “%B“©M“SVL•)“©0“@“é%9”)YK=QB13) ‘M“Algebraically proven Mixture conditions (antithesis) limits (Root B=979)

Q.380. •ù8•ù8“%_@ “%_@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.l’> ‘©0” ”Ù7“â %_*”Ù0“é/“ù ‘l’> “%_?’iM’y>•)\$. “%_M“Y\$.)\$”Ù0 ‘%_ ‘%_>•ù •ù5•

- A) '©M" . "y/ "‰M'Y'?!?'ùk•" "r \$"™\$ "Y?"iM")G"y#“és”Û.'R 0")* "%BG"™?SML")©%:©“é%”YK=1800 /N(Algebraically)proven Mitter constant and constraint limits (Root-B=1100)

Q.381. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM"y.'l> '©0" "U7"â "%5"é8"U\$"Y?"R 8.) "Ù/"â "*"Ù0'9>")@ •ù5•" '%m'%m\$ "Y?"Ù:

- [illegible]

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Surds, Indices & Continued Fractions, Problem #102): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

383. [English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex Percentages & Depreciation Models, Problem 03): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

Q.384. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM"y.'!> '©0" "Ü7"â "%2"é--"™>"%?, 'EG•%."é('iA'Y>"%&"é0 •ù5•", "é "é0 " #"%

- A) ©M" .y/ "%M'Y?I?üK•" "r \$™\$ Y?"iM")G"y#ésÜ.R 0")* "BG©M?SvLr")©.e@é%9")YK=QB52) /M%aQicay)proven Miedler constraints
C) iM"Y?"%>B 'Y5â ")(" %e"é(iM"Y">">©M" >©M'B ("ü6"UD)ü6ed, %e. 12. A = 726) Determine value of e derived via quadratic method or
boundary equality (Root B=1144)
boundary equality (Root D=1768)

conditions below:

- resolution (Root A=735)

Problem #106). Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below.

387. $y'' + 2y' + 2y = 0$ (Time & Work with Alternating Shifts & Efficiency)

A) %OM "\$u,") "%o@'e>'9 'YK 'CB" > 'Y0'%oG "Y>") > '%o?" M'iG"i>B)/M"Y?%&%'B"Y5G S)(%6d'e&E'&B)FR>+a@M'beoMaBe' resolution (RootA=749)

- [illegible]

C) 'iM"Y?"%>'B 'Y5"â "("%."é"é('iM"Y">" > '©M" >'©M'B ("ù6"UDY%ed,%s.%2.A=750 / Delimit value% derived via quadratic

- A) "%%@'é>'%%.'!>'Y@'©A'yM?ù?'YO'%G'Y>")>•y7"Ü'I.'©0'a *B%CM' "\$U%)")%(=953)'XOpna=>x0%Spa n e t e r s c o n f l i c t i n g b o u n d a r e s
C) 'CM' .'/y' "%MY?!ùK•" "r \$™\$ "Y?"iM')G"y#'\$Ü.R 0)* %Q f m s y M v > B0 'E\$ a %g()Zc el q(M)"Analytically prove B if larger than A and
resolution (Root A=763)

- Q.390.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #110
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Quadratic Equations, Discriminant & Sign of Roots, Problem #110): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1210)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root D=1210)
boundary equality (Root D=1210)
- Q.391.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #111
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Triangle Centres: Incentre, Circumcentre, Orthocentre & Centroid, Problem #111): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•)“ “%@‘é>‘9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?“ M‘iG“i>B) / “iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root A=777)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root D=1887)
boundary equality (Root D=1887)
- Q.392.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #112
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Cyclic Quadrilaterals & Tangent-Chord Angles, Problem #112): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1232)
C) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root A=777)
boundary equality (Root D=1904)
- Q.393.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #113
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Circles: Common External & Transverse Tangents, Problem #113): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) “%@‘é> “%.”l> ‘Y@ ‘©A“yM‘ù? ‘Y0“%G “Y>“) > •y7“Ù ‘l. ‘©0‘â “B“©M“ \$“ù,•)“ “%@‘é>‘9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?“ M‘iG“i>B) / “iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1232)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1243)
boundary equality (Root D=1904)
- Q.394.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #114
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Coordinate Geometry: Straight Lines, Collinearity & Intercepts, Problem #114): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1254)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root D=1938)
boundary equality (Root D=1938)
- Q.395.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #115
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Trigonometric Maxima and Minima & Advanced Identities, Problem #115): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•)“ “%@‘é>‘9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?“ M‘iG“i>B) / “iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root A=805)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root D=1955)
boundary equality (Root D=1955)
- Q.396.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #116
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Heights and Distances with Multi-Point Observations, Problem #116): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1276)
C) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root A=812)
boundary equality (Root D=1972)
- Q.397.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #117
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex 2D Mensuration & Shaded Region Areas, Problem #117): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) “%@‘é> “%.”l> ‘Y@ ‘©A“yM‘ù? ‘Y0“%G “Y>“) > •y7“Ù ‘l. ‘©0‘â “B“©M“ \$“ù,•)“ “%@‘é>‘9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?“ M‘iG“i>B) / “iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1287)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root A=819)
resolution (Root A=819)
- Q.398.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #118
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (3D Mensuration: Frustum, Prism, Pyramid & Tetrahedron, Problem #118): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1298)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root D=1933)
boundary equality (Root D=2006)
- Q.399.** •ù8•ù8“%@ “%@‘É@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.‘l> ‘©0” “Ù7“â %&”“Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #119
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Statistics: Mean, Median, Mode, Variance & Standard Deviation, Problem #119): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•)“ “%@‘é>‘9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?“ M‘iG“i>B) / “iM“Y?“%>‘B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6“ÙBù\$6“ \$%.”)2“%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root B=1233)
C) ‘©M“ .”y/ “%M‘Y?“I?“ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.‘R 0”)“ “%&@76) / “Definite Value Derived Via Quadratic Calculus Resolution (Root D=2023)
boundary equality (Root D=2023)

Q.400. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM"y.'l> '©0" "Ù7"â %%"Ù0"é/"ù 'l> "%%?'iM'y>•)\$: "%%M"Y\$.)\$"Ù0 '%% '%%>•ù •ù5•

A) '©M' '\$ù,•)' '%@é>9 'YK '©B' >'YO'%G 'Y>')>'%'?' M'iG">B)/M'Y?%\$'EB'Y&C S/'%605'8F'-Id FR'o adMtoodHtE'mcEtIn's cónstRin

C) '©M' ".y/ "%oM'Y'?I'?ùK-" "r "\$™\$ "Y?"iM")G"y#"é\$"Û:'R 0")* "%0-%0-14' %5.0>èY'(0%0)2 Y0-1807)%0-ally tReHy·proven in dge'r' còd s'ant'uñk
resolution (Root A=973)
boundary equality (Root D=2363)

resolution (Root A=1043)

Q.440. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM"y.'l> '©0" "Ù7"â %%"Ù0"é/"ù 'l> "%?"iM'y>•)\$: "%M"Y\$.)\$"Ù0 '%% '%%>•ù •ù5•

A) '©M' ".y/ "%oM'Y?'I?ùK~". "r \$"™\$ "Y?"IM")G"y#"é\$Ù.R 0")*(~~BB'©M'.~~SVU.)©@_@%é%g%)Y K=2B89)'MnA(Gicay)proven Mieser Gons(fant\$C
constraint limits (RootB=1760)
©.Z.A=1120)©Dgnite vauk.Cerved)via quadrato carcus"ée'sdUde
bndes.compl. (Root D. 2732)

C) $'\text{C}'\text{M}^{\cdot}$ ".y"/ "%m'Y?'I?ùK." "r \$™\$ "Y?"IM")G"y#"\$U:'R 0")* %c m'o Y'mo>=80 %s a %g(1zc<26q3)M". Analytically proven B(itte)u constraint limits (Root B=1771)<=943 M".

[illegible]

A) '©M' '\$,.)' '%@é>9 'YK '©B' >'YO%G 'Y>')>'%'?' M'iG">B) /IM'Y?e\$>'EB'Y&C's'X'066587-1dFR">adMCo©MBE'mcclmngscnstr(air)
C) '©M' '.y/' %oM'Y?'I?'ùK*' "r \$™\$ "Y?"IM")G'y#"\$U'.R 0")* "%S %©tP' %0>e'x'G©C)2'0+221'S'0/%Galyteah•proveh indger'andseant'unk
resolution (Root A=1141)
boundary equality (Root D=2771)

A) 'OM' . 'y' / '%OM'Y?'I?'ÜK'." 'r '\$""\$ "Y?'IM)"G'Y#"\$S'Ü.'R 0")* "BG'OM' \$WU'."C%>@'ë'g")YK=2B32) /Möhaicaay)pröwen Miedel d'ons(fartätsch)

A) $\frac{1}{2} \times 10^{-10} \text{ m}$ B) 10^{-10} m C) $2 \times 10^{-10} \text{ m}$ D) $4 \times 10^{-10} \text{ m}$ E) $8 \times 10^{-10} \text{ m}$

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Allegation & Advanced Mixtures with Sequential Dilution, Problem #166): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) 'IM'Y?'>'B 'Y5'â "('%.é('iM'Y"> '©M" >'©M'B ("ü6ÜB)S\$P%\$ü,"2"%@1629 Determine value of B given q=ad and G=0.14 (rested)

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Time & Work with Alternating Shifts & Efficiency Differentials, Problem #167): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

Q.448. [English]: In SSC CGL Tier-2 Mathematical Abilities examination (Relative Speed, Circular Tracks & Escalator Motion, Problem #168): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

Q.449. •ù8•ù8"%@ "‰@'É@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM'y.'> '©0" "U7"â %, "™A'©& 'éB"™A " #"%@ •)! '©M" ."/ •ù5•" 8'é. "ù\$ '12, Â 8'é

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Polynomial Roots, Factor Theorem & Symmetric Functions, Problems 1400): Determine the rigorous mathematical analysis for the variable satisficing the advanced bounded & conditions below

resolution (Root A=1183)

- Q.450.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #170)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Quadratic Equations, Discriminant & Sign of Roots, Problem #170): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #170)
C) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #170)
Q.451. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #171)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Triangle Centres: Incentre, Circumcentre, Orthocentre & Centroid, Problem #171): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•) ‘%”@‘é>9 ‘YK ‘©B” > ‘Y0“%G “Y>”) > “%?” M‘iG“i>B) / iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #171)
C) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #171)
Q.452. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #172)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Cyclic Quadrilaterals & Tangent-Chord Angles, Problem #172): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #172)
C) ‘iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #172)
Q.453. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #173)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Circles: Common External & Transverse Tangents, Problem #173): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) “%”@‘é> “%.”l> ‘Y@ ‘©A“yM‘ù? ‘Y0“%G “Y>”) > •y7“Ù ‘l. ‘©0“â “B“©M“ \$“ù,•) ‘%”@‘é>9 ‘YK ‘©B” > ‘Y0“%G “Y>”) > “%?” M‘iG“i>B) / iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #173)
C) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #173)
Q.454. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #174)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Coordinate Geometry: Straight Lines, Collinearity & Intercepts, Problem #174): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #174)
C) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #174)
Q.455. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #175)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Trigonometric Maxima and Minima & Advanced Identities, Problem #175): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•) ‘%”@‘é>9 ‘YK ‘©B” > ‘Y0“%G “Y>”) > “%?” M‘iG“i>B) / iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #175)
C) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #175)
Q.456. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #176)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Heights and Distances with Multi-Point Observations, Problem #176): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #176)
C) ‘iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #176)
Q.457. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #177)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex 2D Mensuration & Shaded Region Areas, Problem #177): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) “%”@‘é> “%.”l> ‘Y@ ‘©A“yM‘ù? ‘Y0“%G “Y>”) > •y7“Ù ‘l. ‘©0“â “B“©M“ \$“ù,•) ‘%”@‘é>9 ‘YK ‘©B” > ‘Y0“%G “Y>”) > “%?” M‘iG“i>B) / iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #177)
C) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #177)
Q.458. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #178)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (3D Mensuration: Frustum, Prism, Pyramid & Tetrahedron, Problem #178): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #178)
C) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #178)
Q.459. •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“â %”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #179)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Statistics: Mean, Median, Mode, Variance & Standard Deviation, Problem #179): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•) ‘%”@‘é>9 ‘YK ‘©B” > ‘Y0“%G “Y>”) > “%?” M‘iG“i>B) / iM“Y?“%>B ‘Y5“â “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >©M‘B (“ù6”ÙBù\$6“ \$%.”)2“%”&”Ù5“ù “é\$ “%.”> “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #179)
C) ‘©M“ .”y/ “%M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “â “Ù5“ù “Ù\$‘Y0, “%.”%M‘ù> #179)

Q.460. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM"y.'l> '©0" "Ù7"â %%"Ù0"é/"ù 'l> "%%?'iM'y>•)\$: "%%M"Y\$.)\$"Ù0 '%% '%%>•ù •ù5•

A) 'C'M' .y/ '%M'Y?'I?ÜK• " r"\$™\$ "Y?"IM")G"y#"\$S"Ü.R 0")*(BQ'C'M'SVU,"%B@e"%%)Y K=QB4Q)/MnAqicaay)proven Mieser constraints (art.9 C)

C) 'iM"Y?"%>'B 'Y5"â ") ("%.é""é 'iM"Y">" > 'C'M' >'C'M'B ('ü6"UDüfö6,98...Z.A=1200)Ölymte van de Gerved via quadratische Lösung

A) "%@'é> "%.'l> 'Y@ '©A'yM'ù> 'Y0'%%G 'Y>'> •y7"Ü 'l. '©0'â *B%©K' \$'%)")2d@3079)'YQp0Ba>ext0mEparâmeter'Conflicting'boundâes'€

Q.462. •ù8•ù8"%%@ "%@'E@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM"y."l> '©0" "U7"â % " #" A "è\$'é 'R "Y "%\$'B -"ù("U(, "%."%M"ù> #182): '%

A) 'iM"Y?'%>'B 'Y5"â ") ("%. "é""é('iM"Y>"> '©M" >'©M'B ("ù6"UB)ù\$M" \$%,.)2"%@2749/ Define a value of the variable `q` as a quadratic polynomial (es decir,

Q.463. •ù8•ù8“%o@ “%o@‘E@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.l’> ‘©0” “U7“â %o “ù?” “*” “U0‘l?‘i\$‘!> •ù5• “Y.”)2”U/, .”TM!““A 8’é8”U/“â 3 f2

A) '©M" \$“ù,•)' "%@'é>'9 'YK '©B" > 'Y0'%G "Y>")> "%?" M'iG"i>B))iM("Y?%\$>EB"Y5C# "X" %66é34-44FR"> adMcoodMBr("meeting\$constrain)

Q.464. •ù8•ù8“%o@ “%o@‘E@•ù2 ‘ù?‘ù0-2 ‘y#“ù\$” / ‘YM“y.l’> ‘©0” “U7“ã %o2‘é--“TM>‘%o?, ‘EG•%o.“é(‘iA‘Y>‘%o&“é0 •ù5”, “é “é0 “ #‘%o

A) '©M' ".y/ '">%M'Y?'I?'ùK•" "r \$"TM\$ "Y?"IM")G"y#"^é\$"Û.R 0")* "%BG'©M?'\$U(•)©%@^é%9")YK=2B92 'YAnalogically'proven Miesler-constant\$(\$

Q.465. •ù8•ù8"%@ "%@'E@•ù2 'ù?'ù0-2 'y#"ù\$"/ 'YM"y.l> '©0" "U7"â "% "YM" 5"9&"U" "ô","U"/é (•Y0"U"Y">" M"y?'Rù\$"U0"%."é8"ù

A) "%@'é> "%.'l> 'Y@ '©A"yM'ù? 'Y0'%G "Y>")> •y7"Ù 'l. '©0'â *B%'©M' \$"(%)")2d@-145)'YQp0a>ext0%<parameter?conflicting)bound&\$ (

Q.466. •ù8•ù8“%@@ “%@‘E@•ù2‘ù?‘ù0-2‘y#“ù\$”/‘YM“y.‘l>‘©0”“U7“â%.“ù6”U0‘2“Y‘YM“.“ù“%>•)&”U0‘l>‘©0“ù5“M‘l(,“%.“%.

A) 'iM"Y?"%>'B 'Y5"â ") ("%. "é""é('iM"Y>"> '©M">'©M'B ("ù6"ùB)ù\$M" \$%,.)2"%@302"/ Define a value of the variable `q` as a draw from the `USresid` distribution.

Q.467. •ù8•ù8%00@ "%0@'E@•ù2'ù?'ù0-2 'y#"\$ù\$' / 'YM"y.l"> '©0" "U7"â %0 "e0"U/•ù5•" 8'e/ (•ù "é'lo'©"> " @•ù5•" &'YM"y\$â •ù\$' /

A) '©M" \$“ù,•)“ %o@'é>'9 'YK '©B" > 'Y0%oG "Y>")> %o?" M'iG"i>B))iM("Y?%>{eB"Y5C# \$(“%6Cé“eF“iM“FR" > aM“côM“hâ(“i“e“t“i“g“ô“p“s“k“i“n“

Q.468. •ù8•ù8"%@ "%@'E@•ù2'ù?'ù0-2'y#"ù\$"/'YM"y.l>'©0"U7"â%8"é*y"U7'©>"A5"9\$U\$"/'ùM"H'R"Y"•ù8"U"y2"y"

A) '©M" .y/ "%M'Y?'I?'ùK•" r \$™\$ "Y?"iM")G"y#é\$`Ü.'R 0")* "%BG'©M?'MUr.)©%@é%a")YK=2B44) /Analogically proven Miesler-cons(ante)nd

Q.469. •ù8•ù8%oo@ "‰@'E@•ù2 'ù?•ù0-2 'y#""ù\$"/ 'YM"y.l> '©0" "U7"â‰‰, "YM'A'©&'éB""A'" #‰‰)! '©M" .y/•ù5" 8'é."ù\$ '12,A 8'é

A) "%@'é> "%.'l> 'Y@ '©A'yM'ù? 'Y0'%%G "Y>")> •y7"Ù 'l. '©0'â *B%©M' \$"Ù%")%2D-3213)'Y0p0Ba>xt0%Éparâmetr?Configuring boundaries é

Q.480. •ù8•ù8“%_@ “%_@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.l’> ‘©0” ”Ù7“â %_*”Ù0“é/“ù ‘l’> “%_?’iM’y>•)\$. “%_M“Y\$.)\$”Ù0 ‘%_ ‘%_>•ù •ù5•

Q.500. •ù8•ù8“%_@ “%_@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.l’> ‘©0” ”Ù7“â %_*”Ù0“é/“ù ‘l’> “%_?’iM’y>•)\$. “%_M“Y\$.)\$”Ù0 ‘%_ ‘%_>•ù •ù5•

- Q.510.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”&”Ù5“ù “é\$ “%.” “ # •ù5•” 5“ù5“ù “Ù\$‘Y0, “%.”%”M‘ù> #230)
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Quadratic Equations, Discriminant & Sign of Roots, Problem #230): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%”>B ‘Y5“à “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6”ÙBù\$6“”\$%.”)2“%”&689) Define value derived via quadratic equation (Root D=2530)
C) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root D=3927)
- Q.511.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”\$“U0“ù” “YG ‘Y•G•)&”U0: •Y ‘l ‘Y•G•)&”U0, ‘©0“ù “y ‘iM“ Á
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Triangle Centres: Incentre, Circumcentre, Orthocentre & Centroid, Problem #231): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•) “%”@‘é>9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?” M‘iG“i>B) /iM“Y?“%”<EB“Y5G#SC “%”&689) Define value derived via quadratic equation (Root A=1617)
C) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root D=3927)
- Q.512.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”\$“YM “@ 0 ‘IA M‘UA‘Á “Y “%”M‘©0“Ù6” G‘i>-‘É@“Y> ‘YK
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Cyclic Quadrilaterals & Tangent-Chord Angles, Problem #232): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root B=2552)
C) ‘iM“Y?“%”>B ‘Y5“à “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6”ÙBù\$6“”\$%.”)2“%”&689) Define value derived via quadratic equation (Root D=3944)
- Q.513.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”\$“9\$“U\$: •TM-ù(“ù7”Ù •Y(“ 8“Ù““ M“b “Y ‘l?“ M‘ù “%”M‘©0
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Circles: Common External & Transverse Tangents, Problem #233): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) “%”@‘é> “%.”l> ‘Y@ ‘©A“yM‘ù? ‘Y0“%G “Y>“) > •y7“Ù ‘l. ‘©0“à “%”%”&689) Define value derived via quadratic equation (Root A=1631)
C) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root B=2563)
- Q.514.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %” (“ù0“U&“y6“é ‘R “Ù/“é.”ù\$“ó “%”0“““ 0“y “é•” “Y •Y ‘l‘i’ Á
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Coordinate Geometry: Straight Lines, Collinearity & Intercepts, Problem #234): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%”>B ‘Y5“à “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6”ÙBù\$6“”\$%.”)2“%”&689) Define value derived via quadratic equation (Root B=2574)
C) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root D=3978)
- Q.515.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”\$“U0“ù” “#‘é?‘l@‘ò ‘©M‘©?“yM ‘Ù(“ù.”Ù(“ù7”Ù •ù5•” 8“ M
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Trigonometric Maxima and Minima & Advanced Identities, Problem #235): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•) “%”@‘é>9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?” M‘iG“i>B) /iM“Y?“%”<EB“Y5G#SC “%”&689) Define value derived via quadratic equation (Root A=1645)
C) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root D=3995)
- Q.516.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”\$“• “é•ù5•” &”)0 “%.”TM-A-‘É?•)&” “*“U0“y “Ù72‘Á 8‘é8“Ù/“
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Heights and Distances with Multi-Point Observations, Problem #236): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root B=2596)
C) ‘iM“Y?“%”>B ‘Y5“à “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6”ÙBù\$6“”\$%.”)2“%”&689) Define value derived via quadratic equation (Root D=4012)
- Q.517.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”&“U5“ù5“ù.” / ‘YM“yG‘iM” “ù\$“ò “Y ‘i>‘ù>•) “ù\$ ‘YM“yG‘iM
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Complex 2D Mensuration & Shaded Region Areas, Problem #237): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) “%”@‘é> “%.”l> ‘Y@ ‘©A“yM‘ù? ‘Y0“%G “Y>“) > •y7“Ù ‘l. ‘©0“à “%”%”&689) Define value derived via quadratic equation (Root B=2607)
C) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root A=1659)
- Q.518.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”\$“U0“ù”Ù, ‘©?“ >‘é?‘ Á “ù(“Ù(R “Y ‘©\$“ 7”Ù+“), “%.”%”
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (3D Mensuration: Frustum, Prism, Pyramid & Tetrahedron, Problem #238): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘iM“Y?“%”>B ‘Y5“à “)(“%.”“é““é(‘iM“Y>“ > ‘©M“ >‘©M‘B (“ù6”ÙBù\$6“”\$%.”)2“%”&689) Define value derived via quadratic equation (Root B=2618)
C) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root D=4046)
- Q.519.** •ù8•ù8“%”@ “%”@‘É@•ù2 ‘ù?’ù0-2 ‘y#“ù\$” / ‘YM“y.”l> ‘©0” “Ù7“à %”\$“8“iM“ù?Y@: ‘é>yM“òÀ “é”Ù/“ù “âÀ, “TM-A“), ‘©M“ 8“ #
[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Statistics: Mean, Median, Mode, Variance & Standard Deviation, Problem #239): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:
A) ‘©M“ \$“ù,•) “%”@‘é>9 ‘YK ‘©B“ > ‘Y0“%G “Y>“) > “%?” M‘iG“i>B) /iM“Y?“%”<EB“Y5G#SC “%”&689) Define value derived via quadratic equation (Root A=1673)
C) ‘©M“ .”y/ “%”M‘Y?“l?‘ùK•” “r \$“TM\$ “Y?“iM“)G“y#“é\$“Ù.”R 0”) * “%”%”&689) Analytically proven integer consistency boundary equality (Root D=4063)

Q.520. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM"y.'l> '©0" "Ù7"â %%"Ù0"é/"ù 'l> "%%?'iM'y>•)\$: "%%M"Y\$.)\$"Ù0 '%% '%%>•ù •ù5•

A) "%@é' %>.%>'Y@ '©A"yMù? 'Y0%G "Y"> >•y7"U 'l. '©0'à %>©M "\$"ü")%2 @-6997)'Y©DÜB>ext©Gpar)net&rbkMfing)boüda\$ é
C) '©M" ."/ %>M'Y?'I?'ÜK•" "r \$™\$ "Y?"IM"G'y#é\$Ü.'R 0)"* %>G IM% M%> B0 Y\$ %>(%)Z©E3 E3)M"Analytically prove B if nuder wshen unit
resolution (A=1687)

A) 'iM'Y?'%>'B 'Y5'â "(" (%. "é"e('iM'Y"> '©M' >©M'B ("û6'ÛB) \$B4' \$û, %2' %>6649/ Define value for GeV via q=Adw(GeV) /y (rests) (

A) '©M' '\$,.)' "%@é>9 'YK '©B' >'Y0%'G "Y">' %?'" MiG">B)/M/Y?8\$>'EB'Y5G#>'2%0.86>'E(FI&VFR è 'eiffo' c©mfilBa(ü)ë'ttùg' éon'Stra)'
C) '©M' ".y/' %©M'Y?'I?'ùK#" "r \$™\$ "Y?"IM")G"y#"és"Û.R 0)"* %06 %©n&ar' %0>èx'(©P©)Z'0#8259)%Gally'cay'p7vén' iñdégér'èò'sé'ñt'uñ'
resolution (Root A=1701)
boundary sensitivity (Root C=1131)

A) '©M' .y/ '%mY?'I?ÜK" "r \$™\$ "Y?"IM")G"y# "é\$ "Ü.R 0")* "%B?©M?SWr. ©B@ "é%g")YK=3B72) /Mn@l@ca#y)proven Mieser-Gonsch(artsh(

C) "©M" .\y/ %mM'Y?'I?'ùK" " r \$ "™\$ "Y?'IM")G'y# "é\$ "Ü.R 0")* "constraint limits (Reot B=2695)

A) 'iM'Y'?%>'B 'Y5'â "("%."é"e('iM'Y"> '>'@M'>'@M'B ("ü6"ÜB)S64'%'ü."2'%-@7229/ Defining value of parameter via quadratic (C)

C) '@M' ".y/ "%M'Y'?I?'Ü·K" " 'r \$ "™\$ "Y?'iM")G"y#"é\$'Ü·R 0")* "ü6'%'@M'>'@M'B ("ü6"ÜB)S64'%'ü."2'%-@7229/ Defining value of parameter via quadratic (C)

constraint limits (Root-B=2706)

boundary condition (Root-B=1182)

conditions below:

A) 'COM' '\$\dot{u},s\$)' '%@' 'é>9' 'YK' 'CB' > 'Y0' '%G' 'Y>') > '%?' 'M'iG'i>B)' /M'Y'Y%G' ('EB'Y5G#S)' ('%65487' -ad YR' > adM'co'ordinate' meeting's'constrain

C) 'OM' '\$\dot{u},s\$)' '%@' 'é>9' 'YK' 'CB' > 'Y0' '%G' 'Y>') > '%?' 'M'iG'i>B)' /M'Y'Y%G' ('EB'Y5G#S)' ('%65487' -ad YR' > adM'co'ordinate' meeting's'constrain

[illegible]

Problem #249): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

Q.540. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y#"ù\$" / 'YM"y.'l> '©0" "Ù7"â %%"Ù0"é/"ù 'l> "%%?'iM'y>•)\$: "%%M"Y\$.)\$"Ù0 '%% '%%>•ù •ù5•

- [illegible]

- C) '0M' .y/ '%M'Y?I?i'uK' . "r \$ "TM\$ "Y?iM"G"y#E\$U.R 0")* %G lmsY nmo> B0 \$e a %o(12m e3 243M) Analytically prove B if (a) u is a constant with respect to y (Post A-1833)

- C) '0M' .y/'%M'Y?'!ùK•" "r \$™\$ "Y?"iM')G"y"ë\$'Û.R 0")* %& %@#4P%0-9?0-6%0/2 0-84080/%Galytichy-proven integer %005ah"un%

- C) '0M' .y/ "%M'Y?I?ùk•" "r \$™\$ "Y?"iM")G"y#é\$Ù.R 0")* %0\$ %00041%0.0-?00002Y0034190%00lytically proven inoger %0050h%000

- C) $\lim_{Y \rightarrow \infty} \frac{Y^5}{Y^5} = 1$ (The limit of $\frac{Y^5}{Y^5}$ as Y approaches infinity is 1, which is a constant value.)

- constraint limits (Root B=2915).

- A) 'IM'Y?'%>'B 'Y5'a ")(("%".'é"'é('IM'Y>' > 'CM' >'CM'B ('ùò'ÜBü\$ÖP'\$,%,"%)2'%=Q629/ Define value of parameter via quadratic equation by restricting constraint limits (Root B-2926).

- ferentials, Problem #267): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- Problem #268): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- [English]: In SSC CGL Tier-2 Mathematical Abilities examination (Polynomial Roots, Factor Theorem & Symmetric Functions, Problem #269): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

Q.560. •ù8•ù8"%@ "%@'É@•ù2 'ù?'ù0-2 'y#ù\$" / 'YM'y.'l> '©0" "Ù7"â %%"Ù0"é/"ù 'l> "%?"im'y>•)\$. "%M"Y\$•)\$"Ù0 "%o '%o>•ù •ù5•

[English]: In SSC CGL Tier-2 Mathematical Abilities examination (Probability Theory: Independent Events, Conditional & Cards/Dice, Problem #280): Determine the rigorous mathematical value for the variable satisfying the advanced boundary conditions below:

- A) '©M" .y/ "%M'Y?'l?'ùK•" "r \$™\$ "Y?'IM")G"y#é\$"Ù.R 0)"* "%©M'Y?'l?'ùK•"YK=3B40) /M%Qic%y)proven Miedr condantésC constraint limits (Root B=3080)
C) 'im"Y?'%o>'B 'Y5"â ") ("%. "é""é('im"Y>" > '©M" >'©M'B ("ù6"Ù0"é/"ù "YK=3B40) /M%Qic%y)proven Miedr condantésC constraint limits (Root B=3080)

Q.561. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 6 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 6 years at simple interest. Find the annual rate of interest.

- A) 18.0%
B) 15.0%
C) 12.5%
D) 33.3%

Q.562. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 7 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 7 years at simple interest. Find the annual rate of interest.

- A) 28.6%
B) 15.0%
C) 12.5%
D) 18.0%

Q.563. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 8 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 8 years at simple interest. Find the annual rate of interest.

- A) 12.5%
B) 15.0%
C) 25.0%
D) 18.0%

Q.564. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 9 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 9 years at simple interest. Find the annual rate of interest.

- A) 15.0%
B) 22.2%
C) 12.5%
D) 18.0%

Q.565. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 10 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 10 years at simple interest. Find the annual rate of interest.

- A) 20.0%
B) 15.0%
C) 12.5%
D) 18.0%

Q.566. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 11 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 11 years at simple interest. Find the annual rate of interest.

- A) 18.0%
B) 15.0%
C) 12.5%
D) 18.2%

Q.567. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 12 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 12 years at simple interest. Find the annual rate of interest.

- A) 12.5%
B) 15.0%
C) 16.7%
D) 18.0%

Q.568. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 13 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 13 years at simple interest. Find the annual rate of interest.

- A) 15.0%
B) 15.4%
C) 12.5%
D) 18.0%

Q.569. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 14 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 14 years at simple interest. Find the annual rate of interest.

- A) 15.0%
B) 14.3%
C) 12.5%
D) 18.0%

Q.570. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 5 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 5 years at simple interest. Find the annual rate of interest.

- A) 18.0%
B) 15.0%
C) 12.5%
D) 40.0%

Q.571. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 6 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 6 years at simple interest. Find the annual rate of interest.

- A) 33.3%
B) 15.0%
C) 12.5%
D) 18.0%

Q.572. 'YK•, "%0"é6"ò 8"é""é0'2 , "Ù/"é '©0 7 "Y0"Ù7"¹ 'éG•" 8"Ù5'ù 'Y@ 3 'yA'%@ "™K 'É>'l@ "™H-B 5"é0"Ù7"ù 'ÉM'ù>'Â & " "l

[English]: A sum of money becomes 3 times of itself in 7 years at simple interest. Find the annual rate of interest.

- A) 28.6%
B) 15.0%
C) 12.5%
D) 18.0%

