two Diagrams present Mackente Kall 1) Valprair Acid is a 5 host chan fatty acid, use ld in psychrotric Owarders It Acts an GABA levels in the Bran, Blocks Historie teacytlase inhibitor, and Blocks Valtye goted in Channels. GABA is farmed from OK b in the TCA cycle. Valprore Acid inhibits GABA transammer and succente remial behyde Dehydrayer are, which Degrader 10 ABA Valprare Acid Also has Anti-epiteptir properties
By Blocking Valtage gold Sodam & Cathannels, To
Decreating the frequency of from at most
It has Also Been shown to inhibit
HOLD I high - it HDACI which increases the expression of Apaphasis and anti furnar artial.

Kentres VPA :5 a highly Board protein Resulting in law chearace, 3 Mechanins for metabolism, blucorandation, Beta axidation in the nitochantern and Cyp450 netroted axidation (this is among rate howevers). The Valproate blucarante is the moor metabable 30-60% excreted in the wrine. Italsa can Bebranght into the liver was carnitive shuttle. Then through Petaaxidation it is Broken Down to sever-1 met-balites, A slaw hydrolysis by an Acyl CaA throwthere creto 3-040-VPA which is are endproduct from wied By a thealare which turns it into a thial conjugator

pharmaco Oynamers Valprose Acid (VPA) 260H HDA(9 akb PHOACII ADACI guccing atabab frill Juch succente 6ABA Antifumarg one sen: Aldohyte T HSA1 Nat CACNAISGNAP GABR

Cyp 2 C9 I) It is a phase I day metabolizing cytochome Pubo enzyme isoform that is primarily expressed in the liver and has the 2nd highest expression of all CYP isoforms.

It is responsible for 15-20% afall Drays in the phase I of Detarketho I t is known to metabolize To Ibutanise to by tray netry Italbut ande. substrite (RH) Fe<sup>2</sup> + O<sub>2</sub> B NAOPH

Ht pub of the peduction cyto ware

et years QIt stats when Iron Bunk to the Iron the Iron Becomes Peducato fezt. 3 Oz ten Bands to ten enzyne 3 Oz is tem Activated in a Telectra Reduden A single Oz Atom 27 Dissacrated with an Oxidation of Iron and protonation of complex involved theo leaves The penning a Atam 15 highly energets out peaks with substrate, which the leaves

The Dase are sever to Days at 10 to 20 m/kg at Body weg ht is the Hewaper for Dare Tray change the lysasand of the practical.

Hubule-This effects the peabsorption of theany Minerals such as Mgz+ (az+ K+ and Also CASTS, Pharphalipit, and greare in excretm If contined Day we accurs their it can be- As Ank Deend Followe. If high Daves of 40 mg/kg Are of wered from Castical necrosis onto controlly. From occus Rapidly. While futulas Neccosts is evident the Mechanism of how it has pers is unknown though seval mechanisms are failed potentially Books source.
The Damage is warly be parted After it has Been Darre, as taxily his asheret Rapidly As stated it would a rufly Decreve to Pe at sarption in the placement to Dehydiction, polyncia, by poturon, Norman, Fatigue, weakness, 7615 155 because the R190d value of Decreasing Diametrally and you are large sugar cells, B190d, and Forse There are all courself for furtrange Due to 90% at Reabsorbtin according at the proximal consoling tubule; tus 11 cause 1 arg e Distress. large Distress. It is largely elimented By glandidar

1)Ghodke-Puranik Yogita, Thorn Caroline F, Lamba Jatinder K, Leeder J Steven, Song Wen, Birnbaum Angela K, Altman Russ B, Klein Teri E. "Valproic acid pathway: pharmacokinetics and pharmacodynamics" Pharmacogenetics and genomics (2013)

2)Miners JO, Birkett DJ. Cytochrome P4502C9: an enzyme of major importance in human drug metabolism. Br J Clin Pharmacol. 1998;45:525–538.

3)Cytochrome P4502C9: an enzyme of major importance in human drug metabolism Miners J O, Birkett D J in British journal of clinical pharmacology (1998)

4) Marie-Paule Mingeot-Leclercq\* and Paul M. Tulkens Aminoglycosides: Nephrotoxicity Antimicrob. Agents Chemother. May 1999 vol. 43 no. 5 1003-1012

5)Guengerich F P, HumancytochromesP450.In:OrtizdeMontellano PR, editor. Cytochrome P450, structure, mechanism and biochem- istry (3rd Ed.). New York, Kluwer/Plenum Press, 2005. p. 377–530.