Visual cross-check of resting-state clustering

FOR2107 compared with Tozzi et al.

Preliminary comparison · five- and six-cluster descriptive solutions · 1 September 2026

FOR2107 data sample

2,239 participants in the selected FOR2107 cohort

1,130
Healthy reference participants


1,109
Clinical participants

Clinical diagnosis	n	Role in current analysis
MDD	823	Primary clinical group
Anxiety + comorbid MDD	266	Included with clinical group
Anxiety-only	20	Included with clinical group
Total clinical	**1,109**	MDD + anxiety analysis


Clustering input: 2,237 participants (1,108 clinical; 1,129 healthy reference).

Supplementary Figure 2: how convincing is the separation?


Tozzi et al.



FOR2107 · k = 6




FOR2107 · k = 5




Supplementary Figure 5: what do the cluster profiles look like?


Tozzi et al.



FOR2107 · k = 6



FOR2107 · k = 5



Task equivalence and the replication boundary

Tozzi et al.	Relationship	FOR2107
Task-free fMRI scan	**Close**	Resting-state fMRI scan
Conscious Facial Expressions of Emotion Task	**Related**	Face-matching task
Non-conscious Facial Expressions of Emotion Task	**Related**	Subliminal affective-priming task
Go–NoGo Task	**Missing**	No direct equivalent
No equivalent	**Additional**	Episodic memory-encoding task

Without the exact data acquisition protocol, we doubt that the results can be reproduced exactly.