
$P(\hat{s}^* > \text{band})$
Birds
Hihi Tiritiri Matangi Island, New Zealand 0.99
Hihi Karori, New Zealand 0.99
Blue tits La Rouvière, France 0.99
Blue tits Pirio, France 1.00
Blue tits Muro, France 1.00
Great tits Wytham Woods, UK 1.00
Great tits Hoge Veluwe, the Netherlands 0.98
Collared flycatchers Gotland, Sweden 0.98
Song sparrows Mandarte Island, Canada 0.99
Mammals
Snow voles Graubünden, Switzerland 0.38
Red squirrels Kluane, Canada 0.72
Spotted hyenas Ngorongoro Crater, Tanzania 1.00
Meerkats Kalahari, South Africa 0.86
Red deer Isle of Rum, UK 1.00
Soay sheep St. Kilda, UK 0.99
Bighorn sheep Ram Mountain, Canada 0.99
Rhesus macaques Cayo Santiago, Puerto Rico 0.93
Yellow baboons Amboseli National Park, Kenya 0.99
$10^{-4}$ $10^{-3}$ $10^{-2}$ $10^{-1}$ $10^{0}$ $10^{1}$
Mean heterozygous selection coefficient, $s$
Implied $\hat{s}^* = \hat{V}_A / \hat{U}_d$: mode, 50% and 95% HPD
Measured mean $s$ of new nonsynonymous mutations in wild birds or mammals
$\hat{s}^* > 1$: no selection coefficient maintains $\hat{V}_A$