
Superb fairy wrens Canberra, Australia
Hihi Tiritiri Matangi Island, New Zealand
Hihi Karori, New Zealand
Blue tits La Rouvière, France
Blue tits Pirio, France
Blue tits Muro, France
Great tits Wytham Woods, UK
Great tits Hoge Veluwe, the Netherlands
Collared flycatchers Gotland, Sweden
Song sparrows Mandarte Island, Canada
Birds
Snow voles Graubünden, Switzerland
Red squirrels Kluane, Canada
Spotted hyenas Ngorongoro Crater, Tanzania
Meerkats Kalahari, South Africa
Red deer Isle of Rum, UK
Soay sheep St. Kilda, UK
Bighorn sheep Ram Mountain, Canada
Rhesus macaques Cayo Santiago, Puerto Rico
Yellow baboons Amboseli National Park, Kenya
Mammals
10^{-4}
10^{-3}
10^{-2}
10^{-1}
10^{0}
10^{1}
Additive genetic variance in relative fitness, V_A
Maintainable at MSD balance: $U_d s$, $s = 10^{-3}$ to 10^{-2}
Measured: mode, 50% and 95% HPD
Least needed for observed π (multiplicative bound), study species
Least needed for observed π (multiplicative bound), another population