

a
Great tits (Wytham Woods, UK)
Great tits (Hoge Veluwe, the Netherlands)
Collared flycatchers (Gotland, Sweden)
Song sparrows (Mandarte Island, Canada)
Birds
Red deer (Isle of Rum, UK)
Soay sheep (St. Kilda, UK)
Bighorn sheep (Ram Mountain, Canada)
Rhesus macaques (Cayo Santiago, Puerto Rico)
Yellow baboons (Amboseli National Park, Kenya)
Mammals
10^{-5} 10^{-4} 10^{-3} 10^{-2} 10^{-1} 10^{0}
Effective size relative to breeding adults, N_e/N
b
2.0
2.1
3.2
1.6
1.0
1.3
1.1
0.8
0.8
0.6 0.8 1.0
$\bar{\Omega}$, maps only
Unlinked, multiplicative bound: $e^{-4\kappa_{max}\hat{V}_A}$, mode and 95% HPD
With linkage (total): $\bar{\Omega}e^{-4\kappa_{max}\hat{V}_A}$, assigned DFE: mode and 95% HPD
Observed: δ, π of study species
Observed: δ, π from another population
c
Chromosome Ω_i
1.00
0.75
0.50
0.25
0.00
10^{-3} 10^{-2} 10^{-1}
Selective target per Morgan, f_i/M_i
chromosome ≥ 20 Mb
microchromosome < 20 Mb
Birds
Mammals