

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/281277296>

# Amamiichthys, a new genus for the sparid fish *Cheimerius matsubarai* Akazaki 1962, and redescription of the species, with designation of a neotype

Article in *Zootaxa* · August 2015

DOI: 10.11646/zootaxa.4007.2.3

CITATIONS

0

READS

265

2 authors:



Fumiya Tanaka

Miyazaki University

10 PUBLICATIONS 18 CITATIONS

[SEE PROFILE](#)



Yukio Iwatsuki

Miyazaki University

133 PUBLICATIONS 1,212 CITATIONS

[SEE PROFILE](#)



## ***Amamiichthys*, a new genus for the sparid fish *Cheimerius matsubarai* Akazaki 1962, and redescription of the species, with designation of a neotype**

FUMIYA TANAKA<sup>1,3</sup> & YUKIO IWATSUKI<sup>2</sup>

<sup>1</sup>Interdisciplinary Graduate School of Agriculture and Engineering, University of Miyazaki, 1-1 Gakuen-kibanadai-nishi, Miyazaki, 889-2192, Japan. E-mail: nb11003@student.miyazaki-u.ac.jp

<sup>2</sup>Department of Marine Biology and Environmental Sciences, Faculty of Agriculture, University of Miyazaki, 1-1 Gakuen-kibanadai-nishi, Miyazaki, 889-2192, Japan. E-mail: yuk@cc.miyazaki-u.ac.jp

<sup>3</sup>Corresponding author

### **Abstract**

A new genus, *Amamiichthys*, is proposed for the sparid fish *Cheimerius matsubarai* Akazaki 1962. This genus differs from all other genera in the family Sparidae by the following combination of characters: both jaws with an outer row of small molariform teeth and an inner row of similar, even smaller teeth; frontal bone flat, coarse and porous; strongly bifurcate protuberance of upper ethmoid just beneath mid-region of anteriormost portion of frontal bone; head and body pinkish, gradually becoming silvery toward abdomen, with many small blue spots, some pairs overlapping. The type species, *Amamiichthys matsubarai*, is redescribed and a neotype designated.

**Key words:** Sparidae, new genus, *Amamiichthys*, *Cheimerius matsubarai*, Neotype

### **Introduction**

Species of the family Sparidae are widespread in tropical and temperate marine and brackish waters (Smith 1938; Akazaki 1962; Bauchot & Smith 1983; Carpenter 2001). Most are carnivorous, feeding on benthic invertebrates, and many are commercially important (Carpenter 2001; Iwatsuki *et al.* 2015).

The family Sparidae has historically been classified into four or six subfamilies, primarily on the basis of dentition (Smith 1938; Smith & Smith 1986; Akazaki 1962), although such subfamilies have more recently been considered as non-monophyletic on the basis of mitochondrial cytochrome *b* (*cyt-b*) gene and 16S ribosomal RNA data (Hanel & Sturmbauer 2000; Orrell & Carpenter 2002; Orrell *et al.* 2004; Chiba *et al.* 2009). Chiba *et al.* (2009) has also demonstrated that six genera (*Acanthopagrus*, *Cheimerius*, *Dentex*, *Diplodus*, *Pagellus* and *Pagrus*) are also non-monophyletic. Clearly, generic and higher classification of the Sparidae needs to be revised, using both morphological and genetic data. Furthermore, the position of centrarchid species, often included in the family Sparidae (Orrell *et al.* 2002; Orrell & Carpenter 2004; Chiba *et al.* 2009; Santini *et al.* 2014), needs to be resolved.

The genus *Cheimerius* (type species: *Dentex nufar* Valenciennes 1830) was proposed by Smith (1938) and was included in the subfamily Denticinae on the basis of acute, non-molariform teeth (Smith 1938; Smith & Smith 1986). Currently the genus includes two valid species: *C. nufar* from the Western Indian Ocean and *C. matsubarai* Akazaki 1962 known from only Amami-oshima Island, Ryukyu Islands, Japan. *Cheimerius* has been subsequently characterized as having 12 dorsal-fin spines (first and second very short, third to sixth longer and filamentous) and scales absent on the preopercular flange (Smith 1938; Akazaki 1962). Akazaki (1962) included *C. matsubarai* in the genus *Cheimerius* (subfamily: Denticinae) because both species share the above diagnostic characters. However, *C. matsubarai* has small molariform teeth on each jaw.

The combination of recent morphological (this study) and genetic studies has indicated strongly that *C. matsubarai* should be considered as generically distinct from *C. nufar*. In fact, Chiba *et al.* (2009) noted that *C. matsubarai* belonged to the same clade as Indo-West Pacific species of *Argyrops* Swainson 1839, *Evynnis* Jordan &