

## 26 2025-03-28 | Week 10 | Lecture 25

Start reading 6.1, 6.2

Recall that we say that a sequence  $X_1, X_2, \dots$  of random variables is *independent and identically distributed (IID)* if

- The  $X_i$ 's are independent rvs
- Every  $X_i$  has the same probability distribution as  $X_1$

**Theorem 80** (Central limit theorem). *Suppose we have an IID sequence of random variables  $X_1, X_2, \dots$  with finite mean  $\mathbb{E}[X_1] = \mu$  and finite variance  $\text{Var}(X_1) = \sigma^2 > 0$ . Let*

$$S_n = X_1 + \dots + X_n.$$

*Then for any fixed  $-\infty \leq a \leq b \leq +\infty$ , we have*

$$\lim_{n \rightarrow \infty} \mathbb{P} \left[ a \leq \frac{S_n - n\mu}{\sigma\sqrt{n}} \leq b \right] = \Phi(b) - \Phi(a) = \int_a^b \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy \quad (23)$$

*The conclusion of the CLT can be stated in many equivalent ways. One useful way involves the *sample average*  $\bar{X} := \frac{S_n}{n}$ . The CLT says that*

$$\bar{X} \approx \underbrace{\mu + \frac{\sigma}{\sqrt{n}}Z}_{\mathcal{N}(\mu, \frac{\sigma^2}{n})} \quad (24)$$

where  $Z \sim \mathcal{N}(0, 1)$ .

Usually  $n \geq 30$  is big enough for the approximation to be valid.

**Example 81.** A witch decides to make 50 batches of potions. Unfortunately, her hut is not very clean and bugs keep crawling into the brew. Suppose that the amount of bugs contaminating the  $i$ th batch is a random variable  $X_i$  with mean  $\mu = 4g$  and standard deviation  $\sigma = 1.5g$ . Assume the witch prepares her 50 batches independently. Let

$$\bar{X} = \frac{X_1 + \dots + X_{50}}{50}$$

be the average amount of impurity.

**Question:** What is the probability that  $3.5 \leq \bar{X} \leq 3.8$ ?

**Solution:** Here,  $n = 50 > 30$  so we can apply the central limit theorem. By the central limit theorem,  $\bar{X}$  has distribution approximately

$$\begin{aligned} \bar{X} &\approx \mu + \frac{\sigma}{\sqrt{n}}Z \\ &= 4 + \frac{3}{2\sqrt{50}}Z \end{aligned}$$

where  $Z \sim \mathcal{N}(0, 1)$  is a standard normal random variable. Therefore

$$\begin{aligned} \mathbb{P} [3.5 \leq \bar{X} \leq 3.8] &\approx \mathbb{P} \left[ 3.5 \leq 4 + \frac{3}{2\sqrt{50}}Z \leq 3.8 \right] \\ &= \mathbb{P} \left[ -\frac{1}{2} \leq \frac{3}{2\sqrt{50}}Z \leq -\frac{1}{5} \right] \\ &= \mathbb{P} \left[ -\frac{\sqrt{50}}{3} \leq Z \leq -\frac{2\sqrt{50}}{15} \right] \\ &= \mathbb{P} [-2.36 \leq Z \leq -.94] \\ &= \int_{-2.36}^{-.94} \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy \\ &\approx .16 \end{aligned}$$

End of Example 81.  $\square$

**Example 82** (Application of CLT). The human genome consists of a sequence of nucleotides  $(A, T, C, G)$ . For this problem we can think of the human genome as a sequence of nucleotides

CAGGAAGCATTATATAGAGGTAATAATAATTAAATTTAACTTTTTATTATTCTCTATGCCCAAGAGAATGTG...

and the sequence consists of  $n = 3.2 \times 10^9 = 3,200,000,000$  nucleotide letters.

The DNA mutation rate in humans is approximately  $\mu = 2 \times 10^{-8} = 0.00000002$  mutations per nucleotide per generation. That means each time a person is born, each nucleotide either mutates (with probability  $\mu$ ) or doesn't mutate (with probability  $1 - \mu$ ). Assume that nucleotides mutate *independently*, that is, whether one nucleotide undergoes mutation is independent from any other nucleotides.

Let  $S_n$  be the number of new mutations in the genome of a randomly-selected newborn baby. We can write

$$S_n = X_1 + \dots + X_n$$

where

$$X_i = \begin{cases} 1 & : \text{mutation at site } i \\ 0 & : \text{no mutation at site } i \end{cases}$$

Let  $M$  be the number of new mutations in the genome of a randomly selected newborn baby. Then  $M \sim \text{Bin}(n, \mu)$  where the "number of trials" is  $n = 3.2$  billion and the "success probability" is  $\mu = 2 \times 10^{-8}$ .

**Question 1:** On average, how many mutations does a newborn baby have?

**Solution:** We want

$$\begin{aligned} \mathbb{E}[S_n] &= \mathbb{E}[X_1 + \dots + X_n] \\ &= \mathbb{E}[X_1] + \dots + \mathbb{E}[X_n] \\ &= n\mu \\ &= (3.2 \times 10^9)(2 \times 10^{-8}) \\ &= 64. \end{aligned} \tag{25}$$

**Question 2:** What is the probability distribution of  $S_n$ ?

**Solution:**  $S_n$  is a sum of  $n$  "coin flips" each having success probability  $\mu$ . Hence,  $S_n$  is a binomial random variable with number of trials  $n$  and success parameter  $\mu$ .

**Question 3:** What percentage of people have between 56 and 72 mutations?

**Solution:** Use the central limit theorem. First observe that

$$\mathbb{E}[X_1] = 1 \cdot \mu + 0 \cdot (1 - \mu) = \mu \tag{26}$$

and

$$\mathbb{E}[X_1^2] = 1^2 \cdot \mu + 0^2 \cdot (1 - \mu) = \mu$$

Therefore

$$\begin{aligned} \sigma^2 &= \text{Var}(X_1) \\ &= \mathbb{E}[X_1^2] - (\mathbb{E}[X_1])^2 \\ &= \mu - \mu^2. \end{aligned}$$

Since  $\mu$  is very small  $\mu^2$  is VERY VERY small, and hence we have  $\sigma^2 \approx \mu$ . Therefore

$$\sigma \approx \sqrt{\mu} = \sqrt{2} \times 10^{-4}. \tag{27}$$

Finally, we have

$$\sqrt{n} = 40000\sqrt{2}. \tag{28}$$

Therefore, using Eq. (23) from the Central limit theorem, we have

$$\mathbb{P}\left[a \leq \frac{S_n - n\mu}{\sigma\sqrt{n}} \leq b\right] \approx \int_a^b \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy$$

and plugging in the values from Eqs. (25), (27) and (28), we get

$$\mathbb{P}\left[a \leq \frac{S_n - 64}{8} \leq b\right] \approx \int_a^b \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy$$

or equivalently

$$\mathbb{P}[64 + 8a \leq S_n \leq 64 + 8b] \approx \int_a^b \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy$$

Taking  $a = -1$  and  $b = 1$ , we get

$$\mathbb{P}[56 \leq S_n \leq 72] \approx \int_{-1}^1 \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy \approx .68.$$

Conclusion: about 68% of people have between 56 and 72 mutations.

What percentage of people have between 48 and 80 mutations?

End of Example 82.  $\square$