

Lecture Notes for Math 372: Elementary Probability and Statistics

Last updated: August 28, 2026

Probability theory has a right and a left hand. On the right is the rigorous foundational work using the tools of measure theory. The left hand “thinks probabilistically,” reduces problems to gambling situations, coin-tossing, models of a physical particle. – Leo Breiman

Contents

0	Tentative course outline	2
1	2026-08-24 Week 01 Lecture 01	3
2	2026-08-26 Week 01 Lecture 02	4
2.1	Set theory	4
2.1.1	Definitions and set-builder notation	4
2.1.2	Set operations	5
3	2026-08-28 Week 01 Lecture 03	7
3.1	A general framework for probability	7
3.2	Random variable	8
3.3	The enumeration principle	9

0 Tentative course outline

This course is a problem-oriented introduction to the basic concepts of probability and statistics, providing a foundation for applications and further study.

- **Weeks 1-3:** Introduction to probability theory
 - Experiments, events, sets, probabilities, random variables. Equally likely outcomes, counting techniques. Conditional probability. Independence. Bayes' theorem. (Sections: 2.1-2.5)
- **Weeks 4-5:** Discrete random variables (1.5 weeks)
 - Expected values, mean, variance, binomial distribution, Poisson distribution. Moment generating functions. (Sections: 3.1-3.6)
- **End of week 5: Midterm 1 (Friday, September 25)**
- **Weeks 6-7:** Continuous Random variables (1.5 weeks)
 - Uniform, exponential, gamma, and normal distributions. Intuitive treatment of the Poisson process and development of the relationship with gamma distributions. (Sections: 4.1-4.4)
- **Weeks 8-9:** Multivariate distributions
 - Calculation of probability, covariance, correlation, marginals, conditions. Distributions of sums of random variables and sampling distributions. Central limit theorem. (Sections: 1.1, 1.3, 1.4, 5.1-5.7)
- **End of week 9: Midterm 2 (Tentatively: Friday, October 23)**
- **Week 10:** Introduction to statistical estimation
 - Point and confidence interval estimation. Maximum likelihood, optimal, and unbiased estimators. Examples. (Sections 6.1, 6.2)
- **Weeks 11-12:** Large sample inference
 - Estimation (1.5 weeks): Types and comparison of estimators; sampling distributions for means/proportions, and their use in large sample estimation; sample size. (Sections 7.1, 7.2)
 - Hypothesis testing (1.5 weeks): Components of a test; significance and power; p-values; large-sample tests for means and proportions (Sections: 8.1-8.4)
- **Week 13:** Small sample inference
 - t-distribution, with applications to small sample estimation and testing; χ^2 and F distributions, with applications to inference about variances (Sections: 7.3, 7.4, 8.3)
- **Weeks 14-16:** Regression and χ^2 tests
 - Regression (1.5 weeks): Least squares, correlation coefficient, inference (Sections: 12.1-12.5)
 - χ^2 tests: multinomial distributions, contingency tables, goodness-of-fit (Sections: 14.1-14.3)
- **Last day of instruction: Wednesday, December 9**
- **Final Exam: December 18**

1 2026-08-24 | Week 01 | Lecture 01

- give syllabus
- did the pirates worksheet

2 2026-08-26 | Week 01 | Lecture 02

The nexus question of this lecture: What is a set and what are the basic set operations?

Reading assignment: chapter 2.1

2.1 Set theory

The mathematical formalization of modern probability theory rests on the foundation of set theory.

2.1.1 Definitions and set-builder notation

- A **set** is an unordered collection of distinct objects. The objects of a set are called **elements**. We write the elements of a set inside curly braces, $\{\dots\}$.

Examples: $\{1, 4, 2, 3\}$, $\{\text{cat}, \text{fish}, 5\}$, $\mathbf{Z} := \{\dots, -2, -1, 0, 1, 2, \dots\}$, $\mathbf{R} = \{\text{the real numbers}\}$

- Two sets A and B are **equal**, denoted $A = B$, if they share precisely the same elements. The order that the elements are written is irrelevant.

$$\{1, 2, 3\} = \{3, 1, 2\}.$$

Repeated elements are ignored:

$$\{1, 2, 3\} = \{1, 1, 2, 3\}.$$

- A set is **finite** if it contains a finite number of distinct elements. This number is called the **size** of the set, and is denoted $|A|$.

Example: $|\{\text{cat}, 5\}| = 2$.

- The **empty set** is the set which contains no elements. This is denoted as

$$\emptyset \quad \text{or} \quad \{\}$$

- Set-builder notation** describes a set by specifying the properties that its elements must satisfy.

Examples:

The set of integers between 1 and 36:

$$\{n \in \mathbb{Z} : 1 \leq n \leq 36\}$$

The set of ordered pairs (x, y) such that x and y are both positive real numbers:

$$\{(x, y) \in \mathbb{R}^2 : x, y > 0\}$$

The set of even integers:

$$\{n \in \mathbb{Z} : n \text{ is even}\}$$

The set of ordered pairs x, y such that both x and y are integers between 1 and 4:

$$\{(x, y) : x, y \in \{1, 2, 3, 4\}\}$$


In set builder notation, the colon ‘:’ is read as “such that” and the symbol ‘ $\in$ ’ (which looks like an E) is read as “is an element of”

2.1.2 Set operations

Let A, B be sets.


- The **intersection** of A and B , denoted $A \cap B$, is defined by

$$A \cap B = \{x : x \in A \text{ and } x \in B\}.$$




- The **union** of A and B , denoted $A \cup B$, is defined by

$$A \cup B = \{x : x \in A \text{ or } x \in B\}.$$




- The **set difference** of A and B , denoted $A \setminus B$, is

$$A \setminus B = \{x : x \in A \text{ and } x \notin B\}.$$




- We say that A is a **subset** of B , and write $A \subseteq B$, if

$$x \in A \implies x \in B.$$




- If $A \subseteq B$, the *complement of A in B*, denoted A^c , is

$$A^c = B \setminus A = \{x : x \in B \text{ and } x \notin A\}.$$



- We say that A and B are *disjoint* if $A \cap B = \emptyset$.



3 2026-08-28 | Week 01 | Lecture 03

The nexus question of this lecture: What does set theory have to do with probability?
Reading assignment: finish chapter 2.1, start 2.2

3.1 A general framework for probability

We begin with a general framework and some terminology to formalize the notions of probability.

- An **experiment** is an activity or process whose outcome is subject to uncertainty, and about which an observation is made.

Examples include flipping a coin, rolling a dice, measuring the size of a wave, or the amount of rainfall, conducting a poll, performing a diagnostic test, opening a pack of Pokemon cards, asking your crush on a date, etc.

- The **sample space** S of an experiment is the set of all possible outcomes. The elements of the sample space are called **sample points**. Each sample point represents a **unique** outcome of the experiment.

Roll a dice. There are six outcomes:

$$1, 2, 3, 4, 5, 6$$

These are the sample points. The sample space is

$$S = \{1, 2, 3, 4, 5, 6\}.$$

- In probability, we refer to subsets of the sample space as **events**.

an event = a set of outcomes = a subset of S .

We represent events using brackets $[\dots]$, and usually denote them with capital letters.

Examples of events when rolling a dice are:

$$E_1 = [\text{roll a 1}] = \{1\}$$

$$E_2 = [\text{roll a 2}] = \{2\}$$

$$E_3 = [\text{roll a 3}] = \{3\}$$

$$A = [\text{roll an odd number}] = \{1, 3, 5\}$$

$$B = [\text{roll an even number}] = \{2, 4, 6\}$$

$$C = [\text{roll a number less than 5}] = \{1, 2, 3, 4\}$$

$$D = [\text{roll a 2 or a 3}] = \{2, 3\}$$

- The first three events shown are **singleton** sets, containing only one sample point. The book calls these **simple events**. But note

$$1 \neq \{1\}$$

- The empty set $\emptyset$ and the whole sample space S are always both events:

- S is the event that an outcome occurs (“something happens”).
- $\emptyset$ is the event that no outcome occurs. This is called the **null event**.

- Two events are E, F **mutually exclusive** if $E \cap F = \emptyset$ (i.e., the sets E, F are disjoint). This means that the events E and F cannot both happen at the same time.

The events A, B defined above are mutually exclusive, since the dice roll cannot be both even and odd.

- Translating set operations into the language of events,

- $E \cap F$ is the event that both E **and** F occur
- $E \cup F$ is the event that at least one of the events E **or** F occurs
- $E^c = S \setminus E$ is the event that E does **not** occur.

Example 1 (Examples of sample spaces).

- Flip a coin. the sample space is $S = \{T, H\}$
- The amount of rainfall in a day is $S = \{x \in \mathbb{R} : x \geq 0\}$

An example of an event for this sample space is:

$$[\text{between 1 and 2 inches of rain}] = \{x \in \mathbb{R} : 1 \leq x \leq 2\}.$$

- You've got an urn filled with 300,000,000 balls. Exactly one ball is made of gold. Draw a ball out at random. If it's not the gold ball, put it back and keep repeating until you get the gold ball. Once you get the gold ball, you are done. The output of this experiment is *the number of times you drew a ball from the urn*. The sample space is

$$S = \{1, 2, 3, 4, \dots\}.$$

The last two examples show that the sample space need not be finite.

End of Example 1. $\square$

3.2 Random variable

We often associate numbers with outcomes of an experiment.

Definition 2 (Random variables). Let S be a sample space of an experiment. A **random variable** is any rule that associates a number with each outcome in the S . Formally, it is a function whose domain is S and whose range is $\mathbb{R}$.

Most interesting events are defined using random variables.

Example 3 (sum 2d4). Roll a 4-sided dice twice (this is the **experiment**). There are 16 possible **outcomes**. The **sample space** is

$$S = \{(x, y) : x, y \in \{1, 2, 3, 4\}\},$$

which consists of 16 ordered pairs (x, y) , the **sample points**. We can enumerate the sample space as a table:

		Dice 2			
		1	2	3	4
Dice 1	1	(1, 1)	(1, 2)	(1, 3)	(1, 4)
	2	(2, 1)	(2, 2)	(2, 3)	(2, 4)
	3	(3, 1)	(3, 2)	(3, 3)	(3, 4)
	4	(4, 1)	(4, 2)	(3, 4)	(4, 4)

Let X be the sum of the two rolls. This is an example of a **random variable**. We can represent X by the following table:

		Dice 2			
		1	2	3	4
Dice 1	1	2	3	4	5
	2	3	4	5	6
	3	4	5	6	7
	4	5	6	7	8

We can define events using X . For example

$$\begin{aligned}[X \leq 3] &= \{(x, y) \in S : x + y \leq 3\} = \{(1, 1), (1, 2), (2, 1)\} \\[X = 4] &= \{(x, y) \in S : x + y = 4\} = \{(1, 3), (2, 2), (3, 1)\} \\[X \leq 4] &= [X \leq 3] \cup [X = 4] = \{(1, 1), (1, 2), (2, 1), (1, 3), (2, 2), (3, 1)\} \\[X \text{ is even}] &= \{(1, 1), (1, 3), (2, 2), (2, 4), (3, 1), (3, 3), (4, 2), (4, 4)\} . \\ &\text{etc.}\end{aligned}$$

End of Example 3. $\square$

3.3 The enumeration principle

On Monday's worksheet, I had you do a bunch of problems where you computed probabilities by counting outcomes. This is a simple but critically important idea, called the **enumeration principle**.

Enumeration principle: If the sample space is finite and consists of *equally likely outcomes*, then we can compute lots of probabilities easily by listing outcomes and counting them. To be precise, for any event A ,

$$\mathbb{P}[A] = \frac{|A|}{|S|} = \frac{\# \text{ of outcomes in } A}{\# \text{ of outcomes in } S}.$$

So using X from Example 3,

$$\begin{aligned}\mathbb{P}[X = 4] &= \frac{3}{16} \\ \mathbb{P}[X \text{ is even}] &= \frac{8}{16} = \frac{1}{2}.\end{aligned}$$