

# Testing and Relaxing Distributional Assumptions on Random Coefficients in Demand Models<sup>\*</sup>

Max Lesellier<sup>†</sup> Hippolyte Boucher<sup>†</sup> Gökçe Gökkoca<sup>†</sup>

December 2, 2022

[Link to the last version](#)

## Abstract

The BLP demand model for differentiated products is the workhorse model for demand estimation with market-level data. This model uses random coefficients to account for unobserved preference heterogeneity. The shape of the distribution of random coefficients matters greatly for many counterfactual quantities, such as the cost pass-through. In this paper, we develop new econometric tools to test this distribution and improve its estimation under a flexible parametrization. First, we construct new instruments that are designed to detect deviations from the true distribution of random coefficients. Second, we develop a formal moment-based specification test on the distribution of random coefficients. Third, we show that our instruments can be successfully used to estimate a flexible distribution of random coefficients. Finally, we validate our approach with Monte Carlo simulations and an empirical application using data on car purchases in Germany. We also show that these methods extend to the mixed logit demand model with individual-level data.

**Keywords:** demand estimation, specification test, random coefficients

**JEL codes:** C35, C36, L13, C52

---

<sup>\*</sup>The authors would like to thank Christian Bontemps, Nour Meddahi and Mathias Reynaert for their guidance and support as well as Steve Berry, Richard Blundell, Pierre Dubois, Jean-Marie Dufour, Eric Gautier, Cristina Gualdani, Koen Jochmans, Philip Haile, Vishal Kamat, Pascal Lavergne, and Thierry Magnac for helpful comments and discussions. We also thank conference and seminar participants at the TSE Econometrics and the Yale IO workshops as well as the Milan EEA-ESEM, the Brisbane ESAM, the London IAAE and the Tokyo AMES conferences. Finally, we want to thank Kevin Remmy for providing us with a rich data set on car purchases and characteristics in Germany from 2012 to 2018. All remaining errors are our own.

<sup>†</sup>Toulouse School of Economics, max.lesellier@tse-fr.eu, hippolyte.boucher@tse-fr.eu, gokce.gokkoca@tse-fr.eu.

# 1 Introduction

The differentiated product demand model initiated by [Berry \(1994\)](#) and [Berry, Levinsohn, and Pakes \(1995\)](#) has been used in a wide array of empirical studies. It enables researchers to perform demand estimation in markets with differentiated products using either macro-level (market shares) or micro-level (individual purchases) data while allowing for unobserved heterogeneity in preferences as well as price endogeneity. This unobserved heterogeneity in preferences is modeled through the use of random coefficients (RCs) in the utility function. This framework allows researchers to estimate demand functions, price elasticities and counterfactual outcomes. Applications of the BLP model have notably studied the determinants of market power, the welfare effects resulting from a merger or the introduction of a new good and the economic impact of a tax or a subsidy.<sup>1</sup>

The informativeness of the empirical analysis depends on how well the model can reproduce the underlying substitution patterns and approximate the shape of the demand curve, including its slope and curvature. A recent result in [Miravete, Seim, and Thurk \(2022\)](#) shows that the commonly used Gaussian RC on price imposes strong restrictions on the demand’s curvature and thus limits the range of the implied pass-through. The degree of pass-through of taxes and costs is central to answering many questions in economics such as the impact of tariffs or a cost shock on consumer welfare. However, estimating a more flexible demand system with a non-Gaussian distribution of random coefficients is challenging. First, there is a clear trade-off between the degree of flexibility one chooses (for instance, going from a Gaussian to a Gaussian mixture) and the precision of the estimates one obtains. Therefore, it is important to be able to test the specification chosen by the researcher on the distribution of the RC (for instance, a Gaussian RC) and quantify the degree of misspecification before potentially moving to a more flexible specification. Second, to precisely estimate a more flexible distribution of RC, the researcher must choose instruments (or

---

<sup>1</sup>The BLP demand model has been widely utilized in numerous applications. A non-exhaustive list of examples includes: [Barahona, Otero, Otero, and Kim \(2020\)](#), [Berry et al. \(1995\)](#), [Crawford, Shcherbakov, and Shum \(2019\)](#), [Dubois, Griffith, and O’Connell \(2018\)](#), [Durrmeyer \(2022\)](#), [Grennan \(2013\)](#), [Grigolon, Reynaert, and Verboven \(2018\)](#), [Miller, Sheu, and Weinberg \(2021\)](#), [Miller and Weinberg \(2017\)](#), [Miravete, Moral, and Thurk \(2018\)](#), [Nevo \(2000\)](#), [Petrin \(2002\)](#), [Reynaert \(2021\)](#).

equivalently moment conditions) that strongly identify this distribution. The instruments used by the current empirical practice work well with the standard Gaussian RC, but their performance appears to decline as the specification becomes more flexible in the simulation exercises that we perform.

In this paper, we provide novel econometric tools to address these two challenges. In particular, we construct a new set of instruments designed to detect deviations from the true distribution of random coefficients. Building on these instruments, we provide a formal moment-based specification test on the distribution of random coefficients, which can be implemented without having to re-estimate the model under a more flexible parametrization. Our instruments are designed to maximize the power of this test when the distribution of RC is misspecified. We also show how these instruments can strengthen the identifying power of the moment conditions used for estimation, and thus be successful at estimating a flexibly parameterized distribution of RCs. As an example of a flexible parametric distribution, we consider the Gaussian mixture, which can approximate arbitrarily well any continuous distribution on the real line.

This paper consists of three main contributions. First, we construct a new set of instruments that are designed to detect departures from the true distribution of RCs. The intuition we use is the following. Any given distribution of RCs generates a structural error, which, if correctly specified, is mean-independent with respect to a set of exogenous variables. This identifying condition can be transformed into unconditional moments, which can be used to test whether the chosen distribution of RCs is correctly specified. We formally define this test and construct instruments that maximize its power against a fixed alternative. In a first step, we assume that the econometrician knows the fixed alternative and we derive an expression for the first-best instrument. We call this instrument the most powerful instrument (MPI) and show that this specific choice of instrument achieves the consistency of the test. In a second step, we provide two feasible approximations of the MPI that can be derived without knowledge of the fixed alternative. We call these feasible MPIs the interval instruments in reference to the way they approximate the MPI.

Second, we consider the case where the researcher wants to test whether the distribution of

RCs belongs to a given parametric family. For instance, the researcher may be interested in testing if the random coefficient is normally distributed. This is a composite hypothesis, and we must estimate the unknown parameters of the distribution in a first step. In a second step, we choose instruments to test if the distribution evaluated at the estimated parameters is correctly specified. Here, the interval instruments represent a natural choice of instruments as they are designed to detect deviations from the true distribution of RCs. We study the asymptotic properties of our test when the number of markets,  $T$ , goes to infinity and we prove the asymptotic validity of the test under common assumptions. In particular, we account for the statistical uncertainty stemming from the first step estimation, and we control for the magnitude of the approximations that intervene in the estimation of the BLP model. Our asymptotic results complement previous work by [Freyberger \(2015\)](#) on the asymptotic properties of the BLP estimator when the number of markets grows to infinity.

Third, we show that our interval instruments can be successfully used to estimate the model, and particularly so when the distribution of RCs is flexibly parameterized. We do so by exhibiting the connection between the MPI and the classical optimal instruments used for efficient estimation purposes. Specifically, we show that the MPI devoted to testing the specification of the model at the true parameter against any local alternative can be rewritten as a linear combination of the optimal instruments. This relation between the MPI and the optimal instruments helps us understand why the interval instruments, which approximate the MPI, perform so well in our simulations. So far, the literature has exclusively exploited instruments that approximate the optimal instruments ([Gandhi and Houde \(2019\)](#), [Reynaert and Verboven \(2014\)](#)). We refer to these instruments as traditional instruments. These have been shown to work well in the usual Gaussian case. However, our simulations show that their performance declines when we depart from the Gaussian RC.

To evaluate the performance of our test and instruments, we conduct two sets of simulation experiments. First, we compare the performance of the test when using our interval instruments and when using the instruments commonly adopted by practitioners ([Gandhi and Houde \(2019\)](#), [Reynaert and Verboven \(2014\)](#)). We show that the test has the correct empirical size and that the

interval instruments significantly outperform the traditional instruments in terms of power under alternative distributions. Second, we evaluate the performance of the interval instruments in estimating the model when the distribution of RC is flexibly parametrized, and follows a Gaussian mixture. We show that our instruments outperform the traditional instruments in terms of the mean squared error. In the case where the RC is a Gaussian mixture, the three sets of instruments perform equally well.

Finally, we apply the tools developed in this paper to estimate the demand for cars in Germany from 2012 to 2018. The objective of the empirical exercise is to see how well our instruments perform at estimating a flexible distribution of RCs using a real dataset. Given the importance of price to address most empirical questions, we increase the flexibility of the model by estimating a Gaussian mixture for the RC associated with price. Second, we use our specification test to assess how the degree of misspecification decreases when we increase the flexibility in the distribution of RCs. Third, we use our results to study how the shape of the RC on price can modify important counterfactual quantities such as the pass-through. In particular, our empirical results are consistent with the findings in [Miravete et al. \(2022\)](#).

**Related literature.** Our paper contributes to several strands of the literature. First, it contributes to the literature on the flexible estimation of aggregate demand models for differentiated goods. A few recent papers have proposed non-parametric and semi-parametric methods to estimate aggregate demand functions. [Compiani \(2018\)](#) proposes a non-parametric estimator of the demand functions. If relaxing all the parametric assumptions makes this approach conceptually appealing, it also faces significant theoretical and practical difficulties (more stringent data requirements, large curse of dimensionality, limited scope for counterfactual analysis).<sup>2</sup> [Lu, Shi, and Tao \(2021\)](#) and [Wang \(2022\)](#) propose semi-parametric estimators of the distribution of RCs. These approaches are complementary to ours and the instruments we develop in this paper can

---

<sup>2</sup>In particular, [Compiani \(2018\)](#) relaxes the Type 1 Extreme Value assumption on the taste shock. However, it is not clear how restrictive this assumption is. [McFadden and Train \(2000\)](#) shows that a mixed-logit model with flexibly distributed random coefficients can approximate any discrete choice model derived from random utility maximization. On the other hand, the Type 1 Extreme Value assumption generates massive computational gains, which allows for studying sophisticated markets with many products and many characteristics. Thus, the cost-benefit analysis seems to be largely in favor of the logit specification.

be useful to implement their non-parametric IV estimation procedures, which are known to be rather sensitive to the quality of the instruments (Chetverikov and Wilhelm (2017)). Finally, Ho and Pakes (2014), Tebaldi, Torgovitsky, and Yang (2019) suggest deriving bounds directly on the counterfactual quantities.

Our paper also contributes to the literature on the non-parametric identification of the distribution of RCs in demand models (Fox and Gandhi (2011), Fox, il Kim, Ryan, and Bajari (2012), Dunker, Hoderlein, and Kaido (2022), Wang (2022), Berry and Haile (2014)). First, we slightly extend the identification result in Wang (2022) to link it directly to the primitives of the model, without assuming that the demand functions are identified. Second, we provide a practical way of constructing moments that feature high identifying power with respect to the distribution of RCs.

Third, we contribute to the literature that focuses on the practical estimation of the BLP model. First, we show that the interval instruments that we construct in this paper can be successfully used to estimate the distribution of random coefficients, and particularly so under of flexible distribution of RCs. This new set of instruments complements instruments commonly used by practitioners: Reynaert and Verboven (2014) and Gandhi and Houde (2019) (see Conlon and Gortmaker (2020) for a review). Moreover, we provide a new parametrization of the model, which facilitates the estimation when the distribution of RCs is a Gaussian mixture. This new parametrization complements previous papers that aim at improving the estimation of the model (Dubé, Fox, and Su (2012), Lee and Seo (2015), Salanié and Wolak (2019)).

Finally, our paper contributes to the literature on the asymptotic properties of the BLP estimator (Armstrong (2016), Berry, Linton, and Pakes (2004), Freyberger (2015), Ketz (2019)). In particular, we prove the asymptotic normality and the consistency of the BLP estimator in the large market framework under less stringent assumptions than the remainder of the literature.

**Structure of the paper.** In Section 2, we recall the baseline BLP model, define the structural error of the model, and provide conditions under which the distribution of RCs is non-parametrically identified. In Section 3, we derive the most powerful instrument and show how it relates to the classical optimal instruments. In Section 4, we construct two feasible approximations of the MPI. In Section 5, we present our specification test and show its asymptotic validity. In

Section 6, we conduct Monte Carlo simulations to evaluate the consequences of misspecification on quantities of interest, and gauge the performance of our test and instruments. In Section 7, we apply our new tools to estimate the demand for cars in Germany. We conclude the paper in Section 8.

## 2 Model and identification

### 2.1 Indirect utility and moment restrictions

**Indirect utility.** We first describe the indirect utility function that induces the observed market shares. Our setting closely follows the one introduced in the seminal paper [Berry et al. \(1995\)](#). There are  $T$  markets indexed by  $t = 1, \dots, T$ . There is a continuum of consumers indexed by  $i$ . There are  $J_t$  market-specific products in market  $t$ . Each consumer chooses a product  $j \in \{0, 1, \dots, J_t\}$  where  $j = 0$  corresponds to the outside option. For the sake of exposition and without loss of generality, we will assume throughout our analysis that the number of products is constant across markets ( $\forall t, J_t = J$ ). Product  $j$  is characterized by a vector of characteristics  $x_{jt}$ , which includes the price of the good in most empirical settings. Consumer  $i$  derives an indirect utility  $u_{ijt}$  from purchasing good  $j \in \{0, 1, \dots, J\}$  in market  $t$ :

$$u_{ijt} = \underbrace{x'_{1jt}\beta + \xi_{jt}}_{\delta_{jt}} + x'_{2jt}v_i + \varepsilon_{ijt}, \quad (1)$$

with the following:

- $x_{1jt}$  is a vector of product characteristics of dimension  $K_1$  associated with product  $j$  and for which there is no preference heterogeneity;  $\beta$  represents preferences for  $x_{1jt}$ ;
- $\xi_{jt}$  is an unobserved demand shock on product  $j$  in market  $t$ ;
- $\delta_{jt} \equiv x'_{1jt}\beta + \xi_{jt}$  denotes the mean utility for product  $j$ , the part of the utility that is common to all consumers;
- $x_{2jt}$  is a vector of product characteristics of dimension  $K_2$  for which there is preference heterogeneity;  $v_i$  is the associated random coefficient that follows a distribution characterized

by density  $f$  and is independent of all the other variables:  $v_i \perp\!\!\!\perp (x_t, \xi_t, \{\varepsilon_{ijt}\}_{j=1,\dots,J})$ ;

- $\varepsilon_{ijt}$  is a preference shock that follows an Extreme Value type I (EV1) distribution independent of all other variables and across  $i, j, t$ .

For individual  $i$  in market  $t$ , the indirect utility from purchasing the outside option is normalized to  $u_{i0t} = \varepsilon_{i0t}$ . From the random utility functions in (1), we can infer the demand functions for each good  $j$  in market  $t$  denoted  $\rho_{jt}(f, \beta)$ . Each consumer chooses the product that maximizes his or her utility. Let  $y_{ijt}$  equal 1 if individual  $i$  chooses good  $j = 0, 1, \dots, J$  in market  $t = 1, \dots, T$ . We have the following:

$$\begin{aligned}
\forall j \neq 0, \quad \rho_{jt}(f, \beta) &\equiv \mathbb{P}_{f,\beta}(y_{ijt} = 1 | x_t, \xi_t) \\
&= \mathbb{P}_{f,\beta}(\text{good } j \text{ is chosen in market } t \text{ by individual } i | x_t, \xi_t) \\
&= \mathbb{P}_{f,\beta}(u_{ijt} > u_{ikt} \quad \forall k \neq j | x_t, \xi_t) \\
&= \int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{1jt}\beta + \xi_{jt} + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{x'_{1kt}\beta + \xi_{kt} + x'_{2kt}v\}} f(v) dv.
\end{aligned} \tag{2}$$

For the outside option, the demand function is written as follows:

$$\rho_{0t}(f, \beta) = \mathbb{P}_{f,\beta}(y_{i0t} = 1 | x_t, \xi_t) = \int_{\mathbb{R}^{K_2}} \frac{1}{1 + \sum_{k=1}^J \exp\{x'_{1kt}\beta + \xi_{kt} + x'_{2kt}v\}} f(v) dv.$$

Following the EV1 assumption on the idiosyncratic shock on utility, the demand functions take the usual logit form integrated over the distribution of preference heterogeneity. We assume in this paper that the observed market shares are equal to the shares generated by the model above at the true distribution  $f$  and the true preference parameter  $\beta$ :

$$\forall j, \quad \forall t, \quad s_{jt} = \rho_{jt}(f, \beta). \tag{3}$$

**Moment restrictions.** Following the literature, we assume that the unobserved demand shock  $\xi_{jt}$  is mean independent of  $z_{jt}$ , a set of instrumental variables, namely,  $\mathbb{E}[\xi_{jt} | z_{jt}] = 0$  *a.s.*. The set  $z_{jt}$  traditionally consists of the exogenous characteristics of all the products on the market as well as cost shifters, which are meant to instrument for price. Indeed, the price of a good is usually considered to be an endogenous variable since it is correlated with the unobserved demand shock



$\xi_{jt}$  through the profit maximization problem of firms.<sup>3</sup> To estimate the model, the researcher chooses functions of the instruments  $z_{jt}$  to construct a set of unconditional moments. We refer to these functions as estimation instruments and denote them  $h_E(z_{jt})$ . Likewise, in our analysis, we study the functions of the instruments that are designed to test the specification of the model. We refer to these instruments as testing instruments and we denote them  $h_D(z_{jt})$ , where  $D$  stands for detection.

## 2.2 Inverse demand function and structural error

**Inverse demand function.** For any given distribution of random coefficients  $\tilde{f}$ , we define the demand function  $\rho \equiv (\rho_1(\cdot), \dots, \rho_J(\cdot))$  as the function which maps the vector of mean utilities  $\delta$  to the vector of market shares generated by the model under  $\tilde{f}$ :

$$\rho(\cdot, x_{2t}, \tilde{f}) : \mathbb{R}^J \rightarrow [0, 1]^J$$

$$\delta \mapsto \int_{\mathbb{R}^{K_2}} \frac{\exp\{\delta + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_k + x'_{2kt}v\}} \tilde{f}(v) dv.$$

Berry (1994) shows by applying Brouwer's fixed point that for any  $(s_t, x_{2t})$  and for any distribution of random coefficients  $\tilde{f}$  (even when  $\tilde{f}$  is not the true distribution), there exists a unique  $\tilde{\delta} \in \mathbb{R}^J$  such that:

$$s_t = \rho(\tilde{\delta}, x_{2t}, \tilde{f}).$$

We define the solution to the previous system of equations as the inverse demand functions:  $\rho^{-1}(s_t, x_{2t}, \tilde{f}) = \tilde{\delta}$ . Unfortunately, there is no closed form expression for the inverse demand function, which must be recovered numerically.

**Structural error.** From what precedes, we can uniquely define the structural error  $\xi_{jt}(\tilde{f}, \tilde{\beta})$  generated by a distribution of random coefficient  $\tilde{f}$  and a homogeneous parameter  $\tilde{\beta}$ :

$$\xi_{jt}(\tilde{f}, \tilde{\beta}) = \rho_j^{-1}(s_t, x_{2t}, \tilde{f}) - x'_{1jt}\tilde{\beta}. \quad (4)$$

---

<sup>3</sup>To deal with the endogeneity of prices, Berry et al. (1995) also suggests using exogenous own-product characteristics as well as exogenous characteristics from other products. The main idea behind the use of these instruments is to take advantage of the correlation between price and exogenous characteristics implied by profit-maximizing firms. To be precise, Berry et al. (1995) suggests using the sum of the characteristics from other products produced by the same firm and the sum of exogenous characteristics from rival firms' products as instruments.

The non-linear nature of the model is captured by the inverse demand function which enters the expression of the structural error. The absence of an analytical formula for the inverse demand implies that there is no closed form expression for the structural error, which complicates the estimation of the BLP demand model. If we consider a parametric family of distributions  $\tilde{\mathcal{F}} = \{\tilde{f}(\cdot|\tilde{\lambda}) : \tilde{\lambda} \in \tilde{\Lambda}\}$ , then the structural error generated by a specific element in  $\tilde{f}(\cdot|\tilde{\lambda}) \in \tilde{\mathcal{F}}$  and  $\tilde{\beta}$  is defined as follows:

$$\xi_{jt}(\tilde{f}(\cdot|\tilde{\lambda}), \tilde{\beta}) = \rho_j^{-1}(s_t, x_{2t}, \tilde{f}(\cdot|\tilde{\lambda})) - x'_{1jt}\tilde{\beta}.$$

## 2.3 Non-parametric identification

The main objective of this paper is to provide tools to test the specification on the distribution of random coefficients and to improve its estimation under a flexible specification. A natural first step is to study the conditions under which this distribution is non-parametrically identified. The identification of random coefficients in multinomial choice models has been studied extensively in the literature (Allen and Rehbeck (2020), Berry and Haile (2014), Dunker et al. (2022), Fox and Gandhi (2011), Fox et al. (2012), Wang (2022)). We summarize some of these findings in Appendix C.1. In this Section, we build on an important identification result in Wang (2022) to recover a set of sufficient identifying conditions directly on the primitives of the model. We also show that the identification result holds with a less stringent exogeneity assumption than in Wang (2022).

In contrast to the rest of the literature, Wang (2022) adopts all the parametric assumptions in the standard BLP model and looks for a set of sufficient restrictions under which the identification of the demand functions implies the identification of the distribution of random coefficients. This approach allows him to obtain conditions that are less stringent than the rest of the literature. In particular, Wang (2022) makes no special regressor assumption, no full support assumption, and no continuity assumption on the covariates. Specifically, he shows that if the demand functions  $\rho = (\rho_1, \dots, \rho_J)$  are identified on an open set of  $\mathbb{R}^J$ , then the distribution of random coefficients is identified.<sup>4</sup> His proof exploits the real analytic property of the demand functions.<sup>5</sup> In this

---

<sup>4</sup>Identification of demand functions can be achieved using Theorem 1 in Berry and Haile (2014).

<sup>5</sup>In particular, the real analytic property yields that the local identification of  $\rho$  on  $\mathcal{D} \subset \mathbb{R}^J$  implies the iden-

paper, we build on this injectivity result to find sufficient identifying conditions directly on the primitives of the model (without assuming identification of the demand functions). We also show using a random permutation of the indices that we only require the demand shock  $\xi_{jt}$  to be mean independent of the instrumental variables  $z_{jt}$  across products, but we do not require this to hold for each product  $j$  taken separately. Formally, we only require  $\mathbb{E}[\xi_{jt}|z_{jt}] = 0$  *a.s.* and not  $\mathbb{E}[\xi_{jt}|z_{jt}] = 0$  *a.s.* for all product  $j$  as previously. This is less restrictive, as demand shocks can now be on average non-zero for certain products and account for unobserved quality inherent to each product.

Let us formally state the assumptions that we impose to recover the point identification of  $(f, \beta)$ .

**Assumption A.**

- (i) Strict exogeneity:  $\mathbb{E}[\xi_{jt}|z_{jt}] = 0$  *a.s.*;
- (ii) Completeness: for any measurable function  $g$  such that  $\mathbb{E}[|g(s_t, x_t)|] < \infty$ , if  $\mathbb{E}[g(s_t, x_t)|z_{jt}] = 0$  *a.s.*, then  $g(s_t, x_t) = 0$  *a.s.*;
- (iii) The distribution of the data  $(s_t, x_{2t}, x_{1t}, z_t)$  is fully observed by the econometrician and market shares  $s_t$  are generated by the demand model defined in Section 2.1 by equations (1) and (3);
- (iv) Detectable difference in distributions: we say  $f$  and  $\tilde{f}$  differ (and write  $f \neq \tilde{f}$ ) if there exists  $\bar{v} \in \mathbb{R}^{K_2}$  such that  $F(\bar{v}) \neq \tilde{F}(\bar{v})$ ;
- (v) Let  $x_t = (x_{1t}, x_{2t})$  then  $x_t$  is such that  $\mathbb{P}(x_t'x_t \text{ is positive definite}) > 0 \quad \forall t$ ;
- (vi) There exists  $\bar{x}_t \in \mathcal{X}$  and an open set  $\mathcal{D} \subset \mathbb{R}^J$  such that  $\delta_t = \bar{x}_{1t}\beta_0 + \xi_t$  varies on  $\mathcal{D}$  *a.s.*.

In A(i), we assume that the instruments are strictly exogenous. Assumption A(ii) is a completeness assumption that states that the instruments are strongly relevant with respect to  $(s_t, x_t)$ . This assumption is typical of semiparametric or nonparametric IV models and is equivalent to a full rank assumption in a linear IV model. Intuitively, it means that if the inverse demands are different almost surely, then the instruments will be able to detect the difference. The completeness assumption is a strong assumption that has been widely used in this literature (Berry and Haile (2014), Dunker et al. (2022), Wang (2022)). Assumption A(v) is a standard rank condition.

---

tification of  $\rho$  on  $\mathbb{R}^J$  From the global identification of  $\rho$ , he is then able to show that the random coefficients' distribution is identified under a simple rank condition on  $x_{2t}$ .

Assumption A(vi) is meant to ensure that there is enough variation in  $\delta_t$  to apply the injectivity result in Wang (2022). This assumption indicates that there needs to be sufficient variation in product characteristics across markets in the data to identify  $f$ . In practice, product characteristics are very similar from one market to the other and may not yield sufficient variation. A judicious solution is to create inter-market variation by interacting product characteristics with demographic variables characterizing each market. Let us now state our formal identification result.

**Proposition 1.** Under Assumption A, the distribution of random coefficients  $f$  and the homogeneous preference parameters  $\beta$  are non-parametrically identified:

$$(\tilde{f}, \tilde{\beta}) = (f, \beta) \iff \mathbb{E}[\xi_{jt}(\tilde{f}, \tilde{\beta})|z_{jt}] = \mathbb{E}\left[\rho_j^{-1}(s_t, x_{2t}, \tilde{f}) - x'_{1jt}\tilde{\beta} \middle| z_{jt}\right] = 0 \text{ a.s..}$$

The proof is in Appendix B.1. The identification result above entails that under some fairly weak conditions and in the presence of instruments that generate sufficient variation in the product characteristics, the observed data identifies the distribution of random coefficients non-parametrically. Formally, the model is at the true pair  $(f, \beta)$  if and only if the associated structural error is mean independent of the instrumental variables  $z_{jt}$ . We use this identification result to show the consistency of our test under a specific choice of instruments that we will characterize thereafter.

### 3 Detecting misspecification: the most powerful instrument

The aim of this section is to recover the instrument with the greatest ability to detect misspecification in the distribution of RC. To do so, we consider a setting in which the econometrician wants to test a simple hypothesis of the form  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$ . The upper bar is used to stress the fact that  $\bar{H}_0$  is a simple hypothesis, in contrast to the composite hypothesis  $H_0 : f \in \mathcal{F}_0$  that we study in Section 5. Our approach builds on a simple intuition: if the model under  $\bar{H}_0$  is misspecified, then the structural error will depart from the true demand shock  $\xi_{jt}$ , and our goal is to find the best instrument to pin down this deviation. We proceed as follows. First, we introduce

a moment-based test for  $\bar{H}_0$  and we show its asymptotic validity. Next, we derive an analytical expression for the instrument that maximizes the power of our test against a fixed alternative  $\bar{H}_a : (f, \beta) = (f_a, \beta_a)$ . We call this instrument the most powerful instrument (MPI) and we show how it relates to the classical optimal instruments, derived for efficient estimation purposes. In Section 4, we provide two feasible approximations of the MPI, which have the critical property of being invariant with respect to the alternative  $\bar{H}_a$ .

### 3.1 A moment-based test

We want to test  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$  against  $H_a : (f, \beta) \neq (f_0, \beta_0)$ . For any set of testing instruments  $h_D(z_{jt})$ , we have the following implication:

$$\bar{H}_0 : (f, \beta) = (f_0, \beta_0) \implies \bar{H}'_0 : \mathbb{E}[h_D(z_{jt})\xi_{jt}(f_0, \beta_0)] = 0.$$

We propose to test  $\bar{H}_0$  indirectly through its implication  $\bar{H}'_0$ , which is a set of unconditional moment conditions. We test  $\bar{H}'_0$  with a moment-based test. Our test statistic writes as follows:

$$S_T(h_D, f_0, \beta_0) = TJ \left( \frac{1}{TJ} \sum_{j,t} \xi_{jt}(f_0, \beta_0) h_D(z_{jt}) \right)' \hat{\Omega}_0^{-1} \left( \frac{1}{TJ} \sum_{j,t} \xi_{jt}(f_0, \beta_0) h_D(z_{jt}) \right), \quad (5)$$

with  $\hat{\Omega}_0$  a consistent estimator of  $\Omega_0$  the asymptotic variance-covariance matrix of  $\frac{1}{\sqrt{TJ}} \sum_{j,t} h_D(z_{jt})\xi_{jt}(f_0, \beta_0)$ , that is  $\Omega_0 = \mathbb{E}[\xi_{jt}^2(f_0, \beta_0) h_D(z_{jt}) h_D(z_{jt})']$ . We study the asymptotic properties of our test as the number of markets,  $T$ , goes to infinity. As the focus of this section is on the construction of the most powerful instrument, we postpone the treatment of the specific challenges implied by parameter uncertainty (i.e. when  $\beta_0$  and  $f_0$  must be estimated beforehand) and by the numerical approximations involved in the derivation of the structural error (in practice, the researcher derives a numerical approximation of  $\xi_{jt}(f_0, \beta_0)$ ) to Section 5. Additionally, to keep the results as simple as possible while retaining the key intuitions, we assume independence of the demand shocks in a given market conditional on  $z_{jt}$ . This last assumption is relaxed in the proofs in Appendix B.2 and in Section 5.

**Proposition 2.** Assume that  $(s_t, x_t, z_t)$  are i.i.d. across markets and consistent with the probability model defined by equations (1), (2) and (3) evaluated at  $(f, \beta)$ ,  $\mathbb{E}[\|\xi_{jt}(f_0, \beta_0)h_D(z_{jt})\|^2] < +\infty$ ,  $\Omega_0$  has full rank, and, for  $k \neq j$ ,  $\xi_{jt} \perp \xi_{kt}|z_t$ . We have the following:

- under  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$ ,  $S_T(h_D, f_0, \beta_0) \xrightarrow[T \rightarrow +\infty]{d} \chi^2_{|h_D|_0}$ ,
- under  $H'_a : \mathbb{E}[h_D(z_{jt})\xi_{jt}(f_0, \beta_0)] \neq 0$ ,  $\forall q \in \mathbb{R}^+$ ,  $\mathbb{P}(S_T(h_D, f_0, \beta_0) > q) \xrightarrow[T \rightarrow +\infty]{} 1$ ,

with  $|\cdot|_0$  being the counting norm.

The previous proposition indicates that as long as the testing instruments are functions of  $z_{jt}$ , our test procedure is asymptotically valid for  $\bar{H}_0$ . We are testing  $\bar{H}_0$  by virtue of its implication  $\bar{H}'_0 : \mathbb{E}[h_D(z_{jt})\xi_{jt}(f_0, \beta_0)] = 0$  and, as a consequence, the power properties of our test hinge critically on the choice of the testing instruments  $h_D(z_{jt})$ . This is the focus of the next subsection.

### 3.2 The most powerful instrument (MPI)

The choice of testing instruments  $h_D(z_{jt})$  is key to maximizing the rejection rate of  $\bar{H}_0$  under any alternative  $H_a : (f, \beta) \neq (f_0, \beta_0)$ . To guide our choice of instruments, we first derive the instrument that maximizes the power of our test when the econometrician tests  $\bar{H}_0$  against a fixed alternative  $\bar{H}_a : (f, \beta) = (f_a, \beta_a) \neq (f_0, \beta_0)$ . We refer to this instrument as the most powerful instrument (MPI). In practice, the researcher is often reluctant to fix the alternative. However, the MPI represents a useful first-best solution for which we provide two feasible approximations in Section 4.

**Derivation of the most powerful instrument.** To construct the MPI, we use the following decomposition of the structural error generated under  $\bar{H}_a$ :

$$\xi_{jt}(f_0, \beta_0) = \underbrace{\xi_{jt}(f_a, \beta_a)}_{\text{true error under } \bar{H}_a} + \underbrace{\xi_{jt}(f_0, \beta_0) - \xi_{jt}(f_a, \beta_a)}_{\Delta_{0,a}^{\xi_{jt}}},$$

with  $\Delta_{0,a}^{\xi_{jt}}$  being the correction term due to misspecification under the alternative  $\bar{H}_a$ . Our goal is to compare the ability of our test for different candidates  $h_D(z_{jt})$ , to reject  $\bar{H}_0$  under  $\bar{H}_a$ .

The literature offers many ways to compare the power of competing tests (see [Gourieroux and Monfort \(1995\)](#) for a comprehensive review). First, we distinguish between exact and approximate methods. Exact methods rely on the exact distribution of the test statistic (under  $\bar{H}_0$ ) and allow for comparison in finite sample while asymptotic methods exploit the asymptotic distribution of the test statistic and are informative in larger samples. In our case, the exact distribution of our test is unknown. Thus, we rely on asymptotic methods, which is the most common case in the literature. Second, we divide the methods into local and non-local methods. In parametric tests, local strategies are based on the analysis of the power properties of competing tests under a sequence of local alternatives  $\theta_T$  which converges to  $\theta_0$  at a given rate (usually  $\frac{1}{\sqrt{T}}$ ). The econometrician can compare two competing tests by means of their power functions (or more precisely, the limits of these power functions when sample sizes go to  $+\infty$ ). This is called the direct approach. The dual approach, which is known as Pitman’s relative efficiency, consists of comparing the rates at which the minimal number of observations must increase to ensure a given level of power. The approach we favor in this paper is the non-local approach developed in [Bahadur \(1960\)](#). Here, the econometrician chooses the test with the smallest level  $\alpha$  needed to attain a given power against a fixed alternative and for a given number of observations. In other words, the econometrician chooses the test that minimizes the risk of type I error *ceteris paribus*.

There are several reasons to favor Bahadur’s non-local approach. First, it is better suited for the testing problem we study in this paper. The comparison criterion, known as the asymptotic slope of the test, is in our case straightforward to derive, whereas it is not clear how one should derive Pitman’s efficiency criterion when the test concerns non-parametric objects such as distributions. Moreover, we study the properties of our test against a fixed alternative  $\bar{H}_a : (f, \beta) = (f_a, \beta_a)$  as in Bahadur’s case, which is not necessarily local. Finally, the literature has highlighted many limitations of the local approach. Local criteria are often unable to discriminate between tests even when these tests lead to different decisions (see [Silvey \(1959\)](#)). In addition, as shown in [Dufour and King \(1991\)](#), a locally optimal test in a neighborhood of  $H_0$  may perform very poorly away from  $H_0$ .

Let us now present the intuition for Bahadur’s comparison approach. From [Section 3.1](#), we

have:

$$\text{Under } \bar{H}_0: \quad S_T \equiv S_T(h_D, f_0, \beta_0) \xrightarrow{d} S \quad \text{with } S = \chi^2_{|h_D|_0}.$$

Following the same notations as in [Gourieroux and Monfort \(1995\)](#), we denote:

$$\Lambda(s) = \mathbb{P}_{\bar{H}_0}(S \geq s).$$

The critical value is usually derived using the asymptotic distribution of the test statistic under  $H_0$ . The approximate critical region at a given level  $\alpha$  is then given by:

$$CR_\alpha = \{S_T \geq \Lambda^{-1}(\alpha)\} = \{\Lambda(S_T) \leq \alpha\}.$$

The main idea in Bahadur's approach entails deriving the level of the test if one takes the value of the test statistic as the critical value (this is also known as the p-value). Namely:

$$\alpha_T = \Lambda(S_T).$$

Bahadur suggests preferring the test that displays the lowest level  $\alpha_T$  at least asymptotically. A formal analysis of the asymptotic behavior of  $\alpha_T$  shows that it is better to consider the limit of a transformation of  $\alpha_T$  than the limit of  $\alpha_T$  directly. This gives rise to the concept of the approximate slope of the test.

**Definition 1** (Asymptotic slope of the test).

- (i)  $K_T = -\frac{2}{T} \log(\Lambda(S_T))$  is the approximate slope of the test,
- (ii) Under  $\bar{H}_a$ :  $\text{plim } K_T = c(f_a, \beta_a)$  is the asymptotic slope of the test,

with  $\text{plim}$ , the limit in probability when  $T \rightarrow +\infty$ .

Under the alternative  $\bar{H}_a : (f, \beta) = (f_a, \beta_a)$ , consider two sequences of tests based on  $S_T^1$  and  $S_T^2$  with asymptotic slopes  $c^1(f_a, \beta_a)$  and  $c^2(f_a, \beta_a)$  respectively. The test based on  $S_T^1$  is asymptotically preferred to the test based on  $S_T^2$  in Bahadur's sense if and only if  $c^1(f_a, \beta_a) > c^2(f_a, \beta_a)$ . To derive the asymptotic slopes of our test, we apply an important result in [Geweke \(1981\)](#), which states



that if under  $H_0$ :  $S_T \xrightarrow[T \rightarrow +\infty]{d} \chi_q^2$  (with any  $q \in \mathbb{N}^*$ ), then  $\frac{1}{T}S_T \xrightarrow{a.s.} c(f_a, \beta_a)$  (when the limit exists). In our test, the limiting distribution is chi-squared. Thus, the asymptotic slope of our test with instrument  $h_D(z_{jt})$  writes:

$$c_{h_D}(f_a, \beta_a) = \text{plim} \frac{1}{T} S_T(h_D, f_0, \beta_0) = J\mathbb{E}[\xi_{jt}(f_0, \beta_0)h_D(z_{jt})]' \Omega_0^{-1} \mathbb{E}[\xi_{jt}(f_0, \beta_0)h_D(z_{jt})].$$

Let us note that the asymptotic slope can also be interpreted as a measure of the speed of divergence of the test statistic in terms of population moments, i.e. speed of divergence  $\approx T \times c_{h_D}(f_a, \beta_a)$ . In the next Proposition, we derive an analytical expression for the instrument that maximizes the slope of the test.

**Proposition 3** (Most powerful instrument).

Let  $\mathcal{H}$  be the set of measurable vectorial functions of  $z_{jt}$ . Under any fixed alternative  $\bar{H}_a : (f, \beta) = (f_a, \beta_a)$ , we have the following:

$$(\mathbb{E}[\xi_{jt}(f_0, \beta_0)^2 | z_t])^{-1} \mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}] \in \underset{h_D \in \mathcal{H}}{\operatorname{argmax}} c_{h_D}(f_a, \beta_a).$$

The proof is given in Appendix B.2. The MPI equals the conditional expectation of the correction term  $\Delta_{0,a}^{\xi_{jt}}$  divided by a conditional variance term  $\mathbb{E}[\xi_{jt}(f_0, \beta_0)^2 | z_{jt}]$ . For exposition purposes, we drop the conditional variance term in the subsequent analysis and take the homoskedastic MPI  $h_D^*(z_{jt}) = \mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}]$  as the reference MPI.<sup>6</sup> Methods have been proposed to estimate the conditional variance term non-parametrically and could be adapted to our case. However, it is well known that conditional variance, which also appears in the formulation of the optimal instruments, is difficult to model and estimate in practice. In the BLP framework, the large dimension of  $z_{jt}$  makes the exercise even more difficult. Hence, researchers typically ignore this term or impose a restrictive and ad-hoc structure on the form that it can take (for instance, [Reynaert and Verboven \(2014\)](#)'s approximation of the optimal instruments in the BLP model ignores the variance term). The homoskedastic MPI,  $h_D^*(z_{jt})$ , features other appealing properties including (i) consistency of the associated test and (ii) maximizing correlation with the structural error under the alternative.<sup>7</sup> For simplicity, in what follows, we refer to the homoskedastic MPI as the MPI.

---

<sup>6</sup>This last expression corresponds to the exact formulation of the MPI under homoskedasticity.

<sup>7</sup>The consistency of the test also holds when we keep the conditional variance term.

**(i) Consistency.** By setting  $h_D$  equal to  $h_D^*$ , our moment-based test becomes consistent against any fixed alternative  $\bar{H}_a : (f, \beta) = (f_a, \beta_a) \neq (f_0, \beta_0)$ . Namely, we have the following result:

**Proposition 4** (Consistency of the test with the MPI). Under Assumption A and the same assumptions as in Proposition 2, we have:

$$\bar{H}_a : (f, \beta) = (f_a, \beta_a) \neq (f_0, \beta_0) \implies \forall q \in \mathbb{R}^+, \mathbb{P}(S_T(h_D^*, f_0, \beta_0) > q) \xrightarrow{T \rightarrow +\infty} 1.$$

The proof of this result is given in Appendix B.2.

**(ii) Correlation with the structural error.** Another interesting property of the MPI is to be the function of  $z_{jt}$  which maximizes the correlation with the structural error.

**Proposition 5** (Correlation between the MPI and the structural error).

Let  $\mathcal{H}$  be the set of measurable functions of  $z_{jt}$ , we have under  $\bar{H}_a$ :

$$\forall \alpha \in \mathbb{R}^*, \quad \alpha \mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}] \in \arg \max_{h \in \mathcal{H}} |\text{corr}(\xi_{jt}(f_0, \beta_0), h(z_{jt}))|.$$

The proof is given in Appendix B.2. Intuitively, the MPI  $h_D^*(z_{jt})$  is designed to fully capture the exogenous variation contained in the correction term  $\Delta_{0,a}^{\xi_{jt}}$  implied by the misspecification, which yields the result above.

### 3.3 Connection with the optimal instruments

The MPI maximizes the power of the moment-based test for  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$ . In contrast, the optimal instruments minimize the asymptotic variance-covariance of the GMM estimator when the parameter of interest is identified by conditional moment restrictions. These two problems are seemingly unrelated. However, we show that the MPI devoted to testing the specification of the model at the true parameter against any fixed local alternative can be rewritten as a linear combination of the optimal instruments. Consequently, one can reinterpret the optimal instruments as a local approximation of the MPI devoted to testing the model at the true parameter. This connection between the MPI and the optimal instruments helps us understand why the feasible approximations of the MPI we construct in Section 4 improve the performance of the BLP estimator

in our Monte Carlo simulations when the distribution of RCs is flexible. In this subsection, we first derive the optimal instruments. Then, we exhibit the relation between the optimal instruments and the MPI.

The estimation of the model works as follows. The researcher assumes that  $f$  belongs to a parametric family  $\mathcal{F}_0 = \{f_0(\cdot|\tilde{\lambda}) : \tilde{\lambda} \in \Lambda_0\}$  and wants to estimate the true parameter  $\theta_0 = (\beta'_0, \lambda'_0)'$  under this parametric restriction. In the estimation context that we study here,  $\theta_0$  refers to the true parameter. For now, let us assume that the model is correctly specified:  $f \in \mathcal{F}_0$  and we shorten the notations by removing the dependence of the structural error in  $f_0(\cdot|\tilde{\lambda})$ , which becomes implicit in this context. Namely,  $\xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta})$  becomes  $\xi_{jt}(\tilde{\theta})$ . We further assume that  $\theta_0$  is point identified by the following moment restriction:  $\mathbb{E}[\xi_{jt}(\theta_0)|z_{jt}] = 0$  *a.s.*<sup>8</sup> The researcher must choose the set of instruments  $h_E(z_{jt})$  (or equivalently, the unconditional moments) to include in the GMM objective function:

$$\hat{\theta} = \underset{\tilde{\theta}}{\text{Argmin}} TJ \left( \frac{1}{TJ} \sum_{j,t} \hat{\xi}_{jt}(\tilde{\theta}) h_E(z_{jt}) \right)' \hat{W} \left( \frac{1}{TJ} \sum_{j,t} \hat{\xi}_{jt}(\tilde{\theta}) h_E(z_{jt}) \right).$$

**Optimal instruments in the BLP demand model.** Traditionally, the instruments  $h_E(z_{jt})$  are chosen to minimize the asymptotic variance-covariance of the estimator  $\hat{\theta}$ . The instruments that reach this objective are called the optimal instruments. The resulting estimator is said to be efficient in the sense that its asymptotic variance cannot be reduced by using additional moment conditions. There is a large body of literature on the derivation of optimal instruments in econometric models (Amemiya (1974), Chamberlain (1987), Newey (1990, 2004)). The BLP estimator  $\hat{\theta}$  is a non-linear GMM estimator and classical results in Chamberlain (1987) and Amemiya (1974) show that the optimal instruments in this case write:

$$h_E^*(z_{jt}) = \mathbb{E}[\xi_{jt}(\theta_0)^2|z_{jt}]^{-1} \mathbb{E} \left[ \frac{\partial \xi_{jt}(\theta_0)}{\partial \tilde{\theta}} \middle| z_{jt} \right],$$

---

<sup>8</sup>The identification conditions in the parametric case are less stringent than the conditions for the non-parametric identification in Assumption A.

The corresponding efficiency bound (obtained by setting  $h_E = h_E^*$ ) writes:

$$V^* = \mathbb{E} \left[ \mathbb{E} \left[ \frac{\partial \xi_{jt}(\theta_0)}{\partial \tilde{\theta}} \middle| z_{jt} \right] \mathbb{E} \left[ \frac{\partial \xi_{jt}(\theta_0)}{\partial \tilde{\theta}} \middle| z_{jt} \right]' \mathbb{E}[\xi_{jt}(\theta_0)^2 | z_{jt}]^{-1} \right].$$

For the sake of exhaustivity, we show this result in Appendix B.2.1. As for the MPI, the formulation of the optimal instruments above is obtained under the assumption of conditional independence of demand shocks  $\xi_{jt}$  in the same market:  $k \neq j$ ,  $\xi_{jt} \perp \xi_{kt} | z_t$ . In Appendix B.2.1, we derive the expression for the optimal instruments under weaker assumptions on the demand shock.<sup>9</sup> Consistent with what we did in the case of the MPI, we drop the conditional variance term  $\mathbb{E}[\xi_{jt}(\theta_0)^2 | z_{jt}]$ .

**Connection between the MPI and the optimal instruments.** Let  $\theta_0$  the true parameter. Under the parametric assumption  $f \in \mathcal{F}_0$ , the simple hypothesis  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$  we studied previously becomes  $\bar{H}_0 : \theta = \theta_0$ . It is straightforward to show that, in the parametric case, the associated MPI against a fixed alternative  $\bar{H}_a : \theta = \theta_a$  writes:  $h_D^*(z_{jt}) = \mathbb{E} \left[ \Delta_{\theta_0, \theta_a}^{\xi_{jt}} | z_{jt} \right]$  with  $\Delta_{\theta_0, \theta_a}^{\xi_{jt}} = \xi_{jt}(\theta_0) - \xi_{jt}(\theta_a)$ . By taking a Taylor expansion of  $\xi_{jt}(\theta_a)$  around  $\theta_0$ , we obtain the following:

$$\Delta_{\theta_0, \theta_a}^{\xi_{jt}} = \frac{\partial \xi_{jt}(\theta_0)}{\partial \tilde{\theta}} (\theta_0 - \theta_a) + o(\|\theta_0 - \theta_a\|_2).$$

We see that when  $\theta_a$  is in a neighborhood of  $\theta_0$ , the MPI,  $h_D^*(z_{jt})$ , against this fixed alternative is a linear combination of the optimal instruments  $h_E^*(z_{jt})$ :

$$h_D^*(z_{jt}) = \mathbb{E} \left[ \Delta_{\theta_0, \theta_a}^{\xi_{jt}} | z_{jt} \right] \approx \underbrace{\mathbb{E} \left[ \frac{\partial \xi_{jt}(\theta_0)}{\partial \tilde{\theta}} \middle| z_{jt} \right]'}_{h_E^*(z_{jt})} (\theta_0 - \theta_a).$$

It follows that classical optimal instruments can be interpreted as an approximation of the MPI devoted to testing  $H_0 : \theta = \theta_0$  against any fixed local alternative.<sup>10</sup> Moreover, let us note that the connection between the MPI and the optimal instruments holds if we keep the conditional variance term in both cases.

---

<sup>9</sup>We allow for unrestricted forms of correlation between demand shocks within a given market.

<sup>10</sup>This interpretation of the optimal instruments only holds when the model is well specified i.e.  $f \in \mathcal{F}_0$ , and thus, in general, the optimal instruments shouldn't be used to test the specification of the model.

## 4 A feasible most powerful instrument

The MPI is the most powerful instrument to reject  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$  against a fixed alternative  $\bar{H}_a : (f, \beta) = (f_a, \beta_a)$ . Its derivation requires the knowledge of the alternative while in practice the econometrician typically wants to remain agnostic about the alternative. Moreover, the MPI is defined as a conditional expectation of a non-linear function with respect to a large dimension vector  $z_{jt}$ , and thus, even if the alternative  $\bar{H}_a$  is known, the MPI can be difficult to compute. In this section, we remain in the same configuration, where the econometrician wants to test  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$  against a fixed alternative  $\bar{H}_a : (f, \beta) = (f_a, \beta_a)$ . However now, we assume that this alternative is unknown to the econometrician. We provide two feasible approximations of the MPI, which do not depend on  $\bar{H}_a$ , and that, unlike the MPI, can be computed in practice. To do so, we show that the MPI can be approximated by a linear combination of known functions of  $z_{jt}$ . We call these interval instruments in reference to the way these functions are derived. Our feasible MPI is simply the vector of the interval instruments. The cost to incur for feasibility is that the properties we established for the MPI do not carry over to the feasible MPI. Nevertheless, our Monte Carlo simulations in Section 6 show that the interval instruments perform very well in practice.

By construction, in the BLP demand model, the correction term writes:

$$\begin{aligned}\Delta_{0,a}^{\xi_{jt}} &= x'_{1jt}(\beta_a - \beta_0) + \rho_j^{-1}(s_t, x_{2t}, f_0) - \rho_j^{-1}(s_t, x_{2t}, f_a) \\ &= x'_{1jt}(\beta_a - \beta_0) + \Delta_j(s_t, x_{2t}, f_0, f_a).\end{aligned}\tag{6}$$

The previous equation shows that the correction term is the sum of a linear part, which is standard, and a non-linear part which is specific to the BLP demand model.

**Linear part.** The linear part of the MPI writes:  $\mathbb{E}[x_{1jt}|z_{jt}]'(\beta_a - \beta_0) = \mathbb{E}[x_{1jt}|z_{jt}]'\gamma$ . Thus, for its linear part, the MPI is a linear combination of the conditional expectation of  $x_{1jt}$  with respect to the exogenous variables with unknown weights. If one is interested in specifically testing that  $\beta = \beta_0$ , informative instruments simply consist of the variables in  $\mathbb{E}[x_{1jt}|z_{jt}]$ .

**Non-linear part.** The non-linear part,  $\Delta_j(s_t, x_{2t}, f_0, f_a)$ , is the part which is implied by the misspecification on the distribution of RCs and for which we need to recover a feasible approximation. Equation (6) indicates that the non-linear part is the difference between the inverse demand functions generated by  $f_0$  and  $f_a$ . We now go one step further and derive two analytical approximations of  $\Delta_j(s_t, x_{2t}, f_0, f_a)$  which we then use as building blocks to construct our feasible approximations of the MPIs. The first approximation is based on a local expansion around  $f_0$ . The second approximation is based on an identity that is valid everywhere. The first approximation is more precise locally whereas the second one is more robust to large deviations from  $f_0$ .

## 4.1 Local approximation

First, we consider a local approximation of  $\Delta_j(s_t, x_{2t}, f_0, f_a)$ . This approximation corresponds to the first order term in the expansion of  $\Delta(s_t, x_{2t}, f_0, f_a)$  “around  $f_0$ ”, which is recovered by exploiting the properties of the inverse demand function, which is both  $\mathcal{C}^\infty$  and bijective in  $s_t$ .

### Proposition 6.

A first order expansion of  $\Delta(s_t, x_{2t}, f_0, f_a)$  around  $f_0$  writes:

$$\Delta(s_t, x_{2t}, f_0, f_a) = \left( \frac{\partial \rho(\delta_t^0, x_{2t}, f_0)}{\partial \delta} \right)^{-1} \int_{\mathbb{R}^{K_2}} \left[ \frac{\exp\{\delta_t^0 + x_{2t}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v\}} - \rho(\delta_t^0, x_{2t}, f_0) \right] f_a(v) + \mathcal{R}_0,$$

with  $\delta_t^0 = \rho^{-1}(s_t, x_{2t}, f_0)$  and  $\mathcal{R}_0 = o\left(\int_{\mathbb{R}^{K_2}} |f_a(v) - f_0(v)| dv\right)$ .

The proof is in Appendix B.3.1. We first observe that for any density  $f_0$ , we can construct artificial market shares  $s_t^0$  such that  $\rho^{-1}(s_t, x_{2t}, f_a) = \rho^{-1}(s_t^0, x_{2t}, f_0)$ . Then, we recover the final result by taking a Taylor expansion of  $\rho^{-1}(s_t^0, x_{2t}, f_0)$  around  $s_t$  and showing that the remainder is bounded.<sup>11</sup> This approximation is local by design: it works best when  $f_a$  is a local deviation from  $f_0$ , even if it can be used more generally. To make this expression useful in practice, we must still overcome two difficulties. The distribution  $f_a$  is unknown to the econometrician. In addition, some variables such as  $\delta_{jt}^0$  are endogenous. However, notice that the previous expression may be particularly useful if the econometrician is interested in testing  $\bar{H}_0$  against a fixed and known alternative as we did in the previous section.

---

<sup>11</sup>The expansion is taken around  $s_t$  because  $s_t^0$  depends on  $f_a$  and is thus unknown to the researcher.

**Discretizing the integral.** To solve for the fact  $f_a$  is unknown to the econometrician, we replace the integral in which  $f_a$  appears by a finite Riemann approximation. Namely,

$$\int_{\mathbb{R}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v\}} f_a(v) dv \approx \sum_{l=1}^L \omega_l(f_a) \frac{\exp\{x'_{2jt}v_l\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v_l\}},$$

with  $\{v_l\}_{l=1,\dots,L}$  the points chosen in the domain of definition of  $f_a$ , and  $\{\omega_l(f_a)\}_{l=1,\dots,L}$  the associated weights.<sup>12</sup> We provide more details on how to choose the points in Appendix C.4. It is important to observe that in the Riemann approximation, only the weights depend on the alternative  $f_a$ . This approximation can also be interpreted as approaching a continuous distribution with a discrete one, where each point in  $\{v_l\}_{l=1,\dots,L}$  represents a specific consumer type with an associated probability  $w_l(f_a)$ . The non-linear part of the MPI can thus be approximated as follows:

$$\mathbb{E}[\Delta_j(s_t, x_{2t}, f_0, f_a) | z_{jt}] \approx \sum_{l=1}^L \omega_l(f_a) \mathbb{E}[\pi_{j,l}(s_t, x_t) | z_{jt}],$$

with  $\pi_{j,l}(s_t, x_t) = \left( \frac{\partial \rho(\delta_t^0, x_{2t}, f_0)}{\partial \delta} \right)^{-1} \left[ \frac{\exp\{\delta_t^0 + x_{2t}v_l\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v_l\}} - \rho(\delta_t^0, x_{2t}, f_0) \right]_j$ .

**Approximating the conditional expectation.** Ideally, we would like to estimate the conditional expectation of  $\pi_{j,l}(s_t, x_t)$  with respect to  $z_{jt}$ . The endogenous variables are  $\{\delta_{jt}^0\}_{j=1,\dots,J}$ , and the potential endogenous variables in  $\{x_{2jt}\}_{j=1,\dots,J}$ , which often include prices. In practice, computing the conditional expectation is challenging because the dimension of  $z_{jt}$  can be very large and the functions  $\pi_{j,l}(\cdot)$  are highly non-linear and non-separable in the endogenous variables. This makes it unappealing to use standard non-parametric estimation methods.<sup>13</sup> In the same spirit as [Reynaert and Verboven \(2014\)](#), we first project the endogenous variables on the space spanned by a relevant subset of  $z_{jt}$ . We mark the projected endogenous variables with a hat and

<sup>12</sup>In the usual Riemann sum, the weights correspond to density evaluated at point  $v_l$ :  $f_a(v_l)$  times the width of the interval around  $v_l$ .

<sup>13</sup>For instance, a Sieve nonparametric estimator of the conditional mean. The dimension of  $z_{jt}$  makes this approach of little relevance in practice.

we plug them into our functions  $\pi_{j,l}(\cdot)$ . Namely, we have the following approximation for every interval instrument  $l$ :

$$\mathbb{E}[\pi_{j,l}(s_t, x_t)|z_{jt}] \approx \hat{\pi}_{j,l}(z_{jt}) = \left( \frac{\partial \rho(\hat{\delta}_t^0, \hat{x}_{2t}, f_0)}{\partial \delta} \right)^{-1} \left[ \frac{\exp\{\hat{\delta}_t^0 + \hat{x}_{2t}v_l\}}{1 + \sum_{k=1}^J \exp\{\hat{\delta}_{kt}^0 + \hat{x}'_{2kt}v_l\}} - \rho(\hat{\delta}_t^0, \hat{x}_{2t}, f_0) \right]_j.$$

We show in Appendix C.2 that this strategy yields an estimator of the conditional expectation that converges faster to a first order approximation of the conditional expectation.

**Test procedure.** From what precedes, the MPI (for its non-linear part) can be approximated as follows:  $h_D^*(z_{jt}) \approx \sum_{l=1}^L \omega_l(f_a) \hat{\pi}_{j,l}(z_{jt})$ . As we don't know the weights  $\omega_l(f_a)$ , we propose to take the vector  $\hat{\pi}_j(z_{jt}) = (\hat{\pi}_{j,1}(z_{jt}), \dots, \hat{\pi}_{j,L}(z_{jt}))'$  as our testing instruments. We call them interval instruments in reference to the way we divide the support into several intervals to construct this approximation. Following the test procedure presented in Section 3.1, we perform a moment based test for  $\bar{H}_0 : \mathbb{E}[\hat{\pi}_j(z_{jt})\xi_{jt}(f_0, \beta_0)] = 0$ . Under the same assumptions as in Proposition 2 and setting  $h_D(z_{jt}) = \hat{\pi}_j(z_{jt})$ , we have the following:

$$\text{Under } H_0 : S_T(h_D, f_0, \beta_0) \xrightarrow[T \rightarrow +\infty]{d} \chi_L^2.$$

This approach has the advantage of being feasible since we can construct the vector of interval instruments  $\hat{\pi}_j(z_{jt})$ , while remaining completely agnostic about  $f_a$ . The price to pay is that we lose the optimality properties of the MPI. We further discuss the properties of the feasible MPI in Appendix C.7. Moreover, the infeasible MPI,  $h_D^*(z_{jt})$ , is of dimension one and its test statistic is distributed as  $\chi_L^2$  asymptotically. In contrast, the feasible MPI is of dimension  $L$  and its asymptotic distribution is a  $\chi_L^2$ . This increase in the number of degrees of freedom may lead to some loss of power. An alternative approach would consist in letting the researcher choose the weights  $\{\hat{\omega}_l\}_{l=1,\dots,L}$  and recover an instrument of dimension one. However, for this approach to work well and retain good power properties, the econometrician must choose the weights so that they approximately match the real weights  $\{\omega_l(f_a)\}_{l=1,\dots,L}$ . This requires a good prior knowledge of the cumulative distribution function of the alternative distribution  $f_a$ . Nevertheless, our Monte Carlo simulations in Section 6 show that the feasible MPIs that we propose perform very well in practice.



## 4.2 Global approximation

Second, we consider a global approximation that is based on an identity which is valid everywhere and not only when  $f$  is close to  $f_a$ . Simple algebraic operations (see Appendix B.3.2) allow us to derive the following expression for  $\Delta_j(s_t, x_{2t}, f_0, f_a)$ . Let  $\delta_{jt}^0 = \rho_j^{-1}(s_t, x_{2t}, f_0)$  and  $\delta_{jt}^a = \rho_j^{-1}(s_t, x_{2t}, f_a)$ . We have:

$$\Delta_j(s_t, x_{2t}, f_0, f_a) = \log \left( \frac{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^a + x'_{2kt}v\}} f_a(v) dv}{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v\}} f_0(v) dv} \right).$$

As for the local approximation, we cannot directly exploit this formula as some quantities such as  $f_a$  and  $\delta_{jt}^a$  are unknown and some variables such as  $\delta_{jt}^0$  are endogenous. To remedy these two difficulties, we apply the same methods as previously described: we discretize the integral, and we project the endogenous variables onto the space spanned by a relevant subset of  $z_{jt}$ . To solve for the fact that the mean utility  $\delta_{jt}^a$  under the alternative is unknown, we replace it with the mean utility under the null  $\delta_{jt}^0$ . This should not alter the approximation too much given that  $\delta_{jt}^a$  only enters the expression at the denominator within a sum, which averages out the differences between  $\delta_{jt}^a$  and  $\delta_{jt}^0$  across products. In the end, we are able to provide the following approximation for the non-linear part of the MPI:

$$\mathbb{E}[\Delta_j(s_t, x_{2t}, f_0, f_a) | z_{jt}] \approx \log \left( \sum_{l=1}^L \bar{\omega}_l(f_a) \hat{\pi}_{j,l}(z_{jt}) \right) \text{ with } \hat{\pi}_{j,l}(z_{jt}) = \frac{\frac{\exp\{x'_{2jt}v_l\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v_l\}}}{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v\}} f_0(v) dv},$$

where  $\{\bar{\omega}_l(f_a)\}_{l=1,\dots,L}$  correspond to the unknown weights and the  $\hat{\pi}_{j,l}(z_{jt})$  are set of global interval instruments. The MPI can thus be approximated by the logarithm of a weighted sum of known functions of  $z_{jt}$ . As we did previously, we use  $\hat{\pi}_j(z_{jt}) = (\hat{\pi}_{j,1}(z_{jt}), \dots, \hat{\pi}_{j,L}(z_{jt}))'$  as instruments to test  $\bar{H}_0$ . All the weights are positive and sum to one, which entails that the non-linear part of the correction term is an increasing function of our instruments. This approximation is said to be global because contrary to the second approximation we study, it does not require  $f_0$  to be close to  $f_a$ . Nevertheless, if  $f_a$  is close to  $f_0$ , then the fraction  $\kappa$  inside the logarithm is close to 1 and the well-known approximation  $\log(\kappa) \approx \kappa - 1$  allows us to directly rewrite the MPI as a linear combination of our instruments.

Overall, the feasible MPIs that we derive in this section allows us to approximate the most powerful instrument against a fixed alternative while remaining agnostic about this alternative.

### 4.3 Feasible MPIs for estimation

In the estimation framework, the researcher stipulates that  $f$  belongs to a parametric family  $\mathcal{F}_0 = \{f_0(\cdot|\tilde{\lambda}) : \tilde{\lambda} \in \Lambda_0\}$  and wants to estimate the true parameter  $\theta_0 = (\beta'_0, \lambda'_0)'$  under this parametric restriction. From the connection between the MPI and the local instruments that we present in Section 3.3, we can infer that good estimation instruments  $h_E(z_{jt})$  ought to approximate the MPI devoted to testing  $H_0 : \theta = \theta_0$  against any local alternative. If we have an initial estimator of  $\theta_0$ , we can directly use the interval instruments presented previously to approximate the MPI devoted to testing  $H_0 : \theta = \theta_0$  against an unknown alternative. The fact that the feasible MPIs do not depend on the alternative is key for estimation. Moreover, the transformation of the MPI into a vector of instruments of dimension  $L \geq |\lambda_0|$  is necessary for estimation as the number of instruments must be greater than the dimension of the parameter to estimate.<sup>14</sup> In Appendix C.5, we propose a version of the interval instruments that does not require a first step estimate of  $\theta_0$  and that can be computed directly from the logit specification.

## 5 Composite hypothesis

In the traditional estimation procedure, which encompasses almost all the applications of the BLP model, the econometrician must make a parametric assumption on the distribution of random coefficients to estimate the model. Formally, the econometrician assumes  $f$  belongs to a parametric family  $\mathcal{F}_0 = \{f_0(\cdot|\tilde{\lambda}) : \tilde{\lambda} \in \Lambda_0\}$ , where  $\tilde{\lambda}$  is a parameter that must be estimated. In applied work, researchers typically assume that  $f$  is normally distributed. This parametric choice is rarely grounded in economic theory and, if too restrictive, is likely to impose arbitrary restrictions on some key counterfactual quantities such as the pass-through. In this section, we develop a formal specification test for  $H_0 : f \in \mathcal{F}_0$ . In comparison to the test in Section 3.1, we must now estimate the parameters of the distribution  $\theta_0 = (\beta'_0, \lambda'_0)'$  in a first step, which generates parameter

---

<sup>14</sup>The linear parameter  $\beta_0$  has its own instruments, which are simply the variables in  $x_{1jt}$ .

uncertainty. Moreover, we propose a rigorous treatment of the numerical approximations involved in the derivation of the structural error  $\xi_{jt}(\tilde{\theta})$ . We organize this section as follows. First, we define the pseudo-true value associated with a given specification and the first stage estimator. Second, we define our test procedure and its implementation in practice. Finally, we study the asymptotic properties of our test.

## 5.1 Pseudo-true value and first stage estimator

To estimate the BLP model, researchers must make three choices. They must choose the parametric family  $\mathcal{F}_0$ , the instruments  $h_E(z_{jt})$  to estimate the model, and a weighting matrix  $W$ , which weights the different moments included in the objective function. Given these three choices, we can define the BLP pseudo-true value  $\theta(\mathcal{F}_0, h_E, W) \equiv \theta_0 = (\beta'_0, \lambda'_0)'$  as follows:<sup>15</sup>

$$\theta(\mathcal{F}_0, h_E, W) \in \underset{\tilde{\theta}}{\text{Argmin}} \mathbb{E} \left[ \xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) h_E(z_{jt}) \right]' W \mathbb{E} \left[ h_E(z_{jt}) \xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) \right].$$

If the model is well-specified ( $f \in \mathcal{F}_0$ ) and the pseudo-true value is unique, then the pseudo-true value is the true value:  $\theta_0 = \theta$ . Under misspecification,  $\theta_0$  is a parameter whose value depends on  $(\mathcal{F}_0, h_E, W)$ . For exposition purposes, we omit this dependence in the subsequent analysis. Moreover, here we remain general and do not impose that  $W$  must be equal to the usual optimal weighting matrix. It is often the case in practice, that the researchers choose the identity matrix or regularize the weighting matrix.

**First stage estimator  $\hat{\theta}$ .** The first stage estimator is an empirical counterpart of the BLP pseudo-true value defined previously. The minimization is done with respect to sample analogs. Additionally, we know that there is no closed form expressions for the structural error  $\xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta})$ , and thus, we must use a feasible counterpart  $\hat{\xi}_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta})$  instead.

$$\hat{\theta}(\mathcal{F}_0, h_E, \hat{W}) \equiv \hat{\theta} = \underset{\hat{\theta}}{\text{Argmin}} \left( \sum_{j,t} \hat{\xi}_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) h_E(z_{jt}) \right)' \hat{W} \left( \sum_{j,t} \hat{\xi}_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) h_E(z_{jt}) \right). \quad (7)$$

---

<sup>15</sup>Our definition of a pseudo-true value is closely related to the approach in [White \(1982\)](#) in the context of maximum likelihood. In his case, the pseudo true value minimizes the Kullback-Leibler distance between the assumed likelihood and the true likelihood, whereas in our case, the pseudo-true value minimizes a weighted sum of population moments.

The construction of the feasible structural error  $\hat{\xi}_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta})$  requires the following 3 numerical approximations:

1. The econometrician does not observe a continuum of consumers as in the theoretical model but only empirical averages  $\hat{s}_{jt}$  over the  $n_t$  individuals in market  $t$ .

$$\hat{s}_{jt} = \frac{1}{n_t} \sum_{i=1}^{n_t} y_{ijt}, \quad (8)$$

where  $y_{ijt} \in \{0; 1\}$  are i.i.d. choices over the  $i = 1, \dots, n_t$ .

2. There is no closed form for  $\rho_j(\cdot, x_{2t}, f_0(\cdot|\tilde{\lambda}))$ , the integral has to be computed through numerical integration. A prominent example is Monte Carlo integration:

$$\hat{\rho}_j(\delta, x_{2t}, f_0(\cdot|\tilde{\lambda})) = \frac{1}{R} \sum_{r=1}^R \frac{\exp\{\delta_j + x'_{2jt} v_r\}}{1 + \sum_{k=1}^{J_t} \exp\{\delta_k + x'_{2kt} v_r\}}, \quad (9)$$

with  $v_r$  iid draws from  $f_0(\cdot|\tilde{\lambda})$ .

3. There is no analytical way to recover the inverse of the demand functions  $\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))$ . The most popular way to derive the inverse demand is by solving the following contraction:

$$C : (\cdot, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) : \delta \mapsto \delta + \log(s_t) - \log(\rho(\delta, x_{2t}, f_0(\cdot|\tilde{\lambda}))).$$

This solution has given rise to the popular nested fixed point GMM procedure.<sup>16</sup>

In Section 5.3, we explicitly state the assumptions that allow us to neglect these approximations asymptotically.

## 5.2 Test procedure

Under Assumption A, and assuming  $h_E(z_{jt})$  and  $W$  are such that the pseudo-true value  $\theta_0$  is unique, the following equivalence holds:

$$\begin{aligned} H_0 : f \in \mathcal{F}_0 &\iff \bar{H}_0 : (f, \beta) = (f_0(\cdot|\lambda_0), \beta_0) \\ &\iff \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)|z_{jt}] = 0 \text{ a.s..} \end{aligned}$$

---

<sup>16</sup>Another solution that has gained traction in the literature is the MPEC procedure (Dubé et al. (2012)) that replaces the BLP inversion at each step of the minimization by imposing equilibrium constraints on the minimization program.

The pseudo true value reduces the dimensionality of the problem by allowing us to move from a composite hypothesis  $H_0 : f \in \mathcal{F}_0$  to the simple hypothesis  $\bar{H}_0 : (f, \beta) = (f_0(\cdot|\lambda_0), \beta_0)$  studied previously. As we did in Section 2, we propose a moment-based test of  $H_0$ .<sup>17</sup> Under  $H_0$ , for every set of testing instruments  $h_D(z_{jt})$ , the following moment conditions must hold:

$$H_0 : f \in \mathcal{F}_0 \iff \bar{H}_0 : (f, \beta) = (f_0(\cdot|\lambda_0), \beta_0) \implies \bar{H}'_0 : \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)h_D(z_{jt})] = 0.$$

We now develop a procedure to test  $\bar{H}'_0$ . In comparison to the test in Section 3.1, we must now account for the fact that the pseudo-true value needs to be estimated to derive the test statistic, which generates parameter uncertainty. Moreover, we propose a rigorous treatment of the numerical approximations involved in the derivation of the structural error.

**Test statistic.** For any choice of testing instruments  $h_D(z_{jt})$ , our objective is to test  $\bar{H}'_0 : \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)h_D(z_{jt})] = 0$  where  $\theta_0 = (\beta'_0, \lambda'_0)'$  is the pseudo-true value associated with the parametric family  $\mathcal{F}_0$ .<sup>18</sup> In order to test  $\bar{H}_0$ , we consider the following Wald test statistic:

$$S_T(h_D, \mathcal{F}_0, \hat{\theta}) = TJ \left( \frac{1}{TJ} \sum_{j,t} \hat{\xi}_{jt}(f_0(\cdot|\hat{\lambda}), \hat{\beta})h_D(z_{jt}) \right)' \hat{\Sigma} \left( \frac{1}{TJ} \sum_{j,t} \hat{\xi}_{jt}(f_0(\cdot|\hat{\lambda}), \hat{\beta})h_D(z_{jt}) \right).$$

where  $\hat{\Sigma}$  is a weighting matrix chosen by the econometrician and  $\hat{\theta} = (\hat{\beta}, \hat{\lambda})$  is a consistent estimator of  $\theta_0$ . The number of markets  $T$  is the dimension that we let grow to infinity to the asymptotic properties of our test. We motivate this choice in Appendix C.3. Under some regularity conditions

---

<sup>17</sup>Other testing approaches could have been considered. First, one could use the previous equivalence to directly test  $H_0$  via an integrated conditional moment test. We do not follow this route for at least two reasons. First, this test will contain no information on the nature of the misspecification (it could be completely unrelated to the distribution of RC). Second, in practice the dimension of  $z_{jt}$  is often very large, which substantially reduces the power of this kind of test. Another testing approach would have entailed testing  $H_0 : f \in \mathcal{F}_0$  against a larger class of densities that encompasses  $\mathcal{F}_0$ . For instance, if  $\mathcal{F}_0$  is the family of normal distributions, encompassing families are mixtures of Gaussians with a larger number of components. We do not follow this route for two reasons. First, it is not desirable to restrict the alternative to a class of distributions that encompass the null as the econometrician does not know a priori the misspecification. Second, estimating the BLP model with a more flexible parametrization is challenging. An advantage of our test procedure is that it doesn't require estimating the model with a more flexible parametrization.

<sup>18</sup>Remember that under an alternative specification, the pseudo true value also depends on the estimation instruments  $h_E(z_{jt})$  and the weighting matrix.

that we make explicit in the following section, the asymptotic distribution of the test statistic under  $\bar{H}'_0$  is as follows:

$$S_T(h_D, \mathcal{F}_0, \hat{\theta}) \xrightarrow{d} Z' \Sigma Z, \quad (10)$$

$$\text{with } \frac{1}{\sqrt{T}} \sum_{t=1}^T \sum_{j=1}^J \hat{\xi}_{jt}(f_0(\cdot|\hat{\lambda}), \hat{\beta}) h_D(z_{jt}) \xrightarrow{d} Z \sim \mathcal{N}(0, \tilde{\Omega}_0). \quad (11)$$

$\Sigma$  is the probability limit of  $\hat{\Sigma}$ . We make  $\tilde{\Omega}_0$  explicit in the next subsection (in particular, the derivation of  $\tilde{\Omega}_0$  takes into account parameter uncertainty). Given that  $\hat{\Sigma}$  is chosen by the econometrician and it is possible to derive a consistent estimator of  $\tilde{\Omega}_0$ , the econometrician can always simulate the asymptotic distribution of the test statistic. In some polar cases, which we present hereafter, the asymptotic distribution of our test statistic is pivotal chi-square distribution that does not require to be simulated.

**Two polar cases.** For the sake of exposition, let us now describe two polar cases where the asymptotic distributions are pivotal chi-square distributions, which do not require to be simulated. Denote by  $|\cdot|_0$  the counting norm.

1. **Sargan-Hansen J test:** If the set of estimation instruments and the set of testing instruments are the same ( $h_E = h_D$ ), if  $\hat{W}$  is the 2-step GMM optimal weighting matrix and if  $\hat{\Sigma} = \hat{W}^{-1}$ , then our test boils down to the usual Sargan-Hansen J test and we have under  $\bar{H}'_0$ :

$$S_T(h_D, \mathcal{F}_0, \hat{\theta}) \xrightarrow{d} \chi^2_{|h_E|_0 - |\theta|_0}.$$

2. **Non-redundant  $h_D$  and  $h_E$ :** if  $\tilde{\Omega}_0$  has full rank and if the econometrician sets  $\hat{\Sigma} = \hat{\tilde{\Omega}}_0^{-1}$ , then our test statistic has the following asymptotic distribution under  $\bar{H}'_0$ :<sup>19</sup>

$$S_T(h_D, \mathcal{F}_0, \hat{\theta}) \xrightarrow{d} \chi^2_{|h_D|_0}.$$

---

<sup>19</sup>If  $\Omega_0$  is singular, one can always use directly the asymptotic distribution in 10 or apply the singularity-robust procedure proposed in Andrews and Guggenberger (2019).

**Choice of the testing instruments.** As previously indicated, the power properties of our test hinge critically on the choice of testing instruments  $h_D(z_{jt})$ . We established that the MPI and its feasible counterparts, the interval instruments, feature attractive properties in testing  $\bar{H}_0 : (f, \beta) = (f_0(\cdot|\lambda_0), \beta_0)$  against any fixed alternative. Thus, it is natural to use these instruments for the specification test above. In particular, we show that the consistency of the test with the MPI carries over to the general specification test above in Appendix B.5.

### 5.3 Asymptotic validity

We now study the asymptotic properties of our test when the number of markets  $T$  goes to infinity. To establish the asymptotic validity and consistency of our test, we exploit classical results on the asymptotic normality of the non-linear GMM estimator (Hansen (1982), Newey (1990)) as well on the large- $T$  asymptotics of the BLP estimator (Freyberger (2015)). The main challenge here is to control the magnitude of the approximations that intervene in the derivation of the structural error so that they can be neglected asymptotically. Contrary to Freyberger (2015), we do not assume the convergence of any moments ex-ante and we allow for the approximation error between demand and observed market shares to be non-zero.

#### Assumption B.

- (i)  $(s_t, x_t, z_t)_{t=1}^T$  are i.i.d. across markets and are consistent with the probability model defined by equations (1), (2) and (3) evaluated at  $(f, \beta)$ ;
- (ii) Strong Exogeneity:  $\mathbb{E}[\xi_{jt}(f, \beta)|z_{jt}] = 0$  a.s.;
- (iii) Finite moment conditions:  $x_{2t}$  has bounded support and  $x_{1t}$  has finite 4th moments.

In B(i), we assume that the data are i.i.d. across markets, an assumption which we could relax slightly (technically, only certain moments need to be identical across markets), and that the data are generated by the BLP demand model at a given pair  $(f, \beta)$ . In B(ii), we assume exogeneity of our instrumental variables. Let us stress that to show the asymptotic validity of our specification test, we do not require  $(f, \beta)$  to be non-parametrically identified, as we just need parametric identification under  $H_0$ . In particular, we do not need all the assumptions in A. B(iii) is

a necessary condition to recover the asymptotic normality of the BLP estimator.  $x_{1t}$  having finite 4th moments is standard.  $x_{2t}$  having bounded support has two purposes. First, it implies that the structural error has a finite 4th moment, [Compiani \(2018\)](#) makes the same assumption on price for this purpose. Second, it ensures that the mapping used in the nested fixed point algorithm is a proper smooth contraction, which allows us to prove that the NFP algorithm converges (without truncating the contraction mapping as in [Berry \(1994\)](#) and [Berry et al. \(1995\)](#)) and control for the NFP approximation bias.

### Assumption C.

$\mathcal{F}_0$  is such that :

- (i)  $\lambda_0$  belongs to the interior of  $\Lambda_0$  with  $\Lambda_0$  compact;
- (ii)  $\tilde{\lambda} \mapsto \rho(\delta, x_{2t}, f_0(\cdot|\tilde{\lambda}))$  is well defined and continuously differentiable on  $\Lambda_0$ .

In [C\(i\)](#), we assume that, for any given DGP, the associated pseudo-true-value  $\lambda_0$  associated with the family  $\mathcal{F}_0$  lies in a compact space  $\Lambda_0$ . This condition is standard in establishing the consistency and asymptotic normality of M-estimators. Second in [C\(ii\)](#), we impose that the demand function and its derivative with respect to  $\lambda$  should both be well defined and continuous.

Next, we impose conditions on the instruments that are used for estimation  $h_E(z_{jt})$  and for testing  $h_D(z_{jt})$  and on the BLP estimator itself.

### Assumption D.

For a given  $\mathcal{F}_0$  that satisfies Assumption [C](#) and for some weighting matrix  $W$  and  $\Sigma$ , the following conditions must hold:

- (i) Finite moments for instruments:  $h_E(z_{jt})$  and  $h_D(z_{jt})$  are not perfectly colinear and have finite 4th moments;
- (ii) Global identification of  $\theta_0$ :  $\exists! \theta_0$  such that  $\forall \tilde{\theta} \neq \theta_0$ :

$$\mathbb{E} \left[ \sum_j \xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) h_E(z_{jt})' \right] W \mathbb{E} \left[ \sum_j h_E(z_{jt}) \xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) \right] > \mathbb{E} \left[ \sum_j \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) h_E(z_{jt})' \right] W \mathbb{E} \left[ \sum_j h_E(z_{jt}) \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) \right];$$



(iii) Local identification:  $\Gamma(\mathcal{F}_0, \theta_0, h_E) = \mathbb{E} \left[ \sum_j h_E(z_{jt}) \frac{\partial \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)}{\partial \theta'} \right]$  and  $\Gamma(\mathcal{F}_0, \theta_0, h_D)$  have full column rank;

(iv)  $W$  and  $\Sigma$  are symmetric positive definite and  $\hat{W} \xrightarrow{P} W$ ,  $\hat{\Sigma} \xrightarrow{P} \Sigma$ ;

(v)  $\hat{\theta}$  minimizes objective function (7) and satisfies the FOC of the minimization problem:

$$\left( \sum_{j,t} \frac{\partial \hat{\xi}_{jt}(f(\cdot|\hat{\lambda}), \hat{\beta})}{\partial \theta} h_E(z_{jt}) \right)' \hat{W} \left( \sum_{j,t} \hat{\xi}_{jt}(f(\cdot|\hat{\lambda}), \hat{\beta}) h_E(z_{jt}) \right) = 0.$$

Assumption D restricts the class of instruments which can be used for estimation and for testing. More specifically, D(i) and D(iii) are common regularity conditions necessary to establish asymptotic results whereas D(ii) is an identification condition which ensures that the pseudo true value  $\theta_0$  is uniquely defined, which is critical to show the consistency of the BLP estimator. Finally, Assumptions D(iv) and D(v) impose regularity conditions on the weighting matrix as well as on the BLP estimator itself.

The next assumptions ensure that the numerical approximations involved in the derivation of the structural error do not interfere with the asymptotic theory.

### Assumption E.

(i) Let  $n_t$  be the number of individuals in market  $t$ ,  $(n_t)_{t=1}^T$  is i.i.d. and independent from all other variables. First it must be that  $\forall t \quad \sqrt{T} \mathbb{E}(n_t^{-1/2}) \xrightarrow{T \rightarrow +\infty} 0$ . Second observed market share  $\hat{s}_t$  in market  $t$  must write:

$$\hat{s}_{jt} = \frac{1}{n_t} \sum_{i=1}^{n_t} y_{ijt},$$

with  $(y_{ijt})_{i=1}^{n_t}$  i.i.d. draws generated by the BLP demand model at a given pair  $(f, \beta)$  conditional on  $(x_t, \xi_t)$ .

(ii) Let  $R$  be the number of simulations, then the simulated demand for product  $j$  writes:

$$\hat{\rho}_{jt}(\delta, x_{2t}, f_0(\cdot|\tilde{\lambda})) = \frac{1}{R} \sum_r \frac{\exp\{\delta_j + x'_{2jt} v_r\}}{1 + \sum_k \exp\{\delta_k + x'_{2kt} v_r\}},$$

where  $v_r \stackrel{iid}{\sim} f_0(\cdot|\tilde{\lambda})$ , and  $\frac{T}{R} \xrightarrow{T \rightarrow +\infty} 0$ .

(iii) Let  $H$  be the stopping time for the contraction (which depends on  $T$ ) and  $\epsilon$  the fixed Lipschitz

constant of the contraction mapping used to invert the demand function, then it must be that  $\sqrt{T}\epsilon^H \xrightarrow{T \rightarrow +\infty} 0$ .

A sufficient condition for **E(i)** to hold is that the minimum number of individuals observed in any market is of higher order than the total number of markets. This condition can be checked in practice.<sup>20</sup> Assumptions **E(ii)** and **E(iii)** can also be checked in practice and are more manageable because  $R$  and  $H$  are chosen by the researcher and can always be increased so that these assumptions hold.

Given our assumptions, we derive the asymptotic distribution of our test statistic under the null, and show that the test is consistent.

**Theorem 5.1.** Let  $\hat{\theta} = \hat{\theta}(\mathcal{F}_0, \hat{W}, h_E)$  be the BLP estimator associated with distributional assumption  $\mathcal{F}_0$ , weighting matrix  $\hat{W}$ , estimating instruments  $h_E$ . Under assumptions **B-E**,

- Under  $\bar{H}'_0 : \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)h_D(z_{jt})] = 0$ ,

$$S_T(h_D, \mathcal{F}_0, \hat{\theta}) \xrightarrow{T \rightarrow +\infty} Z'\Sigma Z, \quad Z \sim \mathcal{N}(0, \tilde{\Omega}_0),$$

where 
$$\tilde{\Omega}_0 = \begin{pmatrix} I_{|h_D|_0} & G \end{pmatrix} \begin{pmatrix} \Omega(\mathcal{F}_0, h_D) & \Omega(\mathcal{F}_0, h_D, h_E) \\ \Omega(\mathcal{F}_0, h_D, h_E)' & \Omega(\mathcal{F}_0, h_E) \end{pmatrix} \begin{pmatrix} I_{|h_D|_0} \\ G' \end{pmatrix},$$

$$\Omega(\mathcal{F}_0, h_D, h_E) = cov\left(\sum_j \xi_{jt}(f(\cdot|\lambda_0), \beta_0)h_D(z_{jt}), \sum_j \xi_{jt}(f(\cdot|\lambda_0), \beta_0)h_E(z_{jt})\right),$$

$$G = -\Gamma(\mathcal{F}_0, \theta_0, h_D) [\Gamma(\mathcal{F}_0, \theta_0, h_E)'W\Gamma(\mathcal{F}_0, \theta_0, h_E)]^{-1} \Gamma(\mathcal{F}_0, \theta_0, h_E)'W.$$

- Under  $H'_a : \mathbb{E}[h_D(z_{jt})\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)] \neq 0$ ,

$$\forall q \in \mathbb{R}^+, \quad \mathbb{P}(S_T(h_D, \mathcal{F}_0, \hat{\theta}) > q) \xrightarrow{T \rightarrow +\infty} 1.$$

---

<sup>20</sup>Note that by making stronger assumptions on the higher moments and the support of the observed characteristics, it is possible to find milder conditions on the number of individuals relative to the number of markets.

The proof of Theorem 5.1 is in Appendix B.4 and comprises three main steps. First, we show that under the assumptions in E, the numerical approximation becomes asymptotically negligible. Second, we show the consistency and asymptotic normality of the BLP estimator. Finally, we derive the asymptotic distribution of the test statistic, taking into account parameter uncertainty ( $\theta_0$  is estimated and not observed). The apparent complexity of the asymptotic variance-covariance matrix  $\Omega_0$  is a direct consequence of parameter uncertainty.

## 6 Monte Carlo experiments

In this section, we conduct three distinct sets of Monte Carlo experiments. First, we implement a simple simulation exercise to assess the effects of incorrectly specifying the distribution of random coefficients on quantities of interest such as price elasticities or cross-price elasticities, which are known to play a key role in shaping the counterfactuals. In a second set of Monte Carlo experiments, we study the finite sample performances of the specification test developed in Section 5 with different sets of testing instruments. We first examine the size of our test in finite sample. Then, we investigate the power properties of our test under alternative specifications (with alternatives including Gaussian mixtures, gamma distributions and local alternatives). We show that our test with the interval instruments significantly outperforms the traditional J-test with the usual instruments. Finally, in the last Monte Carlo exercise, we study the performance of the interval instruments to estimate the parameters of the model by means of comparison with the commonly used instruments in the literature.

### 6.1 Simulation design

For the sake of exposition, we will keep the same simulation design for all the simulation experiments. The simulation design closely follows the simulation design used in Dubé et al. (2012), Reynaert and Verboven (2014). The market includes  $J = 12$  products, which are characterized by 3 exogenous product attributes  $x_a$ ,  $x_b$  and  $x_c$  that follow a joint normal distribution. The price  $p$  is endogenous and depends on the observed and unobserved characteristics and on some cost shifters  $c_1$  and  $c_2$ . Consumer heterogeneity is present only in  $x_c$ , and the random coefficient  $v_i$

associated with  $x_c$  follows various distributions depending on the simulation exercise. The sample size  $T$  varies between 50, 100 and 200 markets. We can summarize the DGP as follows:

$$u_{ijt} = 2 + x_{ajt} + 1.5x_{bjt} - 2p_{jt} + x_{cjt}v_i + \xi_{jt} + \varepsilon_{ijt} \quad \xi_{jt} \sim \mathcal{N}(0, 1), \varepsilon_{ijt} \sim EV1,$$

$$\text{and } \begin{bmatrix} x_{a,j} \\ x_{b,j} \\ x_{c,j} \end{bmatrix} \sim N \left( \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 & -0.8 & 0.3 \\ -0.8 & 1 & 0.3 \\ 0.3 & 0.3 & 1 \end{bmatrix} \right),$$

$$p_{jt} = 1 + \xi_{jt} + u_{jt} + \sum_{k=a}^c x_{kjt} + c_{1jt} + c_{2jt} \quad \text{with } u_{j,t} \sim U[-4, -2], \quad c_{1jt} \sim U[2, 4] \text{ and } c_{2jt} \sim U[3, 5].$$


Market shares are generated by integrating over 20,000 consumers. This allows us to essentially remove the approximation error between the observed and theoretical market shares.

## 6.2 Counterfactuals under an alternative distribution

We now present a simple exercise to illustrate how the misspecification of random coefficients can affect the estimation of quantities of interest such as price elasticities and cross-price elasticities. To do so, we simulate data using the simulation design introduced above and we take various distributions for the random coefficient  $v_i$  (respectively: Gaussian mixture, Uniform, Chi-square, Exponential, Student, Gamma). We ensure that all the distributions have the same mean and variance (3 and 3, respectively). For each distribution, we simulate  $T = 100$  markets of data and we estimate the model either assuming no heterogeneity (simple logit) or assuming that  $v_i$  is normally distributed. We replicate the same exercise 500 times for each distribution. This allows us to recover the mean estimate for the parameters as well as to construct 95% “confidence bins” (by trimming the observations below the 2.5% quantile and above the 97.5% quantile). We plot the true densities and their estimated counterparts under the normal and logit assumptions in Figure 1. We observe that the estimated logit parameters and the estimated means of the normal distributions always coincide and are close to 3 for all the distributions. However, there is some

variation between the different specifications. For instance, the estimated means are larger with the exponential distribution. The estimated variances also vary from one specification to the other. The estimated variances for the exponential distribution are smaller, while they are larger for the student distribution.

Figure 1: True densities and estimated densities under normal and logit specifications



In a second stage, we simulate  $N = 5,000$  draws from the true distributions as well as from the estimated logit and normal approximations to compute the demand, the price-elasticity and the cross-price elasticity for the product  $j^*$  with the highest value for  $x_c$ .<sup>21</sup> The cross-price elasticity is arbitrarily taken for product  $j = 1$  with respect to  $p_{j^*}$ . We derive the quantities of interest for 100 equally spaced values of  $p_{j^*}$  ranging in  $]0, 10[$ . We plot the elasticities in Figure 2 and cross-price elasticities in Figure 3 generated by the true distribution as well as those generated by the logit and normal approximations, respectively. We proceed similarly with the demand functions. We see in Figure 9 in Appendix).

One can observe that, as expected, the logit specification poorly replicates the substitution pat-

<sup>21</sup>The expressions for both price-elasticities and the cross-price elasticities are in Appendix D.1.

terns. In particular, it consistently overstates the magnitude of the elasticities and cross-elasticities with respect to the true ones. The absence of consumer heterogeneity on characteristic  $c$  implies that consumers can “renounce” more easily to product  $j^*$  when its price increases. By introducing some heterogeneity, the normal approximation somewhat attenuates this issue. However, significant discrepancies in the shape of elasticities and cross-price elasticities remain. As most counterfactual analyzes rely on the substitution patterns generated by the model, these differences will inevitably create significant biases.

Figure 2: Price elasticities





Figure 3: Cross-price elasticities



### 6.3 Finite sample performance of the specification test

We now study the empirical size and power of our test under different sample sizes and for different sets of testing and estimating instruments. Once again, the data are generated according to the simulation design exhibited previously for various distributions of  $v_i$ . The assumption made throughout the simulations is  $H_0 : f \in \mathcal{F}_0$ , where  $\mathcal{F}_0$  is the family of normal distributions. In other words, we always assume that the random coefficient is normally distributed and we test this hypothesis. We set the nominal size to 5%. We study the finite sample performances of the specification test that we presented in Section 5 using different sets of estimation and testing instruments. For estimation, we take the instruments commonly adopted by practitioners: the differentiation instruments of [Gandhi and Houde \(2019\)](#) and the "optimal" instruments of [Reynaert and Verboven \(2014\)](#). Both of these sets are approximations of the classical optimal instruments. Second, we compare the performance of the test when performing the standard Sargan-Hansen J test (i.e. when we use the same instruments for testing and estimation) and when we use the global and local approximations of the MPI that we constructed in Sections 4.2 and 4.1. We denote the

latter tests as I Local and I Global respectively. The BLP estimator is computed following the NFP GMM procedure described in Section 5.1. For the optimization, only an analytic Jacobian is provided. We ensure that the number of tested restrictions is of the same magnitude across the different sets of instruments. More details on the exact sets of instruments and on the estimation procedure for this specific set of simulations are given in Appendix D.2.

### 6.3.1 Empirical size

The size is the probability of rejecting the null hypothesis when the null is true, so we compute the empirical size by counting and averaging the number of times we reject the null for nominal size 5% over the 1,000 simulations when the random coefficient  $v_i$  is normally distributed. Below in Table 1, we report the empirical sizes of the test with the different sets of instruments described above for the different sample sizes  $T \in \{50, 100, 200\}$  and for different distributions of the RC such that  $v_i \sim f \in \mathcal{F}_0$ .

Table 1: Empirical size for nominal size 5% (1000 replications)

Number of markets	T=50						T=100						T=200					
Estimation instruments	“Differentiation”			“Optimal”			“Differentiation”			“Optimal”			“Differentiation”			“Optimal”		
Test type	J	I	Global	I	local		J	I	Global	I	local		J	I	Global	I	local	
$v_i \sim \mathcal{N}(-1, 0.5^2)$	0.294	0.083	0.091	0.145	0.078	0.063	0.138	0.078	0.058	0.094	0.084	0.047	0.08	0.052	0.053	0.064	0.05	0.04
$v_i \sim \mathcal{N}(0, 0.75^2)$	0.293	0.084	0.085	0.148	0.081	0.071	0.137	0.061	0.06	0.1	0.059	0.05	0.074	0.053	0.045	0.062	0.048	0.036
$v_i \sim \mathcal{N}(1, 1^2)$	0.287	0.084	0.083	0.142	0.084	0.073	0.142	0.055	0.054	0.098	0.053	0.047	0.079	0.042	0.03	0.058	0.035	0.025
$v_i \sim \mathcal{N}(2, 2^2)$	0.288	0.087	0.077	0.145	0.071	0.072	0.138	0.069	0.051	0.099	0.053	0.056	0.077	0.044	0.041	0.069	0.037	0.044
$v_i \sim \mathcal{N}(3, 3^2)$	0.287	0.089	0.071	0.137	0.075	0.066	0.145	0.074	0.06	0.098	0.06	0.061	0.076	0.044	0.037	0.061	0.046	0.046

We observe that with a moderate sample size ( $T = 50, J = 12$ ), all the tests are over-sized. This is within expectations and due to the approximations inherent to the estimation of the BLP models as described in Section 5 and the relatively large number of instruments used for estimation and testing purposes.<sup>22</sup> However, we notice that the Sargan-Hansen J tests are much more over-sized than the tests with the interval instruments: the rejection rate is above 25% for

<sup>22</sup>The number of over-identifying restrictions lies between 6 and 8. The Sargan-Hansen J tests are known to suffer from size distortions as the number of instruments increases.



the Sargan-Hansen J test with differentiation instruments vs around 8% for the I test. Increasing the sample size improves the tests' empirical levels and shifts them towards the nominal level, which is a good indication of the validity of our test. Even with a relatively large number of markets ( $T = 200$ ), the Sargan-Hansen J tests remain slightly oversized (rejection rate is still slightly above 5%). On the other hand, for the test with interval instruments, the empirical size appears to match the nominal level for all but two configurations, where it seems to be slightly undersized.

### 6.3.2 Empirical power

Power is the probability of rejecting the null hypothesis under an alternative. We compute the empirical power by counting and averaging the number of times we reject the null for the test of nominal size 5% over the 1000 simulations when the distribution of random coefficients is misspecified. The simulation setup remains the same as previously with the only modification being that the true distribution of  $v_i$  is now either a mixture of normals or a Gamma. We report the power against the different alternatives in the subsequent tables. The main takeaway from our results is that the test with the interval instruments as testing instruments (I global and I local) largely outperforms the traditional Sargan-Hansen J-test against all the alternative distributions considered in our simulations.

**Power against Gaussian mixture alternatives.** We simulate data with the random coefficients distributed according to the Gaussian mixtures described below. We plot the true distributions in Figure 4. We report the results in Table 2. We observe that the test with the interval instruments has great power against all the mixtures tested. The rejection rates go to 1 very quickly in comparison to the Sargan-Hansen J tests.

$$v = Dv_1 + (1 - D)v_2, \quad \mathbb{P}(D = 1) = p, \quad \mathbb{P}(D = 0) = 1 - p,$$

$$v_1 \sim \mathcal{N}\left(-\sqrt{\frac{3p}{1-p}} + 2, 1\right) \quad v_2 \sim \mathcal{N}\left(\sqrt{\frac{3(1-p)}{p}} + 2, 1\right),$$

with  $p \in \{0.1; 0.2; 0.3; 0.4; 0.5\}$ .

Figure 4: Densities, Gaussian mixture alternatives




Table 2: Empirical power, Gaussian mixture alternatives (1000 replications)

Number of markets	T=50								T=100								T=200							
Estimation instruments	“Differentiation”				“Optimal’				“Differentiation”				“Optimal’				“Differentiation”				“Optimal’			
Test type	J	I	Global	I Local	J	I	Global	I Local	J	I	Global	I Local	J	I	Global	I Local	J	I	Global	I Local	J	I	Global	I Local
Mixture 1	0.533	0.991	0.987	0.987	0.719	0.989	0.989	0.989	0.604	1	1	1	0.967	1	1	1	0.829	1	1	1	1	1	1	1
Mixture 2	0.626	0.996	0.998	0.998	0.613	0.997	0.998	0.998	0.723	1	1	1	0.905	1	1	1	0.933	1	1	1	1	1	1	1
Mixture 3	0.629	0.992	0.995	0.995	0.43	0.996	0.997	0.997	0.741	1	1	1	0.7	1	1	1	0.941	1	1	1	0.977	1	1	1
Mixture 4	0.601	0.983	0.982	0.982	0.275	0.981	0.981	0.981	0.713	1	0.999	0.999	0.368	1	1	1	0.921	1	1	1	0.672	1	1	1
Mixture 5	0.56	0.907	0.904	0.904	0.157	0.9	0.906	0.906	0.635	0.993	0.995	0.995	0.124	0.995	0.996	0.996	0.855	1	1	1	0.146	1	1	1

**Power against Gamma alternatives.** We simulate data with the random coefficients distributed according to the Gamma distribution described below. We plot the true distributions in Figure 5. We report the results in table 3. We observe that the test with interval instruments has great power against all the Gamma distributions tested except for the first one, which we can see on the plot has a distribution that is relatively close to a normal distribution. Even for the first Gamma distribution, it still outperforms the traditional sets of instruments. For all the other Gamma distributions, the rejection rates go to 1 very quickly in comparison to the Sargan-Hansen J-tests. This confirms the superiority of the interval instruments in detecting misspecification in

the distribution of random coefficients. In Appendix D.2, we also study the power properties of our test against local alternatives.

$$v \sim \Gamma(2, k) \quad \text{with } k \in \{0.25, 0.5, 0.75, 1, 1.5\}$$

Figure 5: Densities, Gamma alternatives




Table 3: Empirical power, Gamma alternatives (1000 replications)

Number of markets	T=50						T=100						T=200					
Estimation instruments	“Differentiation”			“Optimal”			“Differentiation”			“Optimal”			“Differentiation”			“Optimal”		
Test type	J	I	Global	I	Local		J	I	Global	I	Local		J	I	Global	I	Local	
Gamma 1	0.293	0.106	0.093	0.142	0.082	0.074	0.154	0.083	0.073	0.094	0.092	0.08	0.118	0.155	0.139	0.066	0.156	0.138
Gamma 2	0.516	0.747	0.752	0.14	0.781	0.77	0.562	0.983	0.978	0.095	0.982	0.98	0.492	1	1	0.08	1	1
Gamma 3	0.607	0.96	0.962	0.157	0.963	0.969	0.693	0.998	1	0.156	1	1	0.922	1	1	0.161	1	1
Gamma 4	0.622	0.97	0.99	0.207	0.962	0.995	0.748	0.999	1	0.263	1	1	0.933	1	1	0.412	1	1
Gamma 5	0.687	0.991	0.999	0.371	0.988	0.999	0.812	1	1	0.585	1	1	0.976	1	1	0.865	1	1

## 6.4 Finite sample performance of interval instruments for estimation

In our last simulation exercise, we evaluate the performance of our interval instruments in estimating the parameters associated with the RC when the distribution of random coefficients is flexibly parametrized. To do so, we simulate data with a distribution of random coefficients following

a mixture of Gaussians and we estimate the parameters of this mixture. We provide a method to estimate the parameters when the distribution of the RC is a mixture in Section C.6 of the Appendix. In particular, we provide a new parametrization of the model, which yields substantial practical gains and may be of interest to researchers independent of the rest of the paper. The simulation design remains the same as previously. We assume that the random coefficient  $v_i$  is distributed according to the following mixture:  $v_i \sim D_i \mathcal{N}(-2, 0.5) + (1 - D_i) \mathcal{N}(4, 0.5)$  with  $\mathbb{P}(D_i = 1) = 0.25$ . Thus, there are 5 parameters associated with the distribution of RC: the means and variances of each component of the mixture and the mixing probability. Our objective is to compare the performance of the global and local interval instruments with the instruments commonly used by practitioners: the differentiation instruments from [Gandhi and Houde \(2019\)](#) and the “optimal instruments” from [Reynaert and Verboven \(2014\)](#). In Table 4, we report the empirical biases and the square root of the MSE for the estimators of the non-linear parameters for each set of instruments and for the different sample sizes. In Appendix D.3, we report the same information for the linear parameters (see Tables 14, 15, and 17) as well as the distribution of the empirical distribution of the non-linear estimates. Table 4 allows us to directly compare the performances of the three sets of instruments in estimating the non-linear parameters. We first observe that for all the sets of instruments, the empirical biases and  $\sqrt{MSE}$  of the estimators decrease when the sample size increases, which is reassuring. Furthermore, it appears clearly that the differentiation instruments perform worse than the “optimal instruments” and the interval instruments. The empirical  $\sqrt{MSE}$  of the estimators with the differentiation instruments is up to 12 times larger than with the interval instruments and up to 6 times larger than with the “optimal instruments”. We reach the same conclusions when we study empirical biases. The interval instruments appear to perform better than the “optimal instruments” even if the difference is less significant than with the differentiation instruments. For the sake of conciseness, we do not report the results obtained with a mixture of 3 components but the observations we make with two components are even more exacerbated. In Appendix D.3, as a means of comparison, we perform the same exercise when the distribution of random coefficients is a simple Gaussian and here, we do not observe any significant differences between the different sets of instruments, which confirms that the interval instruments make a difference when the distribution of RCs is flexible.

Table 4: Estimation non-linear parameters of the mixture (1000 replications)

Instruments		Differentiation					"Optimal"					Interval Global					Interval Local				
Parameter		$\beta_{3L}$	$\sigma_{3L}$	$\beta_{3H}$	$\sigma_{3H}$	$p_L$	$\beta_{3L}$	$\sigma_{3L}$	$\beta_{3H}$	$\sigma_{3H}$	$p_L$	$\beta_{3L}$	$\sigma_{3L}$	$\beta_{3H}$	$\sigma_{3H}$	$p_L$	$\beta_{3L}$	$\sigma_{3L}$	$\beta_{3H}$	$\sigma_{3H}$	$p_L$
Sample size	true	-2	0.5	4	0.5	0.25	-2	0.5	4	0.5	0.25	-2	0.5	4	0.5	0.25	-2	0.5	4	0.5	0.25
T=50, J=12	bias	0.214	0.184	-0.022	-0.045	0.027	0.076	0.059	0.026	-0.111	0.01	0.017	0	-0.045	0.004	0.005	-0.006	-0.005	-0.039	-0.001	0.003
	$\sqrt{MSE}$	0.633	0.734	0.281	0.35	0.075	0.361	0.483	0.212	0.281	0.036	0.277	0.391	0.227	0.259	0.024	0.251	0.34	0.214	0.244	0.019
T=50, J=20	bias	0.189	0.347	0.022	-0.081	0.025	0.074	0.11	0.028	-0.089	0.01	0.013	0.042	-0.018	-0.003	0.004	0.019	0.033	-0.023	0.01	0.003
	$\sqrt{MSE}$	0.566	0.887	0.184	0.291	0.059	0.328	0.563	0.163	0.228	0.033	0.248	0.415	0.166	0.22	0.021	0.228	0.38	0.15	0.184	0.018
T=100, J=12	bias	0.233	0.226	0.02	-0.066	0.027	0.054	0.037	0.019	-0.066	0.007	0.004	-0.012	-0.027	0.005	0.002	0	0	-0.028	0.007	0.001
	$\sqrt{MSE}$	0.592	0.703	0.256	0.305	0.072	0.279	0.4	0.154	0.211	0.028	0.167	0.282	0.157	0.201	0.013	0.127	0.225	0.143	0.164	0.005
T=100, J=20	bias	0.198	0.423	0.047	-0.101	0.025	0.074	0.107	0.033	-0.074	0.01	-0.009	-0.005	-0.008	-0.009	0.001	-0.003	0.004	-0.01	0.004	0.001
	$\sqrt{MSE}$	0.552	0.89	0.164	0.27	0.055	0.311	0.52	0.129	0.194	0.034	0.115	0.264	0.115	0.169	0.005	0.104	0.226	0.103	0.125	0.004
T=200, J=12	bias	0.184	0.167	0.011	-0.049	0.019	0.026	0.011	0.021	-0.061	0.004	-0.006	-0.027	-0.015	-0.001	0.001	0.002	-0.007	-0.016	0.006	0.001
	$\sqrt{MSE}$	0.466	0.601	0.176	0.262	0.053	0.184	0.313	0.113	0.172	0.018	0.088	0.219	0.108	0.164	0.003	0.091	0.174	0.099	0.123	0.003

## 7 Empirical application

The objective of the empirical exercise is twofold. First, we want to verify how well our instruments perform at estimating a flexible distribution of RCs using a real data set. Second, we want to study how the shape of the distribution of RCs can modify key counterfactual quantities such as the price elasticities or the pass-through, and check whether the results we obtain are consistent with the findings in [Miravete et al. \(2022\)](#). To do so, we estimate demand for cars using data on new car registrations in Germany from 2012 to 2018.<sup>23</sup> There are many reasons to focus on the car market. First, cars are highly differentiated products, which makes the BLP framework particularly adapted to this market. As a result, the BLP demand model has been widely applied to study the car industry (e.g., [Berry et al. \(1995\)](#), [Grigolon et al. \(2018\)](#), [Petrin \(2002\)](#)) and one can easily compare our results with previous results obtained in the literature under different specifications. Second, there are many policy-relevant questions related to this market. In particular, the role of road transport in air pollution is significant and many countries have implemented tax policies to reduce the CO2 emissions generated by car transportation.<sup>24</sup> An important strand of the litera-

<sup>23</sup>The dataset was kindly provided to us by Kevin Remmy <https://kevinremmy.com/research/>.

<sup>24</sup>In 2017, road transport was responsible of approximately 19% of total greenhouse has emissions in EU-28 Retrieved from <https://www.eea.europa.eu/data-and-maps/indicators/transport-emissions-of-greenhouse-gases/transport-emissions-of-greenhouse-gases-12> on Octo-

ture has investigated the performance of these different taxation schemes (Alberini and Horvath (2021), Allcott and Wozny (2014), D’Haultfœuille, Givord, and Boutin (2014), Durrmeyer (2022), Durrmeyer and Samano (2018), Gillingham and Houde (2021), Grigolon et al. (2018), Huse and Koptuyug (2022), Kunert (2018)). Other policy-relevant questions include the impact of import tariffs (Miravete et al. (2018)) and the determinants of market power (Berry et al. (1995), Grieco, Murry, and Yurukoglu (2022)). To answer these questions, the researcher must often estimate the demand for cars. The credibility of the implied analysis depends critically on how well the model can reproduce the underlying substitution patterns and the shape of the demand curve. To this end, it is essential to have a demand system that is sufficiently flexible, and particularly so with respect to the random coefficient on price. In this section, we use our instruments to estimate a Gaussian mixture as the random coefficient associated with price. Moreover, we use our test to assess how moving from the usual Gaussian RC to the Gaussian mixture decreases the degree of misspecification. Finally, we compare the counterfactual quantities under a Gaussian mixture and the traditional specifications (Gaussian RC and logit). In line with the findings in Miravete et al. (2022), our results indicate that the Gaussian mixture yields higher pass-through rates and curvatures.

## 7.1 Data

The data set includes state-level new car registrations, publicly available by the German Federal Motor Transport Authority (KBA) from 2012 to 2018. This gives us 112 markets defined by state-year pairs. Data on car characteristics and price are scraped from General German Automobile Club and include horsepower, engine type, size, weight, fuel cost, CO2 emission, number of doors, segment, and body type. The data set is at a granular level where every car is uniquely identified by its manufacturer and its type key code (HSN/TSN) that is defined according to the characteristics of the car. Following the literature, we aggregate products with the same brand, model, engine type, and body combination (e.g. BMW-1 Series-Diesel-Hatchback).<sup>25</sup> Likewise, we follow the literature and define the market size as the number of households in the market. To construct

---

ber 21, 2022.

<sup>25</sup>In aggregating the products from the HSN/TSN level, we use the characteristics of the car with the highest sales.

market shares, we simply divide new car registrations of a given product by the market size. The data set is complemented by information on demographics such as the number of households or the average income per household at the state-year level and yearly average gas price data from ADAC.<sup>26</sup>

**Summary statistics.** The shares of products sold by engine type are presented in Table 5. We focus our analysis on combustion engine vehicles as in our sample period electric-vehicle cars constitute a small market share (always less than 5% of the sold vehicles) and can be seen as a distinct market. Between diesel and gasoline cars, we observe that the market share for diesel cars decreases over time, starting from 2016. The timing is in line with the emissions scandal, known as the Dieselgate, which started in September 2015.

Table 5: Shares (%) of new registrations by engine type

Fuel Type	Year						
	2012	2013	2014	2015	2016	2017	2018
Diesel	46.8	46.1	46.3	46.4	43.9	36.2	30.0
Gasoline	52.6	52.9	52.6	52.3	54.4	60.8	66.5
Battery EV	0.1	0.2	0.3	0.4	0.3	0.7	1.1
Hybrid EV	0.5	0.8	0.7	0.6	1	1.4	1.6
Plug-in hybrid EV	0	0	0.1	0.3	0.4	0.9	0.9

Table 6 provides sales-weighted averages for prices and observed characteristics. We observe that the difference in fuel consumption and resulting fuel costs steadily ranks diesel above gasoline. However, the average price of diesel cars sold is higher than gasoline cars. This implies a potential trade-off in terms of the costs of car ownership at the time of purchase. With a fixed mileage in mind, a consumer with high sensitivity to fuel costs might be willing to pay a higher price for a more fuel-efficient car. We also observe that the horsepower and the size of the newly registered cars increase over time.

<sup>26</sup>State level income [https://ec.europa.eu/eurostat/web/products-datasets/-/nama\\_10r\\_2hhinc](https://ec.europa.eu/eurostat/web/products-datasets/-/nama_10r_2hhinc)

Table 6: Summary Statistics (Sales weighted)

	Year						
	2012	2013	2014	2015	2016	2017	2018
<b><u>Diesel</u></b>							
Price/income	0.74	0.72	0.73	0.72	0.71	0.69	0.68
Size (m2)	8.31	8.31	8.32	8.36	8.42	8.48	8.53
Horsepower (kW/100)	1.09	1.07	1.11	1.11	1.14	1.16	1.21
Fuel cost (euros/100km)	7.90	7.18	6.63	5.53	4.94	5.25	5.83
Fuel cons. (Lt./100km)	5.19	4.98	4.89	4.73	4.61	4.61	4.71
CO2 emission (g/km)	136.19	130.50	127.69	123.58	120.42	120.49	123.27
Nb. of products/market	133	138	146	150	151	149	143
<b><u>Gasoline</u></b>							
Price/income	0.46	0.46	0.46	0.46	0.46	0.45	0.43
Size (m2)	7.23	7.27	7.28	7.30	7.36	7.46	7.53
Horsepower (kW/100)	0.79	0.78	0.80	0.82	0.85	0.88	0.91
Fuel cost (euros/100km)	9.48	8.61	8.11	7.27	6.69	7.06	7.40
Fuel cons. (Lt./100km)	5.76	5.47	5.40	5.31	5.25	5.34	5.38
CO2 emission (g/km)	135.80	128.18	125.27	122.89	121.22	122.86	123.26
Nb. of products/market	157	171	179	185	186	193	188

Note: Provided statistics are sales weighted averages across products. Total number of markets (State\*Year) is 112 .

**Inter-market variation.** Our dataset contains both geographical variation and time variation, as we observe the sales in every state in Germany over the period 2012-2018. States in Germany differ significantly in terms of income per inhabitant, population density and average distance driven.<sup>27</sup> It is fundamental to take this inter-market variation into account in our empirical specification for two reasons. First, our model postulates that consumers’ preferences are the same across markets. However, we observe that the market shares vary from one state to the other even if the choice set remains the same. This feature of the data can only be explained if

<sup>27</sup>For the population density 2019 (inh/km<sup>2</sup>): 69 (Mecklenburg-Vorpommern) to 4118 (Berlin) (from Federal Statistical Office of Germany (Destatis)), GDP per capita 2019: 28.9k (Mecklenburg-Vorpommern) to 67k (Hamburg) (retrieved from <https://www.ceicdata.com/en/germany/esa-2010-gdp-per-capita-by-region/gdp-per-capita-bremen> on 05 November 2022). For average driving distance in 2019: 13079 km (Mecklenburg-Vorpommern) to 9531 (Berlin) retrieved from <https://de.statista.com/statistik/daten/studie/644381/umfrage/fahrleistung-privater-pkw-in-deutschland-nach-bundesland/> on 19 September 2022.



we let the preferences vary from one market to the other. Second, in Section 2.3, we saw that there needs to be sufficient variation in the product characteristics across markets to identify the distribution of RCs. By interacting product characteristics with state demographics, we achieve both objectives: we shift the preferences to a more common representation and we create variation in the product characteristics across markets. To choose which interaction terms to include in the utility function, we first create market specific sales-weighted characteristics for the following variables: price, fuel cost, size, horsepower, height, gasoline dummy, and foreign dummy (equal to one if the manufacturer of the car is not German). Then, we regress these quantities on the demographics of interest: average income, population density, and a time trend. Last, we select the interaction terms that explain the best the variation in sales-weighted characteristics (namely, the variables with a p-value below  $1e^{-10}$ ). The results of these regressions are presented in Table 7. They suggest that income shifts positively the preferences for price, size, and horsepower (i.e. higher income is associated with larger cars, and higher horsepower). In contrast, income shifts negatively the preferences for foreign status, height, and gasoline status.<sup>28</sup> Although weaker, a similar pattern is observed for the effect of population density on car characteristics.

Table 7: Linear regressions of sales-weighted car characteristics on demographic characteristics

	Income(/1000)	Population density (/100)	Time trend
Price( $\times 1000$ )	0.138** (0.013)	0.069* (0.011)	0.286* (0.059)
Fuel cost (euros/100km)	-0.0069 (0.0063)	-0.0036 (0.0056)	0.3587** (0.0296)
Size( $m^2$ )	0.0058** (0.00079)	0.0018* (0.00070)	0.0176* (0.00371)
Horsepower (KW/100)	0.0028** (0.00028)	0.0012* (0.00025)	0.0129** (0.00132)
Foreign	-0.0050** (0.00052)	-0.0014* (0.00046)	0.0295** (0.00246)
Height(m)	-0.00051** (0.000061)	-0.00043** (0.000054)	0.00181* (0.000286)
Gasoline	-0.0067** (0.00059)	-0.0024* (0.00053)	0.0131* (0.00280)

Note: \* p-value lower than 0.01, \*\* p-value lower than  $1e^{-10}$ .

---

<sup>28</sup>In the main analysis, we use price/income to capture the income effect.

**Instruments for the endogeneity of price.** To instrument for price, we use a combination of variables on the intensity of competition and cost shifters. To measure the intensity of competition, we consider the number of competing products of the same class and engine type in a given market, and the number of competing products of the same engine type in a given market. As for cost shifters, we use three complementary datasets: the mean hourly labor cost, the price of steel (interacted with the weight of the car), and exchange rates between Germany and the country of assembly.

1. Labor cost: we use the mean nominal hourly labor cost per employee in the manufacturing sector of the country of assembly of the models. The data on labor costs come from International Labor Organization Statistics (ILOSTAT).<sup>29</sup>
2. Price of steel: we collect the price of steel futures in January of each year.
3. Exchange rates: we construct the exchange rates between Germany and the country of assembly of each car model using exchange rate data from OECD.<sup>30</sup>

## 7.2 Empirical specification

The indirect utility of consumer  $i$ , purchasing product  $j$  in market  $t$  (defined as a state-year pair) is given by:

$$u_{ijt} = \underbrace{x'_{1jt}\beta + \xi_{jt}^*}_{\delta_{jt}} + x'_{2jt}\alpha_i + \varepsilon_{ijt}.$$

The mean utility  $\delta_{jt} = x'_{1jt}\beta + \xi_{jt}^*$  captures homogeneous preferences. The variables in  $x_{1jt}$  consist of the product characteristics for which we assume that there is no preference heterogeneity and the interaction terms that explain the best the geographical variation observed in Table 7.<sup>31</sup>

The demand shock on product  $j$  is decomposed as follows:

$$\xi_{jt}^* = Brand_j + State_t + Year_t + \xi_{jt},$$

---

<sup>29</sup>Retrieved from [https://www.ilo.org/ilostat-files/Documents/Excel/INDICATOR/LAC\\_4HRL\\_ECO\\_CUR\\_NB\\_A\\_EN.xlsx](https://www.ilo.org/ilostat-files/Documents/Excel/INDICATOR/LAC_4HRL_ECO_CUR_NB_A_EN.xlsx)

<sup>30</sup>Retrieved from <https://data.oecd.org/conversion/exchange-rates.htm>

<sup>31</sup>The choice of the variables that display preference heterogeneity is based on our understanding of the car market and follows current empirical practices for this specific market. However, we understand the limitations of this approach, and we are working on an iterative procedure to select the variables that display consumer heterogeneity.

where  $Brand_j$  is a brand fixed effect that captures the unobserved quality of the brand of product  $j$ ,  $State_t$  captures state specific demand shocks that are fixed across time and products and  $Year_t$  captures year-specific demand shocks. Therefore,  $State_t$  and  $Year_t$  play a role in explaining the variation in the overall demand for cars (or equivalently, in the share of the outside option).

The variables in  $x_{2jt}$  are the product characteristics that display preference heterogeneity and which we augment with a RC. In our specification, we include the price, the size, and the gasoline dummy in  $x_{2jt}$ . We estimate the model assuming different specifications for the distribution of RCs. First, we estimate the model without any consumer heterogeneity. Second, we assume that all the RCs are normally distributed. Finally, we consider a Gaussian mixture on price to increase flexibility with respect to the preferences on price. For each different specification, we perform the specification test developed in Section 5 to see how the degree of misspecification evolves as we increase flexibility on the distribution of RCs.

### 7.3 Estimation

**Estimation conditional logit (no heterogeneity).** First, we estimate the logit model, and we report the results in Table 8.<sup>32</sup> As expected, we find a negative effect of price and fuel cost on the utility. The interaction terms indicate that the utility derived from size, horsepower, foreign status and gasoline all decrease with income. Moreover, we observe that the aversion to fuel cost decreases over time, which is likely an artifact implied by increasing fuel cost over the years. In contrast, the utility derived from horsepower appears to increase with time. However, these time effects are smaller in comparison with the heterogeneity due to income. To facilitate the interpretation of these results, we consider a household with a €47,000 income in 2018. This corresponds to the mean income in 2018. For this household, the implied effect of size on the utility is negative, whereas a positive utility is derived from higher horsepower, the car’s brand being German, height, and gasoline engines.

---

<sup>32</sup>In Appendix E, we provide results for baseline specifications including the simple conditional logit and the nested logit (with and without state and year fixed effects).

Table 8: Logit estimation

	Baseline		$\times$ Income (/1000)		$\times$ Pop. density(/100)		$\times$ Time trend	
Homogeneous Preferences	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E
Price/income	-2.4	1.3e-01	-	-	-	-	-	-
Fuel Cost	-0.25	8.6e-03	-	-	-	-	0.014	1.7e-03
Size( $m^2$ )	0.15	4.2e-02	-0.0055	8.5e-04	-	-	-	-
Horsepower(KW/100)	2.7	1.8e-01	-0.019	2.4e-03	-	-	-0.081	7e-03
Foreign	0.18	7.1e-02	-0.017	1.4e-03	-	-	-	-
Height(m)	3.5	2.3e-01	-0.0015	4.6e-03	-0.036	4.7e-03	-	-
Gasoline	1.1	6.3e-02	-0.011	1.2e-03	-	-	-	-

Note: Brand, Year and State FE's are included.

**Estimation with Gaussian random coefficients.** We now increase the flexibility in the *traditional* manner, by assuming that the RCs on the price, the size and the gasoline indicator follow a Gaussian distribution. We report the estimates obtained under this new specification in Table 9. The signs for the homogeneous preference parameters in  $x_{1jt}$  remain the same and the magnitude of the effects do not change significantly. The sign associated with the mean effect of price remains negative. In contrast, the sign on the mean effects of the size and the gasoline dummy are inverted with respect to the logit specification. This last observation illustrates an important empirical finding: average effects are not invariant to the introduction of preference heterogeneity. In other words, the logit estimates do not necessarily match the means, when we introduce a Gaussian RC. Moreover, the three RCs display high variances and particularly so for the gasoline dummy, which indicate a high level of heterogeneity with respect to these three characteristics.<sup>33</sup>

<sup>33</sup>The estimation is performed using the parametrization proposed in Ketz (2019), which avoids boundary issues at 0 for the variances of the RCs.

Table 9: Traditional BLP (Gaussian RC)

	Baseline		× Income (/1000)		× Pop. density(/100)		× Time trend	
Homogeneous Preferences	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E
Price/income	-	-	-	-	-	-	-	-
Fuel Cost	-0.29	5.1e-03	-	-	-	-	0.031	9.2e-04
Size( $m^2$ )	-	-	-0.0053	3.1e-04	-	-	-	-
Horsepower(KW/100)	0.77	1.5e-02	0.0078	6.8e-04	-	-	-0.12	5.6e-03
Foreign	0.21	5.4e-02	-0.019	1.1e-03	-	-	-	-
Height(m)	3.4	1.1e-02	-0.0088	1.2e-03	-0.032	3.6e-04	-	-
Gasoline	-	-	-0.0028	8.6e-04	-	-	-	-
Gaussian RC	$\hat{\beta}$	S.E	$\hat{\sigma}$	S.E				
Price/income	-2.4	2e-02	0.96	5.9e-03	-	-	-	-
Size( $m^2$ )	-0.37	1.5e-02	0.43	3.6e-03	-	-	-	-
Gasoline	-2.3	4.4e-02	4	4.1e-04	-	-	-	-

Note: Brand, Year and State FE's are included.

**Estimation with a Gaussian mixture on the price.** Finally, we increase the flexibility of the model, by replacing the Gaussian RC on the price variable with a Gaussian mixture of 2 components. We focus on the price as the literature shows that the distribution of price sensitivity is absolutely key for many quantities of interest in IO, including the price elasticities and the pass-through. We report the estimates obtained under this new specification in Table 10. The results point out the presence of two distinct modes in the distribution of the RC associated with price. The two modes reveal the presence of two groups of consumers: the first one with high price sensitivity (with the mean component at -9.6) and the second one with low price sensitivity (with the mean component at -2.5). Moreover, the distribution is heavily asymmetric with the probability of the first mode being 0.9, which entails that the majority of consumers are highly sensitive to price. This last feature is completely absent in the logit and Gaussian specifications, which seem to capture only the first mode of the distribution as we can see in Figure 6. Once again the homogeneous parameters are relatively unchanged with respect to the previous specifications. The Gaussian RC on the gasoline still displays a high variance (the standard deviation of the RC

equals 2.8).


Table 10: Estimation Gaussian mixture on Price

	Baseline		× Income (/1000)		× Pop. density(/100)		× Time trend	
Homogeneous Preferences	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E	$\hat{\beta}$	S.E
Price/income	-	-	-	-	-	-	-	-
Fuel Cost	-0.23	5.8e-03	-	-	-	-	0.026	1e-03
Size( $m^2$ )	-	-	-0.0055	3.7e-04	-	-	-	-
Horsepower(KW/100)	1.8	3.6e-02	-0.0016	1.1e-03	-	-	-0.1	7e-03
Foreign	0.26	6.1e-02	-0.021	1.2e-03	-	-	-	-
Height(m)	3.5	1.1e-02	-0.012	1.2e-03	-0.032	3.7e-04	-	-
Gasoline	-	-	-0.026	1.3e-03	-	-	-	-
Gaussian RC	$\hat{\beta}$	S.E	$\hat{\sigma}$	S.E				
Size( $m^2$ )	0.5	1.9e-02	0.1	6.7e-02	-	-	-	-
Gasoline	-0.45	3.8e-03	2.8	9.1e-03	-	-	-	-
Gaussian Mixture	$\hat{\beta}_1$	S.E	$\hat{\sigma}_1$	S.E	$\hat{\beta}_2$	S.E	$\hat{\sigma}_2$	S.E
Price/income	-9.6	1.8e-02	0.1	1.8e-03	-2.5	1.8e-02	0.35	5.2e-04
Probability	0.9	6.8e-05						

Note: Brand, Year and State FE's are included.

In Figure 6, we plot the estimated distribution of random coefficients under the three specifications we consider. We observe little to no variation in the homogeneous parameters from one specification to the other. The main difference comes from the introduction of the Gaussian mixture on price, which reveals the presence of a large group of highly price sensitive consumers.

Figure 6: Estimated distributions of RCs in the three specifications



**Specification Test.** By increasing the flexibility on the distribution of RCs, we recover less precise estimates and the model becomes more difficult to estimate. Thus, it is important to show that the additional flexibility substantially reduces the misspecification of the model. To quantify the degree of misspecification across the different models, we keep the same set of estimation instruments across the different specifications of RCs and we report the value of the associated Sargan-Hansen J statistics in each case. Moreover, for every model, we follow the procedure developed in Section 5 to test if the distribution of RCs on price is well specified. We use the global interval instruments and we denote this test “Interval test”. We report the values of the test statistics and the degrees of freedom of the chi-square under the null in Table 11. We observe an important decrease in the Sargan-Hansen J statistic when we transition from the logit to the Gaussian RC. However, the decrease in the Sargan-Hansen J statistic is much larger when we transition from the Gaussian RC on price to the Gaussian mixture, which indicates that the Gaussian mixture performs much better than the simple Gaussian at capturing the underlying heterogeneity in price sensitivity. The interval test displays a similar behavior, with the largest decrease in the test statistic stemming from the transition from the Gaussian RC to the Gaussian

mixture.

Table 11: Evolution of misspecification with flexibility

Instruments	Logit			Gaussian RC			Gaussian mixture		
Test	Stat.	Critical val.	DF	Stat.	Critical val.	DF	Stat.	Critical val.	DF
J test	2755.7	40.1	27	2341.7	36.4	24	950.3	33.9	21
Interval test	1331.9	14.1	7	999.4	14.1	7	244.0	14.1	7

## 7.4 Counterfactual quantities

The objective of this subsection is to illustrate how changes in the distribution of the RC associated with price affect many counterfactual quantities of interest in empirical IO, such as the price elasticities, the marginal costs faced by car manufacturers, and the pass-through of cost. In order to compare our empirical results with the findings in [Miravete et al. \(2022\)](#), we also calculate the demand curvature under the different specifications. They show that a large demand curvature is necessary to recover a pass-through larger than one.

In the following, we study the effect of different specifications on the price elasticities, demand curvature, marginal costs and mark-ups, and finally on the pass-through. To recover marginal costs and mark-ups, we assume that multi-product firms pricing under Bertrand-Nash pricing. For the pass-through, we calculate the new equilibrium prices using fixed point iterations and, following the literature, study the effect of increasing the marginal costs of each product by 1% and recomputing the marginal cost. In [Appendix E](#), we provide details on the calculation of counterfactual quantities. In our computations, we use the year 2018, which is the last year of our sample.

**Summary of results.** We report the median values for the five counterfactual quantities of interest in [Table 12](#). Several remarks are in order. First, the Gaussian mixture yields a much lower price elasticity than the two other specifications. This is related to the emergence of a group of highly price sensitive consumers in the mixture specification, which we fail to detect with the logit and Gaussian RC specifications. Moreover, the low price elasticities that we recover in the Gaussian and logit specifications, generate unreasonably low marginal costs (even negative ones



as we can see in Figure 7) and excessive mark-ups. In contrast, this problem does not appear with the Gaussian mixture. Finally, to link our results with the findings in Miravete et al. (2022), we now focus on the demand curvature and the pass-through of cost. As expected, the logit displays a curvature and a pass-through equal to 1. In contrast, we can see that the Gaussian mixture displays a larger demand curvature than the other two specifications. This comes from the skewness that the mixture induces in the distribution of price sensitivity. This last feature implies that the Gaussian mixture yields a pass-through much greater than 1 (1.5 on average). Unfortunately, the negative marginal costs we recover with the Gaussian RC prevent us from computing the pass-through in this case.<sup>34</sup>

Table 12: Median counterfactual quantities under different specifications on RCs


RC distribution on price	Logit	Gaussian	Gaussian Mixture
Own price-elasticity	-1.2	-1.1	-2.6
Demand curvature	1.0	1.2	1.3
Marginal cost	9,366	1,929	20,105
Mark-up	24,048	29,572	11,066
Pass-through	1.0	-	1.5

In Figure 7, we plot the empirical distributions of the counterfactual quantities. We can see in the plot featuring the distribution of marginal costs that the logit and Gaussian specifications generate negative marginal costs for some of the cars. This is an indication that the price elasticities implied by these specifications are too low in absolute value.

---


<sup>34</sup>Our algorithm to compute the new equilibrium prices after the change in cost does not converge.

Figure 7: Empirical distribution of counterfactual quantities under different specifications



Finally, in Figure 8, we plot the elasticity functions implied by the different specifications for the 15 most popular cars in our sample. We observe important differences in the elasticities. The Gaussian mixture generates lower price elasticities than the other two specifications. We do the same exercise with the demand curves in Appendix E.

Figure 8: Estimated elasticities under different specifications



## 8 Conclusion

In this paper, we develop novel econometric tools to parsimoniously increase the flexibility of the distribution of random coefficients in the BLP demand model initiated by [Berry et al. \(1995\)](#). Specifically, we construct novel instruments designed to detect deviations from the true distribution of random coefficients. Building on these instruments, we provide a formal moment-based specification test on the distribution of random coefficients, which allows researchers to test the chosen specification without having to re-estimate the model under a more flexible parametrization. Our instruments are designed to maximize the power of the test when the distribution of

RC is misspecified. By exploiting the duality between estimation and testing, we show that these instruments can also improve the estimation of the BLP model under a flexible parametrization. Our Monte Carlo simulations confirm that the interval instruments we develop in this paper outperform the traditional instruments both for testing and estimating purposes. Finally, we apply these new tools to flexibly estimate the demand for cars in Germany. We show that these tools can be applied to the equally popular mixed logit demand model with individual-level data.

In future work, we plan to see if we can generalize these instruments to other non-linear moment-based models, as well as to the general problem of testing distributional assumptions in structural models. From a broader perspective, our paper is part of an existent discussion on the most effective way to model unobserved preference heterogeneity in structural models. Most empirical frameworks feature a clear trade-off between the degree of flexibility one chooses and the precision of the estimates one obtains. It is thus critical to understand how misspecification on the unobserved heterogeneity affects the counterfactual quantities of interest. In the case of the BLP demand model, our paper and others show that misspecification in the distribution of random coefficients substantially distorts the substitution patterns as well as the shape of the demand curve and, thus, is likely to significantly alter the counterfactual quantities.

# Bibliography

- ALBERINI, A. AND M. HORVATH (2021): “All car taxes are not created equal: evidence from Germany,” *Energy Economics*, 100.
- ALLCOTT, H. AND N. WOZNY (2014): “Gasoline prices, fuel economy, and the energy paradox,” *Review of Economics and Statistics*, 96, 779–795.
- ALLEN, R. AND J. REHBECK (2020): “Identification of random coefficient latent utility models,” *arXiv preprint: 2003.00276*.
- AMEMIYA, T. (1974): “The nonlinear two-stage least-squares estimator,” *Journal of Econometrics*, 2, 105–110.
- ANDREWS, D. W. AND P. GUGGENBERGER (2019): “Identification and singularity-robust inference for moment condition models,” *Quantitative Economics*, 10, 1703–1746.
- ARMSTRONG, T. B. (2016): “Large market asymptotics for differentiated product demand estimators with economic models of supply,” *Econometrica*, 84, 1961–1980.
- BAHADUR, R. R. (1960): “Stochastic comparison of tests,” *The Annals of Mathematical Statistics*, 31, 276–295.
- BARAHONA, N., C. OTERO, S. OTERO, AND J. KIM (2020): “Equilibrium effects of food labeling policies,” *SSRN working paper*.
- BERRY, S. T. (1994): “Estimating discrete-choice models of product differentiation,” *The RAND Journal of Economics*, 25, 242–262.
- BERRY, S. T. AND P. A. HAILE (2009): “Nonparametric identification of multinomial choice demand models with heterogeneous consumers,” *NBER working paper 15276*.
- (2014): “Identification in differentiated products markets using market level data,” *Econometrica*, 82, 1749–1797.

- BERRY, S. T., J. LEVINSOHN, AND A. PAKES (1995): “Automobile prices in market equilibrium,” *Econometrica*, 63, 841–890.
- BERRY, S. T., O. B. LINTON, AND A. PAKES (2004): “Limit theorems for estimating the parameters of differentiated product demand systems,” *The Review of Economic Studies*, 71, 613–654.
- CHAMBERLAIN, G. (1987): “Asymptotic efficiency in estimation with conditional moment restrictions,” *Journal of Econometrics*, 34, 305–334.
- CHETVERIKOV, D. AND D. WILHELM (2017): “Nonparametric instrumental variable estimation under monotonicity,” *Econometrica*, 85, 1303–1320.
- COMPIANI, G. (2018): “Nonparametric demand estimation in differentiated products markets,” *SSRN working paper*.
- CONLON, C. AND J. GORTMAKER (2020): “Best practices for differentiated products demand estimation with pyblp,” *The RAND Journal of Economics*, 51, 1108–1161.
- CRAWFORD, G. S., O. SHCHERBAKOV, AND M. SHUM (2019): “Quality overprovision in cable television markets,” *American Economic Review*, 109, 956–995.
- D’HAULTFŒUILLE, X., P. GIVORD, AND X. BOUTIN (2014): “The environmental effect of green taxation: the case of the French *bonus/malus*,” *The Economic Journal*, 124, F444–F480.
- DUBÉ, J.-P., J. T. FOX, AND C.-L. SU (2012): “Improving the numerical performance of static and dynamic aggregate discrete choice random coefficients demand estimation,” *Econometrica*, 80, 2231–2267.
- DUBOIS, P., R. GRIFFITH, AND M. O’CONNELL (2018): “The effects of banning advertising in junk food markets,” *The Review of Economic Studies*, 85, 396–436.
- DUFOUR, J.-M. AND M. L. KING (1991): “Optimal invariant tests for the autocorrelation coefficient in linear regressions with stationary or nonstationary AR (1) errors,” *Journal of Econometrics*, 47, 115–143.

- DUNKER, F., S. HODERLEIN, AND H. KAIDO (2022): “Nonparametric identification of random coefficients in endogenous and heterogeneous aggregate demand models,” *arXiv preprint: 2201.06140*.
- DURRMEYER, I. (2022): “Winners and losers: the distributional effects of the French feebate on the automobile market,” *The Economic Journal*, 132, 1414–1448.
- DURRMEYER, I. AND M. SAMANO (2018): “To rebate or not to rebate: fuel economy standards vs. feebates,” *The Economic Journal*, 128, 3076–3116.
- FOX, J. T. AND A. GANDHI (2011): “Identifying demand with multidimensional unobservables: a random functions approach,” *NBER working paper 17557*.
- FOX, J. T., K. IL KIM, S. P. RYAN, AND P. BAJARI (2012): “The random coefficients logit model is identified,” *Journal of Econometrics*, 166, 204–212.
- FREYBERGER, J. (2015): “Asymptotic theory for differentiated products demand models with many markets,” *Journal of Econometrics*, 185, 162–181.
- GANDHI, A. AND J.-F. HOUDE (2019): “Measuring substitution patterns in differentiated-products industries,” *NBER working paper 26375*.
- GEWEKE, J. (1981): “The approximate slopes of econometric tests,” *Econometrica*, 49, 1427–1442.
- GILLINGHAM, K. AND S. HOUDE (2021): “Consumer myopia in vehicle purchases: evidence from a natural experiment,” *American Economic Journal: Economic Policy*, 207–238.
- GOURIEROUX, C. AND A. MONFORT (1995): *Statistics and econometric models*, vol. 2, Cambridge University Press.
- GRENNAN, M. (2013): “Price discrimination and bargaining: empirical evidence from medical devices,” *American Economic Review*, 103, 145–177.
- GRIECO, P. L., C. MURRY, AND A. YURUKOGLU (2022): “The evolution of market power in the US automobile industry,” *NBER working paper 29013*.

- GRIGOLON, L., M. REYNAERT, AND F. VERBOVEN (2018): “Consumer valuation of fuel costs and tax policy: evidence from the European car market,” *American Economic Journal: Economic Policy*, 10, 193–225.
- HANSEN, L. P. (1982): “Large sample properties of generalized method of moments estimators,” *Econometrica*, 50, 1029–1054.
- HO, K. AND A. PAKES (2014): “Hospital choices, hospital prices, and financial incentives to physicians,” *American Economic Review*, 104, 3841–3884.
- HUSE, C. AND N. KOPTYUG (2022): “Salience and policy instruments: evidence from the auto market,” *Journal of the Association of Environmental and Resource Economists*, 9, 345–382.
- ICHIMURA, H. AND T. S. THOMPSON (1998): “Maximum likelihood estimation of a binary choice model with random coefficients of unknown distribution,” *Journal of Econometrics*, 86, 269–295.
- KETZ, P. (2019): “On asymptotic size distortions in the random coefficients logit model,” *Journal of Econometrics*, 212, 413–432.
- KUNERT, U. (2018): “Diesel fuel and passenger cars receive preferential tax treatment in Europe: reform of taxation needed in Germany,” *DIW Weekly Report*, 8, 289–298.
- LEE, J. AND K. SEO (2015): “A computationally fast estimator for random coefficients logit demand models using aggregate data,” *The RAND Journal of Economics*, 46, 86–102.
- LEWBEL, A. (2000): “Semiparametric qualitative response model estimation with unknown heteroscedasticity or instrumental variables,” *Journal of Econometrics*, 97, 145–177.
- LU, Z., X. SHI, AND J. TAO (2021): “Semi-nonparametric estimation of random coefficient logit model for aggregate demand,” *SSRN working paper*.
- McFADDEN, D. AND K. TRAIN (2000): “Mixed MNL models for discrete response,” *Journal of Applied Econometrics*, 15, 447–470.
- MILLER, N. H., G. SHEU, AND M. C. WEINBERG (2021): “Oligopolistic price leadership and mergers: the United States beer industry,” *American Economic Review*, 111, 3123–3159.



- MILLER, N. H. AND M. C. WEINBERG (2017): “Understanding the price effects of the Miller-Coors joint venture,” *Econometrica*, 85, 1763–1791.
- MIRAVETE, E. J., M. J. MORAL, AND J. THURK (2018): “Fuel taxation, emissions policy, and competitive advantage in the diffusion of European diesel automobiles,” *The RAND Journal of Economics*, 49, 504–540.
- MIRAVETE, E. J., K. SEIM, AND J. THURK (2022): “Robust pass-through estimation in discrete choice models,” *Working paper*.
- NEVO, A. (2000): “Mergers with differentiated products: the case of the ready-to-eat cereal industry,” *The RAND Journal of Economics*, 31, 395–421.
- NEWKEY, W. K. (1990): “Efficient instrumental variables estimation of nonlinear models,” *Econometrica*, 58, 809–837.
- (2004): “Efficient semiparametric estimation via moment restrictions,” *Econometrica*, 72, 1877–1897.
- PETRIN, A. (2002): “Quantifying the benefits of new products: the case of the minivan,” *Journal of Political Economy*, 110, 705–729.
- REYNAERT, M. (2021): “Abatement strategies and the cost of environmental regulation: emission standards on the European car market,” *The Review of Economic Studies*, 88, 454–488.
- REYNAERT, M. AND F. VERBOVEN (2014): “Improving the performance of random coefficients demand models: the role of optimal instruments,” *Journal of Econometrics*, 179, 83–98.
- ROODMAN, D. (2009): “A Note on the theme of too many instruments,” *Oxford Bulletin of Economics and Statistics*, 71, 135–158.
- SALANIÉ, B. AND F. A. WOLAK (2019): “Fast, “robust”, and approximately correct: estimating mixed demand systems,” *NBER working paper 25726*.
- SILVEY, S. D. (1959): “The Lagrangian multiplier test,” *The Annals of Mathematical Statistics*, 30, 389–407.

- TEBALDI, P., A. TORGOVITSKY, AND H. YANG (2019): “Nonparametric estimates of demand in the California health insurance exchange,” *NBER working paper 25827*.
- TRAIN, K. E. (2008): “EM algorithms for nonparametric estimation of mixing distributions,” *Journal of Choice Modelling*, 1, 40–69.
- WANG, A. (2022): “Sieve BLP: A semi-nonparametric model of demand for differentiated products,” *Journal of Econometrics*.
- WHITE, H. (1982): “Maximum likelihood estimation of misspecified models,” *Econometrica*, 50, 1–25.

# Appendices

## A Extension to the mixed logit demand model

The main difference between the BLP demand model and the mixed logit model is that the latter one assumes that the econometrician observes individual data. Let us consider the baseline mixed logit model with no endogeneity and consumer level data.<sup>35</sup> We observe the choices of  $N$  consumers. The indirect utility function of consumer  $i$  making choice  $j \in \{0, 1, \dots, J\}$  is given by:

$$u_{ij} = x'_{1ij}\beta_0 + x'_{2ij}v_i + \varepsilon_{ij}, \quad (12)$$

where

- $\varepsilon_{ij}$  is a preference shock that follows a type I extreme value distribution independent of all other variables and across  $i, j$ ;
- $x_{1ij}$  is a vector of product characteristics interacted with consumer characteristics of dimension  $K_1$  which display no preference heterogeneity;
- $x_{2ij}$  is a vector of product characteristics interacted with consumer characteristics of dimension  $K_2$  which display preference heterogeneity;
- $v_i$  is a vector of random coefficients of dimension  $K_2$  which jointly follows a distribution characterized by a density  $f$ ;

Each consumer chooses the product that maximizes his or her utility. For any couple  $(\tilde{f}, \tilde{\beta})$ , demand for product  $j$  from consumer  $i$  writes:

$$\forall j \neq 0, \quad \rho_j(x_i, \tilde{f}, \tilde{\beta}) = \int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{1ij}\tilde{\beta} + x'_{2ij}v\}}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\tilde{\beta} + x'_{2ik}v\}} \tilde{f}(v) dv. \quad (13)$$

For the outside option, we have:

$$\text{for } j = 0, \quad \rho_j(x_i, \tilde{f}, \tilde{\beta}) = \int_{\mathbb{R}^{K_2}} \frac{1}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\tilde{\beta} + x'_{2ik}v\}} \tilde{f}(v) dv. \quad (14)$$

---

<sup>35</sup>In the mixed logit case, the absence of endogenous variables is not an unrealistic assumption as the econometrician can always model unobserved product quality by incorporating product fixed effects into the utility function.

**Structural error.** As we did in the case of the BLP demand model, we can define the structural error generated by  $(\tilde{\beta}, \tilde{f})$  as follows. Let  $y_{ij}$  equal to 1 if individual  $i$  chooses good  $j = 0, 1, \dots, J$ , the structural error writes:

$$\xi_{ij}(\tilde{f}, \tilde{\beta}) = y_{ij} - \rho_j(x_i, \tilde{\beta}, \tilde{f}).$$

By construction, at the true  $(f, \beta)$ , we have  $\mathbb{E}[\xi_{ij}(\beta, f)|x_i] = \mathbb{E}[y_{ij}|x_i] - \rho_j(x_i, \beta, f) = 0$  a.s.. The notation  $x_i$  refers to  $(x_{ij})_{j=1, \dots, J}$ .

**Most powerful instrument and approximations.** As in the aggregate demand model, we want to derive the instruments with the greatest ability to detect misspecification in the distribution of RCs. Given that the model displays no endogeneity, the set of exogenous variables is simply  $x_i$ . Our objective is to find the functions of  $x_i$ , which provides the most detection power against a wrong distribution. With this objective in mind, we consider a situation where the econometrician has a candidate  $(f_0, \beta_0)$  and wants to test  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$  against  $H_a : (f, \beta) \neq (f_0, \beta_0)$ . We proceed as in the BLP case and we derive the instrument that maximizes the power of the associated moment based test. Second, we propose feasible approximations of the MPI that do not depend on the fixed alternative.

For any set of testing instruments  $h_D(x_i)$ , we have the following implication:

$$\bar{H}_0 : (f, \beta) = (f_0, \beta_0) \implies \bar{H}'_0 : \mathbb{E}[h_D(x_i)\xi_{ij}(f_0, \beta_0)] = 0.$$

We propose to test  $\bar{H}_0$  indirectly through its implication  $\bar{H}'_0$ , which is a set of unconditional moment conditions. We test  $\bar{H}'_0$  with a moment-based test and the test statistic writes as follows:

$$S_N(h_D, f_0, \beta_0) = NJ \left( \frac{1}{NJ} \sum_{i,j} \xi_{ij}(f_0, \beta_0) h_D(x_i) \right)' \hat{\bar{\Omega}}_0^{-1} \left( \frac{1}{NJ} \sum_{i,j} \xi_{ij}(f_0, \beta_0) h_D(x_i) \right), \quad (15)$$

with  $\hat{\bar{\Omega}}_0$  a consistent estimator of  $\bar{\Omega}_0$  the asymptotic variance-covariance matrix of  $\frac{1}{NJ} \sum_{i,j} \xi_{ij}(f_0, \beta_0) h_D(x_i)$  under  $\bar{H}_0$ :

$$\bar{\Omega}_0 = \mathbb{E} \left[ \left( \frac{1}{\sqrt{J}} \sum_j \xi_{ij}(f_0, \beta_0) h_D(x_i) \right) \left( \frac{1}{\sqrt{J}} \sum_j \xi_{ij}(f_0, \beta_0) h_D(x_i) \right)' \right].$$

Assuming that  $(x_i, y_i)$  are i.i.d. across individuals and consistent with the probability model defined by equations (12, 13, 14) evaluated at  $(f, \beta)$ ,  $\mathbb{E}[\|\xi_{ij}(f_0, \beta_0)h_D(x_i)\|^2] < +\infty$ , and  $\Omega_0$  has full rank, we can show:

- under  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$ ,  $S_N(h_D, f_0, \beta_0) \xrightarrow[T \rightarrow +\infty]{d} \chi_{|h_D|_0}^2$ ,
- under  $H'_a : \mathbb{E}[h_D(x_i)\xi_{jt}(f_0, \beta_0)] \neq 0$ ,  $\forall q \in \mathbb{R}^+$ ,  $\mathbb{P}(S_N(h_D, f_0, \beta_0) > q) \xrightarrow[T \rightarrow +\infty]{} 1$ ,

with  $|\cdot|_0$  being the counting norm. The proof is almost identical to the proof of Proposition 2 and thus, we omit it. Following the same steps as the proof of Proposition 3, the expression for the Most Powerful Instrument (that maximizes the slope of the test) writes:

$$h_D^*(x_i) = \mathbb{E}[\xi_{ij}(f_0, \beta_0)^2 | x_i]^{-1} \Delta(x_i, f_0, \beta_0, f_a, \beta_a),$$

where each component  $j$  of the correction term  $\Delta(x_i, f_0, \beta_0, f_a, \beta_a)$  writes:

$$\begin{aligned} \Delta(x_i, f_0, \beta_0, f_a, \beta_a)_j &= \rho_j(x_i, \beta_0, f_0) - \rho_j(x_i, \beta_a, f_a) \\ &= \int_{\mathbb{R}} \left[ \rho_j(x_i, \beta_0, f_0) - \frac{\exp\{x'_{1ij}\beta_a + x'_{2ij}v\}}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\beta_a + x'_{2ik}v\}} \right] f_a(v). \end{aligned}$$

Several remarks are in order. First, contrary to the BLP case, the correction term  $\Delta_{0,a}^{\xi_j}$  is a function of the exogenous variables  $x_i$ , and thus we don't need to compute its conditional expectation as in the BLP model. The conditional variance term can be estimated, even if it is challenging in practice. For the sake of exposition, we drop this term in the rest of the analysis. As we did for the BLP case, we propose two feasible approximations of the MPI, which don't require the knowledge  $(\beta_a, f_a)$  and, which can be computed in practice.

- **Local approximation.** First, we provide a local approximation, which is accurate when  $f_0$  is close to the true density  $f_a$ . To derive this local approximation, we need to impose additional restrictions on  $\beta_0$  and  $\beta_a$  so that  $\|\beta_a - \beta_0\| = O\left(\int_{\mathbb{R}^{K_2}} |f_0(v) - f_a(v)| dv\right)$ . This is the purpose of Assumption 1

**Assumption 1.** We assume that  $\beta_0 = \beta_0^*$  and  $\beta_a = \beta_a^*$  where  $(\beta_0^*, \beta_a^*)$  are both pseudo true values, which maximize the conditional expectation of their respective population log-likelihoods. Namely,

$$\beta_0^* = \operatorname{argmax}_{\tilde{\beta} \in \mathbb{R}^{K_1}} \mathbb{E}[L(x_i, y_i, \tilde{\beta}, f_0) | x_i] \text{ with } L(x_i, y_i, \tilde{\beta}, f_0) = \sum_{j=0}^J \mathbf{1}\{y_{ij} = 1\} \log(\rho_j(x_i, \tilde{\beta}, f_0))$$

$$\beta_a^* = \operatorname{argmax}_{\tilde{\beta} \in \mathbb{R}^{K_1}} \mathbb{E}[L(x_i, y_i, \tilde{\beta}, f_a) | x_i] \text{ with } L(x_i, y_i, \tilde{\beta}, f_a) = \sum_{j=0}^J \mathbf{1}\{y_{ij} = 1\} \log(\rho_j(x_i, \tilde{\beta}, f_a))$$

Now we can derive the following first order approximation of the  $\Delta_j(x_i, f_0, \beta_0, f_a, \beta_a)$  around  $f_0$ .

**Proposition 7.**

Under Assumption 1, a first order expansion of  $\Delta_j(x_i, f_0, \beta_0, f_a, \beta_a)$  around  $f_0$  writes:

$$\Delta_j(x_i, f_0, \beta_0, f_a, \beta_a) = \int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{1ij}\beta_0 + x'_{2ij}v\}}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\beta_0 + x'_{2ik}v\}} (f_0(v) - f_a(v)) dv + \left. \frac{\partial \rho_j(x_i, \beta, f_a)}{\partial \beta} \right|_{\beta=\beta_0} (\beta_a - \beta_0) + \mathcal{R}_0$$

with  $\mathcal{R}_0 = o\left(\int_{\mathbb{R}^{K_2}} |f_0(v) - f_a(v)| dv\right)$ .

The proof is in Appendix B. Building on this approximation, we can discretize the integrals as we did in the BLP case to circumvent the fact that we do not know  $f_a$ .

$$\begin{aligned} \mathbb{E}[\Delta_j(x_i, f_0, f_a) | x_i] &\approx \sum_{l=1}^L \bar{\omega}_{1l}(f_a) \underbrace{\left[ \rho_j(x_i, \beta_0, f_0) - \frac{\exp\{x'_{1ij}\beta_0 + x'_{2ij}v_l\}}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\beta_0 + x'_{2ik}v_l\}} \right]}_{\pi_{1,j,l}(x_i)} \\ &+ \sum_{l=1}^L \bar{\omega}_{2l}(f_a) \underbrace{\frac{\partial}{\partial \beta} \left\{ \frac{\exp\{x'_{1ij}\beta_0 + x'_{2ij}v_l\}}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\beta_0 + x'_{2ik}v_l\}} \right\}}_{\pi_{2,j,l}(x_i)}, \end{aligned}$$

with  $\{v_l\}_{l=1,\dots,L}$   $L$  points chosen in the domain of definition of  $f_a$ , and  $\bar{\omega}_l(f_a)$  the unknown weights associated with each point. The local interval instruments, in the mixed logit case, write:  $(\pi_{1,j,l}(x_i), \pi_{2,j,l}(x_i))$ .

- **Global approximation.** As we did, in the BLP case, we can also write a global approximation of the corection term. To do so, we replace the unknown  $\beta_a$  by a known substitute  $\beta_0$ .<sup>36</sup> To circumvent the fact that  $f_a$  is unknown, we replace the integral with a finite sum. Namely, we have:

$$\mathbb{E}[\Delta_j(x_i, f_0, f_a)|x_i] \approx \sum_{l=1}^L \omega_l \underbrace{\left[ \rho_j(x_i, \beta_0, f_0) - \frac{\exp\{x'_{1ij}\beta_0 + x'_{2ij}v_l\}}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\beta_0 + x'_{2ik}v_l\}} \right]}_{\bar{\pi}_{j,l}(x_i)}$$

with  $\{v_l\}_{l=1,\dots,L}$   $L$  points chosen in the support of  $f_a$ , and  $\omega_l(f_a)$  the unknown weights associated with each point. The local interval instruments, in the mixed logit case, write:  $(\bar{\pi}_{1,j,l}(x_i))$

**Composite hypothesis.** In practice, as in the BLP model, the researcher must make a parametric assumption on the distribution of random coefficients to estimate the model. Formally, the econometrician assumes  $f$  belongs to a parametric family  $\mathcal{F}_0 = \{f_0(\cdot|\tilde{\lambda}) : \tilde{\lambda} \in \Lambda_0\}$ , where  $\tilde{\lambda}$  is a parameter that must be estimated. In applied work, researchers typically assume that  $f$  is normally distributed. The researcher may be interested in testing the validity of the specification. The mixed logit is often estimated by conditional MLE. To test the validity of the specification, the research must follow the same steps as the ones highlighted in Section 5. First, the researcher must estimate a pseudo-true value  $\theta_0 = (\beta'_0, \lambda'_0)' \in \mathbb{R}^{|\theta|_0}$ , which maximizes the conditional expectation of their respective population log-likelihoods under  $H_0 : f \in \mathcal{F}_0$ . Namely:

$$\theta_0 = \underset{\tilde{\theta} \in \mathbb{R}^{|\theta|_0}}{\operatorname{argmax}} \mathbb{E}[L(x_i, y_i, \tilde{\beta}, f_0(\cdot|\tilde{\lambda}))|x_i] \text{ with } L(x_i, y_i, \tilde{\beta}, f_0(\cdot|\tilde{\lambda})) = \sum_{j=0}^J \mathbf{1}\{y_{ij} = 1\} \log(\rho_j(x_i, \tilde{\beta}, f_0(\cdot|\tilde{\lambda})))$$

Next, we test  $H'_0 : \mathbb{E}[h_D(x_i)\xi_{ij}(f_0(\cdot|\tilde{\lambda}), \beta_0)] = 0$  with the moment based test exhibited above. To derive the asymptotic distribution of  $S_N(h_D, f_0(\cdot|\hat{\lambda}, \hat{\beta}))$ , we must now take into account the parameter uncertainty stemming from the first stage estimation. As in the BLP demand model, the integrals must be numerically computed to recover the theoretical probabilities implied by

---

<sup>36</sup>in our simulations, we find that the homogeneous parameters are usually close to each other even when the distributions are somewhat distant from each other.

the model, and compute the conditional likelihood. Thus, one must also take into account the numerical approximations in the derivation of the asymptotic distribution.

## B Proofs

### B.1 Identification

In this subsection, we prove that under Assumption A, the distribution of random coefficients  $f$  is non-parametrically point identified.

*Proof.* Proof of Proposition 1

We want to show that under Assumptions A, the following implication holds:

$$\begin{aligned} (\tilde{f}, \tilde{\beta}) = (f, \beta) &\iff \mathbb{E}[\xi_{jt}(\tilde{f}, \tilde{\beta})|z_{jt}] = 0 \text{ a.s.} \\ &\iff \mathbb{E}\left[\rho_j^{-1}(s_t, x_{2t}, \tilde{f}) - x'_{1jt}\tilde{\beta} \middle| z_{jt}\right] = 0 \text{ a.s..} \end{aligned}$$

**Step 1:** First, we show that for any random permutation of indexes  $j \rightarrow j'$ , the following equivalence holds:

$$\mathbb{E}[\xi_{jt}|z_{jt}] = 0 \text{ a.s.} \iff \mathbb{E}[\xi_{jt}|z_{j't}] = 0 \text{ a.s.} \quad \forall j'.$$

As the new indexation is done exogenously, we have for any  $j'$ :

$$\mathbb{E}[\xi_{jt}(\tilde{f}, \tilde{\beta})|z_{jt}] = \mathbb{E}[\xi_{jt}(\tilde{f}, \tilde{\beta})|z_{jt}, j \rightarrow j'] \equiv \mathbb{E}_{j'}[\xi_{j't}(\tilde{f}, \tilde{\beta})|z_{j't}] \text{ a.s.,}$$

with  $j \rightarrow j'$  indicates index  $j$  has been changed into  $j'$ . Consequently, we have:

$$\mathbb{E}[\xi_{jt}(\tilde{f}, \tilde{\beta})|z_{jt}] = 0 \text{ a.s.} \iff \forall j' \quad \mathbb{E}_{j'}[\xi_{j't}(\tilde{f}, \tilde{\beta})|z_{j't}] = 0 \text{ a.s.}$$

This last equivalence allows us to come back to the exogeneity condition assumed in [Berry and Haile \(2014\)](#) and in [Wang \(2022\)](#):  $\forall k, \quad \mathbb{E}[\xi_{jt}|z_{jt}, j = k] = 0 \text{ a.s.}$ . The only difference being that here  $j'$  is determined completely randomly. Intuitively, the exogeneity condition required for non-parametric identification of the demand functions is stronger than the one needed for the non-parametric identification of the distribution of RC.



**Step 2:** We now need to show the following equivalence:

$$(\tilde{f}, \tilde{\beta}) = (f, \beta) \iff \forall j', \quad \mathbb{E}_{j'}[\xi_{j't}(\tilde{f}, \tilde{\beta})|z_{j't}] = 0 \text{ a.s.}$$

Given the random permutation  $j \rightarrow j'$ , which is market dependent, we must redefine our matrices and vectors as follows:  $\hat{x}_t = M_t x_t$  with  $(M_t)_{i,k} = \mathbf{1}\{i = j_t, k = j'_t\}$ . Likewise  $\hat{s}_t = M_t s_t$ .  $M_t$  is a random matrix. It is straight forward to show the direct implication.

$$(\tilde{f}, \tilde{\beta}) = (f, \beta) \implies \forall j', \quad \mathbb{E}_{j'} \left[ \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - x'_{1j't} \tilde{\beta} \middle| z_{j't} \right] = \mathbb{E}_{j'}[\xi_{j't}(f, \beta)|z_{j't}] = 0 \text{ a.s.}$$

The reverse implication is much more intricate to prove and we will exploit other results in the literature. We want to show:

$$(\tilde{f}, \tilde{\beta}) \neq (f, \beta) \implies \exists j' \quad \left| \quad \mathbb{E}_{j'} \left[ \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - \hat{x}'_{1j't} \tilde{\beta} \middle| z_{j't} \right] = 0 \text{ a.s.} \right. \text{ does not hold.}$$

**Case 1:** First, let us assume that  $\tilde{f} = f$  and  $\tilde{\beta} \neq \beta$ , then we have:

$$\rho^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - \hat{x}_{1t} \tilde{\beta} = \underbrace{\rho^{-1}(\hat{s}_t, \hat{x}_{2t}, f) - x_{1t} \beta}_{\xi_t(f, \beta)} + \hat{x}_{1t}(\beta - \tilde{\beta})$$

By assumption, we have:  $P(x'_{1t} x_{1t} \text{ } dp) > 0$ .  $M_t$  is symmetric, idempotent and full rank. As a consequence,

$$P(\hat{x}'_{1t} \hat{x}_{1t} \text{ } dp) = P(x'_{1t} M_t x_{1t} \text{ } dp) = P(x'_{1t} x_{1t} \text{ } dp) > 0$$

Therefore, we have  $\forall \gamma \neq 0 \in \mathbb{R}^K$ ,

$$\begin{aligned} P(\gamma' \hat{x}'_{1t} \hat{x}_{1t} \gamma > 0) &> P(\hat{x}'_{1t} \hat{x}_{1t} \text{ } dp) > 0 \iff P(\|\hat{x}_{1t} \gamma\|^2 > 0) > 0 \\ &\iff P(\hat{x}_{1t} \gamma \neq 0) > 0 \end{aligned}$$

Thus,  $\exists j' \mid x'_{1j't}(\beta - \tilde{\beta}) = 0 \text{ a.s.}$  does not hold. To conclude, there exists  $j'$  such that:

$$\mathbb{E}_{j'}[\rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, f) - x'_{1j't} \tilde{\beta} | z_{j't}] = \underbrace{\mathbb{E}_{j'}[\xi_{j't}(f, \beta) | z_{j't}]}_{=0} + \underbrace{\mathbb{E}_{j'}[x'_{1j't}(\beta - \tilde{\beta}) | z_{j't}]}_{=0 \text{ a.s. does not hold from the completeness}}$$

**Case 2:** Now let us assume that  $\tilde{f} \neq f$  and we want to show that  $\forall \tilde{\beta} \in \mathbb{R}^k, \exists j'$  such that:

$$\mathbb{E}_{j'} \left[ \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - x'_{1j't} \tilde{\beta} \middle| z_{jt} \right] = 0 \text{ a.s. does not hold.}$$

First, let us observe that  $\forall j'$ ,

$$\mathbb{E}_{j'} [\rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - x'_{1j't} \tilde{\beta} | z_{j't}] = \underbrace{\mathbb{E}_{j'} [\xi_{j't}(f, \beta) | z_{j't}]}_{=0} + \mathbb{E}_{j'} [\rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, f) - x'_{1j't}(\tilde{\beta} - \beta) | z_{j't}].$$

As a consequence, we need to show that  $\exists j'$  such that  $\mathbb{E}_{j'} [\rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, f) - x'_{1j't}(\tilde{\beta} - \beta) | z_{j't}] = 0$  a.s. does not hold. From the completeness condition, a sufficient condition is:  $\exists j'$  such that  $\rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, f) - x'_{1j't}(\tilde{\beta} - \beta) = 0$  a.s. does not hold.

Let  $\gamma = (\tilde{\beta} - \beta)$ . By contradiction, it can be easily be shown that  $\rho(\hat{\delta}_t, \hat{x}_{2t}, f) - \rho(\hat{\delta}_t + \hat{x}_{1t}\gamma, \hat{x}_{2t}, \tilde{f}) \neq 0 \implies \exists j' \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) \neq \rho_j^{-1}(\hat{s}_t, \hat{x}_{2t}, f) + \gamma' x_{1j't}$ . Indeed, assume that  $\rho(\hat{\delta}_t, \hat{x}_{2t}, f) - \rho(\hat{\delta}_t + \hat{x}_{1t}\gamma, \hat{x}_{2t}, \tilde{f}) \neq 0$  and  $\forall j' \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) = \rho_j^{-1}(\hat{s}_t, \hat{x}_{2t}, f) + \gamma' x_{1j't}$ . Then, we have:  $\rho(\rho^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}), \hat{x}_{2t}, \tilde{f}) = \rho(\rho^{-1}(\hat{s}_t, \hat{x}_{2t}, f) + \hat{x}_{1t}\gamma, \hat{x}_{2t}, \tilde{f}) = \rho(\hat{\delta}_t + \hat{x}_{1t}\gamma, \hat{x}_{2t}, \tilde{f}) \neq \rho(\hat{\delta}_t, \hat{x}_{2t}, f) = \hat{s}_t$ . Therefore, we have a contradiction.

Hence, the next step is to show that  $\forall \gamma, \tilde{f} \neq f \implies \rho(\hat{\delta}_t, \hat{x}_{2t}, f) - \rho(\hat{\delta}_t + \hat{x}_{1t}\gamma, \hat{x}_{2t}, \tilde{f}) = 0$  a.s. does not hold.

To this end, we are going to exploit the identification result shown by Wang (2022). Following the notations in this paper, we define  $\mu_i = \hat{x}_{1t}\Gamma + \hat{x}_{2t}v_i = \hat{x}_t\mathbf{v}$  with  $\mathbf{v}_i = (\Gamma, v_i)$ . Here  $\Gamma$  is a degenerate random variable characterized by constant  $c$  such that  $P(\Gamma = c) = 1$ . Let  $G_{\mu|\hat{x}_t}$  the distribution of  $\mu_i|\hat{x}_t$  under  $f^\dagger = (c = 0, f)$  and  $G_{\tilde{\mu}|\hat{x}_t}$  the distribution of  $\mu_i|\hat{x}_t$  under  $\tilde{f}^\dagger = (c = \gamma, \tilde{f})$ . The following result is shown in Wang (2022): for any  $\hat{x}_t \in \text{Supp}(\hat{x}_t)$ ,

$$\exists j' \mid \rho_{j'}(\hat{\delta}_t, G_{\mu|\hat{x}_t}) - \rho_{j'}(\hat{\delta}_t, G_{\tilde{\mu}|\hat{x}_t}) = 0 \text{ on open set } \mathcal{D} \subset \mathbb{R}^J \implies G_{\mu|\hat{x}_t} = G_{\tilde{\mu}|\hat{x}_t}.$$

Thanks to the real analytic property of the demand functions  $\rho$ , Wang (2022) does not require a full support assumption on  $\hat{\delta}_t$ .

Fix the value of  $\hat{x}_t$  as follows:  $\hat{x}_t = \bar{M}_t \bar{x}_t = \hat{\hat{x}}_t$ . By assumption, there exists  $\bar{x}_t \in \text{Supp}(x_t)$  such that  $\bar{x}_t' \bar{x}_t$  is dp and  $\delta_t = \bar{x}_{1t} \beta + \xi_t$  varies on an open set  $\bar{D}$  almost surely. These properties naturally transmit to  $\hat{\hat{x}}_t$  Now. The chosen permutation  $\bar{M}_t$  doesn't matter. Given the result in Wang (2022), in order to prove that  $\rho(\hat{\delta}_t, \hat{x}_{2t}, f_0) - \rho(\hat{\delta}_t + \hat{x}_{1t} \gamma, \hat{x}_{2t}, f) = 0$  a.s. does not hold, we just need to prove that  $\forall \gamma, \tilde{f} \neq f \implies G_{\tilde{\mu}|\hat{\hat{x}}_t} \neq G_{\mu|\hat{\hat{x}}_t}$ . By definition (see assumption A (iv)),  $\tilde{f} \neq f \implies \exists v^* \in \mathbb{R}^{K_2} \tilde{F}(v^*) \neq F(v^*)$ . Take  $x^* = (0_{K_1}, \hat{\hat{x}}_{2t} v^*)' = \hat{\hat{x}}_t(0_{K_1}, v^*)'$ :

$$\begin{aligned} G_{\mu|\hat{\hat{x}}_t}(x^*) &= P(x_t \mathbf{v}_i \leq x^* | x_t = \hat{\hat{x}}_t) = P((x_t' x_t)^{-1} x_t' x_t \mathbf{v}_i \leq (x_t' x_t)^{-1} x_t' \hat{\hat{x}}_t(0_{K_1}, v^*)' | x_t = \hat{\hat{x}}_t). \\ &= (1_{K_1}, P(v_i \leq v^* | x_t = \hat{\hat{x}}_t))' = (1_{K_1}, F(v^*))' \end{aligned}$$

The last equality comes from independence of  $v_i$  and  $x_t$ . Likewise,  $G_{\tilde{\mu}|\hat{\hat{x}}_t}(x^*) = (1\{\gamma > 0\}, \tilde{F}(v^*))'$

Therefore,  $\exists x^*, \forall \gamma \ G_{\tilde{\mu}|\hat{\hat{x}}_t}(x^*) \neq G_{\mu|\hat{\hat{x}}_t}(x^*)$ . Following the result in Wang (2022), we have that for all  $\gamma \in \mathbb{R}^{K_1}$ ,  $\rho(\hat{\delta}_t, \hat{x}_{2t}, f) - \rho(\hat{\delta}_t + \hat{x}_{1t} \gamma, \hat{x}_{2t}, \tilde{f}) = 0$  a.s. does not hold, which in turn implies that for all  $\gamma \in \mathbb{R}^{K_1}$ ,  $\exists j' \ \rho_j^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - \rho_j^{-1}(\hat{s}_t, \hat{x}_{2t}, f) + \hat{x}_{1jt}' \gamma = 0$  a.s. does not hold.

To conclude:  $\forall \beta \in \mathbb{R}^k$ , there exists  $j'$  such that:

$$\rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, \tilde{f}) - \rho_{j'}^{-1}(\hat{s}_t, \hat{x}_{2t}, f) - x_{1j't}'(\tilde{\beta} - \beta) = 0 \text{ a.s. does not hold,}$$

which is what we wanted to show. □

In Section 5, we used the following equivalence between the composite hypothesis and the pseudo-true value to construct the specification test.

**Corollary 2.1.** Under Assumption A, and assume  $h_E(z_{jt})$  and  $W$  are such that the pseudo-true value  $\theta_0$  is unique, then we have:

$$H_0 : f \in \mathcal{F}_0 \iff \bar{H}_0 : (f, \beta) = (f_0(\cdot | \lambda_0), \beta_0).$$

*Proof.* Proof of Corollary 2.1

Let us assume that under specification  $\mathcal{F}_0$ , instruments  $h_E(z_{jt})$  and weighting matrix  $W$ , the pseudo true value is unique.

- Under  $H_0 : f \in \mathcal{F}_0$  and there exists  $\lambda$  such that  $f = f_0(\cdot|\lambda)$ . By the mean independence assumption on the unobserved quality  $\xi_{jt}$ , we have at the true  $\theta = (\beta, \lambda)$ :

$$\xi_{jt}(f_0(\cdot|\lambda), \beta) = \rho_j^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda)) - x'_{1jt}\beta = \xi_{jt} \implies \mathbb{E}[(\xi_{jt}(f_0(\cdot|\lambda), \beta)h_E(z_{jt}))] = 0.$$

Thus,  $\theta$  is solution to the previous minimization problem and as the solution is unique:  $\theta_0 = \theta$ . As a consequence,  $\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) = \xi_{jt}$  and  $\mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)|z_{jt}] = 0$  a.s..

- Under an alternative specification:  $f \notin \mathcal{F}_0$ , we know from the Proposition 1 that  $\forall \tilde{\theta} = (\tilde{\beta}, \tilde{\lambda})$ ,

$$\mathbb{E}\left[\rho_j^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) - x'_{1jt}\tilde{\beta} \middle| z_{jt}\right] = 0 \text{ a.s. does not hold.}$$

In particular, the last equation doesn't hold at the true value  $\tilde{\theta} = \theta_0$ .

□

## B.2 Detecting misspecification: the most powerful instrument

*Proof of Proposition 2.*

- Under  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$ . By assumption, the data are i.i.d. across markets,  $\mathbb{E}[\|\xi_{jt}(f_0, \beta_0)h_D(z_{jt})\|^2] = \frac{1}{J}\mathbb{E}[\sum_j \|\xi_{jt}(f_0, \beta_0)h_D(z_{jt})\|^2] < +\infty$ , the CLT applies:

$$\frac{1}{\sqrt{TJ}} \sum_{j,t} h_D(z_{jt})\xi_{jt}(f_0, \beta_0) = \frac{1}{\sqrt{TJ}} \sum_{j,t} h_D(z_{jt})\xi_{jt} \xrightarrow{T \rightarrow +\infty} \mathcal{N}(0, \tilde{\Omega}_0),$$

with:

$$\begin{aligned} \tilde{\Omega}_0 &= \mathbb{E} \left[ \left( \frac{1}{\sqrt{J}} \sum_{j=1}^J h_D(z_{jt})\xi_{jt} \right) \left( \frac{1}{\sqrt{J}} \sum_{j=1}^J h_D(z_{jt})\xi_{jt} \right)' \right] \\ &= \frac{1}{J} \mathbb{E} \left[ \sum_{j=1}^J h_D(z_{jt})h_D(z_{jt})'\xi_{jt}^2 + \sum_{j=1}^J \sum_{k \neq j}^J h_D(z_{jt})h_D(z_{kt})'\xi_{jt}\xi_{kt} \right] \\ &= \frac{1}{J} \mathbb{E} \left[ \sum_{j=1}^J h_D(z_{jt})h_D(z_{jt})'\xi_{jt}^2 \right] + \frac{1}{J} \sum_{j=1}^J \sum_{k \neq j}^J \mathbb{E} \left[ h_D(z_{jt})h_D(z_{kt})' \underbrace{\mathbb{E}[\xi_{jt}\xi_{kt}|z_{jt}, z_{kt}]}_{=0} \right] \\ &= \mathbb{E} [h_D(z_{jt})h_D(z_{jt})'\xi_{jt}^2] \\ &= \Omega_0. \end{aligned}$$

Third line comes from  $\xi_{jt} \perp \xi_{kt}|z_t$ . By assumption,  $\Omega_0$  has a full rank. Thus, we have by the CMT:

$$S_T(h_D, f_0, \beta_0) = TJ \left( \frac{1}{TJ} \sum_{j,t} \xi_{jt}(f_0, \beta_0) h_D(z_{jt}) \right)' \hat{\Omega}_0^{-1} \left( \frac{1}{TJ} \sum_{j,t} \xi_{jt}(f_0, \beta_0) h_D(z_{jt}) \right) \xrightarrow[T \rightarrow +\infty]{d} \chi^2_{|h_D|_0}.$$

- Under  $H'_a : \mathbb{E}[h_D(z_{jt})\xi_{jt}(f_0, \beta_0)] \neq 0$ . The data are i.i.d. across markets, by the law of large numbers:  $\frac{1}{TJ} \sum_{j,t} h_D(z_{jt})\xi_{jt}(f_0, \beta_0) \xrightarrow{P} \mathbb{E} \left[ \frac{1}{J} \sum_j h_D(z_{jt})\xi_{jt}(f_0, \beta_0) \right]$ . It follows by the continuous mapping theorem:

$$\begin{aligned} \frac{S_T(h_D, f_0, \beta_0)}{T} &\xrightarrow{P} J \mathbb{E} \left[ \frac{1}{J} \sum_j h_D(z_{jt})\xi_{jt}(f_0, \beta_0) \right]' \Omega_0^{-1} \mathbb{E} \left[ \frac{1}{J} \sum_j h_D(z_{jt})\xi_{jt}(f_0, \beta_0) \right] \\ &= J \underbrace{\mathbb{E}[h_D(z_{jt})\xi_{jt}(f_0, \beta_0)]' \Omega_0^{-1} \mathbb{E}[h_D(z_{jt})\xi_{jt}(f_0, \beta_0)]}_{\kappa(h_D, f_0, \beta_0)} \end{aligned}$$

Under  $H'_a$ ,  $\kappa(h_D, f_0, \beta_0)$  is strictly positive because  $\Omega_0$  is positive definite. Thence,

$$\begin{aligned} \forall q \in \mathbb{R}, \quad \lim_{T \rightarrow \infty} \mathbb{P}(S_T(h_D, f_0, \beta_0) > q) &= \lim_{T \rightarrow \infty} \mathbb{P} \left( \frac{S(h_D, f_0, \beta_0) - q}{T} > 0 \right) \\ &= \mathbb{P}(J\kappa(h_D, f_0, \beta_0) > 0) \\ &= 1, \end{aligned}$$

where the second equality holds because convergence in probability implies convergence in distribution. □

*Proof of Proposition 3.* To shorten notations, let  $\xi_{jt0} \equiv \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)$ ,  $\xi_{jta} \equiv \xi_{jt}(f_a, \beta_a)$  and  $\xi_{t0}$  and  $\xi_{ta}$  their stacked versions over  $j$ . Likewise, we define  $h_D(z_t) = (h_D(z_{1t}), \dots, h_D(z_{Jt}))'$ . Under  $\bar{H}_a : (f, \beta) = (f_a, \beta_a)$ , the asymptotic slope of the test writes:

$$\begin{aligned} c_{h_D}(f_a, \beta_a) &= \mathbb{E} \left( \sum_j \xi_{jt0} h_D(z_{jt}) \right)' \mathbb{E} \left( \left( \sum_j \xi_{jt0} h_D(z_{jt}) \right) \left( \sum_{j'} \xi_{j't0} h_D(z_{j't}) \right)' \right)^{-1} \mathbb{E} \left( \sum_j \xi_{jt0} h_D(z_{jt}) \right) \\ &= \mathbb{E}(\xi'_{t0} h_D(z_t)) \mathbb{E}(h_D(z_t)' \xi_{t0} \xi'_{t0} h_D(z_t))^{-1} \mathbb{E}(h_D(z_t)' \xi_{t0}) \\ &= \mathbb{E}(\Delta_{0,a}^{\xi_t}{}' h_D(z_t)) \mathbb{E}(h_D(z_t)' \mathbb{E}(\xi_{t0} \xi'_{t0} | z_t) h_D(z_t))^{-1} \mathbb{E}(h_D(z_t)' \Delta_{0,a}^{\xi_t}). \end{aligned}$$

Third line comes from  $\mathbb{E}(\Delta_{0,a}^{\xi_t}{}' h_D(z_t)) = \mathbb{E}((\xi_{t0} - \xi_{ta})' h_D(z_t)) = \mathbb{E}(\xi_{t0}' h_D(z_t))$  because  $\xi_{ta}$  is the true structural error. Then the slope of the test taking  $h_D^*(z_t) = \mathbb{E}(\xi_{t0} \xi_{t0}' | z_t)^{-1} \mathbb{E}(\Delta_{0,a}^{\xi_t} | z_t)$  is equal to:

$$c_{h_D^*}(f_a, \beta_a) = \mathbb{E} \left( \mathbb{E}(\Delta_{0,a}^{\xi_t} | z_t)' \mathbb{E}(\xi_{t0} \xi_{t0}' | z_t)^{-1} \mathbb{E}(\Delta_{0,a}^{\xi_t} | z_t) \right)$$

To finish the proof, we must show that for any set of instruments  $h_D$ , we have:  $c_{h_D^*}(f_a, \beta_a) \geq c_{h_D}(f_a, \beta_a)$ .

Denote  $\tilde{h}_D(z_t) = \mathbb{E}(\xi_{t0} \xi_{t0}' | z_t)^{1/2} h_D(z_t)$  and  $\tilde{h}_D^*(z_t) = \mathbb{E}(\xi_{t0} \xi_{t0}' | z_t)^{1/2} h_D^*(z_t)$ . With these new notations, we have:

$$\begin{aligned} c_{h_D^*}(f_a, \beta_a) - c_{h_D}(f_a, \beta_a) &= \mathbb{E} \left( \tilde{h}_D^*(z_t)' \tilde{h}_D^*(z_t) \right) - \mathbb{E} \left( \tilde{h}_D^*(z_t)' \tilde{h}_D(z_t) \right) \mathbb{E} \left( \tilde{h}_D(z_t)' \tilde{h}_D(z_t) \right)^{-1} \mathbb{E} \left( \tilde{h}_D(z_t)' \tilde{h}_D^*(z_t) \right) \\ &= G' \begin{pmatrix} \mathbb{E} \left( \tilde{h}_D^*(z_t)' \tilde{h}_D^*(z_t) \right) & \mathbb{E} \left( \tilde{h}_D^*(z_t)' \tilde{h}_D(z_t) \right) \\ \mathbb{E} \left( \tilde{h}_D(z_t)' \tilde{h}_D^*(z_t) \right) & \mathbb{E} \left( \tilde{h}_D(z_t)' \tilde{h}_D(z_t) \right) \end{pmatrix} G \\ &= G' \mathbb{E} \left( \tilde{H} \tilde{H}' \right) G \geq 0, \end{aligned}$$

$$\text{with } \tilde{H} = (\tilde{h}_D^*(z_t), \tilde{h}_D(z_t))' \text{ and } G = \left( 1, -\mathbb{E} \left( \tilde{h}_D^*(z_t)' \tilde{h}_D(z_t) \right) \mathbb{E} \left( \tilde{h}_D(z_t)' \tilde{h}_D(z_t) \right)^{-1} \right)'.$$

**Special case:** when we assume for  $k \neq j$ ,  $\xi_{jt} \perp \xi_{kt} | z_t$ , and take  $\hat{\Omega}_0 = \frac{1}{JT} \sum_{j,t} \xi_{jt0}^2 h_D(z_{jt}) h_D(z_{jt})'$  as our weighting matrix (as we do for illustrations purposes in the main text), we find that the slope under  $\bar{H}_a$  writes:

$$c_{h_D}(f_a, \beta_a) = \mathbb{E}(\Delta_{0,a}^{\xi_{jt}} h_D(z_{jt}))' \mathbb{E}(h_D(z_{jt}) h_D(z_{jt})' \mathbb{E}(\xi_{jt0}^2 | z_{jt}))^{-1} \mathbb{E}(h_D(z_{jt}) \Delta_{0,a}^{\xi_{jt}}).$$

Using the same arguments as previously, one can show that a maximizer of the slope of the test is obtained by taking  $h_D^*(z_{jt}) = \mathbb{E}(\xi_{jt}^2 | z_{jt})^{-1} \mathbb{E}(\Delta_{0,a}^{\xi_{jt}} | z_{jt})$ .

□

*Proof of Proposition 4.*

Under Assumption A, Proposition 1 implies the following:

$$\begin{aligned}
\bar{H}_a : (f, \beta) = (f_a, \beta_a) \neq (f_0, \beta_0) &\implies \mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}] \neq 0 \text{ a.s.} \\
&\implies \mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}]^2 > 0 \text{ a.s.} \\
&\implies \mathbb{E}[\mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}]^2] > 0 \\
&\implies \mathbb{E}[\mathbb{E}[\xi_{jt}(f_0, \beta_0)\mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}]]] > 0 \\
&\implies \mathbb{E}[\xi_{jt}(f_0, \beta_0)\mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}]] > 0 \\
&\implies \bar{H}'_a : \mathbb{E}[\xi_{jt}(f_0, \beta_0) \underbrace{\mathbb{E}[\Delta_{0,a}^{\xi_{jt}}|z_{jt}]}_{h_D^*(z_{jt})}] \neq 0.
\end{aligned}$$

Under the same assumptions as 2, we have the following:

$$\bar{H}'_a : \mathbb{E}[\xi_{jt}(f_0, \beta_0)h_D^*(z_{jt})] \neq 0 \implies \forall q \in \mathbb{R}^+, \mathbb{P}(S_T(h_D^*, \mathcal{F}_0, \hat{\theta}) > q) \rightarrow 1.$$

□

*Proof of Proposition 5.*

Let  $\mathcal{H}$  the set of measurable functions of  $z_{jt}$ , we want to show under  $\bar{H}_a$ :

$$\forall \alpha \in \mathbb{R}^*, \alpha \mathbb{E}[\Delta_{0,a}^{\xi_{jt}}|z_{jt}] \in \arg \max_{h \in \mathcal{H}} \text{corr}(\xi_{jt}(f_0, \beta_0), h(z_{jt})).$$

We proceed in 2 steps. First, we derive the upper bound by showing that for any  $h \in \mathcal{H}$ , we have:

$$\text{corr}(\xi_{jt}(f_0, \beta_0), h(z_{jt})) \leq \sqrt{\frac{\text{var}(\mathbb{E}[\Delta_{0,a}^{\xi_{jt}}|z_{jt}])}{\text{var}(\xi_{jt}(f_0, \beta_0))}}.$$

To do so, we use the definition of the conditional expectation and the Cauchy Schwarz inequality. First notice that we have:  $\mathbb{E}[\Delta_{0,a}^{\xi_{jt}}|z_{jt}] = \mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}]$ . By definition of the conditional expectation, we have for any  $h \in \mathcal{H}$ ,

$$\mathbb{E}[h(z_{jt})\xi_{jt}(f_0, \beta_0)] = \mathbb{E}[h(z_{jt})\mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}]].$$

It follows that:

$$|\text{cov}(h(z_{jt}), \xi_{jt}(f_0, \beta_0))| = \text{cov}(h(z_{jt}), \mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}]) \leq \sqrt{\text{var}(h(z_{jt}))\text{var}(\mathbb{E}[\xi_{jt}(f_0, \beta_0)|z_{jt}])}.$$

The inequality comes from the Cauchy Schwarz inequality. The result follows by using the definition of the correlation coefficient.

Second, we show that the upper bound is reached by taking for any  $\alpha \in \mathbb{R}^*$ ,  $h_D^*(z_{jt}) = \alpha \mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}]$ .

$$\begin{aligned} \text{cov} \left( \xi_{jt}(f_0, \beta_0), \alpha \mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}] \right) &= \alpha \text{cov} \left( \Delta_{0,a}^{\xi_{jt}}, \mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}] \right) \\ &= \alpha \text{var} \left( \mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}] \right). \end{aligned}$$

Consequently,

$$\text{corr}(\xi_{jt}(f_0, \beta_0), h_D^*(z_{jt})) = \frac{\alpha}{\sqrt{\alpha^2}} \sqrt{\frac{\text{var}(\mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}])}{\text{var}(\xi_{jt}(f_0, \beta_0))}} \implies |\text{corr}(\xi_{jt}(f_0, \beta_0), h_D^*(z_{jt}))| = \sqrt{\frac{\text{var}(\mathbb{E}[\Delta_{0,a}^{\xi_{jt}} | z_{jt}])}{\text{var}(\xi_{jt}(f_0, \beta_0))}}.$$

□

### B.2.1 Connection with optimal instruments

In the parametric case and assuming that the model is well specified ( $f \in \mathcal{F}_0$ ) the BLP parameter  $\theta_0$  is identified by the following non-linear conditional moment restriction  $\mathbb{E}[\xi_{jt}(\theta_0) | z_{jt}] = 0$ . The derivation of the optimal instruments in this context has been studied by [Amemiya \(1974\)](#). For an arbitrary choice of  $h_E(z_{jt})$ , the GMM estimator with the 2-step efficient weighting matrix has the following asymptotic distribution:

$$\sqrt{T}(\hat{\theta} - \theta_0) \xrightarrow{d} \mathcal{N}(0, (\Gamma(\mathcal{F}_0, \theta, h_E)' \Omega(\mathcal{F}_0, h_E)^{-1} \Gamma(\mathcal{F}_0, \theta, h_E))^{-1}),$$

with the same notations as previously:

$$\begin{aligned} \Omega(\mathcal{F}_0, h_E) &= \mathbb{E} \left[ \left( \sum_j \xi_{jt}(\theta) h_E(z_{jt}) \right) \left( \sum_j h_E(z_{jt}) \xi_{jt}(\theta) \right)' \right] \\ \Gamma(\mathcal{F}_0, \theta_0, h_E) &= \mathbb{E} \left[ \sum_j h_E(z_{jt}) \frac{\partial \xi_{jt}(\theta_0)}{\partial \tilde{\theta}'} \right]. \end{aligned}$$

For the sake of exposition, we will assume that unobserved demand shock  $\xi_{jt}$  is independent across observations, namely:  $\mathbb{E}[\xi_{jt}(\theta_0) \xi_{j't}(\theta) | z_t] = 0$  for  $j \neq j'$ . The general case extends naturally. The optimal instrument  $h_E^*(z_{jt})$  are chosen to minimize the asymptotic variance covariance matrix.



We derive the form of the optimal instruments in the context of BLP by applying well known results in [Chamberlain \(1987\)](#) and [Amemiya \(1974\)](#)

**Lemma 2.2.** Optimal instruments in the BLP model.

In our setting and assuming  $f \in \mathcal{F}_0$ , the optimal instruments  $h_E^*(z_{jt})$  write:

$$h_E^*(z_{jt}) = \mathbb{E}[\xi_{jt}(\theta_0)^2 | z_{jt}]^{-1} \mathbb{E} \left[ \frac{\partial \xi_{jt}(\theta_0)}{\partial \theta} \middle| z_{jt} \right].$$

and the corresponding efficiency bound (obtained by setting  $h_E = h_E^*$ ) writes:

$$V^* = \mathbb{E} \left[ \sum_j \mathbb{E} \left[ \frac{\partial \xi_{jt}(\theta_0)}{\partial \theta} \middle| z_{jt} \right] \mathbb{E} \left[ \frac{\partial \xi_{jt}(\theta_0)}{\partial \theta} \middle| z_{jt} \right]' \mathbb{E}[\xi_{jt}(\theta)^2 | z_{jt}]^{-1} \right]^{-1}$$

*Proof.* To shorten the notations, we denote:  $\sigma^2(z_{jt}) = \mathbb{E}[\xi_{jt}(\theta_0)^2 | z_{jt}]$  and  $d(z_{jt}) = \mathbb{E} \left[ \frac{\partial \xi_{jt}(\theta_0)}{\partial \theta} \middle| z_{jt} \right]$ . Likewise, we define:

$$\Omega_0(h_E) = \mathbb{E} \left[ \sum_j \mathbb{E}[\xi_{jt}(\theta_0)^2 | z_{jt}] h_E(z_{jt}) h_E(z_{jt})' \right].$$

We want to prove that for any set of instruments  $h_E(z_{jt})$  that  $V^*(z_{jt}) - \Gamma_0(h_E)' \Omega_0(h_E)^{-1} \Gamma_0(h_E)$  matrix is semi definite positive.

$$\begin{aligned} V^*(z_{jt}) - \Gamma_0(h_E) \Omega_0(h_E)^{-1} \Gamma_0(h_E)' &= \\ &= \mathbb{E} \left[ \sum_j d(z_{jt}) d(z_{jt})' \sigma^2(z_{jt}) \right] - \mathbb{E} \left[ \sum_j \frac{\partial \xi_{jt}(\theta_0)}{\partial \theta} h_E(z_{jt})' \right] \Omega_0(h_E)^{-1} \mathbb{E} \left[ \sum_j \frac{h_E(z_{jt}) \partial \xi_{jt}(\theta_0)'}{\partial \theta} \right] \\ &= \mathbb{E} \left[ \sum_j d(z_{jt}) d(z_{jt})' \sigma^{-2}(z_{jt}) \right] - \mathbb{E} \left[ \sum_j d(z_{jt}) h_E(z_{jt})' \right] \mathbb{E} \left[ \sum_j \sigma^2(z_{jt}) h_E(z_{jt}) h_E(z_{jt})' \right]^{-1} \mathbb{E} \left[ \sum_j h_E(z_{jt}) d(z_{jt})' \right] \\ &= \mathbb{E} \left[ \tilde{D}(z_{jt})' \tilde{D}(z_{jt}) \right] - \mathbb{E} \left[ \tilde{D}(z_{jt})' \tilde{H}_E(z_{jt}) \right] \mathbb{E} \left[ \tilde{H}_E(z_{jt})' \tilde{H}_E(z_{jt}) \right]^{-1} \mathbb{E} \left[ \tilde{H}_E(z_{jt})' \tilde{D}(z_{jt}) \right]. \end{aligned}$$

The second line comes from law of iterated expectations. The third line is a matricial way to rewrite the second line.  $\tilde{D}(z_{jt})$  a matrix which stacks  $d(z_{jt})/\sigma(z_{jt})$  over the set of products (each line corresponds to one product  $j$ ). Likewise, let  $\tilde{H}_E(z_{jt})$  a matrix which stacks  $h_E(z_{jt})\sigma(z_{jt})$  over

the set of products (each line corresponds to one product  $j$ ). Now let us define the following matrices.

$$\tilde{X} = \begin{pmatrix} \tilde{D}(z_{jt}) & \tilde{H}_E(z_{jt}) \end{pmatrix} \quad \text{and} \quad \tilde{M} = \begin{pmatrix} I_{|\theta_0|} & -\mathbb{E} \left[ \tilde{D}(z_{jt})' \tilde{H}_E(z_{jt}) \right] \mathbb{E} \left[ \tilde{H}_E(z_{jt})' \tilde{H}_E(z_{jt}) \right]^{-1} \end{pmatrix}'$$

We have:  $V^*(z_{jt}) - \Gamma_0(h_E) \Omega_0(h_E)^{-1} \Gamma_0(h_E) = \tilde{M}' \mathbb{E}[\tilde{X}' \tilde{X}] \tilde{M}$ .

The matrix above is clearly semi definite positive.  $\square$

## B.3 Feasible most powerful instrument

### B.3.1 Local approximation of the MPI

*Proof of Proposition 6.* First, we define  $s_t^0 = \rho(\delta_t, x_{2t}, f_0(\cdot|\lambda_0))$  with  $\delta_t$  the true mean utility. From lemma 2.4  $\rho^{-1}$  is  $\mathcal{C}^\infty$  and in particular,  $\rho^{-1}$  is  $\mathcal{C}^1$ . Thus, the Taylor expansion of  $\rho^{-1}(s_t^0, x_{2t}, f_0(\cdot|\lambda_0))$  around  $s_t$  writes:

$$\begin{aligned} \rho^{-1}(s_t^0, x_{2t}, f_0(\cdot|\lambda_0)) &= \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0)) + \left. \frac{\partial \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))}{\partial s} \right|_{s=s_t} (s_t^0 - s_t) + o(\|s_t^0 - s_t\|) \\ \delta_t &= \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0)) + \left. \frac{\partial \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))}{\partial s} \right|_{s=s_t} (s_t^0 - s_t) + o(\|s_t^0 - s_t\|) \end{aligned}$$

We now derive an expression for the first derivative of the inverse function. We make use of lemma 2.5: for any  $\delta \in \mathbb{R}^J$ ,  $\frac{\partial \rho(\delta, x_{2t}, f)}{\partial \delta}$  is invertible.

$$\begin{aligned} \frac{\partial \rho(\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0)), x_{2t}, f_0(\cdot|\lambda_0))}{\partial s} = I_J &\iff \frac{\partial \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))}{\partial s} \left( \frac{\partial \rho(\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0)), x_{2t}, f_0(\cdot|\lambda_0))}{\partial \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))} \right) = I_J \\ &\iff \frac{\partial \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))}{\partial s} = \left( \frac{\partial \rho(\delta_t^0, x_{2t}, f_0(\cdot|\lambda_0))}{\partial \delta} \right)^{-1} \end{aligned}$$

with  $\delta_t^0 = \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))$ . Consequently,

$$\underbrace{\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0)) - \delta_t}_{\Delta(s_t, x_{2t}, f_0, f_a)} = - \left( \frac{\partial \rho(\delta_t^0, x_{2t}, f_0(\cdot|\lambda_0))}{\partial \delta} \right)^{-1} (s_t^0 - s_t) + o(\|s_t^0 - s_t\|) \quad (16)$$

with  $\delta_t^0 = \rho_j^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))$

Now let us show that there exists a constant  $M$  such that  $\|s_t^0 - s_t\| \leq M\tau(f_0(\cdot|\lambda_0) - f_a)$ . with  $\tau(f_0 - f_a) = \int_{\mathbb{R}^{K_2}} |f_0(v|\lambda_0) - f_a(v)|dv$ . Norms are equivalent in a finite vectorial space and without loss of generality, we will derive the results with the  $L_1$  norm. By definition:

$$s_t^0 - s_t = \int_{\mathbb{R}^{K_2}} \frac{\exp\{\delta_t + x_{2t}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt} + x'_{2jk}v\}} (f_0(v|\lambda_0) - f_a(v))dv$$

Taking the  $L_1$  norm of this vector:

$$\begin{aligned} \|s_t^0 - s_t\|_1 &= \sum_{j=1}^J \left| \int_{\mathbb{R}^{K_2}} \frac{\exp\{\delta_{jt} + x_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt} + x'_{2jk}v\}} (f_0(v|\lambda_0) - f_a(v))dv \right| \\ &\leq \sum_{j=1}^J \int_{\mathbb{R}^{K_2}} \underbrace{\left| \frac{\exp\{\delta_{jt} + x_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt} + x'_{2jk}v\}} \right|}_{\leq 1} |f_0(v|\lambda_0) - f_a(v)|dv. \\ &\leq J \int_{\mathbb{R}^{K_2}} |f_0(v|\lambda_0) - f_a(v)|dv = J\tau(f_0(\cdot|\lambda_0) - f_a). \end{aligned}$$

This proves the statement. As a consequence, we have:  $\|s_t^0 - s_t\|_1 = O(\tau(f_0(\cdot|\lambda_0) - f_a))$  and  $o(\|s_t^0 - s_t\|) = o(\tau(f_0(\cdot|\lambda_0) - f_a))$ .

The problem with the term  $s_t^0 - s_t$  is that it is an expression of  $\delta_t$  which we do not know under misspecification. As we want to be able to compute this approximation of the error term, it is not convenient in practice to have an expression which depends on  $\delta_t$ . On the other hand, we know  $\delta_t^0$  and thus, the simple idea that we exploit is to take a Taylor expansion of the term above around  $\delta_t^0$ . First, let us remark that from equation 16, we have that:

$$\|\delta_t - \delta_t^0\| = \|\delta_t - \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))\| = O(\|s_t^0 - s_t\|) = O(\tau(f_0(\cdot|\lambda_0) - f_a)).$$

Now let us take the Taylor expansion of  $s_t^0 - s_t$  around  $\delta_t^0$ :

$$\begin{aligned}
s_t^0 - s_t &= \int_{\mathbb{R}^{K_2}} \frac{\exp\{\delta_t^0 + x_{2t}v\}}{1 + \sum_{k=1}^J \exp\{\tilde{\delta}_{kt} + x'_{2jk}v\}} (f_0(v|\lambda_0) - f_a(v)) dv \\
&+ \underbrace{\int_{\mathbb{R}^{K_2}} \frac{\partial}{\partial \delta'} \left\{ \frac{\exp\{\delta_t^0 + x_{2t}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2jk}v\}} \right\} (\delta_t - \delta_t^0) (f_0(v|\lambda_0) - f_a(v)) dv}_{B} + o(\|\delta_t - \delta_t^0\|).
\end{aligned}$$

From what precedes, we know that  $o(\|\delta_t - \delta_t^0\|) = o(\tau(f_0(\cdot|\lambda_0) - f_a))$ . Now, let us show that term  $B$  in the previous expansion is also  $o(\tau(f_0(\cdot|\lambda_0) - f_a))$ . Again taking the  $L_1$  norm:

$$\begin{aligned}
\|B\|_1 &= \sum_{j=1}^J \left| \sum_{l=1}^J \int_{\mathbb{R}^{K_2}} \frac{\partial}{\partial \delta_l} \left\{ \frac{\exp\{\delta_{jt}^0 + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\tilde{\delta}_{kt} + x'_{2jk}v\}} \right\} (\delta_{lt} - \delta_{lt}^0) (f_0(v|\lambda_0) - f_a(v)) dv \right| \\
&\leq \sum_{j=1}^J \sum_{l=1}^J \int_{\mathbb{R}^{K_2}} \underbrace{\left| \frac{\partial}{\partial \delta_l} \left\{ \frac{\exp\{\delta_{jt}^0 + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2jk}v\}} \right\} \right|}_{\leq 1} |\delta_{lt} - \tilde{\delta}_{lt}| |f_0(v|\lambda_0) - f_a(v)| dv \\
&\leq J^2 \tau(f_0(\cdot|\lambda_0) - f) O(\tau(f_0(\cdot|\lambda_0) - f_a)) = O(\tau(f_0(\cdot|\lambda_0) - f_a)^2) = o(\tau(f_0(\cdot|\lambda_0) - f_a)).
\end{aligned}$$

Thus,  $\|B\|_1 = o(\tau(f_0(\cdot|\lambda_0) - f_a))$  and by combining all the results together, we get the final result. When  $f_0(\cdot|\lambda_0)$  gets “close “ to  $f_a$ , we have the following approximation:

$$\begin{aligned}
\Delta(s_t, x_{2t}, f_0, f_a) &= \left( \frac{\partial \rho(\delta_t^0, x_{2t}, f_0(\cdot|\lambda_0))}{\partial \delta} \right)^{-1} \int_{\mathbb{R}^{K_2}} \frac{\exp\{\delta_t^0 + x_{2t}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2jk}v\}} (f_a(v) - f_0(v|\lambda_0)) dv \\
&+ o(\tau(f_a - f_0(\cdot|\lambda_0))),
\end{aligned}$$

with  $\delta_t^0 = \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda_0))$  and  $\tau(f_a - f_0(\cdot|\lambda_0)) = \int_{\mathbb{R}^{K_2}} |f_a(v) - f_0(\cdot|\lambda_0)(v)| dv$ .  $\square$

### B.3.2 Global approximation of the MPI

**Lemma 2.3.** Analytical expression for  $\Delta_j(s_t, x_{2t}, f_0, f_a)$ . Let  $\delta_{jt}^0 = \rho_j^{-1}(s_t, x_{2t}, f_0)$  and  $\delta_{jt}^a = \rho_j^{-1}(s_t, x_{2t}, f_a)$ . We have the following:

$$\Delta_j(s_t, x_{2t}, f_0, f_a) = \log \left( \frac{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^a + x'_{2kt}v\}} f_a(v) dv}{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{jt}^0 + x'_{2kt}v\}} f_0(v) dv} \right).$$

*Proof of Lemma 2.3.*

$$\begin{aligned}
1 = \frac{\rho(\delta_{jt}, x_{2t}, f_a)}{\rho(\delta_{jt}^0, x_{2t}, f_0)} &\iff 1 = \frac{\int_{\mathbb{R}^{K_2}} \frac{\exp\{\delta_{jt} + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^a + x'_{2kt}v\}} f_a(v) dv}{\int_{\mathbb{R}^{K_2}} \frac{\exp\{\delta_{jt}^0 + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v\}} f_0(v) dv} \\
&\iff \frac{\exp\{\delta_{jt}^0\}}{\exp\{\delta_{jt}^a\}} = \frac{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x_{2t}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^a + x'_{2kt}v\}} f_a(v) dv}{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v\}} f_0(v) dv} \\
&\iff \Delta_j(s_t, x_{2t}, f_0, f_a) = \log \left( \frac{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^a + x'_{2kt}v\}} f_a(v) dv}{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}^0 + x'_{2kt}v\}} f_0(v) dv} \right).
\end{aligned}$$

□

### B.3.3 Approximation of the MPI in the mixed logit case

*Proof of Proposition 7.* By definition, we have:

$$\begin{aligned}
g_j(x_i, \cdot, f) : \mathbb{R}^{K_1} &\rightarrow [0, 1] \\
\tilde{\beta} &\mapsto \int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{ij1}\tilde{\beta} + x'_{2ij}v\}}{1 + \sum_{k=1}^J \exp\{x'_{ik1}\tilde{\beta} + x'_{2ik}v\}} f(v) dv
\end{aligned}$$

$g$  is  $\mathcal{C}^\infty$  on  $\mathbb{R}^{K_1}$ . Thus, we can take a first order Taylor expansion of  $g_j(x_i, \cdot, f_a)$  around  $\beta_0$ :

$$g_j(x_i, \beta_a, f_a) = g_j(x_i, \beta_0, f_0) + \left. \frac{\partial g(x_i, \tilde{\beta}, f_0)}{\partial \tilde{\beta}} \right|_{\tilde{\beta}=\beta_0} (\beta_a - \beta_0) + o(\|\beta_a - \beta_0\|)$$

This yields immediately,

$$\begin{aligned}
g(x_i, \beta_0, f_0) - g(x_i, \beta_a, f_a) &= \int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{1ij}\beta_0 + x'_{2ij}v\}}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\beta_0 + x'_{2ik}v\}} (f_0(v) - f_a(v)) dv + \\
&\quad \left. \frac{\partial g(x_i, \tilde{\beta}, f_a)}{\partial \tilde{\beta}} \right|_{\tilde{\beta}=\beta_0} (\beta_a - \beta_0) + o(\|\beta_a - \beta_0\|)
\end{aligned}$$

Finally, we need to show  $\|\beta_a - \beta_0\| = o\left(\int_{\mathbb{R}^{K_2}} |f_0(v) - f_a(v)| dv\right)$ . From Assumption 1 by definition of  $\beta_0^*$  and  $\beta_a^*$ , we have  $f_0 = f_a \Rightarrow \beta_0^* = \beta_a^*$ . Going further it can be shown using the Kullback divergence that for any  $\epsilon_1 > 0$  such that  $\int_{\mathbb{R}^{K_2}} |f_0(v) - f_a(v)| dv < \epsilon_1$  there exists

some  $e_2 > 0$  such that  $||\beta_0^* - \beta_a^*|| < e_2$ . In other words if  $\int_{\mathbb{R}^{K_2}} |f_0(v) - f_a(v)| dv$  is small then  $||\beta_a^* - \beta_0^*||$  is also small, thus any small  $o$  of  $||\beta_a - \beta_0|| = ||\beta_a^* - \beta_0^*||$  can be replaced by a small  $o$  of  $\int_{\mathbb{R}^{K_2}} |f_0(v) - f_a(v)| dv$  even if the two quantities are not proportional. Consequently

$$g(x_i, \beta_0, f_0) - g(x_i, \beta_a, f_a) = \int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{1ij}\beta_0 + x'_{2ij}v\}}{1 + \sum_{k=1}^J \exp\{x'_{1ik}\beta_0 + x'_{2ik}v\}} (f_0(v) - f_a(v)) dv + \left. \frac{\partial g(x_i, \tilde{\beta}, f_a)}{\partial \tilde{\beta}} \right|_{\tilde{\beta}=\beta_0} (\beta_a - \beta_0) + o\left(\int_{\mathbb{R}^{K_2}} |f_0(v) - f_a(v)| dv\right)$$

□

## B.4 Specification Test: composite hypothesis

In this section, we prove theorem 5.1, which is the main asymptotic result of the paper. The section is organized as follows. First, we establish the equivalence between the moment condition around which we build our test  $\mathbb{E} \left[ \sum_{jt} \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) h_D(z_{jt}) \right] = 0$  and the one characterizing  $H'_0 : \mathbb{E} [\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) h_D(z_{jt})] = 0$ . Then, we introduce the notations used in the proofs and we decompose  $\hat{\xi}$  according to the BLP approximations. Second we provide technical lemmas which prove that under the assumptions in E, the BLP approximations vanish asymptotically. Third, we prove that the BLP estimator is consistent and asymptotically normal. Finally, we prove the main theorem and we show that under the null the test is pivotal in the 2 polar cases described in the main text.

### B.4.1 Equivalence between moment conditions

Let  $h_D(z_{jt})$  our detection instruments. For conciseness, we omit the dependence in  $f_0$  and denote  $\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) = \xi_{jt}(\theta_0)$ . We want to prove that the following two moment conditions are equivalent:

$$\mathbb{E} [\xi_{jt}(\theta_0) h_D(z_{jt})] = 0 \iff \mathbb{E} \left[ \sum_{j=1}^J \xi_{jt}(\theta_0) h_D(z_{jt}) \right] = 0$$

Let  $R_t$  a categorical random variable which exogenously selects a product  $j$  with probability  $\frac{1}{J}$ . Formally, we have  $(\xi_{jt}(\theta_0), z_{jt}) \perp\!\!\!\perp R_{jt}$ . By construction, we have:

$$\begin{aligned}
\mathbb{E} [\xi_{jt}(\theta_0) h_D(z_{jt})] &= \sum_{k=1}^J \mathbb{E} [\xi_{kt}(\theta_0) h_D(z_{kt}) R_{kt}] = \sum_{k=1}^J \mathbb{E} [\xi_{kt}(\theta_0) h_D(z_{kt})] \mathbb{E}[R_{kt}] \\
&= \frac{1}{J} \mathbb{E} \left[ \sum_{k=1}^J \xi_{kt}(\theta_0) h_D(z_{kt}) \right]
\end{aligned}$$

Second line results from independence of  $(\xi_{jt}(\theta_0), z_{jt})$  and  $R_{jt}$ . This proves the result.

#### B.4.2 Notations

In the proofs, we will adopt the following notations. If the derivations are done under the parametric assumption  $H_0 : f \in \mathcal{F}_0$  then we omit the dependence in  $f_0$  and interchangeably use  $\xi_{jt}(f_0(\cdot|\lambda), \beta)$  and  $\xi_{jt}(\theta)$ . We also omit the dependence of the BLP pseudo true value in  $W$  and  $h_E(z_{jt})$ <sup>37</sup>. Then define the following objectives of the GMM minimization

$$\begin{aligned}
\hat{\mathcal{Q}}_T(\tilde{\theta}) &= \left( \frac{1}{T} \sum_{j,t} \hat{\xi}_{jt}(\tilde{\theta}) h_E(z_{jt}) \right)' \hat{W} \left( \frac{1}{T} \sum_{j,t} \hat{\xi}_{jt}(\tilde{\theta}) h_E(z_{jt}) \right) \\
\mathcal{Q}_T(\tilde{\theta}) &= \left( \frac{1}{T} \sum_{j,t} \xi_{jt}(\tilde{\theta}) h_E(z_{jt}) \right)' \hat{W} \left( \frac{1}{T} \sum_{j,t} \xi_{jt}(\tilde{\theta}) h_E(z_{jt}) \right) \\
\mathcal{Q}(\tilde{\theta}) &= \mathbb{E} \left[ \sum_j \xi_{jt}(\tilde{\theta}) h_E(z_{jt}) \right]' W \mathbb{E} \left[ \sum_j \xi_{jt}(\tilde{\theta}) h_E(z_{jt}) \right]
\end{aligned}$$

We also define the following moments

$$\begin{aligned}
\hat{g}_T(\tilde{\theta}, h) &= \frac{1}{T} \sum_{j,t} \hat{\xi}_{jt}(\tilde{\theta}) h(z_{jt}) \\
g_T(\tilde{\theta}, h) &= \frac{1}{T} \sum_{j,t} \xi_{jt}(\tilde{\theta}) h(z_{jt}) \\
g(\tilde{\theta}, h) &= \mathbb{E} \left[ \sum_j \xi_{jt}(\tilde{\theta}) h(z_{jt}) \right]
\end{aligned}$$

And recall the definition of  $\Gamma(\mathcal{F}_0, \tilde{\theta}, h)$  which is used interchangeably with  $\Gamma(\tilde{\theta}, h)$

$$\hat{\Gamma}_T(\tilde{\theta}, h) = \frac{1}{T} \sum_{j,t} h(z_{jt}) \frac{\partial}{\partial \theta} \hat{\xi}_{jt}(\tilde{\theta})'$$

---

<sup>37</sup>The BLP pseudo true value depends on  $W$  and  $h_E(z_{jt})$  when the model is misspecified

$$\Gamma_T(\tilde{\theta}, h) = \frac{1}{T} \sum_{j,t} h(z_{jt}) \frac{\partial}{\partial \theta} \xi_{jt}(\tilde{\theta})'$$

$$\Gamma(\tilde{\theta}, h) = \mathbb{E} \left[ \sum_j h(z_{jt}) \frac{\partial}{\partial \theta} \xi_{jt}(\tilde{\theta})' \right]$$

Furthermore, unless specified, all limits are taken with respect to  $T$ ; Additionally, we denote by the expression  $X = o_P(T^\kappa)$  a random variable or statistic which is asymptotically degenerate of order  $T^\kappa$ , ie  $X = o_P(T^\kappa) \Leftrightarrow \forall e > 0 \mathbb{P}(|X|T^{-\kappa} > e) \xrightarrow{T \rightarrow \infty} 0$ , and denote by  $X = O_P(T^\kappa)$  a random variable which is (bounded in probability) of order  $T^\kappa$ , ie  $\forall e_1 > 0 \exists e_2 > 0, \exists T_N : \forall T \geq T_N \mathbb{P}(|X|T^{-\kappa} > e_2) < e_1$ . Properties of  $o_P(1)$  and  $O_P(1)$  random variables are used throughout these proofs.

#### B.4.3 Feasible Structural Error and BLP approximations

We now decompose the difference between the true structural error  $\xi_{jt}(\tilde{\theta})$  and the feasible structural error  $\hat{\xi}_{jt}(\tilde{\theta})$  in terms of the different approximations involved in the derivation of the feasible structural error  $\hat{\xi}_{jt}(\tilde{\theta})$ . In market  $t$  given an assumption  $\mathcal{F}_0$ , a parameter  $\tilde{\lambda}$ , market shares  $s_t$  and product characteristics with preference heterogeneity  $x_{2t}$  there exists a unique  $\delta_t \in \mathbb{R}^J$  such that  $s_t = \rho(\delta_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))$  (Brouwer's fixed point theorem, see [Berry \(1994\)](#)) so that  $\delta_t = \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))$ . There is no closed form for  $\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))$  so the NFP algorithm is used. Denote as  $C$  the contraction used to find the mean utilities which solve the demand equal market share constraint

$$C(\cdot, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) : \delta \in \mathbb{R}^J \mapsto \delta + \log(s_t) - \log(\rho(\delta, x_{2t}, f_0(\cdot|\tilde{\lambda})))$$

So that for some starting mean utility  $\delta_0 \in \mathcal{B} \subset \mathbb{R}^J$  where  $\mathcal{B}$  is bounded, the mean utility obtained via NFP at the limit is equal to the unique vector which solves the constraint

$$\delta_t(f_0(\cdot|\tilde{\lambda})) = \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) = \lim_{H \rightarrow \infty} C^{(H)}(\delta_0, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))$$

Similarly the error generated by  $(f_0(\cdot|\tilde{\lambda}, \tilde{\beta}))$  can be obtained from NFP at the limit

$$\xi_t(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) = \delta_t(f_0(\cdot|\tilde{\lambda})) - x_{1t}\tilde{\beta} = \lim_{H \rightarrow \infty} C^{(H)}(\delta_0, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) - x_{1t}\tilde{\beta}$$



This way we obtain a vector of mean utilities for each market  $t$ . There are 3 approximations to consider, market shares are not truly observed, the demand integral has to be simulated, and the contraction is never taken to its limit, so define  $\widehat{\xi}(f_0, \tilde{\lambda})$ ,  $\widehat{\delta}(f_0, \tilde{\lambda})$  and  $\hat{C}$  for some starting value  $\delta_0$

$$\begin{aligned}\widehat{\xi}_t(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) &= \hat{C}^{(H)}(\delta_0, \hat{s}_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) - x_{1jt}\tilde{\beta}, & \widehat{\delta}(f_0, \tilde{\lambda}) &= \hat{C}^{(H)}(\delta_0, \hat{s}_t, x_{2t}, f_0, \tilde{\lambda}) \\ \hat{C} : \delta &\mapsto \delta + \log(\hat{s}_t) - \log(\hat{\rho}(\delta, x_{2t}, f_0(\cdot|\lambda_0)))\end{aligned}$$

Consequently we decompose the difference between the error generated by  $(f_0(\cdot|\tilde{\lambda}), \tilde{\beta})$  and its feasible approximation into 3 differences

$$\begin{aligned}\xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) - \widehat{\xi}_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) &= \delta_{jt}(f_0(\cdot|\tilde{\lambda})) - \widehat{\delta}_{jt}(f_0(\cdot|\tilde{\lambda})) \\ &= \lim_{H \rightarrow \infty} C_j^{(H)}(\delta_0, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) - \hat{C}_j^{(H)}(\delta_0, \hat{s}_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) \\ &= \lim_{H \rightarrow \infty} C_j^{(H)}(\delta_0, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) - C_j^{(H)}(\delta_0, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) \\ &\quad + C_j^{(H)}(\delta_0, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) - C_j^{(H)}(\delta_0, \hat{s}_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) \\ &\quad + C_j^{(H)}(\delta_0, \hat{s}_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) - \hat{C}_j^{(H)}(\delta_0, \hat{s}_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) \\ &\equiv \rho_j^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})) - D_j(\rho, s_t, \tilde{\lambda}) \\ &\quad + D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, \hat{s}_t, \tilde{\theta}) \\ &\quad + D_j(\rho, \hat{s}_t, \tilde{\theta}) - D_j(\hat{\rho}, \hat{s}_t, \tilde{\theta})\end{aligned}$$

In the fourth line, we simply introduce shortened notations for the same objects.

#### B.4.4 Technical Lemmas

The 1st and 2nd lemma establish the smoothness of  $\rho^{-1}$  and the invertibility of the Jacobian matrix of  $\rho$  with respect to  $\delta$ . In the 3rd lemma, we derive the Lipschitz constant of the contraction and we prove that it is bounded away from 0 and 1. The 4th lemma ensures that for key moments and quantities the BLP approximations can be ignored uniformly asymptotically.

**Lemma 2.4.**  $\rho^{-1}$  is  $\mathcal{C}^\infty$

*Proof.* We know that the demand function  $\rho$  is  $\mathcal{C}^\infty$  and invertible on  $\mathbb{R}^J$ . Moreover,  $\forall \delta \in \mathbb{R}^J$ ,  $\frac{\partial \rho(\delta, x_{2t}, f)}{\partial \delta} \neq 0$ . As a consequence,  $\rho^{-1} : [0, 1]^J \rightarrow \mathbb{R}^J$  the inverse demand function is also  $\mathcal{C}^\infty$ .  $\square$

**Lemma 2.5.** For any  $\delta \in \mathbb{R}^J$ ,  $\frac{\partial \rho(\delta, x_{2t}, f)}{\partial \delta}$  is invertible.

*Proof.*  $\frac{\partial \rho}{\partial \delta}$  is a  $J \times J$  matrix such that  $(\frac{\partial \rho}{\partial \delta})_{j,k}$  is:

$$\frac{\partial \rho_j(\delta_t, x_{2t}, f)}{\partial \delta_{kt}} = \begin{cases} \int \mathcal{T}_{jt}(v) (1 - \mathcal{T}_{kt}(v)) f(v) dv & \text{if } j = k \\ - \int \mathcal{T}_{jt}(v) \mathcal{T}_{kt}(v) f(v) dv & \text{if } j \neq k \end{cases}$$

$$\text{with } \mathcal{T}_{jt}(v) \equiv \frac{\exp\{\delta_{jt} + x'_{2jt}v\}}{1 + \sum_{j'=1}^J \exp\{\delta_{j't} + x'_{2j't}v\}}$$

One can easily check that  $\frac{\partial \rho}{\partial \delta}$  is strictly diagonally dominant. Indeed for each row  $j$ :

$$\left| \frac{\partial \rho_j(\delta_t, x_{2t}, f)}{\partial \delta_{kt}} \right| - \sum_{k \neq j} \left| \frac{\partial \rho_j(\delta_t, x_{2t}, f)}{\partial \delta_{kt}} \right| = \int \mathcal{T}_{jt}(v) \underbrace{\left( 1 - \sum_{k=1}^J \mathcal{T}_{kt}(v) \right)}_{>0} f(v) dv > 0$$

□

**Lemma 2.6** (Contraction Mapping Lipschitz Constant).

Given parametric assumption  $\mathcal{F}_0$ , under assumptions **B-E**, assume that starting mean utility  $\delta_0$  is in  $\mathcal{B}$  where  $\mathcal{B}$  is compact, then without loss of generality there exists some  $(\underline{a}, \bar{a}) \in \mathbb{R}^2$  with  $\bar{a} > \underline{a}$  such that for any  $b \in \mathcal{B}$  for any  $j = 1, \dots, J$   $\underline{a} \leq b_j \leq \bar{a}$ , furthermore denote by  $\mathcal{X}$  the compact support of  $x_{2jt}$ . Then on  $\mathcal{B}$  the map  $C(\cdot, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}_0))$  is a contraction with Lipschitz constant

$$\epsilon = \max_{j=1, \dots, J} \sup_{a \in \mathcal{B}, b \in [0; \bar{a} - \underline{a}]^J, x_2 \in \mathcal{X}, \tilde{\lambda} \in \Lambda_0} 1 - \frac{\int \frac{\exp\{a_j + b_j + x'_{2j}v\}}{(1 + \sum_k \exp\{a_k + b_k + x'_{2k}v\})^2} f_0(v|\tilde{\lambda}) dv}{\int \frac{\exp\{a_j + b_j + x'_{2j}v\}}{1 + \sum_k \exp\{a_k + b_k + x'_{2k}v\}} f_0(v|\tilde{\lambda}) dv}$$

which is in  $(0; 1)$

*Proof.* This proof is inspired by the proof of the Theorem in Appendix 1 of [Berry et al. \(1995\)](#). Let  $C_j(\cdot) \equiv C(\cdot, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}_0))$ , we first determine the partial derivative of  $C_j(\cdot)$

$$\begin{aligned} \frac{\partial C_j(a)}{\partial a_j} &= 1 - \frac{1}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} \int \frac{\exp\{a_j + x'_{2jt}v\} (1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}) - \exp\{2(a_j + x'_{2jt}v)\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv \\ &= \frac{1}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} \int \frac{\exp\{2(a_j + x'_{2jt}v)\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv \\ \frac{\partial C_j(a)}{\partial a_{j'}} &= \frac{1}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} \int \frac{\exp\{a_j + x'_{2jt}v\} \exp\{a_{j'} + x'_{2j't}v\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv \end{aligned}$$

Note that for any  $j = 1, \dots, J$  all partial derivatives of  $C_j(\cdot)$  are strictly positive and that the sum of its derivatives evaluated in  $a$  equals

$$\begin{aligned} \sum_{k=1}^J \frac{\partial C_j(a)}{\partial a_k} &= \frac{1}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} \int \frac{\exp\{a_j + x'_{2jt}v\} \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv \\ &= \frac{1}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} \int \frac{\exp\{a_j + x'_{2jt}v\} (1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\} - 1)}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv \\ &= 1 - \frac{\int \frac{\exp\{a_j + x'_{2jt}v\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} \end{aligned}$$

For any  $(a_1, a_2) \in \mathcal{B}^2$  let  $\tilde{a} = (\|a_1 - a_2\|_\infty, \dots, \|a_1 - a_2\|_\infty) \in \mathbb{R}^J$  then

$$\begin{aligned} C_j(a_1) - C_j(a_2) &= C_j(a_2 + a_1 - a_2) - C_j(a_2) \leq C_j(a_2 + \tilde{a}) - C_j(a_2) \\ &\leq \int_{0^J}^{\|a_1 - a_2\|_\infty^J} \frac{\partial C_j(a_2 + b)}{\partial a} db \\ &\leq \|a_1 - a_2\|_\infty \sup_{a \in \mathcal{B}, b \in [0; \bar{a} - \underline{a}]^J} \sum_{k=1}^J \frac{\partial C_j(a + b)}{\partial a_k} \\ &\leq \|a_1 - a_2\|_2 \max_{j=1, \dots, J} \sup_{a \in \mathcal{B}, b \in [0; \bar{a} - \underline{a}]^J, x_2 \in \mathcal{X}, \tilde{\lambda} \in \Lambda_0} \sum_{k=1}^J \frac{\partial C_j(a + b)}{\partial a_k} \\ &\equiv \|a_1 - a_2\|_2 \epsilon \end{aligned}$$

where the 1st inequality holds because  $C_j(\cdot)$  is increasing in all its inputs, the 2nd inequality holds by the fundamental theorem of calculus and by the total derivative formula, the 3rd and 4th inequalities hold by properties of norms.

We now prove that  $\sup_{a \in \mathcal{B}, b \in [0; \bar{a} - \underline{a}]^J, \tilde{\lambda} \in \Lambda_0} \sum_{k=1}^J \frac{\partial C_j(a+b)}{\partial a_k} \in (0; 1)$  which will imply that  $\epsilon \in (0; 1)$ .

To do so we have to prove that  $\sum_{k=1}^J \frac{\partial C_j(a, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))}{\partial a_k}$  is continuous in  $(a, x_{2t}, \tilde{\lambda})$  and takes values in  $(0; 1)$  almost surely, this way because  $\mathcal{B}$ ,  $\mathcal{X}$  and  $\Lambda_0$  are compact by Weierstrass' Extreme Value Theorem the sum of partial derivatives will also take values in a compact which is inside  $(0; 1)$ , then the supremum will become a maximum which can be attained and which is inside  $(0; 1)$ . The sum of partial derivatives is almost surely in  $(0; 1)$  because

$$\int \frac{\exp\{a_j + x'_{2jt}v\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv - \rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))$$

$$\begin{aligned}
&= \int \frac{\exp\{a_j + x'_{2jt}v\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv - \int \frac{\exp\{a_j + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}} f_0(v|\tilde{\lambda}) dv \\
&= - \int \frac{\exp\{a_j + x'_{2jt}v\} \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv < 0 \\
&\Rightarrow \frac{\int \frac{\exp\{a_j + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}} f_0(v|\tilde{\lambda}) dv}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} < 1 \\
&\Rightarrow \sum_{k=1}^J \frac{\partial C_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))}{\partial a_k} = 1 - \frac{\int \frac{\exp\{a_j + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}} f_0(v|\tilde{\lambda}) dv}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} > 0 \\
&\Rightarrow - \frac{\int \frac{\exp\{a_j + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}} f_0(v|\tilde{\lambda}) dv}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} < 0 \\
&\Rightarrow \sum_{k=1}^J \frac{\partial C_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))}{\partial a_k} = 1 - \frac{\int \frac{\exp\{a_j + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}} f_0(v|\tilde{\lambda}) dv}{\rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))} < 1
\end{aligned}$$

Continuity of the sum of the partial derivatives in  $(a, x_{2t})$  is trivial, continuity in  $\tilde{\lambda}$  also holds because  $f_0(\cdot|\tilde{\lambda})$  must be continuously differentiable via Assumption D.  $\forall e_1 > 0, \exists e_2 : \forall(\lambda_1, \lambda_2) : \|\lambda_1 - \lambda_2\|_2 \leq e_2$  implies  $|f_0(v|\lambda_1) - f_0(v|\lambda_2)| < e_1$  for all  $v$  which in turn implies

$$\begin{aligned}
\forall x_2 \in \mathcal{X}, \forall a \in \mathcal{B} \quad &\left| \int \frac{\exp\{a_j + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}} (f_0(v|\lambda_1) - f_0(v|\lambda_2)) dv \right| \\
&\leq \int \frac{\exp\{a_j + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\}} |f_0(v|\lambda_1) - f_0(v|\lambda_2)| dv \\
&\leq e_1
\end{aligned}$$

Thus both  $\tilde{\lambda} \mapsto \rho_j(a, x_{2t}, f_0(\cdot|\tilde{\lambda}))$  and  $\tilde{\lambda} \mapsto \int \frac{\exp\{a_j + x'_{2jt}v\}}{(1 + \sum_{k=1}^J \exp\{a_k + x'_{2kt}v\})^2} f_0(v|\tilde{\lambda}) dv$  are continuous and so is their ratio.  $\square$

**Lemma 2.7** (Uniform Convergence of Objective Function wrt BLP Approximations).

Given parametric assumption  $\mathcal{F}_0$ , under assumptions B-E and  $\forall h$  which satisfies D

$$\begin{aligned}
&\sup_{\tilde{\theta} \in \Theta_0} \sqrt{T} \|\hat{g}_T(\tilde{\theta}, h) - g_T(\tilde{\theta}, h)\|_2 \xrightarrow{P} 0 \\
&\sup_{\tilde{\theta} \in \Theta_0} \|\hat{\Gamma}_T(\tilde{\theta}, h) - \Gamma_T(\tilde{\theta}, h)\|_2 \xrightarrow{P} 0 \\
&\sup_{\tilde{\theta} \in \Theta_0} |\hat{\mathcal{Q}}_T(\tilde{\theta}) - \mathcal{Q}(\tilde{\theta})| \xrightarrow{P} 0
\end{aligned}$$

*Proof.* Parts of this proof are inspired from Freyberger (2015). We prove the 3 statements of the Lemma in order

1. Using the properties of the  $\sup$ , the fact that  $\forall(A, B)$  rv,  $\forall e > 0, \forall \alpha \in (0, 1), \mathbb{P}(A + B > e) \leq \mathbb{P}(A > \alpha e) + \mathbb{P}(B > (1 - \alpha)e)$  and the previous decomposition of the difference between  $\xi$  and  $\hat{\xi}$  we can find an upper bound on the probability that the difference between  $\hat{g}_T(\cdot)$  and  $g_T(\cdot)$  is above a deviation: For any  $e_1 > 0$

$$\begin{aligned} \mathbb{P}(\sup_{\hat{\theta}} \sqrt{T} \|\hat{g}_T(\theta, h) - g_T(\theta, h)\|_2 > e_1) &= \mathbb{P}(\sup_{\hat{\theta}} \sqrt{T} \frac{1}{T} \left\| \sum_{j,t} (\hat{\xi}_t(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) - \xi_t(f_0(\cdot|\tilde{\lambda}), \tilde{\beta})) h(z_{jt}) \right\|_2 > e_1) \\ &\leq \mathbb{P}(\sup_{\tilde{\lambda}} \sqrt{T} \left\| \frac{1}{T} \sum_{j,t} (\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}_0)) - D_j(\rho, s_t, \tilde{\lambda})) h(z_{jt}) \right\|_2 > \frac{e_1}{3}) \\ &\quad + \mathbb{P}(\sup_{\tilde{\lambda}} \sqrt{T} \left\| \frac{1}{T} \sum_{j,t} (D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, \hat{s}_t, \tilde{\lambda})) h(z_{jt}) \right\|_2 > \frac{e_1}{3}) \\ &\quad + \mathbb{P}(\sup_{\tilde{\lambda}} \sqrt{T} \left\| \frac{1}{T} \sum_{j,t} (D_j(\rho, \hat{s}_t, \tilde{\lambda}) - D_j(\hat{\rho}, \hat{s}_t, \tilde{\lambda})) h(z_{jt}) \right\|_2 > \frac{e_1}{3}) \end{aligned}$$

Then we can prove that each element of the upper bound converges to 0

- (a) By properties of contractions and using Lemma 2.6 we have

$$|\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}_0)) - D_j(\rho, s_t, \tilde{\lambda})| \leq \epsilon^H |\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}_0)) - \delta_0| \leq \epsilon^H \kappa$$

for some constant  $\kappa$  which exists due to the compactness of  $\Lambda_0$ ,  $\mathcal{X}$  and  $\mathcal{B}$ . Thus using the iid nature of the data ??(i), the speed of the NFP algorithm Assumption E(iii), the triangle inequality, Markov inequality and Cauchy-Schwarz inequality the 1st element converges to 0

$$\begin{aligned} &\mathbb{P}(\sup_{\tilde{\lambda}} \sqrt{T} \left\| \frac{1}{T} \sum_{j,t} (\rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}_0)) - D_j(\rho, s_t, \tilde{\lambda})) h(z_{jt}) \right\|_2 > \frac{e_1}{3}) \\ &\leq \mathbb{P}(\sqrt{T} \epsilon^H \kappa \left\| \frac{1}{T} \sum_{j,t} h(z_{jt}) \right\|_2 > \frac{e_1}{3}) \leq \mathbb{P}(\sqrt{T} \epsilon^H \frac{1}{T} \sum_{j,t} \|h(z_{jt})\|_2 > \frac{e_1}{3}) \\ &\leq \frac{3\kappa}{e_1} \sqrt{T} \epsilon^H \sum_j \sqrt{\mathbb{E}(\|h(z_{jt})\|_2^2)} \xrightarrow{T \rightarrow \infty} 0 \end{aligned}$$

- (b) Note that  $D_j$  is continuously differentiable in  $s \in (0; 1)$  so that it is uniformly continuous in  $s$ . Indeed  $C$  is  $\mathcal{C}^\infty$  in  $s$  so that

$$\frac{\partial D(\rho, s_t, \tilde{\lambda})}{\partial s} = \prod_{h=1}^H \frac{\partial C(C^{(h-1)}(\delta_0, s_t, x_{2t}, f_0(\cdot|\tilde{\lambda})), s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))}{\partial s}$$

Next because  $\Lambda_0$  is compact it can be covered by some finite union of closed balls in  $\mathbb{R}^{K_2}$ , ie  $\Lambda_0 \subset \cup_{c=1}^N \Lambda_{0,c}^N$  with  $\forall c = 1, \dots, N \quad \Lambda_{0,c}^N = \{\tilde{\lambda} : \|\tilde{\lambda} - \lambda_c\|_2 \leq r_N\}, \lambda_c \in \Lambda_0$  and

$r_N \xrightarrow{N \rightarrow \infty} 0$ . Consequently

$$\begin{aligned}
& \mathbb{P}(\sup_{\tilde{\lambda}} \frac{1}{\sqrt{T}} \|\sum_{j,t} (D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, \hat{s}_t, \tilde{\lambda})) h_E(z_{jt})\|_2 > \frac{e_1}{3}) \\
& \leq \mathbb{P}(\max_{c=1, \dots, N} \sup_{\tilde{\lambda} \in \Lambda_{0,c}^N} \frac{1}{\sqrt{T}} \|\sum_{j,t} (D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, \hat{s}_t, \tilde{\lambda})) h_E(z_{jt})\|_2 > \frac{e_1}{3}) \\
& \leq \sum_{c=1}^N \mathbb{P}(\sup_{\tilde{\lambda} \in \Lambda_{0,c}^N} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, \hat{s}_t, \tilde{\lambda})| \|h_E(z_{jt})\|_2 > \frac{e_1}{3}) \\
& \leq \sum_{c=1}^N \mathbb{P}(\frac{1}{\sqrt{T}} \|\sum_{j,t} (D_j(\rho, s_t, \lambda_c) - D_j(\rho, \hat{s}_t, \lambda_c)) h_E(z_{jt})\|_2 > \frac{e_1}{9}) \\
& \quad + \sum_{c=1}^N \mathbb{P}(\sup_{\tilde{\lambda} \in \Lambda_{0,c}^N} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, s_t, \lambda_c)| \|h_E(z_{jt})\|_2 > \frac{e_1}{9}) \\
& \quad + \sum_{c=1}^N \mathbb{P}(\sup_{\tilde{\lambda} \in \Lambda_{0,c}^N} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, \hat{s}_t, \lambda_c) - D_j(\rho, \hat{s}_t, \tilde{\lambda})| \|h_E(z_{jt})\|_2 > \frac{e_1}{9})
\end{aligned}$$

where the last inequality was obtained using the triangle inequality. Then by uniform continuity of  $D_j$  in  $s$  it follows that  $\exists e_2 > 0$  such that  $\forall c \quad \frac{1}{\sqrt{T}} \|\sum_{j,t} (D_j(\rho, s_t, \lambda_c) - D_j(\rho, \hat{s}_t, \lambda_c)) h_E(z_{jt})\|_2 > \frac{e_1}{9}$  implies  $\frac{1}{\sqrt{T}} \|\sum_{j,t} (s_t - \hat{s}_t)\|_2 > e_2$  thence letting  $\mathbb{P}^* = \mathbb{P}(\cdot | n_t, x_t, \xi_t)$

$$\begin{aligned}
& \mathbb{P}^*(\frac{1}{\sqrt{T}} \|\sum_{j,t} (D_j(\rho, s_t, \lambda_c) - D_j(\rho, \hat{s}_t, \lambda_c)) h_E(z_{jt})\|_2 > \frac{e_1}{9}) \leq \mathbb{P}^*(\frac{1}{\sqrt{T}} \|\sum_{j,t} (s_t - \hat{s}_t)\|_2 > e_2) \\
& \leq \frac{J \sum_t \mathbb{E}^*(\|s_t - \hat{s}_t\|_2)}{e_2 \sqrt{T}} = \frac{J \sum_t \mathbb{E}^*(\sqrt{\sum_j (s_{jt} - \hat{s}_{jt})^2})}{e_2 \sqrt{T}} \leq \frac{J \sum_t \sqrt{\sum_j \mathbb{E}^*((s_{jt} - \hat{s}_{jt})^2)}}{e_2 \sqrt{T}} \\
& \leq \frac{J \sum_t \sqrt{\sum_j \mathbb{E}^*((\frac{1}{n_t} \sum_{i=1}^{n_t} y_{ijt} - \mathbb{E}^*(y_{ijt}))^2)}}{e_2 \sqrt{T}} = \frac{J \sum_t \sqrt{\sum_j \text{Var}^*(\frac{1}{n_t} \sum_{i=1}^{n_t} y_{ijt})}}{e_2 \sqrt{T}} \\
& \leq \frac{J \sum_t \sqrt{\sum_j \frac{1}{n_t} \text{Var}^*(y_{ijt})}}{e_2 \sqrt{T}} \leq \frac{J^{3/2}}{e_2} \frac{1}{\sqrt{T}} \sum_t \frac{1}{\sqrt{n_t}}
\end{aligned}$$

where Markov inequality, Jensen inequality, the fact that  $y_{ijt} \in \{0, 1\}$ , that  $\varepsilon_{ijt}$  is iid extreme-value type 1 distributed across  $i, j$  and  $t$ , and the fact that  $n_t$  is iid and independent of all other variables have been used. Then taking the expectations and summing over  $N$  on both sides implies by Assumption **E**(i)

$$\sum_{c=1}^N \mathbb{P}(\frac{1}{\sqrt{T}} \|\sum_{j,t} (D_j(\rho, s_t, \lambda_c) - D_j(\rho, \hat{s}_t, \lambda_c)) h_E(z_{jt})\|_2 > \frac{e_1}{9}) \leq \frac{J^{3/2} N}{e_2} \sqrt{T} \mathbb{E}(n_t^{-1/2}) \xrightarrow{T \rightarrow \infty} 0$$

Next using continuity of  $D_j$  in  $\tilde{\lambda}$  it must be that for any  $e_1 > 0$  there exists some  $N$  such that  $\forall \tilde{\lambda} \in \Lambda_{0,c}^N$  such that  $\|\tilde{\lambda} - \lambda_c\|_2 \leq r_N$  implies

$$\frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, s_t, \lambda_c)| \|h_E(z_{jt})\|_2 \leq \frac{e_1}{9}$$

because  $r_N \xrightarrow{N \rightarrow \infty} 0$ . By definition of the supremum it also implies that

$$\sup_{\tilde{\lambda} \in \Lambda_{0,c}} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, s_t, \lambda_c)| \|h_E(z_{jt})\|_2 \leq \frac{e_1}{9}$$

The contraposition is that

$$\sup_{\tilde{\lambda} \in \Lambda_{0,c}} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, s_t, \lambda_c)| \|h_E(z_{jt})\|_2 > \frac{e_1}{9}$$

implies  $\forall \tilde{\lambda} \in \Lambda_{0,c}^N \quad \|\tilde{\lambda} - \lambda_c\|_2 > r_N$  which is impossible by definition of  $\Lambda_{0,c}^N$ . Consequently

$$\begin{aligned} & \sum_{c=1}^N \mathbb{P} \left( \sup_{\tilde{\lambda} \in \Lambda_{0,c}} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, s_t, \tilde{\lambda}) - D_j(\rho, s_t, \lambda_c)| \|h_E(z_{jt})\|_2 > \frac{e_1}{9} \right) \\ & \leq \sum_{c=1}^N \mathbb{P}(\cap_{\tilde{\lambda} \in \Lambda_{0,c}^N} \|\tilde{\lambda} - \lambda_c\|_2 > r_N) = 0 \end{aligned}$$

Similarly

$$\sum_{c=1}^N \mathbb{P} \left( \sup_{\tilde{\lambda} \in \Lambda_{0,c}} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, \hat{s}_t, \tilde{\lambda}) - D_j(\rho, \hat{s}_t, \lambda_c)| \|h_E(z_{jt})\|_2 > \frac{e_1}{9} \right) = 0$$

(c) With the same arguments as in (b)

$$\begin{aligned} & \mathbb{P} \left( \sup_{\tilde{\lambda}} \frac{1}{\sqrt{T}} \left\| \sum_{j,t} (D_j(\rho, \hat{s}_t, \tilde{\lambda}) - D_j(\hat{\rho}, \hat{s}_t, \tilde{\lambda})) h_E(z_{jt}) \right\|_2 > \frac{e_1}{3} \right) \\ & \leq \sum_{c=1}^N \mathbb{P} \left( \frac{1}{\sqrt{T}} \left\| \sum_{j,t} (D_j(\rho, \hat{s}_t, \lambda_c) - D_j(\hat{\rho}, \hat{s}_t, \lambda_c)) h_E(z_{jt}) \right\|_2 > \frac{e_1}{9} \right) \\ & \quad + \sum_{c=1}^N \mathbb{P} \left( \sup_{\tilde{\lambda} \in \Lambda_{0,c}^N} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, \hat{s}_t, \tilde{\lambda}) - D_j(\rho, \hat{s}_t, \lambda_c)| \|h_E(z_{jt})\|_2 > \frac{e_1}{9} \right) \\ & \quad + \sum_{c=1}^N \mathbb{P} \left( \sup_{\tilde{\lambda} \in \Lambda_{0,c}^N} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\hat{\rho}, \hat{s}_t, \lambda_c) - D_j(\hat{\rho}, \hat{s}_t, \tilde{\lambda})| \|h_E(z_{jt})\|_2 > \frac{e_1}{9} \right) \\ & = \sum_{c=1}^N \mathbb{P} \left( \frac{1}{\sqrt{T}} \left\| \sum_{j,t} (D_j(\rho, \hat{s}_t, \lambda_c) - D_j(\hat{\rho}, \hat{s}_t, \lambda_c)) h_E(z_{jt}) \right\|_2 > \frac{e_1}{9} \right) \end{aligned}$$

where  $D_j(\rho, s_t, \lambda_c) = C^{(H)}(\delta_0, s_t, x_{2t}, f_0(\cdot|\lambda_c))$ .  $D_j$  is  $\mathcal{C}^\infty$  in  $\rho \in (0; 1)$ , moreover  $\rho_j(\delta_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))$  and  $\hat{\rho}_j(\delta_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))$  are continuously differentiable in  $\Lambda_0$ . Therefore there exists some  $e_2 > 0$  such that

$$\frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, \hat{s}_t, \lambda_c) - D_j(\hat{\rho}, \hat{s}_t, \lambda_c)| \|h_E(z_{jt})\|_2 > \frac{e_1}{9}$$

implies  $\sup_{a \in \mathcal{B}} \frac{1}{\sqrt{T}} \sum_{j,t} \|\rho(a, x_{2t}, f_0(\cdot|\lambda_c)) - \hat{\rho}(a, x_{2t}, f_0(\cdot|\lambda_c))\|_2 > e_2$ , and as  $\mathcal{B}$  is compact we can cover it by  $\tilde{N}$  closed balls  $\mathcal{B}_b^{\tilde{N}} = \{a \in \mathcal{B} : \|a - a_b\| \leq r_{\tilde{N}}\}$  with  $a_b \in \mathcal{B}$  for any  $b = 1, \dots, \tilde{N}$  so that

$$\begin{aligned} & \sum_{c=1}^N \mathbb{P}(\frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, \hat{s}_t, \lambda_c) - D_j(\hat{\rho}, \hat{s}_t, \lambda_c)| \|h_E(z_{jt})\|_2 > \frac{e_1}{9}) \\ & \leq \sum_{c=1}^N \mathbb{P}(\sup_{a \in \mathcal{B}} \frac{1}{\sqrt{T}} \sum_{j,t} \|\rho(a, x_{2t}, f_0(\cdot|\lambda_c)) - \hat{\rho}(a, x_{2t}, f_0(\cdot|\lambda_c))\|_2 > e_2) \\ & \leq \sum_{c,b} \mathbb{P}(\sup_{a \in \mathcal{B}_b^{\tilde{N}}} \frac{1}{\sqrt{T}} \sum_{j,t} \|\rho(a, x_{2t}, f_0(\cdot|\lambda_c)) - \hat{\rho}(a, x_{2t}, f_0(\cdot|\lambda_c))\|_2 > e_2) \\ & = \sum_{c,b} \mathbb{P}(\frac{1}{\sqrt{T}} \sum_{j,t} \|\rho(a_b, x_{2t}, f_0(\cdot|\lambda_c)) - \hat{\rho}(a_b, x_{2t}, f_0(\cdot|\lambda_c))\|_2 > e_2) \end{aligned}$$

where the last equality was obtained reusing arguments from (b). As a consequence let  $F_{jt}(v) = \frac{\exp\{a_{bj} + x'_{2jt}v\}}{1 + \sum_k \exp\{a_{bk} + x'_{2kt}v\}}$  and  $\mathbb{P}^*(\cdot) = \mathbb{P}(\cdot|x_t, \xi_t)$  then using Markov inequality and Cauchy-Schwarz inequality

$$\begin{aligned} & \mathbb{P}^*(\frac{1}{\sqrt{T}} \sum_{j,t} \|\rho(a_b, x_{2t}, f_0(\cdot|\tilde{\lambda})) - \hat{\rho}(a_b, x_{2t}, f_0(\cdot|\tilde{\lambda}))\|_2 > e_2) \\ & \leq \frac{J \sum_t \mathbb{E}^*(\|\hat{\rho}(a_b, x_{2t}, f_0(\cdot|\tilde{\lambda})) - \rho(a_b, x_{2t}, f_0(\cdot|\tilde{\lambda}))\|_2)}{e_2 \sqrt{T}} \\ & \leq \frac{J \sum_t \sqrt{\sum_j \mathbb{E}^* \left( \left( \frac{1}{R} \sum_{r=1}^R F_{jt}(v_r) - \mathbb{E}^*(F_{jt}(v_r)) \right)^2 \right)}}{e_2 \sqrt{T}} = \frac{J \sum_t \sqrt{\sum_j \text{Var}^* \left( \frac{1}{R} \sum_{r=1}^R F_{jt}(v_r) \right)}}{e_2 \sqrt{T}} \\ & \leq \frac{J^{3/2}}{e_2} \sqrt{\frac{T}{R}} \end{aligned}$$

where the fact that  $v_r$  are iid draws from  $f_0(\cdot|\tilde{\lambda})$  independent from all other variables has been used. It follows by taking the expectation and summing over  $N$  and  $\tilde{N}$  that

$$\mathbb{P}(\sup_{\tilde{\lambda}} \frac{1}{\sqrt{T}} \sum_{j,t} |D_j(\rho, \hat{s}_t, \tilde{\lambda}) - D_j(\hat{\rho}, \hat{s}_t, \tilde{\lambda})| \|h_E(z_{jt})\|_2 \xrightarrow{T \rightarrow \infty} 0)$$

by Assumption **E(i)**.



2. The 2nd statement is not formally proven as it largely builds on the proof of the 1st statement. To see why recall that

$$\hat{\Gamma}_T(\tilde{\theta}, h) - \Gamma_T(\tilde{\theta}, h) = \frac{1}{T} \sum_{jt} h(z_{jt}) \frac{\partial}{\partial \theta} (\hat{\xi}(\tilde{\theta}) - \xi_{jt}(\tilde{\theta}))'$$

More precisely let  $e'_j = (0 \dots 0 \underbrace{1}_{\text{j-th coordinate}} 0 \dots 0)$  then

$$\frac{\partial \xi_{jt}(\tilde{\theta})}{\partial \beta} = -x_{1jt}, \quad \frac{\partial}{\partial \lambda} \xi_{jt}(\tilde{\theta}) = -e'_j \left( \frac{\partial \rho(\delta_t(\tilde{\lambda}), x_{2t}, f_0(\cdot|\tilde{\lambda}))}{\partial \delta} \right)^{-1} \int \frac{\exp\{\delta_{jt}(\tilde{\lambda}) + x'_{2jt}v\}}{1 + \sum_{k=1}^J \exp\{\delta_{kt}(\tilde{\lambda}) + x'_{2kt}v\}} \frac{\partial}{\partial \lambda} f_0(v|\tilde{\lambda}) dv$$

Thus the columns of the matrix  $\hat{\Gamma}_T(\tilde{\theta}, h) - \Gamma_T(\tilde{\theta}, h)$  associated to the derivative in  $\beta$  are equal to 0. Furthermore using an uniform continuity argument  $\left| \frac{\partial \xi_{jt}(\tilde{\theta})}{\partial \lambda} - \frac{\partial \xi_{jt}(\tilde{\theta})}{\partial \lambda} \right| > e_1$  is implied by  $\|\hat{\delta}_t(\tilde{\lambda}) - \delta_t(\tilde{\lambda})\|_2 > e_2$  for some  $e_2 > 0$ . Using the compactness of  $\Lambda_0$  and Assumption **E** it is straightforward that  $\sup_{\tilde{\lambda}} \|\hat{\Gamma}_T(\tilde{\theta}, h) - \Gamma_T(\tilde{\theta}, h)\|_2 \xrightarrow{P} 0$  for any  $h$  which satisfies the conditions in Assumption **D**.

3. The 3rd statement follows from the 1st. Indeed using Cauchy-Schwarz and properties of the supremum

$$\begin{aligned} \sup_{\tilde{\theta} \in \Theta_0} |\hat{\mathcal{Q}}_T(\tilde{\theta}) - \mathcal{Q}_T(\tilde{\theta})| &= |(\hat{g}_T(\tilde{\theta}, h_E) - g_T(\tilde{\theta}, h_E))' \hat{W} (\hat{g}_T(\tilde{\theta}, h_E) - g_T(\tilde{\theta}, h_E)) \\ &\quad - 2(\hat{g}_T(\tilde{\theta}, h_E) - g_T(\tilde{\theta}, h_E))' \hat{W} g_T(\tilde{\theta}, h_E)| \\ &\leq \sup_{\tilde{\theta} \in \Theta_0} \|(\hat{g}_T(\tilde{\theta}, h_E) - g_T(\tilde{\theta}, h_E))\|_2^2 \bar{\mu}(\hat{W}) \\ &\quad + 2 \sup_{\tilde{\theta} \in \Theta_0} \|(\hat{g}_T(\tilde{\theta}, h_E) - g_T(\tilde{\theta}, h_E))\|_2 \sup_{\tilde{\theta} \in \Theta_0} \|g_T(\tilde{\theta}, h_E)\|_2 \bar{\mu}(\hat{W}) \end{aligned}$$

where  $\bar{\mu}(\cdot)$  maps a square matrix towards its maximum eigenvalue. By **D(iv)** and definition of the  $L_2$  matrix norm,  $\bar{\mu}(\hat{W}) \xrightarrow{P} \bar{\mu}(W)$ . Then we apply Jennrich's ULLN: the data is iid,  $\Theta_0$  is compact, and  $g_T(\tilde{\theta}, h_E) = \sum_j \xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) h_E(z_{jt})$  has an envelope with finite absolute 1st moment because  $\xi_{jt}(f_0(\cdot|\tilde{\lambda}), \tilde{\beta}) = \rho^{-1}(s_t, x_{2t}, \tilde{\lambda}) - x'_{1jt} \tilde{\beta}$  and  $\rho^{-1}(\cdot)$  has a maximum because it is continuous and its input are in a compact and because  $\tilde{\beta}$  is in a compact and  $x_{1jt}$  has finite 4th moments, see Assumption **B**; Thus by the CMT  $\sup_{\tilde{\theta} \in \Theta_0} \|g_T(\tilde{\theta}, h_E)\|_2 \xrightarrow{P} \sup_{\tilde{\theta} \in \Theta_0} \|g(\tilde{\theta}, h_E)\|_2$ ; Finally using the 1st statement we have  $\|(\hat{g}_T(\tilde{\theta}, h_E) - g_T(\tilde{\theta}, h_E))\|_2 \xrightarrow{P} 0$  therefore by the CMT

$$\sup_{\tilde{\theta} \in \Theta_0} |\hat{\mathcal{Q}}_T(\tilde{\theta}) - \mathcal{Q}_T(\tilde{\theta})| \xrightarrow{P} 0$$

□

#### B.4.5 Asymptotic Properties of the BLP estimator

**Lemma 2.8** (Consistency of BLP Estimator).

Given parametric assumption  $\mathcal{F}_0$  and under assumptions B-E,

$$\hat{\theta} \xrightarrow{P} \theta_0$$

*Proof.* We prove consistency using arguments for the consistency of M-estimators. For any  $e_1 > 0$  such that  $|\hat{\theta} - \theta_0| > e_1$  then by Assumption D(iii) there exists some  $e_2 > 0$  such that  $Q(\hat{\theta}) - Q(\theta_0) > e_2$  as  $\theta_0$  is the unique minimizer of the objective. Thence for any  $e_1 > 0$ ,  $\exists e_2 > 0$  such that

$$\begin{aligned} \mathbb{P}(|\hat{\theta} - \theta_0| > e_1) &\leq \mathbb{P}(Q(\hat{\theta}) - Q(\theta_0) > e_2) \\ &= \mathbb{P}(\hat{Q}_T(\theta_0) - Q(\theta_0) + Q(\hat{\theta}) - \hat{Q}_T(\hat{\theta}) + \hat{Q}_T(\hat{\theta}) - \hat{Q}_T(\theta_0) > e_2) \\ &\leq \mathbb{P}(\hat{Q}_T(\theta_0) - Q(\theta_0) + Q(\hat{\theta}) - \hat{Q}_T(\hat{\theta}) > e_2) \\ &\leq \mathbb{P}(\hat{Q}_T(\theta_0) - Q(\theta_0) > (1 - \alpha)e_2) + \mathbb{P}(Q(\hat{\theta}) - \hat{Q}_T(\hat{\theta}) > \alpha e_2) \end{aligned}$$

where  $\alpha \in (0; 1)$ , the 2nd inequality comes from the fact that  $\hat{Q}_T(\hat{\theta}) - \hat{Q}_T(\theta_0)$  is almost surely negative by definition of  $\hat{\theta}$ , and the 3rd inequality is obtained by utilizing properties of indicator functions. Then by a direct implication of Lemma 2.7 the right-hand-side converges to 0.  $\square$

**Lemma 2.9** (Asymptotic Normality of BLP Estimator).

Given parametric assumption  $\mathcal{F}_0$ , under assumptions B-E and under  $H_0 : f \in \mathcal{F}_0$

$$\sqrt{T}(\hat{\theta} - \theta_0) = (\Gamma'(\theta_0, h_E)W\Gamma(\theta_0, h_E))^{-1} \sqrt{T}\Gamma'(\theta_0, h_E)Wg_T(\theta_0, h_E) + o_P(1)$$

Furthermore under  $H_0; f \in \mathcal{F}_0$

$$\begin{aligned} \sqrt{T}(\hat{\theta} - \theta_0) &\xrightarrow{d} \mathcal{N}(0, (\Gamma'(\theta_0, h_E)W\Gamma(\theta_0, h_E))^{-1} \Gamma'(\theta_0, h_E)W\Omega(\mathcal{F}_0, h_E)W\Gamma(\theta_0, h_E) \\ &\quad (\Gamma'(\theta_0, h_E)W\Gamma(\theta_0, h_E))^{-1}) \end{aligned}$$

*Proof.* We prove asymptotic normality using arguments from M-estimators asymptotics. From Taylor's theorem there exists some  $\tilde{\theta}$  such that  $\|\tilde{\theta} - \theta_0\|_2 \leq \|\hat{\theta} - \theta_0\|_2$  and

$$\hat{g}_T(\hat{\theta}, h_E) = \hat{g}_T(\theta_0, h_E) + \hat{\Gamma}_T(\tilde{\theta}, h_E)(\hat{\theta} - \theta_0)$$

$$\begin{aligned} \Rightarrow \sqrt{T}\hat{\Gamma}'_T(\hat{\theta}, h_E)\hat{W}\hat{g}_T(\hat{\theta}, h_E) &= \sqrt{T}\hat{\Gamma}'_T(\hat{\theta}, h_E)\hat{W}\hat{g}_T(\theta_0, h_E) + \hat{\Gamma}'_T(\hat{\theta}, h_E)\hat{W}\hat{\Gamma}_T(\tilde{\theta}, h_E)\sqrt{T}(\hat{\theta} - \theta_0) = 0 \\ \Leftrightarrow \sqrt{T}(\hat{\theta} - \theta_0) &= - \left( \hat{\Gamma}'_T(\hat{\theta}, h_E)\hat{W}\hat{\Gamma}_T(\tilde{\theta}, h_E) \right)^{-1} \sqrt{T}\hat{\Gamma}'_T(\hat{\theta}, h_E)\hat{W}\hat{g}_T(\theta_0, h_E) \end{aligned}$$

where the 1st implication is due to the FOC Assumption **D(v)**. Then, we apply the CMT to  $(A, B) \mapsto (A'BA)^{-1}A'B$  which is a continuous mapping if  $A$  and  $B$  are full rank so that when taking  $A = \hat{\Gamma}_T(\hat{\theta}, h_E)$  and  $B = \hat{W}$  we obtain:

$$\sqrt{T}(\hat{\theta} - \theta_0) = - (\Gamma'(\theta_0, h_E)W\Gamma(\theta_0, h_E))^{-1} \sqrt{T}\Gamma'(\theta_0, h_E)Wg_T(\theta_0, h_E) + o_P(1)$$

To prove that  $\text{plim } \hat{\Gamma}_T(\hat{\theta}, h_E) = \text{plim } \hat{\Gamma}_T(\tilde{\theta}, h_E) = \Gamma(\theta_0, h_E)$  we make the following decomposition

$$\hat{\Gamma}_T(\hat{\theta}, h_E) - \Gamma(\theta_0, h_E) = \hat{\Gamma}_T(\hat{\theta}, h_E) - \Gamma_T(\hat{\theta}, h_E) + \Gamma_T(\hat{\theta}, h_E) - \Gamma(\hat{\theta}, h_E) + \Gamma(\hat{\theta}, h_E) - \Gamma(\theta_0, h_E)$$

where the 1st difference is  $o_P(1)$  by Lemma 2.7, the 3rd difference is  $o_P(1)$  by the CMT and the consistency of  $\hat{\theta}$ , see Lemma 2.8, and the 2nd difference is  $o_P(1)$  by Jennrich's ULLN. The ULLN can be applied if and only if  $\sum_j h_E(z_{jt}) \frac{\partial \xi_{jt}(\theta)}{\partial \theta}$  has an envelope with finite 1st absolute moments:  $\xi_{jt}(\theta) = \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\lambda)) - x'_{1jt}\beta$  and  $\frac{\partial \xi_{jt}(\theta)}{\partial \beta} = x_{1jt}$  with  $x_{1jt}$  has finite moments of order 4 by Assumption **B(iv)**, whereas  $\frac{\partial \xi_{jt}(\theta)}{\partial \lambda} = \frac{\partial \rho^{-1}(s_t, x_{2t}, f_0(\cdot|\tilde{\lambda}))}{\partial \lambda}$  and  $\rho^{-1}$  is  $C^\infty$  with arguments  $(s_t, x_{2t}, \lambda)$  which take values in a compact thus  $\frac{\partial \rho^{-1}}{\partial \lambda}$  has bounds.

Thence  $\text{plim } \hat{\Gamma}_T(\hat{\theta}, h_E) = \text{plim } \hat{\Gamma}_T(\tilde{\theta}, h_E) = \Gamma(\theta_0, h_E)$  which is full rank by Assumption **D(ii)**,  $\text{plim } \hat{W} = W$  which is full rank by Assumption **D(iv)**, and by Lemma 2.7  $\text{plim } \sqrt{T}(\hat{g}_T(\theta_0, h_E) - g_T(\theta_0, h_E)) = 0$  so we can apply the aforementioned CMT and by the CLT which can be applied because  $g(\theta_0, h_E) = 0$  under the null

$$\begin{aligned} \sqrt{T}(\hat{\theta} - \theta_0) &= - (\Gamma'(\theta_0, h_E)W\Gamma(\theta_0, h_E))^{-1} \sqrt{T}\Gamma'(\theta_0, h_E)Wg_T(\theta_0, h_E) + o_P(1) \\ &\xrightarrow{d} \mathcal{N}(0, (\Gamma'(\theta_0, h_E)W\Gamma(\theta_0, h_E))^{-1}\Gamma'(\theta_0, h_E)W\Omega(\mathcal{F}_0, h_E)W\Gamma(\theta_0, h_E) \\ &\quad (\Gamma'(\theta_0, h_E)W\Gamma(\theta_0, h_E))^{-1}) \end{aligned}$$

□

#### B.4.6 Asymptotic distribution of the test statistic

##### Proof of Theorem 5.1

*Proof.* This proof leans heavily on the proof of Lemma 2.9. By Taylor's theorem there exists  $\tilde{\theta}$  such that  $\|\tilde{\theta} - \theta_0\|_2 \leq \|\hat{\theta} - \theta_0\|_2$

$$\begin{aligned}\sqrt{T}\hat{g}_T(\hat{\theta}, h_D) &= \sqrt{T}\hat{g}_T(\theta_0, h_D) + \hat{\Gamma}_T(\tilde{\theta}, h_D)\sqrt{T}(\hat{\theta} - \theta_0) \\ &= (I_{|h_D|_0} - \Gamma(\theta_0, h_D)(\Gamma'(\theta_0, h_D)W\Gamma(\theta_0, h_D))^{-1}\Gamma'(\theta_0, h_D)W)\sqrt{T} \begin{pmatrix} g_T(\theta_0, h_D) \\ g_T(\theta_0, h_E) \end{pmatrix} + o_P(1) \\ &\equiv (I_{|h_D|_0} - G)\sqrt{T} \begin{pmatrix} g_T(\theta_0, h_D) \\ g_T(\theta_0, h_E) \end{pmatrix} + o_P(1)\end{aligned}$$

The second equality is obtained by relying on the proof of Lemma 2.9 to express  $\sqrt{T}(\hat{\theta} - \theta_0)$  as a function of moments, by relying on Lemma 2.7 so that  $\text{plim } \sqrt{T}\hat{g}_T(\theta_0, h_D) = \text{plim } \sqrt{T}g_T(\theta_0, h_D)$  and  $\text{plim } \hat{\Gamma}_T(\tilde{\theta}, h_D) = \text{plim } \Gamma_T(\theta_0, h_D)$ , and by using the CMT.

- Under  $H_0 : f \in \mathcal{F}_0$  then  $\mathbb{E} \left[ \sum_j h_D(z_{jt})\xi_{jt}(\theta_0) \right] = 0$  by LIE. So using the CLT and Slutsky's Lemma we obtain

$$\sqrt{T}\hat{g}_T(\hat{\theta}, h_D) \xrightarrow{d} Z \sim \mathcal{N}(0, \Omega_0)$$

where

$$\Omega_0 = \begin{pmatrix} I_{|h_D|_0} & G \end{pmatrix} \begin{pmatrix} \Omega(\mathcal{F}_0, h_D) & \Omega(\mathcal{F}_0, h_D, h_E) \\ \Omega(\mathcal{F}_0, h_D, h_E)' & \Omega(\mathcal{F}_0, h_E) \end{pmatrix} \begin{pmatrix} I_{|h_D|_0} \\ G' \end{pmatrix}$$

with

$$\begin{aligned}\Omega(\mathcal{F}_0, h_D) &= \mathbb{E} \left[ \left( \sum_j \xi_{jt}(f(\cdot|\lambda_0), \beta_0)h_D(z_{jt}) \right) \left( \sum_j h_D(z_{jt})\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) \right)' \right] \\ \Omega(\mathcal{F}_0, h_D, h_E) &= \mathbb{E} \left[ \left( \sum_j \xi_{jt}(f(\cdot|\lambda_0), \beta_0)h_D(z_{jt}) \right) \left( \sum_j h_E(z_{jt})\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) \right)' \right] \\ G &= -\Gamma(\theta_0, h_D) [\Gamma(\theta_0, h_E)'W\Gamma(\theta_0, h_E)]^{-1} \Gamma(\theta_0, h_E)'W\end{aligned}$$

Thence by the continuous mapping theorem:

$$S(h_D, \mathcal{F}_0, \hat{\theta}) = \hat{g}_T(\hat{\theta}, h_D)' \hat{\Sigma} \hat{g}_T(\hat{\theta}, h_D) \xrightarrow{d} Z' \Sigma Z$$

□

- Under  $H'_a : \mathbb{E} \left[ \sum_j h_D(z_{jt}) \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) \right] \neq 0$ , we have by Lemma 2.7, by consistency of  $\hat{\theta} \xrightarrow{P} \theta_0$  and the CMT:

$$\hat{g}_T(\hat{\theta}, h_D) = g_T(\theta_0, h_D) + o_P(1)$$

Thus by Assumption D(iv) and the CMT

$$\frac{S(h_D, \mathcal{F}_0, \hat{\theta})}{T} \xrightarrow{P} \underbrace{\mathbb{E} \left[ \sum_j h_D(z_{jt}) \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) \right]' \Sigma \mathbb{E} \left[ \sum_j h_D(z_{jt}) \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) \right]}_{\kappa(h_D, \mathcal{F}_0, \theta_0)}$$

Under  $H'_a$ ,  $\kappa(h_D, \mathcal{F}_0, \theta_0)$  is strictly positive because  $\Sigma$  is positive definite. Thence,

$$\begin{aligned} \forall q \in \mathbb{R} \quad \lim_{T \rightarrow \infty} \mathbb{P}(S(h_D, \mathcal{F}_0, \hat{\theta}) > q) &= \lim_{T \rightarrow \infty} \mathbb{P} \left( \frac{S(h_D, \mathcal{F}_0, \hat{\theta}) - q}{T} > 0 \right) \\ &= \mathbb{P}(\kappa(h_D, \mathcal{F}_0, \theta_0) > 0) \\ &= 1 \end{aligned}$$

where the 2nd equality holds because convergence in probability implies convergence in distribution.

□

#### B.4.7 Application of Theorem 5.1 to the 2 polar cases

##### 1. Sargan-Hansen J test

If  $h_D = h_E$ , with  $W$  and  $\Sigma$  are set to be equal to the GMM 2-step optimal weighting matrix

$$\Sigma = W = \mathbb{E} \left[ \left( \sum_j \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) h_E(z_{jt}) \right) \left( \sum_j \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) h_E(z_{jt}) \right)' \right]^{-1} = \Omega(\mathcal{F}_0, h_E)^{-1}$$

Then under  $H_0$ :

$$S(h_D, \mathcal{F}_0, \hat{\theta}) \xrightarrow{d} \chi^2_{|h_E|_0 - |\theta|_0}$$

*Proof.* By applying theorem 5.1, we have:

$$S(h_D, \mathcal{F}_0, \hat{\theta}) \xrightarrow{d} Z' \Sigma Z \quad \text{with} \quad Z \sim \mathcal{N}(0, \Omega_0)$$

If  $h_D = h_E$  and  $W = \Omega(\mathcal{F}_0, h_E)^{-1}$  then  $\Omega_0$  simplifies to

$$\begin{aligned}\Omega_0 &= \Omega(\mathcal{F}_0, h_E) - \Gamma(\theta_0, h_E) [\Gamma(\theta_0, h_E)' \Omega(\mathcal{F}_0, h_E)^{-1} \Gamma(\theta_0, h_E)]^{-1} \Gamma(\theta_0, h_E)' \\ &= \Omega(\mathcal{F}_0, h_E)^{1/2} M_{\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)} \Omega(\mathcal{F}_0, h_E)^{1/2}\end{aligned}$$

with  $M_{\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)} \equiv I_{|h_E|_0} - P_{\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)}$  is the orthogonal projection on the space orthogonal to  $\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)$ . Let  $\tilde{Z} \sim \mathcal{N}(0, I_{|h_E|_0})$ , we have by definition:

$$\begin{aligned}Z &= \Omega(\mathcal{F}_0, h_E)^{1/2} M_{\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)} \tilde{Z} \implies \Sigma^{1/2} Z = M_{\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)} \tilde{Z} \\ &\implies Z' \Sigma Z = \tilde{Z}' M_{\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)} \tilde{Z}\end{aligned}$$

Second line comes from symmetry and idempotence of  $M_{\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)}$ . Orthogonal projections have eigenvalues equal to either 0 or 1 with the number of eigenvalues equal to one corresponding to the rank of the space it projects into, which in our case is  $|h_E| - |\theta|_0$ . If we denote by  $V$  the matrix of eigenvectors of  $M_{\Omega(\mathcal{F}_0, h_E)^{-1/2} \Gamma(\theta_0, h_E)}$  then note that  $V' \tilde{Z} \sim \mathcal{N}(0, I_{|h_E|_0})$  so that

$$Z' \Sigma Z = \sum_{k=1}^{|h_E|_0 - |\theta|_0} (V' \tilde{Z})_k^2 \sim \chi_{|h_E|_0 - |\theta|_0}^2$$

□

## 2. Non-redundant $h_D$ and $h_E$

If  $\Omega_0$  is full rank and if the econometrician sets  $\Sigma = \Omega_0^{-1}$ , then our test statistic has the following asymptotic distribution under  $H_0$ :

$$S(h_T, \mathcal{F}_0, \hat{\theta}) \xrightarrow{d} \chi_{|h_D|_0}^2$$

One sufficient condition for  $\Omega_0$  being full rank is  $(\xi_{jt}(f(\cdot|\lambda_0), \beta_0))_{j=1}^J$  is independent across  $j$  and  $(h_E(z_{jt}), h_D(z_{jt}))$  not being perfectly colinear.

*Proof.* The asymptotic result is direct;  $(\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0))_{j=1}^J$  being independent across  $j$  and  $(h_E(z_{jt}), h_D(z_{jt}))$  not being perfectly colinear implies that

$$\Omega(\mathcal{F}_0, h_E, h_D) = \sum_j \mathbb{E} [\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)^2 h_E(z_{jt}) h_D(z_{jt})']$$

$$\Rightarrow \Omega_0 = \sum_j (I_{|h_D|_0} \quad G) \text{Var} \left( \begin{pmatrix} \xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) \\ h_D(z_{jt}) \\ h_E(z_{jt}) \end{pmatrix} \right) \begin{pmatrix} I_{|h_D|_0} \\ G' \end{pmatrix}$$

Thus  $\Omega_0$  is positive definite because it is the sum of positive definite matrices.  $\square$

## B.5 Properties of the MPI in the composite specification test: $f \in \mathcal{F}_0$

**Proposition 8** (Consistency of the test for the composite test with the MPI). Under Assumption **A** and Assumptions **B-E**

$$H_a : f \notin \mathcal{F}_0 \implies \forall q \in \mathbb{R}^+, \quad \mathbb{P}(S(h_D^*, \mathcal{F}_0, \hat{\theta}) > q) \rightarrow 1.$$

*Proof of Proposition 8.*

From corollary 2.1. Under Assumption **A**,

$$\begin{aligned} H_a : f \notin \mathcal{F}_0 &\implies \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)|z_{jt}] \neq 0 \text{ a.s} \\ &\implies \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)|z_{jt}]^2 > 0 \text{ a.s} \\ &\implies \mathbb{E}[\mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)|z_{jt}]^2] > 0 \\ &\implies \mathbb{E}[\mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)\mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0)|z_{jt})|z_{jt}]] > 0 \\ &\implies \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)\mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0)|z_{jt})] > 0 \\ &\implies \forall \alpha \neq 0 \quad H'_a : \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0) \underbrace{\alpha \mathbb{E}[\Delta_{0,a}^{\xi_{jt}}|z_{jt}]}_{h_D^*(z_{jt})}] \neq 0. \end{aligned}$$

From theorem 5.1, under Assumptions **B-E**,

$$H'_a : \mathbb{E}[\xi_{jt}(f_0(\cdot|\lambda_0), \beta_0)h_D^*(z_{jt})] \neq 0 \implies \forall q \in \mathbb{R}^+, \quad \mathbb{P}(S(h_D^*, \mathcal{F}_0, \hat{\theta}) > q) \rightarrow 1.$$

$\square$

## C Additional results and comments

### C.1 Literature on the identification of the distribution of RC

In this section, we briefly summarize some recent findings on the identification of random coefficients in multinomial choice models. In their seminal paper, [Berry and Haile \(2014\)](#) shows the identification of the demand functions  $\rho$  in a framework that encompasses the BLP model but their result does not entail identification of the random coefficients' distribution per se. To achieve their identification result, they require a completeness condition on the instruments as well as additional conditions (eg: connected substitutes) to ensure invertibility of the demand functions. They also need to impose that at least one of the product characteristic has a coefficient that is not random and that is equal to 1. Notice that in BLP model, the structure implied by the logit shock guarantees invertibility of the demand functions.

[Fox et al. \(2012\)](#) provides conditions under which the distribution of random coefficients is identified in a mixed logit model with micro-level data and no endogeneity. Their identification result requires continuous characteristics in  $x_{2t}$  and rules out interaction terms (eg polynomial terms of  $x_{2jt}$ ). Moreover, their result is restricted to distributions of random coefficients with a compact support - excluding for instance a normally distributed random coefficient.

[Fox and Gandhi \(2011\)](#) investigates the identification of the joint distribution of random coefficients  $v_i$  and idiosyncratic shocks  $\varepsilon_{ijt}$  in aggregate demand models without endogeneity. They also consider a setting where endogeneity is introduced in a very restrictive way. They show identification under a special regressor assumption and finite support of the unobserved heterogeneity. The special regressor assumption assumes that a variable in  $x_{1t}$  has full support and has an associated coefficient that is either 1 or -1. This special regressor assumption is very common in the literature on the identification of random coefficients (see [Ichimura and Thompson \(1998\)](#), [Berry and Haile \(2009\)](#), [Matzkin \(2007\)](#) and [Lewbel \(2000\)](#)). Their framework does not nest the standard BLP model as  $\varepsilon_{ijt}$  and  $v_i$  are both assumed to have a finite support but it is more general in other dimensions. They do not exploit the logit distributional assumption on  $\varepsilon_{ijt}$ , they do not impose independence between  $v_i$  and  $\varepsilon_{ijt}$ , their identification argument can be extended to the case where multiple goods are purchased.



In a setting much closer to ours, [Dunker et al. \(2022\)](#) studies the identification of the distribution of random coefficients in endogenous aggregate demand models which includes the BLP model as a special case (in particular, no parametric assumption is made on the idiosyncratic shock  $\varepsilon_{ijt}$ ). They make a clever use of the Radon transform to identify  $f$ . The price they have to incur for flexibility is that they need to make stringent assumptions on the product characteristics: variables in  $x_t$  are required to be continuous and to satisfy a joint full support assumption. The idea is to exploit the variation in the covariates in order to trace out the distribution of rc  $f$ . Unfortunately, these requirements are rarely met in real data sets.

In contrast to the rest of the literature, [Wang \(2022\)](#) adopts all the parametric assumptions assumed in the standard BLP model and looks for the set of minimal assumptions under which the distribution of random coefficients is identified. This approach allows him to obtain sufficient conditions which are much less stringent than the rest of the literature (no special regressor assumption, no full support assumption, no continuity assumption). To be more specific, he shows that if the demand functions are identified on an open set of  $\mathbb{R}^J$ <sup>38</sup>, then the distribution of random coefficients is identified. His proof astutely exploits the real analytic property of the demand functions<sup>39</sup>.

## C.2 Feasible MPI: conditional expectation

In this subsection, we briefly motivate our approach of approximating the conditional expectation by first projecting the endogenous variables on a relevant subset of the exogenous variables. The problem we encounter can be summarized as follows. We want to compute  $\mathbb{E}[g(x_{1t}, x_{2t})|z_{jt}]$ , where  $g$  is highly non-linear,  $x_{1t}$  are endogenous variables and  $x_{2t}$  are exogenous variables. Moreover,  $z_{jt}$  has a large dimension (in the BLP model, its order of magnitude the number of products  $\times$  number of characteristics). Our approach consists in first projecting the endogenous variables  $x_{1t}$  on a relevant subset of  $z_t$ , before plugging them into  $g(x_{1t}, x_{2t})$ . The traditional approach consists in using a non-parametric estimator of  $\mathbb{E}[g(x_{1t}, x_{2t})|z_{jt}]$ . However, given the dimension of  $z_{jt}$ , this

---

<sup>38</sup>which can be achieved using theorem 1 in [Berry and Haile \(2014\)](#)

<sup>39</sup>In particular, the real analytic property yields that the local identification of  $\rho$  on  $\mathcal{D} \subset \mathbb{R}^J$  implies identification of  $\rho$  on  $\mathbb{R}^J$ . From global identification of  $\rho$ , he is then able to show that the random coefficients' distribution is identified under a simple rank condition on  $x_{2t}$

approach is likely to yield poor results in practice because of the huge curse of dimensionality. In contrast, we know that some endogenous variables in  $x_{1t}$  only depend on a subset of  $z_{jt}$ , which we denote  $\tilde{z}_{1jt}$ , then we can use this to our advantage to construct a more precise estimator of  $\mathbb{E}[g(x_{1t}, x_{2t})|z_{jt}]$ . First,  $\hat{x}_{1t} = \mathbb{E}[x_{1t}|\tilde{z}_{1jt}]$  will be much more accurately estimated as we only condition on  $\tilde{z}_{1jt}$  (for instance, the price usually depends on its own cost shifters and its own product characteristics, while the dependence with respect to characteristics of other goods is much weaker and can be ignored empirically). For exposition, we further assume that  $\dim(x_{1t}) = 1$ . We take a second order Taylor expansion of  $\mathbb{E}[g(x_{1t}, x_{2t})|z_{jt}]$  around  $\hat{x}_{1t}$ .

$$\begin{aligned}\mathbb{E}[g(x_{1t}, x_{2t})|z_{jt}] &= \mathbb{E}[g(\hat{x}_{1t}, x_{2t})|z_{jt}] + \mathbb{E}\left[\frac{\partial g(\hat{x}_{1t}, x_{2t})}{\partial x_1}(x_{1t} - \hat{x}_{1t})|z_{jt}\right] + \mathbb{E}\left[\frac{\partial^2 g(\tilde{x}_{1t}, x_{2t})}{\partial x_1^2}(x_{1t} - \hat{x}_{1t})^2|z_{jt}\right] \\ &= g(\hat{x}_{1t}, x_{2t}) + \frac{\partial g(\hat{x}_{1t}, x_{2t})}{\partial x_1} \underbrace{\mathbb{E}[(x_{1t} - \hat{x}_{1t})|z_{jt}]}_{=0} + \mathbb{E}\left[\frac{\partial^2 g(\tilde{x}_{1t}, x_{2t})}{\partial x_1^2}(x_{1t} - \hat{x}_{1t})^2|z_{jt}\right]\end{aligned}$$

with  $\tilde{x}_{1t} \in [x_{1t}; \hat{x}_{1t}]$ . Our approach yields an estimator that converges faster to  $g(\hat{x}_{1t}, x_{2t})$ , which is a first order approximation of  $\mathbb{E}[g(x_{1t}, x_{2t})|z_{jt}]$ .

### C.3 Choice of the large- $T$ asymptotics

In this paper, we study the asymptotics of our test when the number of markets  $T$  grows to infinity. We could also study the asymptotic properties of the BLP estimator and of the test when  $J$  grows to infinity and  $T$  stays fixed. We do not pursue this route for several reasons. First, from an economic point of view, a market with a number of products that grows to infinity is hardly conceivable in industries with imperfect competition and barriers to entry. Second, from a theoretical point of view there is a tension between the identification of demand which require all market shares to be strictly positive, see [Berry and Haile \(2014\)](#), and the large market asymptotics which require all market shares to tend to 0 as  $J$  grows to infinity, see [Berry et al. \(2004\)](#). At the same time it is well established that a many (weak) instruments problem can easily occur in a BLP model with a fixed number of markets and many products especially when using the traditional BLP instruments, see [Armstrong \(2016\)](#).

Consequently only markets with perfect competition and a careful choice of instruments could somehow fit the assumptions necessary for the BLP model to yield consistent estimators and

valid tests with large  $J$ . Yet in the majority of empirical IO papers the markets have imperfect competition, sometimes oligopolies, and use the traditional BLP instruments. Thus we establish our theory with a large number of independent markets, which is a natural setting for empirical IO papers and which is not plagued with the aforementioned theoretical problems.

## C.4 Construction of the interval instruments in practice

We now provide more details on how to construct the interval instruments in practice. The procedure to construct the interval instruments is as follows:

1. Given  $(\mathcal{F}_0, \hat{W}, h_E)$ , the researcher derives the BLP estimator  $\hat{\theta}$
2. Then the researcher chooses  $L$  points  $(v_l)_{l=1}^L \in \mathbb{R}^L$  in the presumed support of  $f_0(\cdot|\hat{\lambda})$ .
3. Finally, the researcher can construct a set of  $L$  interval instruments based on the approximations of the MPI that we develop in sections 4.2 and 4.1.

- Global approximation:  $\{\pi_{j,l}(z_{jt})\}_{l=1,\dots,L}$  interval instruments, which are such that:

$$\mathbb{E}[\Delta_j(s_t, x_{2t}, f_0, f_a)|z_{jt}] \approx \log \left( \sum_{l=1}^L \omega_l \pi_{j,l}(z_{jt}) \right) \text{ with } \pi_{j,l}(z_{jt}) = \frac{\frac{\exp\{x'_{2jt} v_l\}}{1 + \sum_{k=1}^J \exp\{\hat{\delta}_{kt}^0 + x'_{2kt} v_l\}}}{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt} v\}}{1 + \sum_{k=1}^J \exp\{\hat{\delta}_{kt}^0 + x'_{2kt} v\}} f_0(v) dv}$$

with  $\hat{\delta}_t^0$  the linear projection of  $\delta_t^0$  on  $z_{jt}$  (or a carefully chosen subset of  $z_{jt}$ ).

- Local approximation:  $\{\bar{\pi}_{j,l}(z_{jt})\}_{l=1,\dots,L}$  interval instruments such that

$$\mathbb{E}[\Delta_j(s_t, x_{2t}, f_0, f_a)|z_{jt}] \approx \sum_{l=1}^L \bar{\omega}_l \bar{\pi}_{j,l}(z_{jt})$$

$$\text{with } \bar{\pi}_{j,l}(z_{jt}) = \left( \frac{\partial \rho(\hat{\delta}_t^0, x_{2t}, f_0)}{\partial \delta} \right)^{-1} \left[ \frac{\exp\{\hat{\delta}_t^0 + x_{2t} v_l\}}{1 + \sum_{k=1}^J \exp\{\hat{\delta}_{kt}^0 + x'_{2kt} v_l\}} - \rho_j(\hat{\delta}_t^0, x_{2t}, f_0) \right]$$

with  $\hat{\delta}_t^0$  the linear projection of  $\delta_t^0$  on  $z_{jt}$  (or a carefully chosen subset of  $z_{jt}$ ).

**Choice of the  $L$  points in the domain of  $f_a$**  The researcher doesn't know a priori the support of the true density  $f_a$ . Thus, he/she must choose points in the domain of definition of  $f_a$ . If this choice coincides with points of the support where  $|f_0(\cdot|\lambda_0) - f_a|$  is large, then this choice generates

more informative instruments. In practice, one can take points in the high density regions of  $f_0(\cdot|\lambda_0)$  (eg if  $\mathcal{F}_0$  is the Gaussian family, then one can take points around the mean  $\lambda_0$ ). The choice of the number of instruments  $N$  obeys a usual bias variance tradeoff. On the one hand, a large  $L$  allows to better approximate the MPI and thus increases the detection ability of the instruments. On the other hand, it is well-known that a larger number of instruments can induce finite sample bias and can distort asymptotic distributions of estimators and tests such as the over-identification test.<sup>40</sup> Moreover, we observe in our simulations, that when one takes points in the support that are too close to each other, the implied instruments suffer from high levels of colinearity. For these reasons we advise not to use too few or too many interval instruments, in our simulations and application we use between 6 and 10 instruments (in every dimension). We leave a formal analysis of the optimal choice of  $L$  and of the general approximations properties of the interval instruments for future work.

## C.5 Feasible MPIs for estimation

In the estimation framework, the researcher assumes that  $f \in \mathcal{F}_0 = \{f_0(\cdot|\tilde{\lambda}) : \tilde{\lambda} \in \Lambda_0\}$  and wants to estimate the true parameter  $\theta_0 = (\beta'_0, \lambda'_0)'$  under this parametric restriction. From the connection between the MPI and the local instruments that we present in Section 3.3, we infer that good testing instruments  $h_E(z_{jt})$  ought to approximate the MPI devoted to test  $H_0 : \theta = \theta_0$  against any local alternative. If we have an initial estimator of  $\theta_0$ , we can directly use the interval instruments presented in Section 4 to approximate the MPI devoted to test  $H_0 : \theta = \theta_0$ . However, this approach requires to estimate  $\hat{\theta}$  in a first step. Here, we present an alternative approach based on the global approximation of the MPI we derived in Section 4.2, which has the advantage of not requiring a first stage estimate of  $\theta_0$ . it is straightforward to show that for any true parameter  $\theta_0$  and any alternative  $\theta_a$ , we can rewrite the global approximation of the non-linear part of the MPI as follows:

---

<sup>40</sup>See [Roodman \(2009\)](#) for a review on the effect of many possibly weak moments on estimation and testing.

$$\mathbb{E}[\Delta_j(s_t, x_{2t}, \theta_0, \theta_a) | z_{jt}] \approx \log \left( \sum_{l=1}^L \bar{\omega}_l(\theta_0, \theta_a) \hat{\pi}_{j,l}(z_{jt}) \right) \text{ with } \hat{\pi}_{j,l}(z_{jt}) = \frac{\exp\{x'_{2jt} v_l\}}{1 + \sum_{k=1}^J \exp\{\hat{\delta}_{jt}^0 + x'_{2jk} v_l\}}$$

$$\text{and } \bar{\omega}_l(\theta_0, \theta_a) = \frac{\bar{\omega}_l(\theta_a)}{\int_{\mathbb{R}^{K_2}} \frac{\exp\{x'_{2jt} v\}}{1 + \sum_{k=1}^J \exp\{\delta_{jt}^0 + x'_{2jk} v\}} f_0(\cdot | \lambda_0)(v) dv}$$

, with  $\hat{\delta}_{jt}^0$  projected first stage estimates of  $\delta_{jt}^0$ , which can be obtained, for example, under the logit specification.  $\hat{\pi}_{j,l}(z_{jt})$  do not depend on  $f_0$  and can be used for estimation.

## C.6 Estimation procedure when the distribution of RC is a mixture

In this section, we present a procedure to estimate the BLP model when the distribution of RC is parametrized as a mixture. Namely, we perform the estimation under  $H_0 : f \in \mathcal{F}_0$  with  $\mathcal{F}_0$  the family of Gaussian mixtures with  $L$  components. The pdf of a Gaussian mixture writes as follows:

$$\forall x \in \mathbb{R}, f_0(x | \lambda_0) = \sum_{l=1}^L p_{l0} f_l(x | \lambda_{l0}) \quad \sum_{l=1}^L p_{l0} = 1 \quad L \geq 1$$

where  $f_{l0}(\cdot | \lambda_{l0})$  is the pdf of a  $\mathcal{N}(\mu_{l0}, \sigma_{l0}^2)$ .

As long as the means are different ( $\mu_{l0} \neq \mu_{l'0} \forall l \neq l'$ ), the gaussian mixture is uniquely characterized by the vector  $\lambda_0 = (p_{10}, \dots, p_{L0}, \mu_{10}, \dots, \mu_{L0}, \sigma_{10}^2, \dots, \sigma_{L0}^2)$  up to permutations of indexes<sup>41</sup>. The objective of our procedure is to estimate the parameters of the model  $\theta_0 = (\beta_0, \lambda_0)$  where  $\lambda_0$  characterizes the mixture. In general, the problem of estimating a density by a mixture is solved through the use of the well-known Expectation-Maximization (EM) algorithm. In our case, the application of this algorithm is made difficult by two main obstacles. First, we do not observe directly the random coefficients. Second, we do not have individual choice data which would have enabled us to construct a likelihood as in Train (2008). As an alternative, we propose to adapt the BLP estimation procedure to estimate the parameters of a mixture of gaussians instead of the single normal distribution. The mixture affects the derivation of the market shares. The random coefficient  $v_i$  is now a gaussian mixture. Hence,  $v_i = \sum_{l=1}^L \mathbf{1}\{D_i = l\} v_{il}$  where  $(v_{il})_{i=1}^n$  are iid and

---

<sup>41</sup>If for some  $l \neq l'$  we have  $\mu_{l0} = \mu_{l'0}$  then the Gaussian mixture becomes observationally equivalent to an infinite number of other Gaussian mixtures

have density  $f_{l0}(\cdot|\lambda_{l0})$  known up to  $\lambda_{l0}$  for  $l = 1, \dots, L$ , and where  $(D_i)_{i=1}^n$  are iid categorically distributed with pmf  $\mathbb{P}(D_i = l) = p_{l0}$ . For all market  $t$  and product  $j$ , the demand functions are as follows:

$$\begin{aligned}\rho_j(\delta_t, x_{2t}, f_0(\cdot|\lambda_0)) &= \mathbb{P}(j \text{ chosen in market } t \text{ by } i|x_{1t}, x_{2t}, \xi_t) \\ &= \int_{\mathbb{R}} \frac{\exp\{x'_{1jt}\beta_0 + x'_{2jt}v + \xi_{jt}\}}{1 + \sum_{j'=1}^J \exp\{x'_{1j't}\beta_0 + x'_{2j't}v + \xi_{j't}\}} f_0(v|\lambda_0) dv \\ &= \sum_{l=1}^L p_{l0} \int_{\mathbb{R}} \frac{\exp\{\delta_{jt} + x'_{2jt}v\}}{1 + \sum_{j'=1}^J \exp\{\delta_{j't} + x'_{2j't}v\}} f_{l0}(v|\lambda_{l0}) dv\end{aligned}$$

**Reparametrization.** The parameter  $\lambda$  associated with the mixture consists of the means, the standard deviation and the probability of each component. As highlighted by [Ketz \(2019\)](#) in the simple Gaussian case, the way we parametrize the model can greatly affect the asymptotic properties of the estimator as well as the quality of the estimation. In particular, he shows that the standard deviations  $\sigma$  should be reparametrized in order to avoid boundaries issues when  $\sigma$  close to 0. We follow this parametrization and perform the minimization with respect to  $\{(+/-)\sqrt{\sigma_l}\}_{l=1}^L$  instead and  $(\sigma_l)_{l=1}^L$  directly. An additional difficulty in the case of mixtures concerns the estimation of the probabilities associated to each component. These probabilities must all be between 0 and 1 and their sum must be equal to 1. To smoothly integrate these constraints, we perform the optimization with respect to  $\gamma = (\gamma_2, \dots, \gamma_L)$  with  $p = (p_1, p_2, \dots, p_L) = (\frac{1}{1 + \sum_{l=2}^L \exp\{\gamma_l\}}, \frac{\exp\{\gamma_2\}}{1 + \sum_{l=2}^L \exp\{\gamma_l\}}, \dots, \frac{\exp\{\gamma_L\}}{1 + \sum_{l=2}^L \exp\{\gamma_l\}})$ .

**Estimation details.** Apart from the modification in the computation of the market shares and the new parametrization of the model, the estimation procedure with a mixture follows closely the traditional one and the parameters of interest are estimated by minimizing a GMM criterion. Let  $\mathcal{Q}(\theta)$  the GMM objective function:

$$\mathcal{Q}(\theta) = \hat{\xi}(\theta)' h_E(Z) W h_E(Z)' \hat{\xi}(\theta)$$

We now describe the derivation of the Gradient that we provide to the minimization program.

$$\frac{\partial \mathcal{Q}}{\partial \theta} = 2 \left[ \frac{\partial \hat{\xi}(\theta)}{\partial \theta} \right]' h_E(Z) W h_E(Z)' \hat{\xi}(\theta)$$

Where  $\frac{\partial \hat{\xi}(\theta)}{\partial \beta} = -x_1$  and where by the implicit function theorem we have  $\hat{\rho}_j(\delta_t, x_{2t}, \lambda) - s_{jt} = 0 \quad \forall j, t$  which implies:

$$\frac{\partial \hat{\xi}(\theta)}{\partial \lambda} = \frac{\partial \hat{\delta}(\theta)}{\partial \lambda} = - \left[ \frac{\partial \hat{\rho}(\delta, x_2, \lambda)}{\partial \delta} \right]^{-1} \frac{\partial \hat{\rho}(\delta, x_2, \lambda)}{\partial \lambda}$$

- $\frac{\partial \rho}{\partial \delta}$  is a  $JT \times JT$  diagonal by block matrix such that:

$$\frac{\partial \rho_j(\delta_t, x_{2t}, \lambda)}{\partial \delta_{kt}} = \begin{cases} \sum_l p_l \int \mathcal{T}_{jlt}(v) (1 - \mathcal{T}_{klt}(v)) \phi_l(v) dv & \text{if } j = k \\ - \sum_l p_l \int \mathcal{T}_{jlt}(v) \mathcal{T}_{klt}(v) \phi_l(v) dv & \text{if } j \neq k \end{cases}$$

$$\text{with } \mathcal{T}_{jlt}(v) \equiv \frac{\exp\{\delta_{jt} + x'_{2jt} v_l\}}{1 + \sum_{j'=1}^J \exp\{\delta_{j't} + x'_{2j't} v_l\}}$$

- $\frac{\partial \rho}{\partial \lambda}$  is a  $JT \times (3L - 1)$  matrix such that:

$$\begin{aligned} \frac{\partial \rho_j(\delta_t, x_{2t}, \lambda)}{\partial \mu_l} &= p_l \int \mathcal{T}_{jlt} \left( x_{2jt} - \sum_{j'} \mathcal{T}_{j'lt} x_{2j't} \right) \phi(v) dv \\ \frac{\partial \rho_j(\delta_t, x_{2t}, \lambda)}{\partial \sigma_l} &= p_l \int \mathcal{T}_{jlt} \left( x_{2jt} - \sum_{j'} \mathcal{T}_{j'lt} x_{2j't} \right) v \phi(v) dv \\ \frac{\partial \rho_j(\delta_t, x_{2t}, \lambda)}{\partial \gamma_l} &= \sum_{l'=1}^L \zeta(l, l') \int \mathcal{T}_{jlt} \end{aligned}$$

$$\text{With } \zeta(l, l') = \frac{-\exp\{\gamma_l\}}{1 + \sum_{k \neq 1} \exp\{\gamma_k\}} \times \frac{\exp\{\gamma_{l'}\}}{1 + \sum_{k \neq 1} \exp\{\gamma_k\}} + \mathbf{1}\{l = l'\} \frac{\exp\{\gamma_l\}}{1 + \sum_{k \neq 1} \exp\{\gamma_k\}} = -p_l \times p_{l'} + \mathbf{1}\{l = l'\} p_l$$

## C.7 Properties of the feasible approximations of the MPI

So far, we have studied the properties of the MPI, which is an ideal instrument that cannot be derived in practice. Nevertheless, in light of the previous results, the MPI provides a useful upper bound on the power that can be reached using our specification test. More precisely, the

asymptotic slope reached by the MPI can be interpreted as a power envelope on our specification test. Ideally, we want our specification test, with the approximated MPIs as instruments, to achieve slopes close to the ones reached by the MPI. We now distinguish 2 situations.

First, we consider the case where the econometrician tests  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$  against the a known alternative *over*  $\bar{H}_a : (f, \beta) = (f_a, \beta_a)$ . This situation is not interesting in practice as the econometrician usually doesn't know the true alternative and doesn't want to specify an alternative. Nevertheless, it illustrates that in this specific case, we can (in theory) derive a consistent estimator of the MPI. Indeed, in this particular case, we can directly derive an analytical expression for the correction term  $\Delta_{0,a}^{\xi_{jt}}$  either using its definition or the expression in 4.2. Next, we must to compute the conditional expectation of our the correction term with respect to  $z_{jt}$ . This step is quite challenging because the dimension of  $z_{jt}$  is large and because the correction term is heavily non-linear and non-separable with respect to the endogenous variables. In theory, a solution is to perform a Sieve non-parametric estimation of the conditional mean and under standard regularity conditions recover a consistent estimator of  $\mathbb{E}[\Delta_{0,a}^{\xi_{jt}}|z_{jt}]$ . Unfortunately, the rate of converge will be extremely slow given the dimension of  $z_{jt}$  and we don't recommend to do this in practice. Instead, we suggest to use the global approximation and to project the endogenous variables on the space spanned by a relevant subset of  $z_{jt}$ . As we show in Appendix C.2, this strategy yields an estimator which converges faster to a first order approximation of the MPI.

Second, we consider the more realistic situation where the econometrician tests  $\bar{H}_0 : (f, \beta) = (f_0, \beta_0)$  against an unspecified alternative. In this case, we use the interval instruments that we developed in Section 4 as an approximation of the MPI. Due to the different layers of approximations which intervene in the construction of these instruments and the absence of knowledge of  $f_a$ , it is quite difficult to establish conditions under which these instruments can reach the optimal slope of the MPI. A thorough analysis of the properties of these instruments is beyond the scope of this paper and may constitute an interesting starting point for future research. In spite of the lack of theoretical analysis, our Monte Carlo exercises show that the interval instruments perform really well in finite sample.



## D Monte Carlo experiments

### D.1 Counterfactuals under an alternative distribution

For the simulation exercise presented in Section 6.2, we use the following expressions for own- and cross-price elasticities for product  $j \in \{1, 2, \dots, J\}$ . For the sake of simplicity, we drop the market index,  $t$ , in the following expressions.

- Own-price elasticity:

$$\eta_j^j = \frac{p_j}{s_j} \frac{\partial s_j}{\partial p_j} = \frac{p_j}{s_j} \int -\alpha \left( 1 - \frac{\exp\{\delta_j + x_{cj}v_i\}}{1 + \sum_{j'=1}^J \exp\{\delta_{j'} + x_{cj'}v_i\}} \right) \underbrace{\frac{\exp\{\delta_j + x_{cj}v_i\}}{1 + \sum_{j'=1}^J \exp\{\delta_{j'} + x_{cj'}v_i\}}}_{s_{ij}} f_{\theta}(v) dv$$


- Cross-price elasticity ( $k \neq j$ ):

$$\eta_j^k = \frac{p_k}{s_j} \frac{\partial s_j}{\partial p_k} = \frac{p_k}{s_j} \int \alpha \left( \frac{\exp\{\delta_j + x_{cj}v_i\}}{1 + \sum_{j'=1}^J \exp\{\delta_{j'} + x_{cj'}v_i\}} \right) \frac{\exp\{\delta_k + x_{ck}v_i\}}{1 + \sum_{j'=1}^J \exp\{\delta_{j'} + x_{cj'}v_i\}} f_{\theta}(v) dv$$

where  $\alpha = 2$  and  $\delta_j = 2 + x_{aj} + 1.5x_{bj} - 2p_j + \xi_j$  in the DGP.

**Demand functions.** In Figure 9, we plot the demand functions generated under the different specifications (logit and gaussian) of the true densities.

Figure 9: Demand function



## D.2 Finite sample performance of the test

**Practical implementation of the test.** For each setting, we estimate the model for 1000 replications. Minimization is performed with `nloptr` (algorithm: NLOPT-LD-LBFGS). We provide an analytical gradient. The Threshold for the outer loop is  $1e-9$  while the threshold for the inner loop is  $1e-13$ . We use `squrem` and a C++ implementation for the computation of the market shares to speed up the contraction. We also parallelize the contraction over markets using 7 independent cores. Now we formally describe the instruments included in each test.

**Power against local alternatives.** We now assess the local power properties of our test by assuming that the random coefficient  $v_i$  is distributed according to a local alternative. Namely, we assume  $v_i \sim \left(1 - \frac{1}{\sqrt{T}}\right) \mathcal{N}(2, 1) + \frac{1}{\sqrt{T}} Y$  where  $Y$  is an alternative distribution including exponential,

Chi-square, Student, Uniform. We ensure that  $Y$  has mean 2 and variance 1. The results are reported in 13. First, we can observe that except for the uniform local alternative, our test appears to have non-trivial power against all the other local alternatives. For the exponential and chi-square distributions, it is clear that our test with interval instruments outperforms the Sargan-J test with traditional instruments. For the student local alternative, the results seem quite unstable for small sample sizes but as  $T$  increases, interval instruments also seem to perform better. For the uniform alternative, it appears that we don't have power against this local alternative.

Table 13: Empirical power, local alternatives (1000 replications)

Number of markets	T=50				T=100				T=200			
Test type	J	I	J	I Local	J	I Local	J	I	J	I Local	J	I Local
Exponential	0.266	0.704	0.227	0.677	0.222	0.869	0.272	0.868	0.236	0.982	0.394	0.975
Chi-square	0.217	0.219	0.134	0.174	0.13	0.167	0.096	0.151	0.099	0.171	0.086	0.15
Student	0.212	0.139	0.33	0.436	0.115	0.115	0.127	0.093	0.082	0.13	0.134	0.312
Uniform	0.198	0.1	0.126	0.074	0.107	0.062	0.095	0.051	0.073	0.049	0.084	0.044

### D.3 Finite sample performance of Interval instruments for estimation

**Practical implementation of the estimation procedure.** To assess the performance of our instruments in estimating the non-linear parameters with a flexible distribution of random coefficients, we simulate data with a distribution of random coefficients following a mixture of gaussians and we estimate the parameters of this mixture. For each setting, we estimate the model for 1050 replications. We select the replications with an objective function below a certain threshold (in order to avoid local minima). Minimization is performed with nloptr (algorithm: NLOPT-LD-LBFGS). We provide an analytical gradient, which we describe subsequently. The Threshold for the outer loop is 1e-9 while the threshold for the inner loop is 1e-13. We use squarem and a C++ implementation for the computation of the market shares to speed up the contraction. We also parallelize the contraction over markets using 7 independent core. Before we formally define the different sets of instruments, let us present the estimation procedure when the distribution of random coefficients is assumed to be a mixture.

**Instruments** Now we formally describe the instruments present in each different sets used for estimation.

- Differentiation instruments: differentiation instruments + exogenous characteristics (polynomial terms) + cost shifters (20 instruments)
- Optimal instruments are computed in two stages. The first stage instruments consist of differentiation instruments and exogenous characteristics (polynomial terms). Second stage instruments consist of polynomial terms of exogenous characteristics and the approximation of optimal instruments proposed in [Reynaert and Verboven \(2014\)](#) (approximation of  $\mathbb{E} \left[ \frac{\partial \rho_j^{-1}(s_t, x_{2t}, \lambda)}{\partial \lambda} \middle| z_t \right]$ ). The set called optimal instruments includes 15 instruments.
- Interval Instruments are computed in two stages. The first stage instruments consist of differentiation instruments and exogenous characteristics (polynomial terms). Second stage instruments are the interval instruments couples with some exogenous characteristics. A total of 23 instruments. The points in the support to compute the interval instruments are chose as follows: we take equally spaced points in the interval  $\{\beta_{3L} - 0.5(\beta_{3H} - \beta_{3L}), \beta_{3H} + 0.5(\beta_{3H} - \beta_{3L})\}$ .

**Comparison of the performance between the different sets of instruments.** We now report the mean biases and the empirical  $\sqrt{MSE}$  of the estimates for each set of instruments and for different sample sizes. We also plot the distributions of estimates for the non-linear parameters for the different sets of instruments. First, we plot the distribution of estimates obtained when the set of differentiation instruments from [Gandhi and Houde \(2019\)](#) is used with a sample of  $T = 200$  markets and  $J = 12$  products. We observe that despite a relatively large sample, the differentiation instruments perform rather poorly in estimating the non-linear parameters associated with the mixture of Gaussians. In particular, the estimates of the standard deviation parameters associated to each component are very dispersed and a large portion of the estimates are bunched at zero. Second, we plot the distribution of non-linear estimates obtained with the

optimal instruments from [Reynaert and Verboven \(2014\)](#). They tend to perform better than the differentiation instruments as we can see that the estimates are more concentrated around the true value. Yet, it is important to emphasize that the optimal instruments display large failure rates caused by perfect colinearity of the instruments. Finally, we plot the distribution of estimates for the non linear parameters when we use the interval instruments developed in Section 4. It appears clearly that the interval instruments yield a more concentrated distribution of estimates than the two other sets of instruments. For the sake of conciseness, we do not report the results with a mixture with 3 components but the observations we make with two components are even more exacerbated.

Table 14: Estimation mixture with “differentiation” instruments (1000 replications)

Parameter		$\beta_0$	$\alpha$	$\beta_1$	$\beta_2$	$\beta_{3L}$	$\sigma_{3L}$	$\beta_{3H}$	$\sigma_{3H}$	$p_L$
Sample size	true	2	-2	1.5	1	-2	0.5	4	0.5	0.25
T=50, J=12	bias	-0.12	0.022	-0.016	-0.018	0.214	0.184	-0.022	-0.045	0.027
	$\sqrt{MSE}$	0.308	0.06	0.215	0.215	0.633	0.734	0.281	0.35	0.075
T=50, J=20	bias	-0.064	0.011	-0.01	-0.011	0.189	0.347	0.022	-0.081	0.025
	$\sqrt{MSE}$	0.231	0.044	0.165	0.166	0.566	0.887	0.184	0.291	0.059
T=100, J=12	bias	-0.058	0.01	-0.012	-0.012	0.233	0.226	0.02	-0.066	0.027
	$\sqrt{MSE}$	0.204	0.041	0.147	0.148	0.592	0.703	0.256	0.305	0.072
T=100, J=20	bias	-0.04	0.006	-0.007	-0.007	0.198	0.423	0.047	-0.101	0.025
	$\sqrt{MSE}$	0.165	0.032	0.117	0.116	0.552	0.89	0.164	0.27	0.055
T=200, J=12	bias	-0.038	0.007	-0.003	-0.003	0.184	0.167	0.011	-0.049	0.019
	$\sqrt{MSE}$	0.152	0.03	0.11	0.11	0.466	0.601	0.176	0.262	0.053

Table 15: Estimation mixture with “Optimal” instruments(1000 replications)

Parameter		$\beta_0$	$\alpha$	$\beta_1$	$\beta_2$	$\beta_{3L}$	$\sigma_{3L}$	$\beta_{3H}$	$\sigma_{3H}$	$p_L$
Sample size	true	2	-2	1.5	1	-2	0.5	4	0.5	0.25
T=50, J=12	bias	-0.09	0.016	-0.012	-0.013	0.076	0.059	0.026	-0.111	0.01
	$\sqrt{MSE}$	0.296	0.057	0.234	0.232	0.361	0.483	0.212	0.281	0.036
T=50, J=20	bias	-0.046	0.007	0	0.001	0.074	0.11	0.028	-0.089	0.01
	$\sqrt{MSE}$	0.225	0.044	0.178	0.176	0.328	0.563	0.163	0.228	0.033
T=100, J=12	bias	-0.041	0.007	-0.004	-0.003	0.054	0.037	0.019	-0.066	0.007
	$\sqrt{MSE}$	0.202	0.039	0.157	0.158	0.279	0.4	0.154	0.211	0.028
T=100, J=20	bias	-0.029	0.004	-0.003	-0.003	0.074	0.107	0.033	-0.074	0.01
	$\sqrt{MSE}$	0.153	0.03	0.126	0.124	0.311	0.52	0.129	0.194	0.034
T=200, J=12	bias	-0.029	0.005	-0.001	-0.001	0.026	0.011	0.021	-0.061	0.004
	$\sqrt{MSE}$	0.136	0.026	0.111	0.111	0.184	0.313	0.113	0.172	0.018

Table 16: Estimation mixture with Global Interval instruments(1000 replications)

Parameter		$\beta_0$	$\alpha$	$\beta_1$	$\beta_2$	$\beta_{3L}$	$\sigma_{3L}$	$\beta_{3H}$	$\sigma_{3H}$	$p_L$
Sample size	true	2	-2	1.5	1	-2	0.5	4	0.5	0.25
T=50, J=12	bias	-0.154	0.029	-0.043	-0.045	0.017	0	-0.045	0.004	0.005
	$\sqrt{MSE}$	0.341	0.067	0.257	0.258	0.277	0.391	0.227	0.259	0.024
T=50, J=20	bias	-0.092	0.017	-0.02	-0.021	0.013	0.042	-0.018	-0.003	0.004
	$\sqrt{MSE}$	0.245	0.048	0.19	0.19	0.248	0.415	0.166	0.22	0.021
T=100, J=12	bias	-0.07	0.013	-0.017	-0.019	0.004	-0.012	-0.027	0.005	0.002
	$\sqrt{MSE}$	0.2	0.039	0.161	0.161	0.167	0.282	0.157	0.201	0.013
T=100, J=20	bias	-0.047	0.008	-0.006	-0.007	-0.009	-0.005	-0.008	-0.009	0.001
	$\sqrt{MSE}$	0.158	0.031	0.13	0.129	0.115	0.264	0.115	0.169	0.005
T=200, J=12	bias	-0.039	0.007	-0.004	-0.003	-0.006	-0.027	-0.015	-0.001	0.001
	$\sqrt{MSE}$	0.141	0.027	0.109	0.109	0.088	0.219	0.108	0.164	0.003

Table 17: Estimation mixture with Local Interval instruments(1000 replications)

Parameter		$\beta_0$	$\alpha$	$\beta_1$	$\beta_2$	$\beta_{3L}$	$\sigma_{3L}$	$\beta_{3H}$	$\sigma_{3H}$	$p_L$
Sample size	true	2	-2	1.5	1	-2	0.5	4	0.5	0.25
T=50, J=12	bias	-0.134	0.025	-0.023	-0.024	-0.006	-0.005	-0.039	-0.001	0.003
	$\sqrt{MSE}$	0.307	0.059	0.26	0.259	0.251	0.34	0.214	0.244	0.019
T=50, J=12	bias	-0.084	0.016	-0.024	-0.025	0.019	0.033	-0.023	0.01	0.003
	$\sqrt{MSE}$	0.245	0.047	0.188	0.186	0.228	0.38	0.15	0.184	0.018
T=50, J=12	bias	-0.075	0.015	-0.018	-0.016	0	0	-0.028	0.007	0.001
	$\sqrt{MSE}$	0.199	0.039	0.159	0.16	0.127	0.225	0.143	0.164	0.005
T=50, J=12	bias	-0.039	0.007	-0.011	-0.011	-0.003	0.004	-0.01	0.004	0.001
	$\sqrt{MSE}$	0.162	0.032	0.129	0.129	0.104	0.226	0.103	0.125	0.004
T=50, J=12	bias	-0.037	0.007	-0.008	-0.007	0.002	-0.007	-0.016	0.006	0.001
	$\sqrt{MSE}$	0.136	0.026	0.11	0.109	0.091	0.174	0.099	0.123	0.003

Figure 10: Distribution of estimates for non-linear parameters with “Differentiation” instruments ( $T = 200, J = 12$ )




Figure 11: Distribution of estimates for non-linear parameters with “Optimal” instruments ( $T = 200, J = 12$ )




Figure 12: Distribution of estimates for non-linear parameters with “Global Interval” instruments  
 $(T = 200, J = 12)$





Figure 13: Distribution of estimates for non-linear parameters with “Local interval” instruments  
 $(T = 200, J = 12)$





### D.3.1 Estimation with a single Gaussian

Table 18: Estimation with a single Gaussian (1000 replications)

Instruments		Differentiation						"Optimal"						Interval Global						Interval Local					
Parameter		$\beta_0$	$\alpha$	$\beta_1$	$\beta_2$	$\beta_3$	$\sigma_3$	$\beta_0$	$\alpha$	$\beta_1$	$\beta_2$	$\beta_3$	$\sigma_3$	$\beta_0$	$\alpha$	$\beta_1$	$\beta_2$	$\beta_3$	$\sigma_3$	$\beta_0$	$\alpha$	$\beta_1$	$\beta_2$	$\beta_3$	$\sigma_3$
Sample size	true	2	-2	1.5	1	1.5	0.5	2	-2	1.5	1	1.5	0.5	2	-2	1.5	1	1.5	0.5	2	-2	1.5	1	1.5	0.5
T=50, J=12	bias	-0.16	0.032	-0.031	-0.028	-0.032	-0.004	-0.09	0.018	-0.016	-0.014	-0.018	-0.003	-0.15	0.03	-0.028	-0.026	-0.03	-0.004	-0.15	0.03	-0.028	-0.026	-0.03	-0.001
	$\sqrt{MSE}$	0.292	0.057	0.212	0.209	0.138	0.069	0.27	0.053	0.214	0.211	0.138	0.067	0.288	0.056	0.212	0.209	0.138	0.066	0.286	0.056	0.212	0.209	0.138	0.064
T=50, J=20	bias	-0.091	0.018	-0.022	-0.022	-0.015	0.001	-0.047	0.009	-0.013	-0.013	-0.006	0.001	-0.084	0.017	-0.021	-0.021	-0.013	0	-0.086	0.017	-0.021	-0.021	-0.014	0.002
	$\sqrt{MSE}$	0.209	0.041	0.159	0.16	0.106	0.05	0.199	0.039	0.16	0.161	0.106	0.05	0.206	0.041	0.16	0.16	0.106	0.052	0.208	0.041	0.159	0.16	0.106	0.052
T=100, J=12	bias	-0.088	0.017	-0.001	0	-0.027	0.001	-0.052	0.01	0.007	0.007	-0.02	0.001	-0.082	0.016	0	0.001	-0.026	0.001	-0.074	0.014	-0.016	-0.016	-0.013	0.001
	$\sqrt{MSE}$	0.199	0.039	0.146	0.145	0.1	0.045	0.189	0.037	0.148	0.147	0.099	0.047	0.197	0.039	0.146	0.146	0.1	0.044	0.185	0.036	0.151	0.152	0.099	0.044
T=100, J=20	bias	-0.043	0.009	-0.011	-0.012	-0.006	-0.001	-0.021	0.004	-0.007	-0.008	-0.002	-0.001	-0.04	0.008	-0.011	-0.012	-0.006	-0.001	-0.035	0.007	-0.01	-0.009	-0.004	0
	$\sqrt{MSE}$	0.145	0.028	0.115	0.114	0.075	0.035	0.141	0.028	0.115	0.114	0.075	0.035	0.145	0.028	0.115	0.114	0.076	0.035	0.14	0.027	0.116	0.115	0.076	0.035
T=100, J=20	bias	-0.038	0.007	-0.012	-0.012	-0.004	0.001	-0.017	0.003	-0.006	-0.007	-0.001	0	-0.032	0.006	-0.009	-0.01	-0.004	0	-0.033	0.006	-0.009	-0.01	-0.004	0.001
	$\sqrt{MSE}$	0.132	0.026	0.11	0.11	0.073	0.032	0.127	0.025	0.109	0.109	0.069	0.032	0.129	0.026	0.109	0.109	0.069	0.032	0.129	0.026	0.109	0.109	0.069	0.031

## E Empirical application

### E.1 First stage regression: instruments on price

In Table 19 we present the first stage regression for the endogenous variable, price. The explanatory variables include exogenous characteristics as well as the excluded instruments we presented in 7. We find that the excluded instruments are jointly significant with an F-stat of 467.41. As expected, we see that the steel futures price and its interaction with the weight of the car correlate positively with the price. We also see that the higher the exchange rate between the Euro and the local currency at the place of assembly, the lower the price of the car as the cost of production decreases. Moreover, we also see that if the location places a role as the European (country of assembly) dummy is negatively correlated with price. This could point to shipping expenses that are reflected in the price. Note that the effect of the labor costs is not as strong and not of the expected sign. Although we control for missing labor costs in the data with a missing dummy variable, it could still be causing a bias for the coefficient. Finally, the correlation between the competition-related instruments and the price shows that the degree of competition across cars of the same class matters.

Table 19: First stage regression for price

	Price/income
Labor costs (hourly)	−0.0002** (0.0001)
Steel (futures) price	−0.0001*** (0.00001)
Steel (futures) price * Weight	0.00003*** (0.00000)
# Cars by engine-type	0.001*** (0.0001)
# Cars by engine-type and class	−0.002*** (0.00004)
Exchange rate (non European)	−0.0002*** (0.00002)
Europe dummy	−0.018*** (0.003)
Horsepower	0.527*** (0.003)
Gasoline	−0.057*** (0.003)
Fuel cost	−0.003*** (0.001)
Size	0.037*** (0.002)
Foreign	−0.008*** (0.003)
Height	0.062*** (0.009)
Observations	38,999
R <sup>2</sup>	0.896
Adjusted R <sup>2</sup>	0.896
F Statistic	5,167.373*** (df = 65; 38933)

*Note: Brand, Year and State FE's are included.*

\*p<0.1; \*\*p<0.05; \*\*\*p<0.01

## E.2 Baseline specifications: logit and nested logit

Table 20 shows the results from the logit and nested logit specifications. We define the nests by the class of the car, therefore limiting the substitution between the cars that belong to the same

class within a nest.<sup>42</sup> We observe that the estimates are stable across specifications.

Table 20: Estimation results - Logit and Nested Logit

	<i>OLS</i>		<i>IV</i>		
	(1)	(2)	(3)	(4)	(5)
Price/income	-0.354*** (0.041)	-2.907*** (0.133)	-2.356*** (0.124)	-2.729*** (0.053)	-2.615*** (0.052)
log(within market shares)				0.420*** (0.006)	0.407*** (0.006)
Fuel Cost	-0.210*** (0.008)	-0.138*** (0.006)	-0.247*** (0.009)	-0.074*** (0.004)	-0.126*** (0.006)
Size( $m^2$ )	0.031 (0.038)	0.001 (0.040)	0.158*** (0.041)	-0.001 (0.025)	0.104*** (0.026)
Horsepower(KW/100)	0.136 (0.089)	3.151*** (0.183)	2.511*** (0.172)	2.586*** (0.080)	2.431*** (0.078)
Foreign	0.351*** (0.064)	0.083 (0.073)	0.120* (0.070)	-0.106** (0.046)	-0.101** (0.044)
Height(m)	0.870*** (0.216)	1.505*** (0.197)	3.487*** (0.228)	1.121*** (0.125)	2.270*** (0.145)
Gasoline	1.399*** (0.055)	0.625*** (0.061)	1.118*** (0.063)	0.190*** (0.039)	0.422*** (0.041)
Fuel cost $\times$ income	0.020*** (0.002)	-0.002** (0.001)	0.014*** (0.002)	-0.002*** (0.001)	0.007*** (0.001)
Size $\times$ income	-0.005*** (0.001)	-0.002*** (0.001)	-0.006*** (0.001)	0.0003 (0.001)	-0.002*** (0.001)
Horsepower $\times$ income	0.009*** (0.002)	-0.026*** (0.002)	-0.017*** (0.002)	-0.027*** (0.001)	-0.024*** (0.001)
Horsepower $\times$ time	-0.084*** (0.006)	-0.068*** (0.007)	-0.083*** (0.007)	-0.038*** (0.004)	-0.045*** (0.004)
Foreign $\times$ income	-0.019*** (0.001)	-0.015*** (0.001)	-0.016*** (0.001)	-0.008*** (0.001)	-0.008*** (0.001)
Height $\times$ income	-0.006 (0.004)	0.032*** (0.004)	-0.002 (0.005)	0.016*** (0.003)	-0.003 (0.003)
Height $\times$ density	-0.037*** (0.004)	-0.003*** (0.0003)	-0.037*** (0.004)	-0.001*** (0.0002)	-0.021*** (0.003)
Gasoline $\times$ income	-0.016*** (0.001)	-0.003*** (0.001)	-0.010*** (0.001)	0.0004 (0.001)	-0.003*** (0.001)
Gasoline $\times$ Post 2015		-0.024 (0.019)		-0.019 (0.012)	
Constant	-7.937*** (0.167)	-12.482*** (0.149)	-11.171*** (0.167)	-9.144*** (0.092)	-8.506*** (0.102)
State FE/ Year FE	<i>Yes</i>	<i>No</i>	<i>Yes</i>	<i>No</i>	<i>Yes</i>
Observations	39,888	39,888	39,888	39,888	39,888
R <sup>2</sup>	0.385	0.217	0.272	0.686	0.709
Adjusted R <sup>2</sup>	0.384	0.216	0.271	0.686	0.709

\*p<0.1; \*\*p<0.05; \*\*\*p<0.01

Note: Brand FE's are included.

<sup>42</sup>Car classes in the data are: Mini, small, lower-middle, middle, upper-middle, luxury.

### E.3 Counterfactual quantities under different specifications

We define quantities of interest and derive them under the different specifications considered previously. For exposition purposes, we omit the dependence of the market shares in  $\delta_t$ ,  $x_{2t}$  and  $f$ , and simply write  $s_j(\mathbf{p})$  instead of  $\rho_j(\delta_t, x_{2t}; f)$ , where  $\mathbf{p}$  is the price vector.

**Price elasticities.** For the calculation of the price elasticities one can refer to [D.1](#) that writes the quantities for the simulation exercise.

**Demand curvature.** The demand curvature is defined using second derivative of demand as follows:  $\eta_j^2(\mathbf{p}) = s_j(\mathbf{p}) \frac{\partial^2 s_j(\mathbf{p})}{\partial p_j^2} \left( \frac{\partial s_j(\mathbf{p})}{\partial p_j} \right)^{-2}$ .

**Marginal costs and mark-ups.** To recover the marginal costs and the implied mark-ups, we need to make additional assumptions on the supply side. Following the literature, we consider that each multi-product firm  $f \in F$  sets prices for its own products in accordance with a Bertrand-Nash equilibrium. The profit of each firm writes:

$$\Pi_f(\mathbf{p}) = \sum_t \sum_{j \in J_f} (p_j - c_j) M_t s_{jt}(\mathbf{p})$$

where  $J_f$  is the set of goods produced by firm  $f$ ,  $c_j$  is the marginal cost for good  $j$ ,  $M_t$  is the market size and  $s_j(p)$  is the market share of product  $j$ . The first-order condition with respect to price  $p_j$  writes:

$$\sum_t M_t s_{jt}(\mathbf{p}) + \sum_t M_t \sum_{j' \in J_f} (p_{j'} - c_{j'}) \frac{\partial s_{j'}(\mathbf{p})}{\partial p_j} = 0.$$

We gather all the FOCs and rewrite them in matricial form:

$$\mathbf{s}(\mathbf{p}) + (\mathbf{\Delta}(\mathbf{p})) (\mathbf{p} - \mathbf{c}) = 0.$$

where  $\mathbf{\Delta}(\mathbf{p}) = \sum_t M_t \frac{\partial s_{j'}(\mathbf{p})}{\partial p_j}$  if  $j'$  and  $j$  are produced by the same firm and equals to zero otherwise.  $\mathbf{\Delta}(\mathbf{p})$  is known as the ownership matrix. Assuming that the prices are in equilibrium, one can recover the marginal costs using the following equation:

$$\mathbf{c} = \mathbf{p} - (\mathbf{\Delta}(\mathbf{p}))^{-1} \mathbf{s}(\mathbf{p}). \tag{17}$$

The mark-up for product  $j$  simply writes:  $p_j - c_j$ .

**Pass-through** The pass-through of cost is defined as follows. Let us assume that the marginal cost for product  $j$  goes from  $c_j$  to  $c'_j$  (with  $c'_j > c_j$ ), then the cost pass-through equals  $\alpha_j = \frac{p'_j - p_j}{c'_j - c_j}$ , where  $p'_j$  is the new equilibrium price. We calculate the new equilibrium price using Eqn. 17 using fixed point iteration. The pass-through corresponds to the proportion of the cost increase that is transmitted to the price.

Table 21: Counterfactual quantities under different specifications on RCs (20 most popular cars)

Counterfactual quantity		Price elasticity			Curvature			Marginal cost			Mark-up			Pass-through		
Car	Manufacturer	Logit	Gaussian	Mixture	Logit	Gaussian	Mixture	Logit	Gaussian	Mixture	Logit	Gaussian	Mixture	Logit	Gaussian	Mixture
Golf	Volkswagen	-1.09	-0.95	-3.03	1.00	1.14	1.21	1260	-9670	15436	24098	35028	9922	0.92	-	1.30
Polo	Volkswagen	-0.74	-0.70	-2.50	1.00	1.15	1.09	-6643	-14366	9073	23819	31542	8103	1.05	-	1.09
Passat	Volkswagen	-1.43	-1.21	-2.27	1.00	1.17	1.57	9488	-1033	17826	24631	35153	16294	1.02	-	2.65
Corsa	PSA	-0.66	-0.63	-2.28	1.00	1.14	1.07	-8432	-11246	8410	24088	26902	7246	1.02	-	1.12
Fiesta	Ford	-0.62	-0.60	-2.18	1.00	1.15	1.07	-8983	-10806	7657	23487	25310	6847	1.03	-	1.10
Tiguan	Volkswagen	-1.32	-1.14	-2.28	1.00	1.17	1.55	6831	-2919	16211	24118	33868	14738	1.01	-	2.62
Golf	Volkswagen	-1.17	-1.03	-3.12	1.00	1.18	1.27	3128	-7932	16582	23828	34888	10374	0.99	-	1.41
up!	Volkswagen	-0.53	-0.52	-1.92	1.00	1.14	1.05	-11231	-17703	4594	23278	29749	7453	1.04	-	0.96
Tiguan	Volkswagen	-1.34	-1.15	-3.09	1.00	1.19	1.38	7051	-4117	19186	23842	35009	11706	1.01	-	1.66
1er-Reihe	BMW	-1.16	-1.03	-3.09	1.00	1.18	1.28	3845	-769	19179	25138	29753	9805	0.99	-	1.39
Octavia	Volkswagen	-1.23	-1.08	-2.33	1.00	1.17	1.50	4629	-4504	15464	24211	33345	13377	1.01	-	2.34
A4	Volkswagen	-1.56	-1.30	-2.26	1.00	1.19	1.56	13209	1995	20260	25865	37079	18814	1.01	-	2.66
Clio	Renault	-0.73	-0.70	-2.49	1.00	1.16	1.10	-6240	-8684	9817	23120	25563	7063	1.03	-	1.17
T-Roc	Volkswagen	-0.87	-0.81	-2.80	1.00	1.17	1.14	-3645	-12275	11578	23798	32427	8575	1.06	-	1.16
Kuga	Ford	-1.16	-1.03	-3.09	1.00	1.18	1.28	3654	-518	18214	23684	27856	9124	1.03	-	1.39
Golf	Volkswagen	-1.10	-0.99	-2.34	1.00	1.16	1.44	1548	-7284	13678	23929	32762	11799	0.96	-	2.13
A-Klasse	Daimler	-1.28	-1.10	-3.07	1.00	1.19	1.35	6608	562	20662	25066	31112	11013	1.01	-	1.56
Golf	Volkswagen	-1.05	-0.94	-2.33	1.00	1.16	1.42	417	-8115	13135	24177	32710	11460	0.72	-	2.11
Golf	Volkswagen	-1.18	-1.05	-3.15	1.00	1.18	1.27	3202	-8230	16705	23921	35353	10418	0.98	-	1.40
Octavia	Volkswagen	-1.05	-0.95	-3.02	1.00	1.17	1.21	380	-8835	14808	23862	33077	9433	0.78	-	1.30

Figure 14: Estimated demand functions under different specifications

