

Concepts in Abstract Mathematics

MAT246 LEC0101 Winter 2020

Problem Set 2

Solutions

1. (a) Let a and b be relatively prime natural numbers greater than or equal to 2. Prove that $a^{\phi(b)} + b^{\phi(a)} \equiv 1 \pmod{ab}$.
- (b) Find the remainder when $47^{144} + 185^{46} \cdot (46! + 2)^{46}$ is divided by 8695.

Solution.

- (a) Since a and b are relatively prime, Euler's Theorem implies that $a^{\phi(b)} \equiv 1 \pmod{b}$ and $b^{\phi(a)} \equiv 1 \pmod{a}$. Thus, $b \mid a^{\phi(b)} - 1$ and $a \mid b^{\phi(a)} - 1$, so

$$ab \mid (a^{\phi(b)} - 1)(b^{\phi(a)} - 1),$$

which means that $(a^{\phi(b)} - 1)(b^{\phi(a)} - 1) \equiv 0 \pmod{ab}$. Hence, expanding the left-hand side, we get

$$a^{\phi(b)}b^{\phi(a)} - a^{\phi(b)} - b^{\phi(a)} + 1 \equiv 0 \pmod{ab}.$$

Since $a, b \geq 2$, we have $\phi(a), \phi(b) \geq 1$, so $ab \mid a^{\phi(b)}b^{\phi(a)}$ and hence $a^{\phi(b)}b^{\phi(a)} \equiv 0 \pmod{ab}$. Thus, we get $0 - a^{\phi(b)} - b^{\phi(a)} + 1 \equiv 0 \pmod{ab}$, so $a^{\phi(b)} + b^{\phi(a)} \equiv 1 \pmod{ab}$.

- (b) Let $a = 47$ and $b = 185 = 5 \cdot 37$ so that $ab = 8695$. Since for any primes p, q , we have $\phi(p) = p - 1$ and $\phi(pq) = (p - 1)(q - 1)$, we get that $\phi(a) = 46$ and $\phi(b) = 4 \cdot 36 = 144$. Moreover, the canonical factorizations into primes of a and b are $a = 47$ and $b = 5 \cdot 37$, so they have no prime factor in common, and hence they are relatively prime. Thus, $47^{144} + 185^{46} \equiv 1 \pmod{8695}$ by part (a).

Now, a is prime so $a \mid (a - 1)! + 1$ by Wilson's Theorem, and hence

$$ab \mid b \cdot ((a - 1)! + 1).$$

In other words, $185 \cdot (46! + 1) \equiv 0 \pmod{8695}$, so

$$185 \cdot (46! + 2) \equiv 185 \cdot (46! + 1) + 185 \equiv 185 \pmod{8695}.$$

Hence,

$$47^{144} + 185^{46} \cdot (46! + 2)^{46} = 47^{144} + (185 \cdot (46! + 2))^{46} \equiv 47^{144} + 185^{46} \equiv 1 \pmod{8695},$$

so the remainder is 1.

2. (a) Let p and q be primes. Prove that \sqrt{pq} is rational if and only if $p = q$.
 (b) Prove that $\sqrt{57} + \sqrt{n}$ is irrational for all $n \in \mathbb{N}$.

Solution.

- (a) If $p = q$ then $\sqrt{pq} = \sqrt{p^2} = p$ is rational. For the converse, we give two different proofs.

Proof 1. Suppose, by contradiction, that $p \neq q$ and $\sqrt{pq} = \frac{a}{b}$ for some relatively prime numbers $a, b \in \mathbb{N}$. Then, $pqb^2 = a^2$, so $p \mid a^2$ and, since p is prime, this implies that $p \mid a$ (Lemma 7.2.2). Similarly, $q \mid a^2$, so $q \mid a$. Since p and q are distinct primes, they are relatively prime, so $pq \mid a$ (this was proved in Lecture 7 and is also a special case Q6 in PS1). Hence, $a = pqk$ for some $k \in \mathbb{N}$, so $pqb^2 = a^2 = p^2q^2k^2$ and hence $b^2 = pqk^2$. Repeating the same argument, we get that $p \mid b$ and $q \mid b$ so $pq \mid b$. Hence, $pq \mid a$ and $pq \mid b$ contradicting that a and b are relatively prime.

Proof 2. Suppose that \sqrt{pq} is rational. Since the square root of a natural number is rational only if the square root is a natural number (Theorem 8.2.8), we have $\sqrt{pq} = n$ for some $n \in \mathbb{N}$. Hence, $pq = n^2$. Let $n = r_1^{\alpha_1} \cdots r_k^{\alpha_k}$ be the canonical factorization of n , so that $pq = r_1^{2\alpha_1} \cdots r_k^{2\alpha_k}$. Then, we must have that $p = q$, as otherwise, we get two canonical factorizations of the same number, where all the exponents of the first one (i.e. pq) are 1 while all the exponents of the second one (i.e. $r_1^{2\alpha_1} \cdots r_k^{2\alpha_k}$) are even, contradicting uniqueness of canonical factorizations.

- (b) Note that $57 = 3 \cdot 19$ is the product of two distinct primes. (Side note: although 57 is not prime, it is often jokingly called the *Grothendieck prime*.) Hence, $\sqrt{57}$ is irrational by (a). Suppose, by contradiction, that $\sqrt{n} + \sqrt{57} = r \in \mathbb{Q}$. Then, $\sqrt{n} = r - \sqrt{57}$, so $n = (r - \sqrt{57})^2 = r^2 - 2r\sqrt{57} + 57$ and hence $\sqrt{57} = \frac{r^2 + 57 - n}{2r} \in \mathbb{Q}$, contradicting that $\sqrt{57}$ is irrational. Hence, $\sqrt{n} + \sqrt{57}$ is irrational.

3. (a) Prove that if $z, w \in \mathbb{C}$ then $\overline{z + w} = \bar{z} + \bar{w}$.
 (b) Prove that if $z, w \in \mathbb{C}$ then $\overline{zw} = \bar{z}\bar{w}$.
 (c) Prove that if $r \in \mathbb{C}$ is a root of a polynomial with real coefficients, then \bar{r} is also a root of that polynomial.

Solution.

- (a) Let $z = a + bi$ and $w = c + di$, where $a, b, c, d \in \mathbb{R}$. Then,

$$\begin{aligned}\overline{z + w} &= \overline{(a + bi) + (c + di)} \\ &= \overline{(a + c) + (b + d)i} \\ &= (a + c) - (b + d)i \\ &= (a - bi) + (c - di) \\ &= \bar{z} + \bar{w}.\end{aligned}$$

- (b) Let $z = a + bi$ and $w = c + di$, where $a, b, c, d \in \mathbb{R}$. Then,

$$\begin{aligned}\overline{zw} &= \overline{(a + bi)(c + di)} \\ &= \overline{(ac - bd) + (ad + bc)i} \\ &= (ac - bd) - (ad + bc)i \\ &= (ac - (-b)(-d)) + (a(-d) + (-b)c)i \\ &= (a - bi)(c - di) \\ &= \bar{z}\bar{w}.\end{aligned}$$

- (c) Let $p(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0$ be a polynomial with $a_n \in \mathbb{R}$ and let $r \in \mathbb{C}$ be a root of $p(z)$, i.e. $p(r) = 0$. We want to show that $p(\bar{r}) = 0$. By (a), we have $\overline{a_1 z + a_0} = \bar{a}_1 \bar{z} + \bar{a}_0$. Applying (a) again, we get $\overline{a_2 z^2 + a_1 z + a_0} = \overline{a_2 z^2} + \bar{a}_1 \bar{z} + \bar{a}_0 = a_2 \bar{z}^2 + \bar{a}_1 \bar{z} + \bar{a}_0$. Hence, applying (a) n times, we get

$$\overline{a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0} = \overline{a_n z^n} + \overline{a_{n-1} z^{n-1}} + \cdots + \bar{a}_1 \bar{z} + \bar{a}_0$$

Now, by (b), we have $\overline{a_i z^i} = \bar{a}_i \bar{z}^i = \bar{a}_i \bar{z}^i$ for all i . But $a_i \in \mathbb{R}$, so $\bar{a}_i = a_i$ and hence we have shown that

$$\overline{a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0} = a_n \bar{z}^n + a_{n-1} \bar{z}^{n-1} + \cdots + a_1 \bar{z} + a_0,$$

or in other words,

$$\overline{p(z)} = p(\bar{z}).$$

In particular, if $p(r) = 0$, then $p(\bar{r}) = \overline{p(r)} = \bar{0} = 0$.

4. Show that $|\mathbb{R}^n| = |\mathbb{R}|$ for all $n \in \mathbb{N}$.

Solution. We first show that $|\mathbb{R}^2| = |\mathbb{R}|$. Since $|\mathbb{R}| = |[0, 1]|$ (Theorem 10.3.8), we also have $|\mathbb{R}^2| = |[0, 1] \times [0, 1]|$. Indeed, the equality $|\mathbb{R}| = |[0, 1]|$ implies that there is a bijection $f : \mathbb{R} \rightarrow [0, 1]$ and hence the function $F : \mathbb{R}^2 \rightarrow [0, 1] \times [0, 1]$ given by $F(x, y) = (f(x), f(y))$ is also a bijection. Now, we also showed that $|[0, 1] \times [0, 1]| = |\mathbb{R}|$ (Theorem 10.3.33) so we have $|\mathbb{R}^2| = |[0, 1] \times [0, 1]| = |\mathbb{R}|$. Hence, $|\mathbb{R}^2| = |\mathbb{R}|$ (the fact that if $|S| = |T|$ and $|T| = |U|$ then $|S| = |U|$ follows from the fact that if $f : S \rightarrow T$ and $g : T \rightarrow U$ are bijections, then $g \circ f : S \rightarrow U$ is a bijection since $f^{-1} \circ g^{-1}$ is an inverse).

Now, we show by induction on n that $|\mathbb{R}^n| = |\mathbb{R}|$ for all $n \in \mathbb{N}$. The base case $n = 1$ is trivial, since $\mathbb{R}^1 = \mathbb{R}$. Suppose that $|\mathbb{R}^k| = |\mathbb{R}|$ for some $k \in \mathbb{N}$. We want to show that $|\mathbb{R}^{k+1}| = |\mathbb{R}|$. Since $|\mathbb{R}^k| = |\mathbb{R}|$ we have a bijection $f : \mathbb{R}^k \rightarrow \mathbb{R}$. Then, $\mathbb{R}^{k+1} = \mathbb{R}^k \times \mathbb{R}$ and we have a bijection $g : \mathbb{R}^k \times \mathbb{R} \rightarrow \mathbb{R} \times \mathbb{R}$ given by $g(x, y) = (f(x), y)$, so $|\mathbb{R}^k \times \mathbb{R}| = |\mathbb{R} \times \mathbb{R}|$. Hence, $|\mathbb{R}^{k+1}| = |\mathbb{R}^k \times \mathbb{R}| = |\mathbb{R} \times \mathbb{R}| = |\mathbb{R}^2| = |\mathbb{R}|$. To show that g is a bijection, we show that it is both surjective and injective. It is injective since if $g(x_1, y_1) = g(x_2, y_2)$ then $(f(x_1), y_1) = (f(x_2), y_2)$ so $f(x_1) = f(x_2)$ and $y_1 = y_2$. Since f is injective, we have $x_1 = x_2$, so $(x_1, y_1) = (x_2, y_2)$. Now, g is also surjective since if $(x, y) \in \mathbb{R} \times \mathbb{R}$ then, since f is surjective, there exists $x_1 \in \mathbb{R}^k$ such that $f(x_1) = x$ so $g(x_1, y) = (f(x_1), y) = (x, y)$.

5. Find the cardinality of each of those sets.

- (a) The set of lines in the plane.
- (b) The set of circles in the plane whose centre has rational coordinates and whose radius is the square root of a prime number.

Solutions.

- (a) We claim that the cardinality is c , the cardinality of \mathbb{R} .

First, a vertical line is uniquely determined by its intersection with the x -axis, and hence the set of vertical lines is in bijection with \mathbb{R} . A line that is not vertical is of the form $y = ax + b$ for unique $a, b \in \mathbb{R}$, and hence the set of non-vertical lines is in bijection with \mathbb{R}^2 . Hence, the set of all lines in the plane is in bijection with $\mathbb{R} \cup \mathbb{R}^2$. Now, since $|\mathbb{R}| = |[0, 1]|$ (Theorem 10.3.8) we have a bijection $f : \mathbb{R} \rightarrow [0, 1]$ and since $|\mathbb{R}^2| = |\mathbb{R}|$ (Q5) and $|\mathbb{R}| = |(1, 2]|$ (Theorem 10.3.7 and Theorem 10.3.8) we have another bijection $g : \mathbb{R}^2 \rightarrow (1, 2]$. Hence, we can construct a function $h : \mathbb{R} \cup \mathbb{R}^2 \rightarrow [0, 2]$ by defining $h(x) = f(x)$ for $x \in \mathbb{R}$ and $h(y, z) = g(y, z)$ for $(y, z) \in \mathbb{R}^2$. Then, h is a bijection since it has an inverse $k : [0, 2] \rightarrow \mathbb{R} \cup \mathbb{R}^2$ defined by $k(x) = f^{-1}(x)$ if $x \in [0, 1]$ and $k(x) = g^{-1}(x)$ if $x \in (1, 2]$, where f^{-1} and g^{-1} are the inverses of f and g respectively. Hence, $|\mathbb{R} \cup \mathbb{R}^2| = |[0, 2]| = |[0, 1]| = |\mathbb{R}|$.

- (b) Let C be the set of those circles. We claim that the cardinality of C is \aleph_0 . Since C is infinite and \aleph_0 is the smallest infinite cardinality, we have $\aleph_0 \leq |C|$. Hence, by the Cantor-Bernstein Theorem, it suffices to show that $|C| \leq \aleph_0$. In other words, it suffices to show that C is countable.

A circle in C is uniquely specified by a pair of rational numbers $x, y \in \mathbb{Q}$ and a prime number p , where (x, y) are the coordinates of the centre and \sqrt{p} is the radius. Hence, C is in bijection with $\mathbb{Q}^2 \cup \mathbb{P}$, where $\mathbb{Q}^2 = \mathbb{Q} \times \mathbb{Q} = \{(x, y) : x, y \in \mathbb{Q}\}$ and $\mathbb{P} \subseteq \mathbb{N}$ is the set of prime numbers. We begin by proving the following lemma.

Lemma. *If S and T are countable sets, then so is $S \times T$.*

Proof. The set $S \times T$ is the union of the sets $\{s\} \times T$ for $s \in S$. Now, for all $s \in S$, the set $\{s\} \times T$ is in bijection with T , which is countable, so $\{s\} \times T$ is also countable. Since S is countable and the union of a countable number of countable sets is countable (Theorem 10.2.10), we have that $S \times T$ is countable. \square

By this lemma, \mathbb{Q}^2 is countable. Also \mathbb{P} is countable since $\mathbb{P} \subseteq \mathbb{N}$ and a subset of a countable set is countable. Hence, $\mathbb{Q}^2 \cup \mathbb{P}$ is countable since a union of two countable sets is countable (Theorem 10.2.10). So C is countable and infinite, and hence $|C| = \aleph_0$.

6. Show that a set S has infinitely many elements if and only if it has a subset $S_0 \subseteq S$ such that $S_0 \neq S$ and $|S_0| = |S|$.

Solutions. Suppose that S has infinitely many elements. Since \aleph_0 is the smallest infinite cardinality, we have $\aleph_0 \leq |S|$, so there is an injection $f : \mathbb{N} \rightarrow S$. Let $s_i = f(i)$, so that s_1, s_2, s_3, \dots is an infinite sequence of distinct elements of S . Let $S_0 = S \setminus \{s_1\}$. Then, $S_0 \neq S$ since $s_1 \notin S_0$. We claim that $|S_0| = |S|$. Define $g : S \rightarrow S_0$ by $g(s_i) = s_{i+1}$ for all $i \in \mathbb{N}$ and $g(x) = x$ if $x \neq s_i$ for all i . Then, g is bijective since it has an inverse $h : S_0 \rightarrow S$ defined by $h(s_i) = s_{i-1}$ for all $i \geq 2$ and $h(x) = x$ if $x \neq s_i$ for all i . Indeed, if $h(g(s_i)) = h(s_{i+1}) = s_i$ and if $x \neq s_i$ then $h(g(x)) = h(x) = x$. Similarly, $g(h(x)) = x$ for all $x \in S$. Hence, g is a bijection between S and S_0 , so $|S| = |S_0|$.

Conversely, if S is finite and $S_0 \subseteq S$ is a subset such that $S_0 \neq S$, then $|S| = n$ for some $n \in \mathbb{N}$ and S_0 has strictly less elements than S , so $|S_0| = k$ for some $k < n$ and hence $|S_0| < |S|$.