

---

# Hand In 2

## FYSC20

---

*Author*

Max Eriksson

maxerikss@gmail.com

Lund University  
Department of Physics



**LUND**  
UNIVERSITY

November 23, 2024

**Problem:** A spherical shell of radius  $R_a$  carries a surface charge  $\sigma = \alpha \cos \theta$ .

- (a) Calculate its dipole moment with respect to the center of the sphere.
- (b) Calculate the approximate potential at points far away from the sphere.
- (c) The exact solution is given by

$$V(r, \theta) = \frac{\alpha R_a^3}{3\epsilon_0 r^2} \cos \theta \quad (1)$$

What does this imply for the higher poles?

**Solution:** We start by drawing a sketch seen in Fig. 1.




Figure 1: Sketch of the problem setup.

Since we are looking at a spherical shell it is reasonable to use spherical coordinates, as can be seen in the sketch. The cartesian axes are shown as well.

**(a) Solution:** The dipole moment  $\mathbf{p}$  of a continuous distribution centered at the origin is

$$\mathbf{p} = \int_V \mathbf{r}' \rho(\mathbf{r}') d\tau', \quad (2)$$

where  $\rho$  is the volume charge density, and  $V$  is the volume we are integrating over. In our case, since the charge is constricted on the surface, we can use a delta function.

$$\rho(\mathbf{r}) = \delta(r - R_a)\sigma = \delta(r - R_a)\alpha \cos \theta. \quad (3)$$

Using the fact that the vector  $\mathbf{r}$  and the volume element  $d\tau$  in spherical coordinates is  $\mathbf{r} = r\hat{\mathbf{r}}$  and  $d\tau = (dr)(rd\theta)(r \sin \theta d\varphi)$  we can rewrite Eq. (2).