

Metal Accumulation and Toxicity in the Submerged Aquatic Macrophyte *Vallisneria americana*

Author(s): Gretchen K. Bielmyer-Fraser, Ruth Adeyemi and Blessing Akintunde

Source: *Florida Scientist*, 2022, Vol. 85, No. 1 (2022), pp. 10-25

Published by: Florida Academy of Sciences, Inc.

Stable URL: <https://www.jstor.org/stable/10.2307/27186335>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



Florida Academy of Sciences, Inc. is collaborating with JSTOR to digitize, preserve and extend access to *Florida Scientist*

JSTOR

Metal Accumulation and Toxicity in the Submerged Aquatic Macrophyte *Vallisneria americana*

Gretchen K. Bielmyer-Fraser, Ruth Adeyemi, and Blessing Akintunde

Department of Chemistry, Jacksonville University, 2800 University Blvd N., Jacksonville, FL 32211

Abstract Submerged aquatic vegetation (SAV) are ecologically important plant species, providing habitat and food for many aquatic organisms and enhancing water quality. Anthropogenic practices have increased the input of metal pollutants in aquatic systems. SAV are known to accumulate high concentrations of metal pollutants yet less is known about the toxic effects of metals on SAV. In this study, we examined metal accumulation, distribution, and photosynthetic toxicity in the commonly found SAV species *Vallisneria americana* after exposure to copper, cadmium, and lead in the laboratory. Metal content in the root and shoot system of *V. americana* was quantified and photosynthetic parameters were measured after exposure to three exposure concentrations (control, 10 and 50 µg/L) of copper, cadmium, and lead for 4 d. Significant metal accumulation in both leaf and root tissue was observed over time in *V. americana* and the distribution varied with exposure time, concentration, and metal. Photosynthetic impairment corresponded with increasing metal accumulation in *V. americana* with toxicity occurring in the 10 µg/L lead treatment. These findings could be used to help assess tolerance of *V. americana* in aquatic environments impacted by metals.

Keywords: SAV, metals, accumulation, toxicity, *Vallisneria americana*, photosynthesis.

Introduction

Submerged aquatic vegetation (SAV) are important plant species in both freshwater and brackish ecosystems (Pinto et al. 2019; Best 1981). SAV provide vital habitat for many aquatic organisms and food for waterfowl and marine mammals (Best 1981, White et al. 2002). SAV help to anchor sediments, reduce turbidity, and protect organisms from wave energy (Scheffer and Van Nes 2007, Poirrier et al. 2017). Further, SAV enhance water quality with light availability, aid in carbon and nutrient cycling, and absorb contaminants from the environment (Barnett and Schneider 1974, Laughlin 1982, Hauxwell et al. 2004, Li et al. 2007, Scheffer and Van Nes 2007, Camp et al. 2012, Poirrier et al. 2017). While SAV are crucial to the survival and health of many aquatic ecosystems, they have substantially declined over the past 50 years (Li et al. 2007, Orth et al. 2010, Pyati et al. 2012, Goldberg et al. 2018, Pinto et al. 2019, Goldberg and Trent 2020). Pollution, sedimentation, and deteriorating water quality are contributors to this phenomenon, which is exacerbated by the decline in SAV biomass (Alexander et al. 1993, Li et al. 2007, Goldberg and Trent 2020).

Corresponding Author: Gretchen K. Bielmyer-Fraser, Email: gbielmy@ju.edu

Metals enter waterbodies naturally through the weathering of rock, and more substantially via anthropogenic sources like industrial effluents, domestic sewage, stormwater runoff from urban and agricultural lands, and as leachate from metal-based antifoulant paints (Bryan 1974, Guzmán and Jiménez 1992, Sundberg 1998, Jones 2010, Tchounwou 2012). Increased metal discharge into aquatic systems can lead to contamination of the water column and sediments, and cause toxicity to aquatic organisms when levels exceed acceptable limits (Nayar et al. 2004, Jaishankar et al., 2014, Jarvis and Bielmyer-Fraser 2015, Rani Das et al. 2017).

Many aquatic macrophytes have the capacity to accumulate significant concentrations of metals from water and sediment (Keskinan et al. 2004, Peng et al. 2008, Xue et al. 2010, Harguinteguy et al. 2014, Jarvis and Bielmyer-Fraser 2015, Bai et al. 2018). The translocation of metals in plants can be monitored by comparing the root-to-shoot uptake of metals, assessing metal transport, and quantifying distribution of metals in the root and shoot system of aquatic plants (Page et al. 2006). Aquatic macrophytes have been used in remediation of metal-contaminated sites to remove metals and enhance water quality (Miretzky et al. 2006). High biomass of SAV can reduce resuspension of metals from sediment (Zhang et al. 2016). However, if metal concentrations in the aquatic environment exceed certain thresholds, SAV abundance will decline (Keskinan et al. 2004). While metal accumulation in SAV has been well documented in field studies, fewer studies have focused on metal accumulation, distribution, and toxicity in SAV in a controlled laboratory environment. In field situations, measured toxic responses in organisms may be due to a variety of individual or a combination of environmental stressors. Elucidating metal-induced toxicity in the field can be challenging, which prompts the need for more laboratory experiments.

Responses of aquatic plants to metals include increased phytochelatin and chlorophyll production, antioxidant responses, decreased photosynthetic capabilities, and decreased growth (Baker and Walker 1989, Prasad and Strzałka 1999, Wang et al. 2008, Jarvis and Bielmyer-Fraser 2015). Chlorophyll fluorescence has been used to measure photosynthetic parameters in aquatic plants (Jarvis and Bielmyer-Fraser 2015, Siddiqui and Bielmyer-Fraser 2019). Fluorometry is a technique used to differentiate among photon energy captured by a chlorophyll-*a* pigment molecule used to drive photosynthesis, the energy emitted as fluorescence, and energy converted to heat (Schreiber and Bilger 1995, Rosenqvist and Van Kooten 2003). Measuring sub-lethal responses of plants to contaminants can be useful to assess plant health before they decline (Miller et al. 2017). Laboratory toxicity studies can effectively assess organism responses to specific contaminants and results can be applied to field situations.

The objective of this study was to measure metal accumulation, distribution, and toxicity in the perennial submerged aquatic macrophyte, *Vallisneria spiralis* (commonly called wild celery, eelgrass and tape grass) is distributed throughout North America ranging from Canada southward to Texas and Florida in both non-tidal and tidal regions (Moore 2009, USDA NRCS 2020). *V. spiralis* can tolerate a range of salinity from freshwater to brackish (<10 ppt) and are commonly found

throughout river and estuarine systems (Lacoul and Freedman 2006, Dobberfuhl 2007, Sagan 2007, Goldberg et al. 2018). In several waterbodies, abundance of *V. americana* is indicative of ecosystem health and it has been used to monitor sediment-bound metal concentrations in the environment (St-Cyr and Campbell 2000).

Materials and Methods

Experimental organism. *V. americana* was purchased from Buceplant and maintained at 25 °C in a large tank filled with synthetic hard freshwater and continuous aeration. The plants were acclimated to experimental conditions for 48 hours prior to use.

Experimental solutions. Synthetic hard freshwater was made by adding 120 mg/L CaSO₄, 8 mg/L KCl, 120 mg/L MgSO₄, and 192 mg/L NaHCO₃ to ultra-pure Milli-Q water following US EPA guidelines (USEPA 2002). The water was mixed for at least 24 h prior to use (Bielmyer et al. 2012). *V. americana* have been found in aquatic systems with this level of water hardness (100-120 mg CaCO₃/L) (Bielmyer-Fraser et al. 2020). Using synthetic freshwater in controlled toxicity experiments reduces the influence of other variables like contaminants and dissolved organic carbon which could alter metal toxicity (Lauren and McDonald 1986; Playle et al. 1992; Bielmyer-Fraser et al. 2018). Certified stock solutions of 1000 µg/mL copper, as Cu(NO₃)₂, lead, as Pb(NO₃)₂ and cadmium as Cd(NO₃)₂ (Perkin Elmer, Norwalk, CT) were used to make 10 and 100 µg/L Cu, 10 and 100 µg/L Pb, and 10 and 100 µg/L Cd. The metal solutions were transferred into 2-L plastic beakers, each containing a plastic mesh, and equilibrated in their respective containers for 24 h.

Experimental design. There were seven treatments, including the control, each with three replicates and all experimental beakers were arranged in a randomized design. Temperature was maintained at 24.2±0.56 °C using a water bath and recirculating chiller/heater (Model MC-1/4HP, AquaEuro Systems, Los Angeles, California) throughout the 4-day static-renewal experiment. A 16-h light: 8-h darkness cycle was maintained with cool white fluorescent lighting.

At the start of the experiment, one *V. americana* plant was added to each beaker and the plastic mesh was used to anchor the plant. Sediments were not used in this experiment to better characterize direct metal uptake by the plant. A 10 mL water sample was collected daily from each replicate, and the solution was filtered using a 0.45 µm filter, acidified with 50 µL trace metal grade HNO₃, and stored in a 15 mL polypropylene centrifuge tube. New solutions were made, equilibrated for 24 h, and used to renew the experimental solutions throughout the experiment. Three root and three leaf samples were collected from plants in the acclimation tank at the start of the experiment to be analyzed for background metal concentration. One leaf sample was then collected from each replicate at 2 and 4 d in the same manner. Additionally, at the end of the experiment, root samples were also collected for metal analysis. The plant tissue samples were measured for wet mass, dried in an oven at 60°C for 24 h, and then weighed to determine dry mass. Samples were then fully digested with addition of trace metal grade HNO₃ and heated in a 60°C hot water bath for 3 h.

Daily, temperature and dissolved oxygen (DO) were measured using a YSI meter (Model YSI-85; Pentair Aquatic Ecosystems, Apopka, Florida), pH was measured using a calibrated pH meter (WTW, Xylem Inc.), and light intensity was measured using a lux light meter (Extech, Nashua, NH). No algal growth was observed in any of the experimental beakers throughout the length of the experiment. At the end of the experiment, the mean ± standard deviation of DO, temperature, pH, and light intensity values were calculated (Table 1). Following American Public Health Association guidelines, hardness and alkalinity of the base water were measured using titrimetric methods (method 2340 and 2320, respectively; APHA 1985; Table 1).

Photosynthetic parameters. In each replicate, *V. americana* were dark adapted for 30 min and then measured twice in the same location on the leaf for a variety of parameters using a modulated chlorophyll fluorometer (OS5p+; OptiScience, Hudson, NH) and a rapid light curve test (one minute illumination with saturating pulses at every 10 seconds; Genty et al. 1989, Ralph et al. 1999, Beer and Bjork 2000). At 0, 1, 2, and 4 d, the effective quantum yield of photosystem II (amount of energy used

Table 1. Water chemistry (mean ± standard deviation) in experimental solutions. Parameters were measured daily except for alkalinity and hardness which were measured at the start of the experiment in the base water.

Parameter	Measured Value
Dissolved Oxygen	7.78 ± 0.33 mg/L
Temperature	24.2 ± 0.56 °C
pH	8.60 ± 0.20
Light illumination	14501 ± 4628 lux
Alkalinity	128 mg CaCO ₃ /L
Hardness	118 mg CaCO ₃ /L

for photosynthesis; YPSII), the maximum electron transport rate (ETR_{max}), the minimum amount of light necessary to saturate the system (I_k), and the efficiency of the system (α) were measured in the leaves of *V. americana*.

Metal analysis. All the digested root and leaf samples were diluted using ultra-pure 18 mΩ Milli-Q water. The diluted tissue and water samples were analyzed to determine cadmium, copper, and lead concentrations using an atomic absorption spectrophotometer with graphite furnace detection (AAAnalyst 800, Perkin Elmer, Norwalk, CT). Certified metal standards dissolved in 2% HCl, ultra-pure Milli-Q water blanks, and QA/QC samples were analyzed throughout each analysis (detection limits: 0.1 µg/L cadmium, 0.7 µg/L copper, 0.7 µg/L lead). The plant tissue data are presented as µg metal/g dry weight.

Statistical analysis. At the end of the exposure, data from day 4 were analyzed using one-way analysis of variance (one-way ANOVA) to determine differences among treatments. Normality and equality of variance of the data were tested using a Shapiro-Wilk test and Levene’s test, respectively. Because the data were not normally distributed, a Kruskal-Wallis one-way ANOVA on ranks was used followed by a Tukey’s test to determine pairwise comparisons (p ≤ 0.05). The same procedures were used to analyze tissue metal concentrations and photosynthetic data over time.

Results

Experimental conditions. The water chemistry in the experimental solutions is presented in Table 1. The values remained stable and were within acceptable conditions. The measured metal concentrations in the experimental solutions were similar or just below nominal concentrations with one exception (50 µg/L cadmium treatment; Table 2). The highest Cd treatment was approximately double the nominal value, however, the concentration remained stable during the experiment.

Table 2. Measured metal concentrations (mean ± standard deviation) in experimental solutions.

Treatment	Measured Metal Concentration (µg/L)
Control - Cd	0.04 ± 0.04
Control - Cu	1.35 ± 0.98
Control - Pb	2.77 ± 1.51
10 µg/L Cd	9.43 ± 4.27
50 µg/L Cd	122 ± 25.6
10 µg/L Cu	7.20 ± 2.46
50 µg/L Cu	38.5 ± 8.23
10 µg/L Pb	6.91 ± 1.11
50 µg/L Pb	43.6 ± 9.39




Figure 1. A. copper (Cu), B. cadmium (Cd), and C. lead (Pb) accumulation in the leaves of *V. americana* exposed to waterborne metal over 4 d. See Table 2 for measured metal concentrations. Asterisks denote significant difference from the control at a particular time point.

Metal accumulation. Metal significantly accumulated in the leaves of *V. americana* over the duration of the experiment ($P=0.005$; Figure 1). For all metals, accumulation was highest after 2 d of exposure with significant differences observed in leaves of both treatments of cadmium and copper and the highest lead




Figure 1. Continued.

treatment, as compared to control values (Figure 1). Metal accumulation in leaves of both cadmium treatments decreased slightly by 4 d, but cadmium and copper accumulation in both treatments remained significantly elevated, as compared to the control (Figure 1A, B). Alternatively, the lead concentration in the leaves of *V. americana* exposed to the highest lead treatment returned to control levels by 4 d (Figure 1C). The metal accumulation in leaves of the 10 µg/L lead treatment did not significantly differ from control values over the 4-d exposure.

The *V. americana* in the control treatments had no detectable cadmium or lead in the leaves or roots, however, the mean copper concentration in the leaves and roots of the control group were 29.1 and 24.1 µg/g dw, respectively (Figures 1, 2). At the end of the 4-d exposure, root copper concentration was significantly elevated in both copper treatments above control levels, similar to the pattern of copper accumulation in the leaves (Figure 2). No significant differences were detected in the root cadmium concentration in the 10 and 50 µg/L treatments as compared to controls. Alternatively, the root lead concentration was significantly elevated in the 10 µg/L treatment, as compared to control levels (Figure 2).

Photosynthetic parameters. The exposure of *V. americana* to elevated metal concentrations showed notable changes in the photosynthetic parameters (Figure 3). The control values for all parameters did not significantly vary over the duration of the experiment or as compared to background levels (Figure 3). The YPSII and ETR_{max} in *V. americana* were significantly increased after 2 d of exposure to 10 µg/




Figure 2. Cadmium, lead and copper concentrations in the tissues (leaves and roots) of *V. americana* exposed to control, 10, and 50 µg/L waterborne metal for 4 d. See Table 2 for measured metal concentrations. Asterisks indicate a significant difference from the control within a specified tissue type.

L copper (Figure 3A, B). Alternatively, a significant decrease in YPSII was observed in the 50 µg/L copper, 50 µg/L cadmium, and 10 µg/L lead treatments after 4 d of exposure (Figure 3). The Ik was significantly elevated in the 50 µg/L copper and 10 µg/L lead treatments and α was significantly decreased in the 10 µg/L lead treatment (Figure 3C, D).




Fig. 3. A. effective quantum yield of photosystem II (YPSII), B. maximum electron transport rate (ETR_{max}), C. Ik (minimum amount of saturating light), and D. α (efficiency of photosystem II) in *V. americana* after exposure to control, 10 and 100 µg/L Cu, Pb, and Cd over 4 d. Asterisks denote significant difference from the control at a particular time point. "Background" denotes measurements taken at the start of the experiment prior to metal exposure.

Table 3. Shoot: root ratio for *V. americana* exposed to 10 and 100 µg/L cadmium, copper, or lead for 4 d.

Concentration	Lead	Copper	Cadmium
10 µg/L	0.24	1.11	1.74
100 µg/L	1.10	1.15	1.55

Discussion

Metal accumulation. The tissue metal concentrations in *V americana* in this study were similar to those reported in field monitoring studies in various rivers (St-Cyr and Campbell 1994, Hudon et al. 1998, St-Cyr and Campbell 2000). Mean concentrations of lead and copper in various aquatic plants were reported to be 11 and 48 µg/g dw, respectively (Hutchinson 1975). Bielmyer-Fraser et al. (2017) reported plant tissue concentrations of 53–798 µg/g dw copper, 3–95 µg/g dw cadmium, and 7–330 µg/g dw lead from a variety of macrophytes collected from two rivers with varying anthropogenic impact in southern Georgia. The lead concentration in leaf tissues of *Thlaspi praecox* collected from a heavily polluted site near a lead mine was 45 µg/g dw (Vogel-Mikus et al. 2008). Hudon (1980) reported 9.3–28.9 µg/g dw copper, 0.62–3.50 µg/g dw cadmium, and 0.85–2.68 µg/g dw lead in the above ground tissues of *V. americana* collected from St. Lawrence and Ottawa rivers in Canada. St-Cyr and Campbell (1994) reported concentrations of 9.1–39.6 µg/g dw copper, 0.35–5.12 µg/g dw cadmium, and 1.2–8.0 µg/g dw lead in the above-ground parts of *V. americana* collected from Lake St. Pierre, part of the St. Lawrence River. Further research by St-Cyr and Campbell (2000) showed concentrations of 6.2–35.4 and 4.4–39.4 µg/g dw copper, 0.28–4.3 and 0.07–2.35 µg/g dw cadmium, and 0.6–10.9 and 3.2–20.2 µg/g dw lead in the leaves and roots, respectively, of *V. americana*.

Wang et al. (2010) reported concentrations of 396 and 114 µg/g dw copper and 63.8 and 48.0 µg/g dw cadmium in the roots and shoots, respectively, of *V. spiralis* exposed to metal-contaminated sediments for 21 d. *V. spiralis* has been shown to rapidly accumulate copper, cadmium, and lead in the laboratory with copper accumulating evenly between the leaves and the roots; cadmium accumulating to a greater extent in the leaves, and lead accumulating to a higher degree in the roots of *V. spiralis* (Gupta and Chandra 1993, Sinha et al. 1994, Wang et al. 2008). After 4 d of metal exposure in the present study, using similar exposure conditions (e.g. without sediment), a similar pattern of accumulation was observed in *V. americana*. Likewise, Welsh and Denny (1980) reported a similar distribution pattern as compared to the present study for copper and lead in a variety of aquatic macrophytes in the field (English lakes) with shoot: root ratios ranging from 0.74–1.01 for copper and 0.17–0.25 for lead. In this study, shoot: root ratio ranged from 1.11–1.15 for copper, 0.24–1.10 for lead, and 1.55–1.74 for cadmium (Table 3). Metal distribution generally depends on the metal, the exposure concentration, and the plant species (Jarvis and Bielmyer-Fraser 2015, Bai et al. 2018).

Rooted submerged macrophytes can naturally absorb elements through their leaves from the surrounding water (Welsh and Denny 1980) and through their roots

from the sediment interstitial water (St-Cyr and Campbell 2000). Once absorbed, elements can be distributed to various tissues, dependent on biological requirements and/or detoxification (Ximenenz-Embun et al. 2002, Page et al. 2006, Horník et al. 2007). Copper is an essential element necessary for a variety of metabolic processes, including acting as a cofactor for many enzymes (Clemens et al. 2002). Cadmium and lead are nonessential and are therefore presumed in most cases to enter plant cells through essential transporters for other elements, such as zinc (Vogel-Mikus et al. 2008). Welsh and Denny (1980) reported two different pathways for copper and lead transfer from sediments to the shoot of various taxa of submerged aquatic macrophytes. Copper accumulation in these macrophytes was thought to occur primarily by absorption through the roots, translocation to the shoots, and re-translocation to the roots. Copper adsorption into the shoots from the water occurred to a lesser extent (Welsh and Denny 1980). Alternatively, lead accumulation in the shoots mainly occurred through adsorption from the surrounding water (Welsh and Denny 1980). Further, uptake of lead from the water occurred rapidly (Welsh and Denny 1980).

In this study, rapid lead accumulation in the leaves of *V. americana* was observed in the 50 µg/L treatment after 2 d of exposure. However, lead levels also rapidly declined in the leaves by 4 d of exposure. This observed accumulation may have been due to adsorption of lead from the surrounding water, similar to the observations of Welsh and Denny (1980). The significantly increased lead accumulation in the root tissue of *V. americana* exposed to 10 µg/L lead was likely due to uptake via the roots. It is important to note that the leaves and roots of *V. americana* exposed to 10 µg/L had higher lead concentration than those exposed to 50 µg/L lead after 4 d of exposure. Although not significant, the same pattern was observed with cadmium exposure. Jarvis and Bielmyer-Fraser (2015) observed a similar phenomenon in the seaweed, *Ulva lactuca*, after metal exposure. The decreased metal uptake with increasing exposure concentration was attributed to downregulation of specific metal transporters (Wang and Dei 1999, Jarvis and Bielmyer-Fraser 2015). Given that seaweed do not have root systems, further investigation is needed to assess this phenomenon in SAV, as this may be an important consideration if using *V. americana* for metal bioremediation.

For the St. Lawrence River, St-Cyr and Campbell (2000) concluded that roots of *V. americana* were better bioindicator organs than shoots for monitoring metal contamination in sediments. In addition to biota, sediments generally act as a sink for metals entering aquatic systems and therefore concentrations are generally higher in the sediments than the water column (Hall 1989, DiToro et al. 2001, Nayer et al. 2004, Pinto et al. 2019).

Martinez and Shu-Nyamboli (2011) also showed significant correlations between aquatic macrophytes and sediment bioavailable metal concentrations in the Gallinas River. Further, significantly higher metal concentrations were reported in the plant roots as compared to the shoots (Martinez and Shu-Nyamboli 2011). Sediment chemistry can influence the bioavailability of metals (Zhang et al. 2014). In the present study, sediment was eliminated from the metal exposure scenario, to assess metal uptake directly from the water into the plant tissue. It should be noted

that the presence of sediment and the anoxic interstitial water microenvironment could change the metal uptake kinetics in *V. americana* (Zhang et al. 2014). Additionally, developmental stage of the plant and type of root system have been shown to affect metal accumulation (Page et al. 2006). Several species of aquatic plants have mechanisms to protect against metal toxicity including rapid translocation and detoxification strategies (Geng et al. 2019); however, at elevated levels metals may exert toxic effects.

Metal toxicity. Excess copper, cadmium, and lead accumulation in plants can cause physiological effects and growth inhibition in a variety of species (Pålsson 1989, Fernandes and Henriques 1991, Wang et al 2008, 2010, 2013, Jarvis and Bielmyer-Fraser 2015, Zhu et al. 2016). Reduction of chlorophyll in aquatic plants after copper, cadmium, and lead exposure has been shown in several studies (Gupta and Chandra 1993, Prasad et al. 2001, Moferran et al. 2009, Zhu et al 2016). For example, Wang et al. (2010) reported a decreased chlorophyll concentration in *V. spiralis* exposed to copper- and cadmium-contaminated sediments. Cadmium and lead exposure to the plant *Hydrilla verticillata* has been shown to inhibit chlorophyll synthesis enzymes (Singh et al. 2013). Zhu et al. (2016) reported reduced chlorophyll and reduced relative growth rate in *V. natans* exposed to copper-contaminated sediments. Further, the roots of *V. natans* were significantly damaged including stunted roots and poor growth initiation after exposure to increased copper in the sediments (Zhu et al. 2016).

In the current study, increased copper, cadmium and lead exposure to *V. americana* caused decreased YPSII. Decreased photosynthesis has been a reported consequence of metal exposure in other plant species as well (Jarvis and Bielmyer-Fraser 2015, Siddiqui and Bielmyer-Fraser 2019) and is likely a result of the reduction in chlorophyll, as previously mentioned. The decreased energy used for photosynthesis could have been used to upregulate production of phytochelatin (peptides that sequester metals) and antioxidant molecules (e.g. glutathione) which are used for metal detoxification (Wang et al. 2008, Main et al. 2010, Singh et al. 2013, Brock and Bielmyer 2013, Patel and Bielmyer-Fraser 2015). Wang et al. (2008) observed increased cadmium accumulation in leaves and roots, increased phytochelatin concentrations, increased glutathione concentrations, and decreased growth in *V. spiralis* exposed to 18–72 µg/L cadmium for 7 d as compared to controls. Similarly, decreased YPSII was observed in *V. americana* exposed to 50 µg/L cadmium for 4 d in the present study.

Tissue metal concentration correlated with toxicity in *V. americana*. Specifically, the treatments with the highest metal concentrations in the roots (50 µg/L copper and 10 µg/L lead) had the most severe impairment of photosynthetic parameters by 4 d. The 50 µg/L copper and cadmium treatments and 10 µg/L lead treatment had a lower YPSII which corresponded to decreased photosynthetic efficiency and an increase in the minimum amount of light necessary to saturate photosystems in *V. americana*. Decreased photosynthetic capability was likely a result of both metal accumulation in the leaf tissues (where photosynthesis occurs), in addition to metal accumulation in the root system and possibly impairment of water and nutrient uptake (Pålsson 1989, Zhu et al. 2016).

Interestingly, a hormesis effect was observed after 2 d exposure to 10 µg/L copper, where YPSII and ETR_{max} were increased in the *V. americana*. Copper plays an essential role in electron transfer reactions in photosynthesis and may be beneficial in low concentrations over short-term exposure (O'Halloran and Culotta 2000, Clemens 2001, Clemens et al. 2002). However, in excess, the redox activity of copper may lead to the generation of damaging reactive oxygen species and declines in photosynthetic efficiency (Clemens et al. 2002).

This laboratory study demonstrated metal- and concentration-dependent differences in metal accumulation and distribution in *V. americana* as well as photosynthetic toxicity at environmentally realistic metal concentrations. SAV have been used to monitor metals and in bioremediation of metal-contaminated environments (Gupta and Chandra 1998). *V. americana* has the capacity to accumulate copper, cadmium, and lead from surrounding waters and these results highlight the utility of this species for both monitoring and remediation efforts. *V. americana* may be useful for bioremediation of copper- and cadmium-contaminated sites especially, given that many polluted areas have waterborne copper and cadmium concentrations below the 50 µg/L, which was the lowest concentration causing photosynthetic impairment in this study (Bielmyer-Fraser et al. 2020). Furthermore, this information could be used to help manage and protect SAV in aquatic systems impacted by metals.

Acknowledgements The fluorometer was funded by a Jacksonville University EPIC grant awarded to G.K. Bielmyer-Fraser. The authors would also like to thank the Jacksonville University Chemistry department for helping to fund this research and Dr. Nisse Goldberg for constructive scientific dialogue.

References

- Alexander C, Smith R, Calder F, Schropp S, Windom H. 1993. The historical record of metal enrichment in two Florida estuaries. *Estuaries* 16:627–637.
- APHA. 1985. Standard methods for the examination of water and wastewater, 16 ed. American Public Health Association.
- Bai Li, Xiao-Long L, Hu J, Li J, Wang Z-L, Han G, Li S-L, Liu C-Q. 2018. Heavy metal accumulation in common aquatic plants in rivers and lakes in the Taihu Basin. *International Journal of Environmental Research and Public Health* 15:2857.
- Baker AJM, Walker PL. 1989. Physiological responses of plants to heavy metals and the quantification of tolerance and toxicity. *Chemical Speciation and Bioavailability* 1:7–17.
- Barnett BS, Schneider RW. 1974. Fish populations in dense submersed plant communities. *Hyacinth Control Journal* 12:12–14.
- Beer S, Björk M. 2000. Measuring rates of photosynthesis of two tropical seagrasses by pulse amplitude modulated (PAM) fluorometry. *Aquatic Botany* 66:69–76.
- Best RC. 1981. Food and feeding habits of wild and captive sirenia. *Mammal Review* 11:3–29.
- Bielmyer-Fraser GK, Arnold WR, Isely JJ, Klaine SJ, Tomasso J. 2012. Effects of roof and rainwater characteristics on copper concentrations in roof runoff. *Environmental Monitoring and Assessment* 184:2797–2804.
- Bielmyer-Fraser GK, Harper B, Picariello C, Albritton-Ford A. 2018. The influence of salinity and water chemistry on acute toxicity of cadmium to two euryhaline fish species. *Comparative Biochemistry and Physiology C* 214:23–27.

- Bielmyer-Fraser GK, Llazar K, Ramirez J, Ward A, Santiago F. 2020. Spatial and Temporal Water Quality Assessment in the Lower St. Johns River, Florida. *Southeastern Naturalist* 19(3), 582–600.
- Bielmyer-Fraser, GK, Waters MN, Duckworth CG, Patel P, Webster B, Blocker A, Crummey CH, Duncan A, Nwokike S, Picariello C, Ragan JT, Schumacher EL, Tucker R, Tuttle E, Wiggins C. 2017. Assessment of metal contamination in the biota of four rivers experiencing varying degrees of human impact. *Environmental Monitoring and Assessment* 189:23.
- Brock, J., G.K. Bielmyer. 2013. Metal accumulation and sublethal responses in the sea anemone, *Aiptasia pallida* after waterborne exposure to metal mixtures. *Comparative Biochemistry and Physiology Part C* 158:150–158.
- Bryan GW. 1974. Some aspects of heavy metal tolerance in aquatic organisms. Pp. 7–34 in Lockwood APM, ed. *Effects of Pollutants on Aquatic Organisms*. Cambridge University Press, Cambridge.
- Camp EV, Gwinn DC, Pine WE III, Frazer TK. 2012. Changes in submersed aquatic vegetation affect predation risk of a common prey fish, *Lucania parva* (Cyprinodontiformes: Fundulidae), in a spring-fed coastal river. *Fisheries Management and Ecology* 19:245–251.
- Clemens S, Palmgren MG, Kramer U. 2002. A long way ahead: understand and engineering plant metal accumulation. *Trends in Plant Science* 7:309–315.
- Clemens S. 2001. Molecular mechanisms of plant metal homeostasis and tolerance. *Planta* 212:475–486.
- DiToro DM, Allen HE, Bergman HL, Meyer JS, Paquin P, Santore RC. 2001. Biotic ligand model of the acute toxicity of metals. 1. Technical basis. *Environmental Toxicology and Chemistry* 20:2383–2396.
- Dobberfuhl DR. 2007. Light-limiting thresholds for submerged aquatic vegetation in a blackwater river. *Journal of Aquatic Botany* 86:346–352.
- Fernandes JC, Henriques FS. 1991. Biochemical, physiological, and structural effects of excess copper in plants. *The Botanical Review* 57:246–273.
- Geng N, Wu Y, Zhang M, Tsang DCW, Rinklebe J, Xia Y, Lu D, Zhu L, Palansooriya KN, Kim K-H, Ok S. 2019. Bioaccumulation of Potentially Toxic Elements by Submerged Plants and Biofilms: A Critical Review. *Environment International* 131:105015.
- Genty B, Briantais JM, Baker NR. 1989. The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochimica et Biophysica Acta* 990:87–92.
- Goldberg NA, Trent T, Hendrickson J. 2018. Temporal and spatial changes in *Vallisneria americana* Michaux (Tape-grass) beds in the Lower St. Johns River, Florida, from 2002–2011. *Southeastern Naturalist* 17:396–410.
- Goldberg NA, Trent T. 2020. Patterns in submerged aquatic vegetation in the lower St. Johns River, Florida, from 2001 to 2019. *Journal of Aquatic Plant Management* 58:000–000.
- Gupta M, Chandra P. 1994. Lead accumulation and toxicity in *Vallisneria spiralis* (L.) and *Hvdrilla verticillata* (L.f.) Royle, *Journal of Environmental Science and Health Part A: Environmental Science and Engineering and Toxicology* 29:503–516.
- Guzmán HM, Jiménez CE. 1992. Contamination of coral reefs by heavy metals along the Caribbean coast of Central America. *Marine Pollution Bulletin* 24:554–561.
- Hall L. 1989. The effects of dredging and reclamation on metal levels in water and sediments from an estuarine environment off Trinidad, West Indies. *Environmental Pollution* 56:189–207.
- Harguinteguy CA, Cirelli AF, Pignata ML. 2014. Heavy metal accumulation in leaves of aquatic plant *Stuckenia filiformis* and its relationship with sediment and water in the Suquia river (Argentina). *Microchemical Journal* 114:111–118.
- Hauxwell J, Frazer TK, Osenberg CW. 2004. Grazing by manatees excludes both new and established Wild Celery transplants: Implications for restoration in Kings Bay, FL, USA. *Journal of Aquatic Plant Management* 42:49–53.
- Horník M, Pipiška M, Sekáčová J, Augustín J. 2007. Determination of long distance transport of Cs⁺, Co²⁺ and Zn²⁺ ions in vascular plants by autoradiography and gamma-spectrometry. *Nova Biotechnologica* 7:33–40.

- Hudon C. 1998. Metal accumulation in American wild celery (*Vallisneria americana* Michx.) in the St. Lawrence River: effects of water depth and exposure to current. *Canadian Journal of Fisheries and Aquatic Sciences* 55:2317–2328.
- Hutchinson GE. 1975. A Treatise on Limnology, Vol. 3. Limnological Botany. Wiley, New York.
- Jaishankar M, Tseten T, Anbalagan N, Mathew BB, Beeregowda KN. 2014. Toxicity, mechanism and health effects of some heavy metals. *Interdisciplinary Toxicology* 7:60–72.
- Jarvis T, Bielmyer-Fraser G. 2015. Accumulation and effects of metal mixtures in two seaweed species. *Comparative Biochemistry and Physiology* 171:28–33.
- Jones R. 2010. Environmental contamination associated with a marine landfill ('seafill') beside a coral reef. *Marine Pollution Bulletin* 60:1993–2006.
- Keskinkan O, Goksu M, Basibuyuk M, Forster C. 2004. Heavy metal adsorption properties of a submerged aquatic plant (*Ceratophyllum demersum*). *Bioresource Technology* 92:197–200.
- Lacoul P, Freedman B. 2006. Environmental influences on aquatic plants in freshwater ecosystems. *Environmental Reviews* 14:89–136.
- Laughlin RA. 1982. Feeding habits of the Blue Crab, *Callinectes sapidus* Rathbun, in the Apalachicola estuary, Florida. *Bulletin of Marine Science* 32:807–822.
- Lauren D, McDonald D. 1986. Influence of water hardness, pH, and alkalinity on the mechanisms of copper toxicity in juvenile rainbow trout, *Salmo gairdneri*. *Canadian Journal of Fisheries and Aquatic Sciences* 43:1488–1496.
- Li X, Weller D, Gallegos C, Jordan T, Kim H. 2007. Effects of watershed and estuarine characteristics on the abundance of submerged aquatic vegetation in Chesapeake Bay subestuaries. *Estuaries and Coasts* 30:840–854.
- Main, WPL, Ross C, Bielmyer GK. 2010. Copper accumulation and oxidative stress in the sea anemone, *Aiptasia pallida*, after waterborne copper exposure. *Comparative Biochemistry and Physiology Part C* 151:216–221.
- Martinez EA, Shu-Nyamboli C. 2011. Determination of selected heavy metal concentrations and distribution in a southwestern stream using macrophytes. *Ecotoxicology and Environmental Safety* 74:1504–1511.
- Miller RJ, Muller EB, Cole B, Martin T, Nisbet R, Bielmyer-Fraser GK, Jarvis TA, Keller AA, Cherr G, Lenihan HS. 2017. Photosynthetic efficiency predicts toxic effects of metal nanomaterials in phytoplankton. *Aquatic Toxicology* 185:83–93.
- Miretzky P, Saralegui A, Fernández Cirelli A. 2006. Simultaneous heavy metal removal mechanism by dead macrophytes. *Chemosphere* 62:247–254
- Monferrán MV, Agudo JAS, Pignata ML, Wunderlin DA. 2009. Copper-induced response of physiological parameters and antioxidant enzymes in the aquatic macrophyte *Potamogeton pusillus*. *Environmental Pollution* 157:2570–2576.
- Moore K. 2009. Submerged Aquatic Vegetation (SAV) in the Lower St. Johns River and the Influences of Water Quality Factors on SAV. *St. Johns River Water Management District* 9:1–23
- Nayar S, Goh BPL, Chou LM. 2004. Environmental impact of heavy metals from dredged and resuspended sediments on phytoplankton and bacteria assessed in situ mesocosms. *Ecotoxicology and Environmental Safety* 59:349–369.
- O'Halloran TV, Culotta VC. 2000. Metallochaperones, an intracellular shuttle service for metal ions. *Journal of Biological Chemistry* 275:25057–25060.
- Orth RJ, Williams MR, Marion SR, Wilcox DJ, Carruthers TJ, Moore KA, Kemp WM, Dennison WC, Rybicki N, Bergstrom P, Batiuk RA. 2010. Long-term trends in submersed aquatic vegetation (SAV) in Chesapeake Bay, USA, related to water quality. *Estuarine and Coasts* 33:1144–1163.
- Page V, Weisskopf L, Feller U. 2006. Heavy metals in white lupin: uptake, root-to-shoot transfer and redistribution within the plant. *New Phytologist* 171:329–341.
- Påhlsson AMB. 1989. Toxicity of heavy metals (Zn, Cu, Cd, Pb) to vascular plants. *Water, Air, and Soil Pollution* 47:287–319.
- Patel P, Bielmyer-Fraser GK. 2015. The influence of salinity and copper exposure on copper accumulation and physiological impairment in the sea anemone, *Exaiptasia pallida*. *Comparative Biochemistry and Physiology C* 168:39–47.

- Peng K, Luo C, Lou L, Li, X, Shen, Z. 2008. Bioaccumulation of heavy metals by the aquatic plants *Potamogeton pectinatus* L. and *Potamogeton malaianus* Miq. and their potential use for contamination indicators and in wastewater treatment. *Science of the Total Environment* 392:22–29.
- Pinto G, Bielmyer-Fraser GK, Goldberg N, Ouellette A, Le A, Pyati R, Zoellner B, Closmann C. 2019. 2019 State of the River Report for the Lower St. Johns River Basin, Florida: Water Quality, Fisheries, Aquatic Life, and Contaminants (LSJR). City of Jacksonville, Environmental Protection Board.
- Playle R, Gensemer R, Dixon D. 1992. Copper accumulation on gills of fathead minnows: Influence of water hardness, complexation, and pH of the gill micro-environment. *Environmental Toxicology and Chemistry* 11:381–391.
- Poirrier MA, Caputo CE, Franze CD. 2017. Biogeography of submerged aquatic vegetation (SAV) in the Pontchartrain Basin: Species salinity zonation and 1953–2016 Lake Pontchartrain trends. *Southeastern Geographer* 57:273–293.
- Prasad M, Malec P, Waloszek A, Bojko M, Strzalka K. 2001. Physiological responses of *Lemna trisulca* L. (duckweed) to cadmium and copper bioaccumulation. *Plant Science* 161:881–889.
- Prasad M, Strzalka K. 1999. Impact of heavy metals on photosynthesis. Pp. 117–138 in Prasad M, ed. *Heavy Metal Stress in Plants: From Molecules to Ecosystems*. Springer, Berlin.
- Pyati R, Bielmyer GK, Chalk S, McCarthy D, McCarthy H, Pinto G, Sonnenberg L, Welsh P. 2012. Case Study: St. Johns River Basin, USA, Fourth United Nations World Water Development Report. World Water Assessment Programme. UNESCO Publishing, Paris.
- Ralph PJ, Gademan R, Larkun AWD, Schrieber U. 1999. In situ underwater measurements of photosynthetic activity of coral zooxanthellae and other reef dwelling dinoflagellate endosymbionts. *Marine Ecology Progress Series* 180:139–147.
- Rani Das P, Hossain M, Sarker B, Parvin A, Swarna Das S, Moniruzzaman M, Saha B. 2017. Heavy metals in farm sediments, feeds and bioaccumulation of some selected heavy metals in various tissues of farmed *Pangasius hypophthalmus* in Bangladesh. *Fisheries and Aquaculture Journal* 8:1–8.
- Rosenqvist E, Van Kooten O. 2003. Chlorophyll fluorescence: a general description and nomenclature. Pp. 31–74 in Deell J, Toivonen P, eds. *Practical Applications of Chlorophyll Fluorescence in Plant Biology*. Springer, Verlag.
- Sagan JJ. 2007. Interim reports associated with quarterly sampling and groundtruth surveys for the St. Johns River Water Management District, Palatka, FL. SAV Monitoring Project: Interim reports I–V.
- Scheffer M, Van Nes EH. 2007. Shallow lakes theory revisited: various alternative regimes driven by climate, nutrients, depth and lake size. *Hydrobiologia* 584:455–466.
- Schreiber U, Bilger W, Neubauer C. 1995. Chlorophyll fluorescence as a noninvasive indicator for rapid assessment of in vivo photosynthesis. Pp. 49–70 in Schulze E, Caldwell M, eds. *Ecophysiology of Photosynthesis* Springer, Berlin.
- Siddiqui S, Bielmyer-Fraser GK. 2019. Accumulation and effects of dissolved and nanoparticle silver and copper in two marine seaweed species. *Georgia Journal of Science* 77:1.
- Singh A, Kumar CS, Agarwal A. 2013. Effect of lead and cadmium on aquatic plant *Hydrilla verticillata*. *Journal of Environmental Biology* 34:1027–1031.
- St-Cyr L, Campbell PGC. 1994. Trace metals in submerged plants of the St. Lawrence River. *Canadian Journal of Botany* 72:429–439.
- St-Cyr L, Campbell PGC. 2000. Bioavailability of sediment-bound metals for *Vallisneria americana* Michx, a submerged aquatic plant, in the St. Lawrence River. *Canadian Journal of Fisheries and Aquatic Sciences* 57:1330–1341.
- Sundberg R. 1998. The fate of copper released from the Vasa shipyard museum. *Metallurgy* 52:230–231.
- Tchounwou PB, Yedjou CG, Patlolla AK, Sutton DJ. 2012. Heavy metal toxicity and the environment. *Experientia Supplementum* 101:133–164.
- U.S. EPA. 2002. Short-term methods for estimating the chronic toxicity of effluents and receiving waters to freshwater organisms, 4th ed. Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

- US Department of Agriculture Natural Resources Conservation Service (USDA NRCS) Plants National Database. 2017. Profile for *Vallisneria americana* Michx. <http://plants.usda.gov/> Accessed; July 3, 2020.
- Vogel-Mikus K, Regvar M, Mesjasz-Przybylowicz J, Przybylowicz WJ, Simcic J, Pelicon P, Budnar M. 2008. Spatial distribution of cadmium in leaves of metal hyperaccumulating *Thlaspi praecox* using micro-PIXE. *New Phytologist* 179:712–721.
- Wang WX, Dei RCH. 1999. Kinetic measurements of metal accumulation in two marinemacroalgae. *Marine Biology* 135:11–23.
- Wang C, Sun Q, Wang Q. 2008. Cadmium toxicity and phytochelatin production in a rooted-submerged macrophyte *Vallisneria spiralis* exposed to low concentrations of cadmium. *Environmental Toxicology* 24:271–278.
- Wang Q, Li Z, Cheng S, Wu Z. 2010. Influence of humic acids on the accumulation of copper and cadmium in *Vallisneria spiralis* L. from sediment. *Environmental Earth Sciences* 61:1207–1213.
- Wang P, Wang C, Ouyang P, Qian J, Shi R. 2013. Physiological responses of *Vallisneria spiralis* L. induced by different hydraulic conditions when exposed to copper and nitrogen. *African Journal of Biotechnology* 10:7441–7452.
- Welsh RPH, Denny P. 1980. The uptake of lead and copper by submerged aquatic macrophytes in two English lakes. *Journal of Ecology*. 68:443–455.
- White AQ, Pinto GF, Robison AP. 2002. Seasonal Distribution of Manatees, *Trichechus manatus latirostris*, in Duval County and Adjacent Waters, Northeast Florida. *Florida Scientist* 65:208–221.
- Ximénez-Embún P, Rodríguez-Sanz B, Madrid-Albarrán Y, Cámara C. 2002. Uptake of heavy metals by lupin plants in artificially contaminated sand: preliminary results. *International Journal of Environmental Analytical Chemistry* 82:805–813.
- Xue PY, Li GX, Liu WJ, Yan CZ. 2010. Copper uptake and translocation in a submerged aquatic plant *Hydrilla verticillata* (L.f.) Royle. *Chemosphere* 81:1098–1103.
- Zhang Y, Liu X, Qin B, Shi K, Deng J, Zhou Y. 2016. Aquatic vegetation in response to increased eutrophication and degraded light climate in Eastern Lake Taihu: Implications for lake ecological restoration. *Scientific Reports* 6:1–12.
- Zhang C, Yu Z-G, Zeng G-M, Jiang M, Yang Z-Z, Cui F, Zhu M-Y, Shen L-Q, Hu L. 2014. Effects of sediment geochemical properties on heavy metal bioavailability. *Environment International* 73:270-281.
- Zhu Z, Song S, Pengshan L, Jeelani N, Wang P, Yuan H, Zhang J, An S, Leng X. 2016. Growth and physiological responses of submerged plant *Vallisneria natans* to water column ammonia nitrogen and sediment copper. *Peer J* 4:e1953.

Submitted: May 19, 2021

Accepted: July 21, 2021