Package leaflet: Vesperil

Vesperil (ormelteon)

Read all of this document carefully before you start. Keep it: you may need to read it again. If you have further questions, ask your doctor, pharmacist or occupational health service.

1. What Vesperil is and what it is used for

Vesperil is melatonin MT1/MT2 receptor agonist. It is intended for: insomnia associated with shift work disorder in adults.

How it works

Ormelteon binds to the MT1 and MT2 melatonin receptors in the suprachiasmatic nucleus, the brain's master clock. MT1 is linked to falling asleep, MT2 to shifting the phase of the daily rhythm. It has no measurable effect on GABA receptors, so it is not a sedative-hypnotic in the usual sense.

2. What you need to know before you use it

- Common side effects: sleepiness, dizziness, fatigue and nausea.
- Do not combine with fluvoxamine or other strong CYP1A2 inhibitors. They raise ormelteon levels many times over.
- Not recommended in severe liver impairment, and to be used with caution in moderate impairment.
- Changes in prolactin and testosterone have been reported. Consider testing if a patient has absent periods, milk production, reduced libido or fertility problems.
- Patients should not drive or use machinery until they know how Vesperil affects them.

3. Reporting problems

If you notice any side effect or problem, talk to your doctor, pharmacist or occupational health service. You can also report it to your national authority and to NeuroStat at safety@neurostat.com. Reporting helps provide more information on the safety of this product.

4. About this document

Marketing authorisation holder: NeuroStat Holding AG, Thurgauerstrasse 66, 8050 Zürich, Switzerland. First approved 2006. Last revised September 2028.