Package leaflet: Lunavel

Lunavel (corexant)

Read all of this document carefully before you start. Keep it: you may need to read it again. If you have further questions, ask your doctor, pharmacist or occupational health service.

1. What Lunavel is and what it is used for

Lunavel is dual orexin receptor antagonist (DORA). It is intended for: insomnia characterised by difficulty with sleep onset and/or sleep maintenance.

How it works

Orexin neuropeptides promote wakefulness by acting on the OX1 and OX2 receptors. Corexant blocks both receptors and so reduces the wake drive, allowing sleep to occur naturally. Its mechanism differs from GABA-A modulators and benzodiazepines, which broadly depress central nervous system activity.

2. What you need to know before you use it

- Common adverse reactions: somnolence, headache, abnormal dreams and dry mouth.
- Next-day impairment: patients should not drive or perform activities requiring full alertness unless they are fully awake. Risk is higher at higher doses.
- Complex sleep behaviours, sleep paralysis and hypnagogic or hypnopompic hallucinations have been reported.
- Contraindicated in narcolepsy. Worsening of depression and suicidal ideation can occur: monitor patients.
- Avoid with strong CYP3A inhibitors. Reduce the dose with moderate CYP3A inhibitors.

3. Reporting problems

If you notice any side effect or problem, talk to your doctor, pharmacist or occupational health service. You can also report it to your national authority and to NeuroStat at safety@neurostat.example. Reporting helps provide more information on the safety of this product.

4. About this document

Marketing authorisation holder: NeuroStat Holding AG, Thurgauerstrasse 66, 8050 Zürich, Switzerland. First approved 2017. Last revised September 2028.