Package leaflet: Lunavel

Lunavel (corexant)

Read all of this document carefully before you start. Keep it: you may need to read it again. If you have further questions, ask your doctor, pharmacist or occupational health service.

1. What Lunavel is and what it is used for

Lunavel is dual orexin receptor antagonist (DORA). It is intended for: insomnia characterised by difficulty with sleep onset and/or sleep maintenance.

How it works

Orexin neuropeptides keep us awake by acting on the OX1 and OX2 receptors. Corexant blocks both, which lowers the drive to stay awake and lets sleep come naturally. This is a different mechanism from benzodiazepines and other GABA-A modulators, which depress the central nervous system more broadly.

2. What you need to know before you use it

- Common side effects: sleepiness, headache, abnormal dreams and dry mouth.
- Next-day impairment can occur, more often at higher doses. Patients should not drive or do anything that needs full alertness unless they are completely awake.
- Complex sleep behaviours, sleep paralysis and hallucinations on falling asleep or waking have been reported.
- Contraindicated in narcolepsy. Depression and suicidal thoughts can get worse, so patients should be monitored.
- Avoid strong CYP3A inhibitors, and lower the dose with moderate ones.

3. Reporting problems

If you notice any side effect or problem, talk to your doctor, pharmacist or occupational health service. You can also report it to your national authority and to NeuroStat at safety@neurostat.example. Reporting helps provide more information on the safety of this product.

4. About this document

Marketing authorisation holder: NeuroStat Holding AG, Thurgauerstrasse 66, 8050 Zürich, Switzerland. First approved 2017. Last revised September 2028.