Package leaflet: Vesperil

Vesperil (ormelteon)

Read all of this document carefully before you start. Keep it: you may need to read it again. If you have further questions, ask your doctor, pharmacist or occupational health service.

1. What Vesperil is and what it is used for

Vesperil is melatonin MT1/MT2 receptor agonist. It is intended for: insomnia associated with shift work disorder in adults.

How it works

Ormelteon binds with high affinity to the MT1 and MT2 melatonin receptors in the suprachiasmatic nucleus, the brain's master clock. MT1 activity is associated with the onset of sleep and MT2 activity with shifting the phase of the circadian rhythm. Ormelteon has no measurable affinity for GABA receptors, which is why it is not a sedative-hypnotic in the usual sense.

2. What you need to know before you use it

- Common adverse reactions: somnolence, dizziness, fatigue and nausea.
- Do not use with fluvoxamine or other strong inhibitors of CYP1A2, which raise ormelteon exposure many-fold.
- Not recommended in severe hepatic impairment. Use with caution in moderate impairment.
- Reports of changes in prolactin and testosterone levels: consider if amenorrhoea, galactorrhoea, reduced libido or fertility problems occur.
- Patients should not drive or operate machinery until they know how Vesperil affects them.

3. Reporting problems

If you notice any side effect or problem, talk to your doctor, pharmacist or occupational health service. You can also report it to your national authority and to NeuroStat at safety@neurostat.example. Reporting helps provide more information on the safety of this product.

4. About this document

Marketing authorisation holder: NeuroStat Holding AG, Thurgauerstrasse 66, 8050 Zürich, Switzerland. First approved 2006. Last revised September 2028.