

# New characterizations of the summatory function of the Möbius function

Maxie Dion Schmidt

Georgia Institute of Technology

School of Mathematics

Tuesday 23<sup>rd</sup> March, 2021

## Abstract

The Mertens function,  $M(x) := \sum_{n \leq x} \mu(n)$ , is defined as the summatory function of the classical Möbius function for  $x \geq 1$ . The inverse function sequence  $\{g^{-1}(n)\}_{n \geq 1}$  taken with respect to Dirichlet convolution is defined in terms of the strongly additive function  $\omega(n)$  that counts the number of distinct prime factors of any integer  $n \geq 2$ . For large  $x$  and  $n \leq x$ , we associate a natural combinatorial significance to the magnitude of the distinct values of the function  $g^{-1}(n)$  that depends directly on the exponent patterns in the prime factorizations of the integers in  $\{2, 3, \dots, x\}$  viewed as multisets.

We prove an Erdős-Kac theorem analog for the distribution of the unsigned sequence  $|g^{-1}(n)|$  over  $n \leq x$  with a limiting central limit theorem type tendency towards normal as  $x \rightarrow \infty$ . For all  $x \geq 1$ , discrete convolutions of  $G^{-1}(x) := \sum_{n \leq x} \lambda(n) |g^{-1}(n)|$  with the prime counting function  $\pi(x)$  determine exact formulas and new characterizations of asymptotic bounds for  $M(x)$ . In this way, we prove another concrete link of the distribution of  $L(x) := \sum_{n \leq x} \lambda(n)$  with the Mertens function and connect these classical summatory functions with an explicit normal tending probability distribution at large  $x$ . The proofs of these resulting combinatorially motivated new characterizations of  $M(x)$  are rigorous and unconditional.

**Keywords and Phrases:** *Möbius function; Mertens function; Dirichlet inverse; Liouville lambda function; prime omega function; prime counting function; Dirichlet generating function; Erdős-Kac theorem; strongly additive function.*

**Math Subject Classifications (MSC 2010):** *11N37; 11A25; 11N60; 11N64; and 11-04.*

# Index

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Preliminaries . . . . .	3
1.1.1	Definitions . . . . .	3
1.1.2	Properties . . . . .	3
1.1.3	Conjectures on boundedness and limiting behavior . . . . .	4
1.2	A concrete new approach to characterizing $M(x)$ . . . . .	4
1.2.1	Summatory functions of Dirichlet convolutions of arithmetic functions . . . . .	5
1.2.2	An exact expression for $M(x)$ via strongly additive functions . . . . .	5
1.2.3	Formulas illustrating the new characterizations of $M(x)$ . . . . .	7
1.3	Newer analytic methods utilized in the proofs within the article . . . . .	8
1.4	Notation and conventions . . . . .	8
<b>2</b>	<b>Initial elementary proofs of new results</b>	<b>10</b>
2.1	Establishing the summatory function properties and inversion identities . . . . .	10
2.2	Proving the characteristic signedness property of $g^{-1}(n)$ . . . . .	11
2.3	Results on the distribution of exceptional values of $\omega(n)$ and $\Omega(n)$ . . . . .	13
<b>3</b>	<b>Auxiliary sequences expressing the Dirichlet inverse function <math>g^{-1}(n)</math></b>	<b>16</b>
3.1	Definitions and properties of triangular component function sequences . . . . .	16
3.2	Relating the function $C_{\Omega(n)}(n)$ to exact formulas for $g^{-1}(n)$ . . . . .	16
3.3	Another connection to the distribution of the primes . . . . .	17
<b>4</b>	<b>The distributions of <math>C_{\Omega(n)}(n)</math> and <math> g^{-1}(n) </math></b>	<b>19</b>
4.1	Analytic proofs and adaptations of DGF methods for summing additive functions . . . . .	19
4.2	Average order of the unsigned sequences . . . . .	25
4.3	Erdős-Kac theorem analogs for the distributions of the unsigned sequences . . . . .	27
<b>5</b>	<b>Proofs of new formulas and limiting relations for <math>M(x)</math></b>	<b>30</b>
5.1	Establishing initial asymptotic bounds on the summatory function $G^{-1}(x)$ . . . . .	30
5.2	Bounding $M(x)$ by asymptotics for $G^{-1}(x)$ . . . . .	31
<b>6</b>	<b>Conclusions</b>	<b>33</b>
<b>A</b>	<b>Appendix: Asymptotic formulas</b>	<b>37</b>
<b>B</b>	<b>Table: The Dirichlet inverse function <math>g^{-1}(n)</math></b>	<b>40</b>

The conclusion of the proof of Proposition 2.1 in fact implies the stronger result that

$$g^{-1}(n) = \lambda(n) \times \sum_{d|n} \mu^2\left(\frac{n}{d}\right) C_{\Omega(d)}(d),$$

where we adopt the notation that for  $n \geq 2$ ,  $C_{\Omega(n)}(n) = (\Omega(n))! \times \prod_{p^\alpha || n} \frac{1}{\alpha!}$ , where the sequence is taken to be one at  $n := 1$ .

### 2.3 Results on the distribution of exceptional values of $\omega(n)$ and $\Omega(n)$

The next theorems reproduced from [14, §7.4] characterize the relative scarcity of the distributions of  $\omega(n)$  and  $\Omega(n)$  for  $n \leq x$  such that  $\omega(n), \Omega(n) > \log \log x$ . Since  $\mathbb{E}[\omega(n)], \mathbb{E}[\Omega(n)] = \log \log n + B$  for  $B \in (0, 1)$  an absolute constant in each case, these results imply a regular, normal tendency of these additive arithmetic functions towards their respective average orders.

**Theorem 2.2** (Upper bounds on exceptional values of  $\Omega(n)$  for large  $n$ ). *Let*

$$\begin{aligned} A(x, r) &:= \#\{n \leq x : \Omega(n) \leq r \cdot \log \log x\}, \\ B(x, r) &:= \#\{n \leq x : \Omega(n) \geq r \cdot \log \log x\}. \end{aligned}$$

*If  $0 < r \leq 1$  and  $x \geq 2$ , then*

$$A(x, r) \ll x(\log x)^{r-1-r \log r}, \quad \text{as } x \rightarrow \infty.$$

*If  $1 \leq r \leq R < 2$  and  $x \geq 2$ , then*

$$B(x, r) \ll_R x \cdot (\log x)^{r-1-r \log r}, \quad \text{as } x \rightarrow \infty.$$

Theorem 2.3 is a special case analog to the celebrated Erdős-Kac theorem typically stated for the normally distributed values of the scaled-shifted function  $\omega(n)$  over  $n \leq x$  as  $x \rightarrow \infty$  [14, cf. Thm. 7.21] [10, cf. §1.7].

**Theorem 2.3** (Exact limiting bounds on exceptional values of  $\Omega(n)$  for large  $n$ ). *We have that as  $x \rightarrow \infty$*

$$\#\{3 \leq n \leq x : \Omega(n) \leq \log \log n\} = \frac{x}{2} + O\left(\frac{x}{\sqrt{\log \log x}}\right).$$

**Theorem 2.4** (Montgomery and Vaughan). *Recall that we have defined*

$$\widehat{\pi}_k(x) := \#\{n \leq x : \Omega(n) = k\}.$$

*For  $0 < R < 2$  we have that uniformly for all  $1 \leq k \leq R \cdot \log \log x$*

$$\widehat{\pi}_k(x) = \mathcal{G}\left(\frac{k-1}{\log \log x}\right) \frac{x}{\log x} \frac{(\log \log x)^{k-1}}{(k-1)!} \left[1 + O_R\left(\frac{k}{(\log \log x)^2}\right)\right],$$

*where*

$$\mathcal{G}(z) := \frac{1}{\Gamma(z+1)} \times \prod_p \left(1 - \frac{z}{p}\right)^{-1} \left(1 - \frac{1}{p}\right)^z, \quad 0 \leq |z| < R.$$

**Remark 2.5.** We can extend the work in [14] on the distribution of  $\Omega(n)$  to find analogous results bounding the distribution of  $\omega(n)$ . We have that for  $0 < R < 2$

$$\pi_k(x) = \widehat{\mathcal{G}}\left(\frac{k-1}{\log \log x}\right) \frac{x}{\log x} \cdot \frac{(\log \log x)^{k-1}}{(k-1)!} \left[1 + O_R\left(\frac{k}{(\log \log x)^2}\right)\right], \quad \text{unif. for } 1 \leq k \leq R \log \log x. \quad (10)$$

The analogous function to express these bounds for  $\omega(n)$  is defined by  $\widehat{\mathcal{G}}(z) := \widehat{F}(1, z)/\Gamma(1 + z)$  where we take

$$\widehat{F}(s, z) := \prod_p \left(1 + \frac{z}{p^s - 1}\right)^{-1} \left(1 - \frac{1}{p^s}\right)^z, \operatorname{Re}(s) > \frac{1}{2}; |z| \leq R < 2.$$

Let the functions

$$\begin{aligned} C(x, r) &:= \#\{n \leq x : \omega(n) \leq r \log \log x\} \\ D(x, r) &:= \#\{n \leq x : \omega(n) \geq r \log \log x\}. \end{aligned}$$

Then we have the next uniform upper bounds given by

$$\begin{aligned} C(x, r) &\ll x(\log x)^{r-1-r \log r}, \text{ uniformly for } 0 < r \leq 1, \\ D(x, r) &\ll x(\log x)^{r-1-r \log r}, \text{ uniformly for } 1 \leq r \leq R < 2. \end{aligned}$$

With the next corollary, we can accurately approximate asymptotic order of the sums  $\mathcal{A}_\omega(x)$  (defined below) for large  $x$  by only considering the truncated sums  $\mathcal{D}_\omega(x)$  where we have the known uniform bounds on the summands for  $1 \leq k \leq \log \log x$ . This result is cited in the proof of our new result stated in Corollary 4.4 of Section 4 (see Appendix A).

**Corollary 2.6.** *Suppose that for  $x > e$  we define the following functions:*

$$\begin{aligned} \mathcal{N}_\omega(x) &:= \left| \sum_{k > \log \log x} (-1)^k \pi_k(x) \right| \\ \mathcal{D}_\omega(x) &:= \left| \sum_{k \leq \log \log x} (-1)^k \pi_k(x) \right| \\ \mathcal{A}_\omega(x) &:= \left| \sum_{k \geq 1} (-1)^k \pi_k(x) \right|. \end{aligned}$$

As  $x \rightarrow \infty$ , we have that  $\mathcal{D}_\omega(x)/\mathcal{N}_\omega(x) = o(1)$  and  $\mathcal{A}_\omega(x) \sim \mathcal{D}_\omega(x)$ .

*Proof.* First, we sum the main term for the function  $\mathcal{D}_\omega(x)$  by applying the limiting asymptotics for the incomplete gamma function derived in Lemma A.3 to obtain that

$$\begin{aligned} \mathcal{D}_\omega(x) &= \left| \sum_{1 \leq k \leq \log \log x} \frac{(-1)^k \cdot x}{\log x} \cdot \frac{(\log \log x)^{k-1}}{(k-1)!} \right| + O(E_\omega(x)) \\ &= \frac{x}{\sqrt{2\pi \log \log x}} + O(E_\omega(x)), \end{aligned}$$

The error term from the bound in the previous equation is defined according to (10) with  $\widehat{\mathcal{G}}\left(\frac{k-1}{\log \log x}\right) \gg 1$  for all  $1 \leq k \leq \log \log x$  as

$$\begin{aligned} E_\omega(x) &:= \sum_{k \leq \log \log x} \frac{x}{\log x} \cdot \frac{k(\log \log x)^{k-3}}{(k-1)!} \leq \frac{x}{\log x} \times \sum_{1 \leq k \leq \log \log x} \frac{(\log \log x)^{k-2}}{(k-1)!} \\ &\leq \frac{x}{(\log x)(\log \log x)} e^{\log \log x} \leq \frac{x}{\log \log x}. \end{aligned}$$

The right-hand-side expression in the previous equation follows by applying Lemma A.3.

Next, we utilize the notation for and bounds on the function  $D(x, r)$  from Remark 2.5 to bound the function  $\mathcal{N}_\omega(x)$  as follows:

$$\frac{1}{x} \times |\mathcal{N}_\omega(x)| \leq \sum_{k \geq \log \log x} \frac{\pi_k(x)}{x} = \frac{1}{x} \times \sum_{k \geq \log \log x} \#\{2 \leq n \leq x : \omega(n) = k\} \ll 1.$$

Then we see that

$$\left| \frac{\mathcal{D}_\omega(x)}{\mathcal{N}_\omega(x)} \right| = O\left( \frac{1}{\sqrt{\log \log x}} \right) = o(1), \text{ as } x \rightarrow \infty.$$

Equivalently, we have shown that  $\mathcal{D}_\omega(x) = o(\mathcal{N}_\omega(x))$ . The following results from the triangle inequality when  $x$  is large:

$$1 + o(1) = \left( \frac{\mathcal{D}_\omega(x) - \mathcal{N}_\omega(x)}{\mathcal{D}_\omega(x)} \right)^{-1} \ll \frac{\mathcal{D}_\omega(x)}{\mathcal{A}_\omega(x)} \ll \left( \frac{\mathcal{D}_\omega(x) + \mathcal{N}_\omega(x)}{\mathcal{D}_\omega(x)} \right)^{-1} = 1 + o(1).$$

The last equation implies that  $\mathcal{A}_\omega(x) \sim \mathcal{D}_\omega(x)$  as  $x \rightarrow \infty$ . □

## References

- [1] T. M. Apostol. *Introduction to Analytic Number Theory*. Springer–Verlag, 1976.
- [2] P. Billingsley. On the central limit theorem for the prime divisor function. *Amer. Math. Monthly*, 76(2):132–139, 1969.
- [3] P. Erdős and M. Kac. The gaussian errors in the theory of additive arithmetic functions. *American Journal of Mathematics*, 62(1):738–742, 1940.
- [4] C. E. Fröberg. On the prime zeta function. *BIT Numerical Mathematics*, 8:87–202, 1968.
- [5] R. L. Graham, D. E. Knuth, and O. Patashnik. *Concrete Mathematics: A Foundation for Computer Science*. Addison-Wesley, 1994.
- [6] G. H. Hardy and E. M. Wright, editors. *An Introduction to the Theory of Numbers*. Oxford University Press, 2008 (Sixth Edition).
- [7] P. Humphries. The distribution of weighted sums of the Liouville function and Pólya’s conjecture. *J. Number Theory*, 133:545–582, 2013.
- [8] G. Hurst. Computations of the Mertens function and improved bounds on the Mertens conjecture. *Math. Comp.*, 87:1013–1028, 2018.
- [9] H. Hwang and S. Janson. A central limit theorem for random ordered factorizations of integers. *Electron. J. Probab.*, 16(12):347–361, 2011.
- [10] H. Iwaniec and E. Kowalski. *Analytic Number Theory*, volume 53. AMS Colloquium Publications, 2004.
- [11] T. Kotnik and H. té Riele. The Mertens conjecture revisited. *Algorithmic Number Theory*, 7<sup>th</sup> International Symposium, 2006.
- [12] T. Kotnik and J. van de Lune. On the order of the Mertens function. *Exp. Math.*, 2004.
- [13] R. S. Lehman. On Liouville’s function. *Math. Comput.*, 14:311–320, 1960.
- [14] H. L. Montgomery and R. C. Vaughan. *Multiplicative Number Theory: I. Classical Theory*. Cambridge, 2006.
- [15] G. Nemes. The resurgence properties of the incomplete gamma function ii. *Stud. Appl. Math.*, 135(1):86–116, 2015.
- [16] N. Ng. The distribution of the summatory function of the Möbius function. *Proc. London Math. Soc.*, 89(3):361–389, 2004.
- [17] A. M. Odlyzko and H. J. J. té Riele. Disproof of the Mertens conjecture. *J. Reine Angew. Math*, 1985.
- [18] Frank W. J. Olver, Daniel W. Lozier, Ronald F. Boisvert, and Charles W. Clark, editors. *NIST Handbook of Mathematical Functions*. Cambridge University Press, 2010.
- [19] A. Renyi and P. Turan. On a theorem of Erdős-Kac. *Acta Arithmetica*, 4(1):71–84, 1958.
- [20] P. Ribenboim. *The new book of prime number records*. Springer, 1996.
- [21] G. Robin. Estimate of the Chebyshev function  $\theta$  on the  $k^{th}$  prime number and large values of the number of prime divisors function  $\omega(n)$  of  $n$ . *Acta Arith.*, 42(4):367–389, 1983.

- [22] J. B. Rosser and L. Schoenfeld. Approximate formulas for some functions of prime numbers. *Illinois J. Math.*, 6:64–94, 1962.
- [23] M. D. Schmidt. SageMath and Mathematica software for computations with the Mertens function, 2021. <https://github.com/maxieds/MertensFunctionComputations>.
- [24] N. J. A. Sloane. The Online Encyclopedia of Integer Sequences, 2021. <http://oeis.org>.
- [25] K. Soundararajan. Partial sums of the Möbius function. *Annals of Mathematics*, 2009.
- [26] E. C. Titchmarsh. *The theory of the Riemann zeta function*. Clarendon Press, 1951.

## A Appendix: Asymptotic formulas

**Facts A.1** (The incomplete gamma function). The (upper) *incomplete gamma function* is defined by [18, §8.4]

$$\Gamma(s, x) = \int_x^\infty t^{s-1} e^{-t} dt, \operatorname{Re}(s) > 0.$$

The following properties of  $\Gamma(a, x)$  hold at  $z, a > 0$ :

$$\Gamma(a, z) = (a-1)! \cdot e^{-z} \times \sum_{k=0}^{a-1} \frac{z^k}{k!}, a \in \mathbb{Z}^+, z > 0, \quad (26a)$$

$$\Gamma(a, z) \sim x^{a-1} \cdot e^{-z}, \text{ for fixed } a > 0, \text{ as } z \rightarrow +\infty. \quad (26b)$$

Moreover, for real  $z > 0$ , as  $z \rightarrow +\infty$  we have that [15]

$$\Gamma(z, z) = \sqrt{\frac{\pi}{2}} z^{z-\frac{1}{2}} e^{-z} + O(z^{z-1} e^{-z}), \quad (26c)$$

If  $z, a \rightarrow \infty$  with  $z = \lambda a$  for some  $\lambda > 0$  such that  $(\lambda - 1)^{-1} = o(|a|^{1/2})$ , then [15]

$$\Gamma(a, z) \sim z^a e^{-z} \times \sum_{n \geq 0} \frac{(-a)^n b_n(\lambda)}{(z-a)^{2n+1}}, \quad (26d)$$

where the sequence  $b_n(\lambda)$  satisfies the characteristic relation that  $b_0(\lambda) = 1$  and<sup>D</sup>

$$b_n(\lambda) = \lambda(1-\lambda)b'_{n-1}(\lambda) + \lambda(2n-1)b_{n-1}(\lambda), n \geq 1.$$

**Proposition A.2.** Suppose that  $z$  and  $a > 0$  are real parameters. If  $|z| = \lambda a$  for some  $\lambda > 1$ , then as  $z \rightarrow +\infty$  we have that

$$\Gamma(a, z) = \frac{z^{a-1} e^{-z}}{1-\lambda^{-1}} + O_\lambda(z^{a-2} e^{-z}),$$

and if  $a \geq 1$  is integer-valued, then as  $z \rightarrow +\infty$  we have

$$\Gamma(a, -z) \sim \frac{z^{a-1} e^z}{1+\lambda^{-1}} + O_\lambda(z^{a-2} e^{-z}).$$

*Proof.* We can see that for  $\lambda > 1$  and  $n \geq 1$  [5, §6.2]

$$\sum_{k=1}^n [\lambda^k] b_n(\lambda) = \frac{(2n)!}{2^n n!} \sim \sqrt{2} \left( \frac{2n}{e} \right)^n.$$

Thus we conclude that for all large enough  $n$

$$b_n(\lambda) \ll (2n)! \cdot \lambda^n.$$

It follows from (26d) that for  $N := z \left( 1 - \frac{\log \lambda}{\lambda} - \frac{1}{\lambda} \right)$  (cf. [15, §A.1])

$$\Gamma(a, z) \sim z^{a-1} e^{-z} \times \sum_{0 \leq n < N} \frac{(-1)^n b_n(\lambda)}{z^n \cdot (1-\lambda^{-1})^{2n+1}} = z^{a-1} e^{-z} (1 + E(a, z; \lambda)),$$

<sup>D</sup> An exact formula for  $b_n(\lambda)$  is given in terms of the *second-order Eulerian number triangle* [24, A008517] as follows:

$$b_n(\lambda) = \sum_{k=1}^n \left\langle \left\langle \begin{matrix} n \\ k-1 \end{matrix} \right\rangle \right\rangle \lambda^k.$$



where

$$|E(a, z; \lambda)| \ll \sum_{1 \leq n < N} \frac{(2\lambda^2 n)^n}{((\lambda - 1)^2 e z)^n}.$$

For all  $\lambda > 1$ , we have that the  $n^{\text{th}}$  summand in the previous bound is given by  $c_n^n$  where  $0 < c_n < 1$ . We conclude that since  $z \rightarrow \infty$ , we get  $E(a, z; \lambda) = o(1)$ . This argument justifies the main term for  $\Gamma(a, z)$  stated as our proposition.

We obtain the following expansions using a similar justification on which terms in the sum are main terms as above for  $a \in \mathbb{Z}^+$  and for  $N := z \left(1 + \frac{\log \lambda}{\lambda} + \frac{1}{\lambda}\right)$  as  $z \rightarrow -\infty$ :

$$\begin{aligned} \Gamma(a, z) &= z^a e^z \times \sum_{n \geq 0} \frac{(-a)^n b_n(\lambda)}{(z - a)^{2n+1}} \sim z^{a-1} e^z \times \sum_{n \geq 0} \frac{(-1)^n b_n(\lambda)}{z^n \left(1 + \frac{1}{\lambda}\right)^{2n+1}} \\ &= \frac{z^{a-1} e^z}{\left(1 + \frac{1}{\lambda}\right)} + O\left(\sum_{1 \leq n < N} \frac{\lambda^{2n} (2n e^{-1})^n}{(1 + \lambda)^{2n+1}}\right). \end{aligned} \quad \square$$

**Lemma A.3.** *For  $x > e$ , we have that*

$$S_1(x) := \frac{x}{\log x} \times \left| \sum_{1 \leq k \leq \log \log x} \frac{(-1)^k (\log \log x)^{k-1}}{(k-1)!} \right| \sim \frac{x}{\sqrt{2\pi \log \log x}}. \quad (27a)$$

*Proof of (27a).* We set  $t := \log \log x$  and let  $t \rightarrow +\infty$ . We can write

$$\frac{\log x}{x} \cdot S_1(x) = \left| \sum_{0 \leq k < t} \frac{(-t)^k}{k!} \right|.$$

By Taylor's theorem for the exponential function with remainder terms, we have that for some  $0 < s < t$

$$e^{-t} = \sum_{0 \leq k < t} \frac{(-t)^k}{k!} + \frac{(-s)^t}{t!} e^{-s}. \quad (27b)$$

Clearly, we have that the left-hand-side of (27b) corresponds to an  $O\left(\frac{1}{\log x}\right)$  error term. We can also compute that as  $t \rightarrow \infty$  the remainder term in the previous equation satisfies

$$\left| \frac{(-s)^t}{t!} e^{-s} \right| \leq \frac{t^t}{t!} \sim \frac{\log x}{\sqrt{2\pi \log \log x}},$$

by applying Stirlings formula to approximate  $t!$  when  $t$  is sufficiently large. A tight bound on the main term of our sum is argued in cases on the sign of the first sum on the right-hand-side of (27b). Let  $s_t := \text{sgn}\left(\sum_{0 \leq k < t} \frac{(-t)^k}{k!}\right)$ . If  $s_t = -1$ , then we get that

$$\left| \sum_{0 \leq k < t} \frac{(-t)^k}{k!} \right| \leq \frac{\log x}{\sqrt{2\pi \log \log x}} + O\left(\frac{1}{\log x}\right).$$

When  $s_t = +1$ , then for all large  $x$  we must have that the sign on the remainder term from Taylor's theorem in (27b) is negative. Reversing the corresponding inequality yields the symmetric bound that

$$\left| \sum_{0 \leq k < t} \frac{(-t)^k}{k!} \right| \geq \frac{\log x}{\sqrt{2\pi \log \log x}} + O\left(\frac{1}{\log x}\right). \quad \square$$

**Lemma A.4.** *For  $x > e$ , we have that*

$$S_3(x) := \sum_{1 \leq k \leq \log \log x} \frac{(\log \log x)^{k+1/2}}{(2k+1)(k-1)!} = \frac{1}{2}(\log x) \sqrt{\log \log x} + O\left(\frac{\log x}{\sqrt{\log \log x}}\right). \quad (27c)$$

*Proof.* We can sum this series symbolically with *Mathematica* to find that

$$S_3(x) = \frac{1}{2}(\log x)\sqrt{\log \log x} - \frac{\sqrt{\pi}}{4} \operatorname{erfi}\left(\sqrt{\log \log x}\right) - \frac{{}_2F_2\left(1, \frac{3}{2} + \log \log x; 1 + \log \log x, \frac{5}{2} + \log \log x; \log \log x\right)(\log x)(\log \log x)}{2\sqrt{2\pi}(2 \log \log x + 3)}. \quad (27d)$$

We will bound each component term in the above expansion of  $S_3(x)$  to see that the dominant asymptotic order of this function is given by the leading term.

As  $|z| \rightarrow \infty$ , the *imaginary error function*, denoted by  $\operatorname{erfi}(z)$ , has the following asymptotic expansion [18, §7.12]:

$$\operatorname{erfi}(z) = -i + \frac{e^{z^2}}{\sqrt{\pi}} \left( z^{-1} + \frac{z^{-3}}{2} + \frac{3z^{-5}}{4} + \frac{15z^{-7}}{8} + O(z^{-9}) \right).$$

It follows that

$$\frac{\sqrt{\pi}}{4} \operatorname{erfi}\left(\sqrt{\log \log x}\right) = \frac{(\log x)}{4} \left( \frac{1}{\sqrt{\log \log x}} + O\left(\frac{1}{(\log \log x)^{3/2}}\right) \right).$$

By bounding the remaining hypergeometric series term in the expansion of  $S_3(x)$ , we see that

$$\begin{aligned} & \frac{{}_2F_2\left(1, \frac{3}{2} + \log \log x; 1 + \log \log x, \frac{5}{2} + \log \log x; \log \log x\right)}{(2 \log \log x + 3)} \\ &= \frac{1}{2 \log \log x} \times \sum_{k \geq 0} \frac{1}{\left(1 + \frac{2k+3}{2 \log \log x}\right)} \prod_{i=1}^k \left(1 + \frac{i}{\log \log x}\right)^{-1} \\ &= \frac{1}{2 \log \log x} \times \sum_{\substack{k \geq 0 \\ 2k+3 < \log \log x}} \left(1 + \frac{2k+3}{\log \log x}\right)^{-1} \times \prod_{i=1}^k \left(1 + \frac{i}{\log \log x}\right)^{-1} + O\left(\frac{1}{\log \log x} \times \prod_{i=1}^{\log \log x - 1} \left(1 + \frac{i}{\log \log x}\right)^{-1}\right). \end{aligned} \quad (27e)$$

The rightmost sum corresponds to another error term. Indeed, we see that the inner sum for the error term in the second to last line over the bounds  $2k+3 \geq \log \log x$  is convergent to some constant. The leading product factors remaining in the last equation satisfy

$$\prod_{i=1}^{\log \log x - 1} \left(1 + \frac{i}{\log \log x}\right) \geq \frac{1}{2} + \frac{\log \log x}{2},$$

by appealing to a polynomial expansion of the factorial product by the Stirling numbers of the first kind where each component term  $\frac{j}{\log \log x} < 1$  whenever  $1 \leq j < \log \log x$ . It follows that the error term in (27e) is  $O((\log \log x)^{-1})$ . The main term in (27e) is expanded as

$$\begin{aligned} & \sum_{\substack{k \geq 0 \\ 2k+3 < \log \log x}} \left(1 + \frac{2k+3}{\log \log x}\right)^{-1} \times \prod_{i=1}^k \left(1 + \frac{i}{\log \log x}\right)^{-1} \\ & \ll \sum_{\substack{k \geq 0 \\ 2k+3 < \log \log x}} \prod_{i=1}^k \left(1 + \frac{i}{\log \log x} + O\left(\frac{i^2}{(\log \log x)^2}\right)\right) \times \left(1 + \frac{2k+3}{\log \log x} + O\left(\frac{(2k+3)^2}{(\log \log x)^2}\right)\right) \\ &= \sum_{\substack{k \geq 0 \\ 2k \leq \log \log x - 3}} \left[1 + \frac{(2k+1)(2k+2)}{2 \log \log x} + O\left(\frac{k^2}{(\log \log x)^2}\right)\right] \\ & \sim \frac{(\log \log x)^2}{12} + \frac{3(\log \log x)}{8} - \frac{19}{12} + \frac{1}{8 \log \log x}. \end{aligned}$$

Hence, the main term is the first leading term in the expansion of  $S_3(x)$  from (27d).  $\square$