



Maxie Schmidt <maxieds@gmail.com>

arXiv submission submit/7520536

1 message

e-prints@arxiv.org <e-prints@arxiv.org>

Sat, Apr 25, 2026 at 10:45 PM

Reply-To: help@arxiv.org

To: maxieds@gmail.com

Dear Maxie Schmidt,

Thank you for submitting your work to arXiv.

Your submission has been received and is under consideration. The temporary submission number is: submit/7520536.

As with all submissions, this work will go through technical and moderation checks. You will be contacted by arXiv when the work is announced or if any issues are identified.

Our goal is to screen and announce papers as quickly as possible while ensuring that papers meet long-term archival standards. Generally, this process takes two business days, with announcements occurring at 20:00 ET, Sunday through Thursday.

You can make changes and view the current status of the submission from your user dashboard: <https://arxiv.org/user/>.

Below is a copy of the submission information.

Regards,
arXiv Support

\\
arXiv:submit/7520536
From: Maxie Schmidt <maxieds@gmail.com>
Date: Sat, 25 Apr 2026 23:45:11 EST (14kb)

Title: Picking up the partial sums of the $M^{\{o\}}$ bius function problem with probabilistic number theory
 Authors: Dr. Maxie Dion Schmidt
 Categories: math.NT math.PR
 Comments: Keywords and phrases: $M^{\{o\}}$ bius function; Mertens function; Liouville lambda function; prime omega function; Dirichlet inverse; probabilistic number theory; Erdős-Kac theorem
 MSC-class: (2010): 11A25, 11N37, 11N60, 11N64, and 11K65
 License: <http://arxiv.org/licenses/nonexclusive-distrib/1.0/>

\\
 We revisit several hybrid multiplicative-to-additive type functions from a recent preprint article. These functions, $g(n)$ with Dirichlet generating function (DGF) $\zeta(s)^{-1} (1+P(s))^{-1}$ for $\text{Re}(s) > 1$ where $P(s) = \sum_p p^{-s}$ is the prime zeta function, $|g(n)| = \lambda(n) g(n)$ with DGF $\zeta(2s)^{-1} (1-P(s))^{-1}$, and $C_{\Omega}(n)$ with DGF $(1-P(s))^{-1}$. Each of these function variants are defined in terms of the additive (respectively, strongly additive) functions $\omega(n)$ and $\Omega(n)$. These two auxiliary functions are used in the prior manuscript to relate partial sums of the classical $M^{\{o\}}$ bius function, $\mu(n)$, to signed partial sums

involving the prime counting function, $\pi(x)$, and the Liouville lambda function, $\lambda(n) := (-1)^{\Omega(n)}$.

In this article, we explore summing the identities from the first manuscript using several probabilistic assumptions about the independence of the values of $\Omega(n)$ and $\mu^2(n)$ for $n \leq x$ at large x . We recover proofs of the limiting asymptotic growth of $M(x) / \sqrt{x}$ whose hypotheses promise to be substantially more attainable to make rigorous than past results from other authors relying on the Riemann Hypothesis or assumption of the linear independence of the simple, non-trivial zeros of $\zeta(s)$.

\\