

# Factorization theorems and canonical representations for generating functions of special sums

Maxie Dion Schmidt

Georgia Institute of Technology  
School of Mathematics  
Atlanta, GA 30318, USA

Ph.D. Thesis Final Defense  
July 2022

# Thesis Committee



**Georgia Institute  
of Technology**

Dr. Josephine Yu  
School of Mathematics  
*Georgia Tech*

Dr. Matthew Baker  
School of Mathematics  
*Georgia Tech*

Dr. Rafael de la Llave  
School of Mathematics  
*Georgia Tech*

Dr. Jayadev Athreya  
Department of Mathematics  
*University of Washington*

Dr. Bruce Berndt  
Department of Mathematics  
*University of Illinois at Urbana-Champaign*

# Overview of topics

- ▶ Gentle introduction to sequence generating functions (OGFs)
- ▶ Motivate certain “factorized” forms of OGFs for special sums
- ▶ Examples and main results from publications
- ▶ Topics on the frontier of these research topics
- ▶ Questions from the committee and audience

# Overview of topics

- ▶ Gentle introduction to sequence generating functions (OGFs)
- ▶ Motivate certain “factorized” forms of OGFs for special sums
- ▶ Examples and main results from publications
- ▶ Topics on the frontier of these research topics
- ▶ Questions from the committee and audience

# Overview of topics

- ▶ Gentle introduction to sequence generating functions (OGFs)
- ▶ Motivate certain “factorized” forms of OGFs for special sums
- ▶ Examples and main results from publications
- ▶ Topics on the frontier of these research topics
- ▶ Questions from the committee and audience

# Overview of topics

- ▶ Gentle introduction to sequence generating functions (OGFs)
- ▶ Motivate certain “factorized” forms of OGFs for special sums
- ▶ Examples and main results from publications
- ▶ Topics on the frontier of these research topics
- ▶ Questions from the committee and audience

# Overview of topics

- ▶ Gentle introduction to sequence generating functions (OGFs)
- ▶ Motivate certain “factorized” forms of OGFs for special sums
- ▶ Examples and main results from publications
- ▶ Topics on the frontier of these research topics
- ▶ Questions from the committee and audience

# Generating functions are essential tools in discrete mathematics

- For a sequence,  $\mathcal{F} := \{f_n\}_{n \geq 0} \subset \mathbb{C}$ , we define its **ordinary generating function** (OGF) to be

$$F(z) := \sum_{n \geq 0} f_n z^n.$$

- **Notation:** For  $n \geq 0$ ,  $[z^n]F(z) := f_n$  (**coefficient extraction**)
- **Good concise explanation:** *A generating function is a clothesline on which we hang up a sequence of numbers for display (Wilf, [23])*
- We can treat  $F(z)$  using complex analysis or may work with it formally (e.g., disregard convergence; see [10])
- Usually only consider integer sequences (or rational ones over  $\frac{f_n}{n!}$ )



# Generating functions are essential tools in discrete mathematics

- For a sequence,  $\mathcal{F} := \{f_n\}_{n \geq 0} \subset \mathbb{C}$ , we define its **ordinary generating function** (OGF) to be

$$F(z) := \sum_{n \geq 0} f_n z^n.$$

- **Notation:** For  $n \geq 0$ ,  $[z^n]F(z) := f_n$  (**coefficient extraction**)

- **Good concise explanation:** *A generating function is a clothesline on which we hang up a sequence of numbers for display* (Wilf, [23])
- We can treat  $F(z)$  using complex analysis or may work with it formally (e.g., disregard convergence; see [10])
- Usually only consider integer sequences (or rational ones over  $\frac{f_n}{n!}$ )

# Generating functions are essential tools in discrete mathematics

- For a sequence,  $\mathcal{F} := \{f_n\}_{n \geq 0} \subset \mathbb{C}$ , we define its **ordinary generating function** (OGF) to be

$$F(z) := \sum_{n \geq 0} f_n z^n.$$

- **Notation:** For  $n \geq 0$ ,  $[z^n]F(z) := f_n$  (**coefficient extraction**)
- **Good concise explanation:** *A generating function is a clothesline on which we hang up a sequence of numbers for display* (Wilf, [23])
- We can treat  $F(z)$  using complex analysis or may work with it formally (e.g., disregard convergence; see [10])
- Usually only consider integer sequences (or rational ones over  $\frac{f_n}{n!}$ )

# Generating functions are essential tools in discrete mathematics

- For a sequence,  $\mathcal{F} := \{f_n\}_{n \geq 0} \subset \mathbb{C}$ , we define its **ordinary generating function** (OGF) to be

$$F(z) := \sum_{n \geq 0} f_n z^n.$$

- **Notation:** For  $n \geq 0$ ,  $[z^n]F(z) := f_n$  (**coefficient extraction**)
- **Good concise explanation:** *A generating function is a clothesline on which we hang up a sequence of numbers for display* (Wilf, [23])
- We can treat  $F(z)$  using complex analysis or may work with it formally (e.g., disregard convergence; see [10])
- Usually only consider integer sequences (or rational ones over  $\frac{f_n}{n!}$ )

# Generating functions are essential tools in discrete mathematics

- For a sequence,  $\mathcal{F} := \{f_n\}_{n \geq 0} \subset \mathbb{C}$ , we define its **ordinary generating function** (OGF) to be

$$F(z) := \sum_{n \geq 0} f_n z^n.$$

- **Notation:** For  $n \geq 0$ ,  $[z^n]F(z) := f_n$  (**coefficient extraction**)
- **Good concise explanation:** *A generating function is a clothesline on which we hang up a sequence of numbers for display* (Wilf, [23])
- We can treat  $F(z)$  using complex analysis or may work with it formally (e.g., disregard convergence; see [10])
- Usually only consider integer sequences (or rational ones over  $\frac{f_n}{n!}$ )

# Focus of the thesis is on peer-reviewed publications from 2017–2021 (since enrolling at GT)

- ▶ Primary publications summarized in the thesis: [17, 20, 5, 4, 7, 6, 8]
- ▶ Publications focused on *Jabobi-type continued fractions* (*J-fractions*): [11, 14, 12, 16, 15]
- ▶ Other related peer-reviewed publications: [19, 13, 18, 22, 21, 20]

# Focus of the thesis is on peer-reviewed publications from 2017–2021 (since enrolling at GT)

- ▶ Primary publications summarized in the thesis: [17, 20, 5, 4, 7, 6, 8]
- ▶ Publications focused on **Jabobi-type continued fractions (J-fractions)**: [11, 14, 12, 16, 15]
- ▶ Other related peer-reviewed publications: [19, 13, 18, 22, 21, 20]

# Focus of the thesis is on peer-reviewed publications from 2017–2021 (since enrolling at GT)

- ▶ Primary publications summarized in the thesis: [17, 20, 5, 4, 7, 6, 8]
- ▶ Publications focused on **Jabobi-type continued fractions (J-fractions)**: [11, 14, 12, 16, 15]
- ▶ Other related peer-reviewed publications: [19, 13, 18, 22, 21, 20]

# Motivating series expansions for the OGFs of special sums (LGFs)

- For arithmetic functions  $f$  and  $g$ , we define their **Dirichlet convolution at  $n$**  by

$$(f * g)(n) = \sum_{d|n} f(d)g\left(\frac{n}{d}\right), \text{ integers } n \geq 1.$$

- A **Lambert series generating function** (LGF) is an OGF that allows us to generate multiplicative functions expressed via divisor sums of the form  $(f * \mathbb{1})(n)$ :

$$L_f(q) := \sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \sum_{m \geq 1} (f * \mathbb{1})(m)q^m.$$

- OGF relation:  $F(q) = L_{f * \mu}(q)$  (for  $\mu(n)$  the **Möbius function**)



# Motivating series expansions for the OGFs of special sums (LGFs)

- For arithmetic functions  $f$  and  $g$ , we define their **Dirichlet convolution at  $n$**  by

$$(f * g)(n) = \sum_{d|n} f(d)g\left(\frac{n}{d}\right), \text{ integers } n \geq 1.$$

- A **Lambert series generating function** (LGF) is an OGF that allows us to generate multiplicative functions expressed via divisor sums of the form  $(f * \mathbb{1})(n)$ :

$$L_f(q) := \sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \sum_{m \geq 1} (f * \mathbb{1})(m)q^m.$$

- OGF relation:  $F(q) = L_{f * \mu}(q)$  (for  $\mu(n)$  the Möbius function)

# Motivating series expansions for the OGFs of special sums (LGFs)

- For arithmetic functions  $f$  and  $g$ , we define their **Dirichlet convolution at  $n$**  by

$$(f * g)(n) = \sum_{d|n} f(d)g\left(\frac{n}{d}\right), \text{ integers } n \geq 1.$$

- A **Lambert series generating function** (LGF) is an OGF that allows us to generate multiplicative functions expressed via divisor sums of the form  $(f * \mathbb{1})(n)$ :

$$L_f(q) := \sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \sum_{m \geq 1} (f * \mathbb{1})(m)q^m.$$

- OGF relation:  $F(q) = L_{f * \mu}(q)$  (for  $\mu(n)$  the **Möbius function**)

# Examples: Some number theoretic function LGFs

$$\sum_{n \geq 1} \frac{\mu(n)q^n}{1 - q^n} = q \quad (1a)$$

$$\sum_{n \geq 1} \frac{\phi(n)q^n}{1 - q^n} = \frac{q}{(1 - q)^2}, |q| < 1 \quad (1b)$$

$$\sum_{n \geq 1} \frac{n^\alpha q^n}{1 - q^n} = \sum_{m \geq 1} \sigma_\alpha(m) q^m, \alpha \in \mathbb{R} \quad (1c)$$

$$\sum_{n \geq 1} \frac{\lambda(n)q^n}{1 - q^n} = \sum_{m \geq 1} q^{m^2} \quad (1d)$$

$$\sum_{n \geq 1} \frac{\Lambda(n)q^n}{1 - q^n} = \sum_{m \geq 1} \log(m) q^m. \quad (1e)$$

# Definitions – Some standard notation

- ▶ **Iverson's convention:** The symbol  $[\text{cond}]_\delta \in \{0, 1\}$  is one if and only if  $\text{cond}$  is true (cf. [2])
- ▶ The **greatest common divisor (GCD)**:  $(n, m) \equiv \gcd(n, m)$
- ▶ The **(infinite)  $q$ -Pochhammer symbol**:  

$$(a; q)_\infty := \prod_{m \geq 1} (1 - aq^{m-1})$$
- ▶ The **(Euler) partition function**: The number of (unordered) partitions of  $n$  is  $p(n) := [q^n](q; q)_\infty^{-1}$ , with  $p(0) := 1$ , for integers  $n \geq 0$
- ▶ The sequences  $s_e(n, k)$  (and  $s_o(n, k)$ ) denote the the number of  $k$ 's in all partitions of  $n$  into an even (and odd, respectively) number of distinct parts for integers  $1 \leq k \leq n$

# Definitions – Some standard notation

- ▶ **Iverson's convention:** The symbol  $[\text{cond}]_\delta \in \{0, 1\}$  is one if and only if  $\text{cond}$  is true (cf. [2])
- ▶ The **greatest common divisor (GCD)**:  $(n, m) \equiv \gcd(n, m)$
- ▶ The (infinite)  $q$ -Pochhammer symbol:  

$$(a; q)_\infty := \prod_{m \geq 1} (1 - aq^{m-1})$$
- ▶ The (Euler) partition function: The number of (unordered) partitions of  $n$  is  $p(n) := [q^n](q; q)_\infty^{-1}$ , with  $p(0) := 1$ , for integers  $n \geq 0$
- ▶ The sequences  $s_e(n, k)$  (and  $s_o(n, k)$ ) denote the the number of  $k$ 's in all partitions of  $n$  into an even (and odd, respectively) number of distinct parts for integers  $1 \leq k \leq n$

# Definitions – Some standard notation

- ▶ **Iverson's convention:** The symbol  $[\text{cond}]_\delta \in \{0, 1\}$  is one if and only if  $\text{cond}$  is true (cf. [2])
- ▶ The **greatest common divisor (GCD):**  $(n, m) \equiv \gcd(n, m)$
- ▶ The **(infinite)  $q$ -Pochhammer symbol:**  

$$(a; q)_\infty := \prod_{m \geq 1} (1 - aq^{m-1})$$
- ▶ The **(Euler) partition function:** The number of (unordered) partitions of  $n$  is  $p(n) := [q^n](q; q)_\infty^{-1}$ , with  $p(0) := 1$ , for integers  $n \geq 0$
- ▶ The sequences  $s_e(n, k)$  (and  $s_o(n, k)$ ) denote the the number of  $k$ 's in all partitions of  $n$  into an even (and odd, respectively) number of distinct parts for integers  $1 \leq k \leq n$

# Definitions – Some standard notation

- ▶ **Iverson's convention:** The symbol  $[\text{cond}]_\delta \in \{0, 1\}$  is one if and only if  $\text{cond}$  is true (cf. [2])
- ▶ The **greatest common divisor (GCD):**  $(n, m) \equiv \gcd(n, m)$
- ▶ The **(infinite)  $q$ -Pochhammer symbol:**  
 $(a; q)_\infty := \prod_{m \geq 1} (1 - aq^{m-1})$
- ▶ The **(Euler) partition function:** The number of (unordered) partitions of  $n$  is  $p(n) := [q^n](q; q)_\infty^{-1}$ , with  $p(0) := 1$ , for integers  $n \geq 0$
- ▶ The sequences  $s_e(n, k)$  (and  $s_o(n, k)$ ) denote the the number of  $k$ 's in all partitions of  $n$  into an even (and odd, respectively) number of distinct parts for integers  $1 \leq k \leq n$

# Definitions – Some standard notation

- ▶ **Iverson's convention:** The symbol  $[\text{cond}]_\delta \in \{0, 1\}$  is one if and only if  $\text{cond}$  is true (cf. [2])
- ▶ The **greatest common divisor (GCD):**  $(n, m) \equiv \gcd(n, m)$
- ▶ The **(infinite)  $q$ -Pochhammer symbol:**  
 $(a; q)_\infty := \prod_{m \geq 1} (1 - aq^{m-1})$
- ▶ The **(Euler) partition function:** The number of (unordered) partitions of  $n$  is  $p(n) := [q^n](q; q)_\infty^{-1}$ , with  $p(0) := 1$ , for integers  $n \geq 0$
- ▶ The sequences  $s_e(n, k)$  (and  $s_o(n, k)$ ) denote the the number of  $k$ 's in all partitions of  $n$  into an even (and odd, respectively) number of distinct parts for integers  $1 \leq k \leq n$



# Factorization theorems for LGF series

- ▶ Overlapping ideas in publications by MDS and M. Merca: [17, 3]
- ▶ Coauthored work over the next few years: [5, 4, 7, 6]
- ▶ Key idea is to re-write the LGF series as in the following LHS expansion:

$$\sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \frac{1}{(q; q)_\infty} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n \quad (\text{LGF-FT})$$

- ▶ For any  $N \geq 1$ , the matrices  $(s_{n,k})_{1 \leq n, k \leq N}$  are lower triangular with ones on the diagonal (and hence invertible)

# Factorization theorems for LGF series

- ▶ Overlapping ideas in publications by MDS and M. Merca: [17, 3]
- ▶ Coauthored work over the next few years: [5, 4, 7, 6]
- ▶ Key idea is to re-write the LGF series as in the following LHS expansion:

$$\sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \frac{1}{(q; q)_\infty} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n \quad (\text{LGF-FT})$$

- ▶ For any  $N \geq 1$ , the matrices  $(s_{n,k})_{1 \leq n, k \leq N}$  are lower triangular with ones on the diagonal (and hence invertible)

# Factorization theorems for LGF series

- ▶ Overlapping ideas in publications by MDS and M. Merca: [17, 3]
- ▶ Coauthored work over the next few years: [5, 4, 7, 6]
- ▶ Key idea is to re-write the LGF series as in the following LHS expansion:

$$\sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \frac{1}{(q; q)_{\infty}} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n \quad (\mathbf{LGF-FT})$$

- ▶ For any  $N \geq 1$ , the matrices  $(s_{n,k})_{1 \leq n, k \leq N}$  are lower triangular with ones on the diagonal (and hence invertible)

# Factorization theorems for LGF series

- ▶ Overlapping ideas in publications by MDS and M. Merca: [17, 3]
- ▶ Coauthored work over the next few years: [5, 4, 7, 6]
- ▶ Key idea is to re-write the LGF series as in the following LHS expansion:

$$\sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \frac{1}{(q; q)_{\infty}} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n \quad (\mathbf{LGF-FT})$$

- ▶ For any  $N \geq 1$ , the matrices  $(s_{n,k})_{1 \leq n, k \leq N}$  are lower triangular with ones on the diagonal (and hence invertible)

# Factorization theorems for LGF series (cont'd)

- ▶ We prove:  $s_{n,k} = s_o(n, k) - s_e(n, k)$
- ▶ We prove:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- ▶ **Interpretations:** Interesting new ties between OGFs for multiplicative functions and the more additive theory of partitions
- ▶ **Key questions to keep in mind for later:**
  - ▶ Why was the factor of  $(q; q)_{\infty}^{-1}$  in the OGF factorization in equation **(LGF-FT)** so natural?
  - ▶ Collecting common denominators of the partial sums of the RHS yields this OGF factor in the limiting case (algebraic rationale for the choice)
  - ▶ Is there a deeper underlying principle to explain why this factorized form should be the most natural?

# Factorization theorems for LGF series (cont'd)

- ▶ We prove:  $s_{n,k} = s_o(n, k) - s_e(n, k)$
- ▶ We prove:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- ▶ **Interpretations:** Interesting new ties between OGFs for multiplicative functions and the more additive theory of partitions
- ▶ **Key questions to keep in mind for later:**
  - ▶ Why was the factor of  $(q; q)_{\infty}^{-1}$  in the OGF factorization in equation **(LGF-FT)** so natural?
  - ▶ Collecting common denominators of the partial sums of the RHS yields this OGF factor in the limiting case (algebraic rationale for the choice)
  - ▶ Is there a deeper underlying principle to explain why this factorized form should be the most natural?

# Factorization theorems for LGF series (cont'd)

- ▶ We prove:  $s_{n,k} = s_o(n, k) - s_e(n, k)$
- ▶ We prove:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- ▶ **Interpretations:** Interesting new ties between OGFs for multiplicative functions and the more additive theory of partitions
- ▶ **Key questions to keep in mind for later:**
  - ▶ Why was the factor of  $(q; q)_{\infty}^{-1}$  in the OGF factorization in equation **(LGF-FT)** so natural?
  - ▶ Collecting common denominators of the partial sums of the RHS yields this OGF factor in the limiting case (algebraic rationale for the choice)
  - ▶ Is there a deeper underlying principle to explain why this factorized form should be the most natural?

# Factorization theorems for LGF series (cont'd)

- ▶ We prove:  $s_{n,k} = s_o(n, k) - s_e(n, k)$
- ▶ We prove:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k)\mu\left(\frac{n}{d}\right)$
- ▶ **Interpretations:** Interesting new ties between OGFs for multiplicative functions and the more additive theory of partitions
- ▶ **Key questions to keep in mind for later:**
  - ▶ Why was the factor of  $(q; q)_{\infty}^{-1}$  in the OGF factorization in equation **(LGF-FT)** so natural?
  - ▶ Collecting common denominators of the partial sums of the RHS yields this OGF factor in the limiting case (algebraic rationale for the choice)
  - ▶ Is there a deeper underlying principle to explain why this factorized form should be the most natural?



# Factorization theorems for LGF series (cont'd)

- ▶ We prove:  $s_{n,k} = s_o(n, k) - s_e(n, k)$
- ▶ We prove:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- ▶ **Interpretations:** Interesting new ties between OGFs for multiplicative functions and the more additive theory of partitions
- ▶ **Key questions to keep in mind for later:**
  - ▶ Why was the factor of  $(q; q)_{\infty}^{-1}$  in the OGF factorization in equation **(LGF-FT)** so natural?
  - ▶ Collecting common denominators of the partial sums of the RHS yields this OGF factor in the limiting case (algebraic rationale for the choice)
  - ▶ Is there a deeper underlying principle to explain why this factorized form should be the most natural?

# Factorization theorems for LGF series (cont'd)

- ▶ We prove:  $s_{n,k} = s_o(n, k) - s_e(n, k)$
- ▶ We prove:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- ▶ **Interpretations:** Interesting new ties between OGFs for multiplicative functions and the more additive theory of partitions
- ▶ **Key questions to keep in mind for later:**
  - ▶ Why was the factor of  $(q; q)_{\infty}^{-1}$  in the OGF factorization in equation **(LGF-FT)** so natural?
  - ▶ Collecting common denominators of the partial sums of the RHS yields this OGF factor in the limiting case (algebraic rationale for the choice)
  - ▶ Is there a deeper underlying principle to explain why this factorized form should be the most natural?

# LGF factorization theorems – Other results

- ▶ Let the **(normalized) average order** of the function  $f$  be defined by

$$\Sigma_f(x) := \sum_{1 \leq n \leq x} f(n), \text{ for } x \geq 1.$$

- ▶ Let  $a_f(n) := \sum_{1 \leq k \leq n} s_{n,k} f(k)$  where the lower triangular  $s_{n,k}$  are the same as in **(LGF-FT)**

- ▶ **Theorem:** For all  $n \geq 1$

$$\begin{aligned} \Sigma_{f*1}(n+1) = & \sum_{b=\pm 1} \sum_{k=1}^{\left\lfloor \frac{\sqrt{24n+1}-b}{6} \right\rfloor + 1} (-1)^{k+1} \Sigma_{f*1} \left( n+1 - \frac{k(3k+b)}{2} \right) \\ & + \sum_{1 \leq k \leq n} a_f(k+1). \end{aligned}$$

# LGF factorization theorems – Other results

- ▶ Let the **(normalized) average order** of the function  $f$  be defined by

$$\Sigma_f(x) := \sum_{1 \leq n \leq x} f(n), \text{ for } x \geq 1.$$

- ▶ Let  $a_f(n) := \sum_{1 \leq k \leq n} s_{n,k} f(k)$  where the lower triangular  $s_{n,k}$  are the same as in **(LGF-FT)**

- ▶ **Theorem:** For all  $n \geq 1$

$$\begin{aligned} \Sigma_{f*1}(n+1) = & \sum_{b=\pm 1} \sum_{k=1}^{\left\lfloor \frac{\sqrt{24n+1}-b}{6} \right\rfloor + 1} (-1)^{k+1} \Sigma_{f*1} \left( n+1 - \frac{k(3k+b)}{2} \right) \\ & + \sum_{1 \leq k \leq n} a_f(k+1). \end{aligned}$$

# LGF factorization theorems – Other results

- ▶ Let the **(normalized) average order** of the function  $f$  be defined by

$$\Sigma_f(x) := \sum_{1 \leq n \leq x} f(n), \text{ for } x \geq 1.$$

- ▶ Let  $a_f(n) := \sum_{1 \leq k \leq n} s_{n,k} f(k)$  where the lower triangular  $s_{n,k}$  are the same as in **(LGF-FT)**

- ▶ **Theorem:** For all  $n \geq 1$

$$\begin{aligned} \Sigma_{f*1}(n+1) = & \sum_{b=\pm 1} \sum_{k=1}^{\left\lfloor \frac{\sqrt{24n+1}-b}{6} \right\rfloor + 1} (-1)^{k+1} \Sigma_{f*1} \left( n+1 - \frac{k(3k+b)}{2} \right) \\ & + \sum_{1 \leq k \leq n} a_f(k+1). \end{aligned}$$

# LGFs – Other results (cont'd)

- **Notation:**  $\sigma(n) \equiv \sigma_1(n) := \sum_{d|n} d$  is the **(ordinary)**  
**sum-of-divisors function**

- **Corollary:** For all  $x \geq 1$

$$\Sigma_{\sigma}(x+1) = \sum_{s=\pm 1} \left( \sum_{0 \leq n \leq x} \sum_{k=1}^{\lfloor \frac{\sqrt{24n+25}-s}{6} \rfloor} (-1)^{k+1} \frac{k(3k+s)}{2} p(x-n) \right)$$

- **Compare to classical bounds:**  $\Sigma_{\sigma}(x) = \frac{\pi^2 x^2}{12} \times \left(1 + O\left(\frac{\log x}{x}\right)\right)$
- **Improvement (Walfisz, 1964):**

$$\Sigma_{\sigma}(x) = \frac{\pi^2 x^2}{12} \times \left(1 + O\left(\frac{(\log x)^{\frac{2}{3}}}{x}\right)\right)$$

# LGFs – Other results (cont'd)

- **Notation:**  $\sigma(n) \equiv \sigma_1(n) := \sum_{d|n} d$  is the **(ordinary)**  
**sum-of-divisors function**

- **Corollary:** For all  $x \geq 1$

$$\Sigma_{\sigma}(x+1) = \sum_{s=\pm 1} \left( \sum_{0 \leq n \leq x} \sum_{k=1}^{\lfloor \frac{\sqrt{24n+25}-s}{6} \rfloor} (-1)^{k+1} \frac{k(3k+s)}{2} p(x-n) \right)$$

- **Compare to classical bounds:**  $\Sigma_{\sigma}(x) = \frac{\pi^2 x^2}{12} \times (1 + O(\frac{\log x}{x}))$   
 ► **Improvement (Walfisz, 1964):**

$$\Sigma_{\sigma}(x) = \frac{\pi^2 x^2}{12} \times \left( 1 + O\left(\frac{(\log x)^{\frac{2}{3}}}{x}\right) \right)$$

# LGFs – Other results (cont'd)

- **Notation:**  $\sigma(n) \equiv \sigma_1(n) := \sum_{d|n} d$  is the **(ordinary)**  
**sum-of-divisors function**

- **Corollary:** For all  $x \geq 1$

$$\Sigma_{\sigma}(x+1) = \sum_{s=\pm 1} \left( \sum_{0 \leq n \leq x} \sum_{k=1}^{\lfloor \frac{\sqrt{24n+25}-s}{6} \rfloor} (-1)^{k+1} \frac{k(3k+s)}{2} p(x-n) \right)$$

- **Compare to classical bounds:**  $\Sigma_{\sigma}(x) = \frac{\pi^2 x^2}{12} \times (1 + O(\frac{\log x}{x}))$   
 ► **Improvement (Walfisz, 1964):**

$$\Sigma_{\sigma}(x) = \frac{\pi^2 x^2}{12} \times \left( 1 + O\left(\frac{(\log x)^{\frac{2}{3}}}{x}\right) \right)$$



# LGFs – Other results (cont'd)

- **Notation:**  $\sigma(n) \equiv \sigma_1(n) := \sum_{d|n} d$  is the **(ordinary)**  
**sum-of-divisors function**

- **Corollary:** For all  $x \geq 1$

$$\Sigma_{\sigma}(x+1) = \sum_{s=\pm 1} \left( \sum_{0 \leq n \leq x} \sum_{k=1}^{\lfloor \frac{\sqrt{24n+25}-s}{6} \rfloor} (-1)^{k+1} \frac{k(3k+s)}{2} p(x-n) \right)$$

- **Compare to classical bounds:**  $\Sigma_{\sigma}(x) = \frac{\pi^2 x^2}{12} \times (1 + O(\frac{\log x}{x}))$   
 ► **Improvement (Walfisz, 1964):**

$$\Sigma_{\sigma}(x) = \frac{\pi^2 x^2}{12} \times \left( 1 + O\left(\frac{(\log x)^{\frac{2}{3}}}{x}\right) \right)$$

# Branching out from LGFs I

- **Motivation:** We find that

$$\Sigma_f(x) = \sum_{\substack{d|x \\ d>1}} f(d) + \sum_{\substack{1 \leq d \leq x \\ (d,x)=1}} f(d) + \sum_{\substack{1 < d \leq x \\ 1 < (d,x) < x}} f(d), \text{ for } x \geq 1,$$

- The first summations are generated by LGFs as

$$\sum_{\substack{d|x \\ d>1}} f(d) = [q^n] L_f(q) - f(1), \text{ for any } n \geq 1.$$

- What about the next two sum terms?

# Branching out from LGFs I

- **Motivation:** We find that

$$\Sigma_f(x) = \sum_{\substack{d|x \\ d>1}} f(d) + \sum_{\substack{1 \leq d \leq x \\ (d,x)=1}} f(d) + \sum_{\substack{1 < d \leq x \\ 1 < (d,x) < x}} f(d), \text{ for } x \geq 1,$$

- The first summations are generated by LGFs as

$$\sum_{\substack{d|x \\ d>1}} f(d) = [q^n] L_f(q) - f(1), \text{ for any } n \geq 1.$$

- What about the next two sum terms?

# Branching out from LGFs I

- **Motivation:** We find that

$$\Sigma_f(x) = \sum_{\substack{d|x \\ d>1}} f(d) + \sum_{\substack{1 \leq d \leq x \\ (d,x)=1}} f(d) + \sum_{\substack{1 < d \leq x \\ 1 < (d,x) < x}} f(d), \text{ for } x \geq 1,$$

- The first summations are generated by LGFs as

$$\sum_{\substack{d|x \\ d>1}} f(d) = [q^n] L_f(q) - f(1), \text{ for any } n \geq 1.$$

- What about the next two sum terms?

# Branching out from LGFs I

- **Motivation:** We find that

$$\Sigma_f(x) = \sum_{\substack{d|x \\ d>1}} f(d) + \sum_{\substack{1 \leq d \leq x \\ (d,x)=1}} f(d) + \sum_{\substack{1 < d \leq x \\ 1 < (d,x) < x}} f(d), \text{ for } x \geq 1,$$

- The first summations are generated by LGFs as

$$\sum_{\substack{d|x \\ d>1}} f(d) = [q^n] L_f(q) - f(1), \text{ for any } n \geq 1.$$

- What about the next two sum terms?

# Branching out from LGFs II

- Generalized factorization theorems for GCD-type sums in [8]
- For integers  $1 \leq k \leq x$ , we define

$$T_f(x) = \sum_{\substack{d=1 \\ (d,x)=1}}^x f(d), \quad (\text{Type I Sums})$$

$$L_{f,g,k}(x) = \sum_{d|(k,x)} f(d)g\left(\frac{x}{d}\right). \quad (\text{Type II Sums})$$

- The factorization theorems considered are now of the form

$$T_f(x) = [q^x] \left( \frac{1}{(q; q)_\infty} \times \sum_{n \geq 2} \sum_{k=1}^n t_{n,k} f(k) q^n + f(1)q \right),$$

$$g(x) = [q^x] \left( \frac{1}{(q; q)_\infty} \times \sum_{n \geq 2} \sum_{k=1}^n u_{n,k}(f, w) \left( \sum_{m=1}^k L_{f,g,m}(k) w^m \right) q^n \right),$$

- Here,  $w \in \mathbb{C} \setminus \{0\}$  is a non-zero indeterminate parameter inserted to force invertibility of the sequences  $u_{n,k}(f, w)$  above

# Branching out from LGFs II

- ▶ Generalized factorization theorems for GCD-type sums in [8]
- ▶ For integers  $1 \leq k \leq x$ , we define

$$T_f(x) = \sum_{\substack{d=1 \\ (d,x)=1}}^x f(d), \quad (\text{Type I Sums})$$

$$L_{f,g,k}(x) = \sum_{d|(k,x)} f(d)g\left(\frac{x}{d}\right). \quad (\text{Type II Sums})$$

- ▶ The factorization theorems considered are now of the form

$$T_f(x) = [q^x] \left( \frac{1}{(q; q)_\infty} \times \sum_{n \geq 2} \sum_{k=1}^n t_{n,k} f(k) q^n + f(1)q \right),$$

$$g(x) = [q^x] \left( \frac{1}{(q; q)_\infty} \times \sum_{n \geq 2} \sum_{k=1}^n u_{n,k}(f, w) \left( \sum_{m=1}^k L_{f,g,m}(k) w^m \right) q^n \right),$$

- ▶ Here,  $w \in \mathbb{C} \setminus \{0\}$  is a non-zero indeterminate parameter inserted to force invertibility of the sequences  $u_{n,k}(f, w)$  above

# Branching out from LGFs II

- Generalized factorization theorems for GCD-type sums in [8]
- For integers  $1 \leq k \leq x$ , we define

$$T_f(x) = \sum_{\substack{d=1 \\ (d,x)=1}}^x f(d), \quad (\text{Type I Sums})$$

$$L_{f,g,k}(x) = \sum_{d|(k,x)} f(d)g\left(\frac{x}{d}\right). \quad (\text{Type II Sums})$$

- The factorization theorems considered are now of the form

$$T_f(x) = [q^x] \left( \frac{1}{(q; q)_\infty} \times \sum_{n \geq 2} \sum_{k=1}^n t_{n,k} f(k) q^n + f(1)q \right),$$

$$g(x) = [q^x] \left( \frac{1}{(q; q)_\infty} \times \sum_{n \geq 2} \sum_{k=1}^n u_{n,k}(f, w) \left( \sum_{m=1}^k L_{f,g,m}(k) w^m \right) q^n \right),$$

- Here,  $w \in \mathbb{C} \setminus \{0\}$  is a non-zero indeterminate parameter inserted to force invertibility of the sequences  $u_{n,k}(f, w)$  above



# Branching out from LGFs II

- ▶ Generalized factorization theorems for GCD-type sums in [8]
- ▶ For integers  $1 \leq k \leq x$ , we define

$$T_f(x) = \sum_{\substack{d=1 \\ (d,x)=1}}^x f(d), \quad (\text{Type I Sums})$$

$$L_{f,g,k}(x) = \sum_{d|(k,x)} f(d)g\left(\frac{x}{d}\right). \quad (\text{Type II Sums})$$

- ▶ The factorization theorems considered are now of the form

$$T_f(x) = [q^x] \left( \frac{1}{(q; q)_\infty} \times \sum_{n \geq 2} \sum_{k=1}^n t_{n,k} f(k) q^n + f(1)q \right),$$

$$g(x) = [q^x] \left( \frac{1}{(q; q)_\infty} \times \sum_{n \geq 2} \sum_{k=1}^n u_{n,k}(f, w) \left( \sum_{m=1}^k L_{f,g,m}(k) w^m \right) q^n \right),$$

- ▶ Here,  $w \in \mathbb{C} \setminus \{0\}$  is a non-zero indeterminate parameter inserted to force invertibility of the sequences  $u_{n,k}(f, w)$  above

# Examples I: What other types of sums might we want to generate?

## Example ( $\mathcal{A}$ -Set convolutions, ACVL)

For each  $n \geq 1$ , let  $A(n) \subseteq \{1 \leq d \leq n : d|n\}$  be a subset of the divisors of  $n$ . We say that  $n$  is  $A$ -primitive if  $A(n) \equiv \{1, n\}$ . Let the set of  $A$ -primitive positive integers be denoted by

$$\mathcal{A} := \{n \geq 1 : n \text{ is } A\text{-primitive}\}.$$

Then we may consider the following invertible convolutions:

$$S_{1,\mathcal{A}}(f, g; n) := \sum_{\substack{d|n \\ d \in \mathcal{A}}} f(d)g\left(\frac{n}{d}\right),$$

$$S_{2,\mathcal{A}}(f, g; n) := \sum_{\substack{d|n \\ d, \frac{n}{d} \in \mathcal{A}}} f(d)g\left(\frac{n}{d}\right).$$

# Examples II: What other types of sums might we want to generate?

## Example (Unitary convolutions, UCVL)

The *unitary convolution* of  $f$  and  $g$  at integers  $n \geq 1$  is defined by

$$(f \odot g)(n) := \sum_{\substack{d|n \\ (d, \frac{n}{d})=1}} f(d)g\left(\frac{n}{d}\right).$$

# Examples III: What other types of sums might we want to generate?

## Example ( $\mathcal{D}$ -Kernel convolutions, DCVL)

Suppose that  $\mathcal{D} : \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mathbb{C}$  is an **invertible** and **lower triangular kernel function**: i.e.,  $\mathcal{D}(n, k) = 0$  whenever  $k > n$  and  $\mathcal{D}(n, n) \neq 0$  for all  $n \geq 1$ . We want to study a generalized class of  $\mathcal{D}$ -convolution type sums of the form

$$(f \boxdot_{\mathcal{D}} g)(n) := \sum_{1 \leq k \leq n} f(k)g(n+1-k)\mathcal{D}(n, k), \text{ for integers } n \geq 1.$$

# Definitions of generalized factorization theorems

- **Option 1:** For  $n \geq 1$  and multiplier OGFs such that  $\mathcal{C}(0) \neq 0$

$$\sum_{\substack{k \in A_n \\ A_n \subseteq [1, n] \cup \{n\}}} f(k) := [q^n] \left( \frac{1}{\mathcal{C}(q)} \times \sum_{\substack{n \geq 1 \\ 1 \leq k \leq n}} v_{n,k}(\mathcal{A}, \mathcal{C}) f(k) q^n \right)$$

- **Option 2:** For  $n \geq 1$ , weights  $\mathcal{T}_{j,j} \neq 0$  for all  $j \geq 1$ , and multiplier OGFs such that  $\mathcal{C}(0) \neq 0$

$$\sum_{1 \leq k \leq n} \mathcal{T}_{n,k} f(k) := [q^n] \left( \frac{1}{\mathcal{C}(q)} \times \sum_{\substack{n \geq 1 \\ 1 \leq k \leq n}} u_{n,k}(\mathcal{T}, \mathcal{C}) f(k) q^n \right)$$

- These generalized sum types allow us to consider (weighted) forms of the special case LGF and GCD-type OGF factorizations we have seen so far (**Explain, clarify quantifiers**)

# Definitions of generalized factorization theorems

- **Option 1:** For  $n \geq 1$  and multiplier OGFs such that  $\mathcal{C}(0) \neq 0$

$$\sum_{\substack{k \in A_n \\ A_n \subseteq [1, n] \cup \{n\}}} f(k) := [q^n] \left( \frac{1}{\mathcal{C}(q)} \times \sum_{\substack{n \geq 1 \\ 1 \leq k \leq n}} v_{n,k}(\mathcal{A}, \mathcal{C}) f(k) q^n \right)$$

- **Option 2:** For  $n \geq 1$ , weights  $\mathcal{T}_{j,j} \neq 0$  for all  $j \geq 1$ , and multiplier OGFs such that  $\mathcal{C}(0) \neq 0$

$$\sum_{1 \leq k \leq n} \mathcal{T}_{n,k} f(k) := [q^n] \left( \frac{1}{\mathcal{C}(q)} \times \sum_{\substack{n \geq 1 \\ 1 \leq k \leq n}} u_{n,k}(\mathcal{T}, \mathcal{C}) f(k) q^n \right)$$

- These generalized sum types allow us to consider (weighted) forms of the special case LGF and GCD-type OGF factorizations we have seen so far (**Explain, clarify quantifiers**)

# Definitions of generalized factorization theorems

- **Option 1:** For  $n \geq 1$  and multiplier OGFs such that  $\mathcal{C}(0) \neq 0$

$$\sum_{\substack{k \in A_n \\ A_n \subseteq [1, n] \cup \{n\}}} f(k) := [q^n] \left( \frac{1}{\mathcal{C}(q)} \times \sum_{\substack{n \geq 1 \\ 1 \leq k \leq n}} v_{n,k}(\mathcal{A}, \mathcal{C}) f(k) q^n \right)$$

- **Option 2:** For  $n \geq 1$ , weights  $\mathcal{T}_{j,j} \neq 0$  for all  $j \geq 1$ , and multiplier OGFs such that  $\mathcal{C}(0) \neq 0$

$$\sum_{1 \leq k \leq n} \mathcal{T}_{n,k} f(k) := [q^n] \left( \frac{1}{\mathcal{C}(q)} \times \sum_{\substack{n \geq 1 \\ 1 \leq k \leq n}} u_{n,k}(\mathcal{T}, \mathcal{C}) f(k) q^n \right)$$

- These generalized sum types allow us to consider (weighted) forms of the special case LGF and GCD-type OGF factorizations we have seen so far **(Explain, clarify quantifiers)**

# A good open question to ask (LGF case)

- **Recall:** The original factorization theorem expansions for the LGF case are of the form

$$L_f(q) := \frac{1}{(q; q)_\infty} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n.$$

- We proved:  $s_{n,k} = s_{n,k} = s_o(n, k) - s_e(n, k)$
- We proved:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- The matrix entries and inverses are expressed in terms of partition theoretic functions (cf. [1])
- This gives new connections between functions in multiplicative number theory and the theory of partitions



# A good open question to ask (LGF case)

- **Recall:** The original factorization theorem expansions for the LGF case are of the form

$$L_f(q) := \frac{1}{(q; q)_\infty} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n.$$

- We proved:  $s_{n,k} = s_{n,k} = s_o(n, k) - s_e(n, k)$
- We proved:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- The matrix entries and inverses are expressed in terms of partition theoretic functions (cf. [1])
- This gives new connections between functions in multiplicative number theory and the theory of partitions

# A good open question to ask (LGF case)

- **Recall:** The original factorization theorem expansions for the LGF case are of the form

$$L_f(q) := \frac{1}{(q; q)_\infty} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n.$$

- We proved:  $s_{n,k} = s_{n,k} = s_o(n, k) - s_e(n, k)$
- We proved:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- The matrix entries and inverses are expressed in terms of partition theoretic functions (cf. [1])
- This gives new connections between functions in multiplicative number theory and the theory of partitions

# A good open question to ask (LGF case)

- **Recall:** The original factorization theorem expansions for the LGF case are of the form

$$L_f(q) := \frac{1}{(q; q)_\infty} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n.$$

- We proved:  $s_{n,k} = s_{n,k} = s_o(n, k) - s_e(n, k)$
- We proved:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- The matrix entries and inverses are expressed in terms of partition theoretic functions (cf. [1])
- This gives new connections between functions in multiplicative number theory and the theory of partitions

# A good open question to ask (LGF case)

- **Recall:** The original factorization theorem expansions for the LGF case are of the form

$$L_f(q) := \frac{1}{(q; q)_\infty} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k} f(k) \right) q^n.$$

- We proved:  $s_{n,k} = s_{n,k} = s_o(n, k) - s_e(n, k)$
- We proved:  $s_{n,k}^{-1} = \sum_{d|n} p(d-k) \mu\left(\frac{n}{d}\right)$
- The matrix entries and inverses are expressed in terms of partition theoretic functions (cf. [1])
- This gives new connections between functions in multiplicative number theory and the theory of partitions

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ The view of the OGF,  $\mathcal{C}(q) := (q; q)_\infty$ , in the LGF case being “*optimal*” (or somehow encoding the most meaningful hidden information about this sum type) is inherently **qualitative**
- ▶ How can we precisely define a corresponding **quantitative** metric with which we can express the intuition from the special case?

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ The view of the OGF,  $\mathcal{C}(q) := (q; q)_\infty$ , in the LGF case being “*optimal*” (or somehow encoding the most meaningful hidden information about this sum type) is inherently **qualitative**
- ▶ How can we precisely define a corresponding **quantitative** metric with which we can express the intuition from the special case?

# Generalizing “canonically best” OGF factorizations (cont’d)

- **Idea (first approximation):** For  $1 \times N$  vectors  $\vec{a} := (a_1, \dots, a_n)$  and  $\vec{b} := (b_1, \dots, b_N)$ , one standard way to evaluate how well matched these vectors are is given by the **(normalized) correlation statistic**

$$\text{PCorr}(\vec{a}, \vec{b}) := \frac{\frac{1}{N} \times \sum_{1 \leq j \leq N} a_j b_j}{\sqrt{\left( \sum_{1 \leq i \leq N} a_i^2 \right) \left( \sum_{1 \leq j \leq N} b_j^2 \right)}} \in [-1, 1] \quad (\text{PC-STAT})$$

- **Idea (refinement):** Use the correlation statistic in **(PC-STAT)** with infinite sequences in place of the  $N$ -vectors; These sequences should depend on (reflect key features of) the series coefficients of  $\mathcal{C}(q)^{\pm 1}$  and  $\mathcal{D}(n, k)$  (or  $\mathcal{D}^{-1}(n, k)$ ) – *Precise definitions in the thesis manuscript*

# Generalizing “canonically best” OGF factorizations (cont’d)

- **Idea (first approximation):** For  $1 \times N$  vectors  $\vec{a} := (a_1, \dots, a_n)$  and  $\vec{b} := (b_1, \dots, b_N)$ , one standard way to evaluate how well matched these vectors are is given by the **(normalized) correlation statistic**

$$\text{PCorr}(\vec{a}, \vec{b}) := \frac{\frac{1}{N} \times \sum_{1 \leq j \leq N} a_j b_j}{\sqrt{\left( \sum_{1 \leq i \leq N} a_i^2 \right) \left( \sum_{1 \leq j \leq N} b_j^2 \right)}} \in [-1, 1] \quad (\text{PC-STAT})$$

- **Idea (refinement):** Use the correlation statistic in **(PC-STAT)** with infinite sequences in place of the  $N$ -vectors; These sequences should depend on (reflect key features of) the series coefficients of  $\mathcal{C}(q)^{\pm 1}$  and  $\mathcal{D}(n, k)$  (or  $\mathcal{D}^{-1}(n, k)$ ) – *Precise definitions in the thesis manuscript*



# Visualizing the LGF case – High-level procedure I

- Can we visualize the notion of an optimally correlated OGF,

$$\mathcal{C}(q) := 1 + \sum_{n \geq 1} c_n(\mathcal{C}) q^n \in \mathbb{Z}[[q]],$$

to see if taking  $\mathcal{C}(q) := (q; q)_\infty$  is really the best? **(YES!)**

- **Notation:** Let the set of (unsigned) pentagonal numbers be defined as follows:  $\mathcal{N}_{\text{Pent}} := \{G_j : j \geq 0\}$  where

$$G_j := \frac{1}{2} \left\lfloor \frac{j}{2} \right\rfloor \left\lceil \frac{3j+1}{2} \right\rceil \mapsto \{0, 1, 2, 5, 7, 12, 15, 22, \dots\}$$

- For  $1 \leq k \leq n$ , let the correlation component

$$f_{\text{LGF}}[\mathcal{C}](n, k) := \frac{\frac{1}{n} \times [k \in \mathcal{N}_{\text{Pent}}]_\delta \times \mu^2\left(\frac{n}{k}\right) [k|n]_\delta}{\sqrt{2^{\omega(n)} \times \sum_{0 \leq k \leq n} [k \in \mathcal{N}_{\text{Pent}}]_\delta}}.$$

# Visualizing the LGF case – High-level procedure I

- Can we visualize the notion of an optimally correlated OGF,

$$\mathcal{C}(q) := 1 + \sum_{n \geq 1} c_n(\mathcal{C}) q^n \in \mathbb{Z}[[q]],$$

to see if taking  $\mathcal{C}(q) := (q; q)_\infty$  is really the best? **(YES!)**

- **Notation:** Let the set of **(unsigned) pentagonal numbers** be defined as follows:  $\mathcal{N}_{\text{Pent}} := \{G_j : j \geq 0\}$  where

$$G_j := \frac{1}{2} \left\lfloor \frac{j}{2} \right\rfloor \left\lceil \frac{3j+1}{2} \right\rceil \mapsto \{0, 1, 2, 5, 7, 12, 15, 22, \dots\}$$

- For  $1 \leq k \leq n$ , let the correlation component

$$f_{\text{LGF}}[\mathcal{C}](n, k) := \frac{\frac{1}{n} \times [k \in \mathcal{N}_{\text{Pent}}]_\delta \times \mu^2\left(\frac{n}{k}\right) [k|n]_\delta}{\sqrt{2^{\omega(n)} \times \sum_{0 \leq k \leq n} [k \in \mathcal{N}_{\text{Pent}}]_\delta}}.$$

# Visualizing the LGF case – High-level procedure I

- Can we visualize the notion of an optimally correlated OGF,

$$\mathcal{C}(q) := 1 + \sum_{n \geq 1} c_n(\mathcal{C}) q^n \in \mathbb{Z}[[q]],$$

to see if taking  $\mathcal{C}(q) := (q; q)_\infty$  is really the best? **(YES!)**

- **Notation:** Let the set of **(unsigned) pentagonal numbers** be defined as follows:  $\mathcal{N}_{\text{Pent}} := \{G_j : j \geq 0\}$  where

$$G_j := \frac{1}{2} \left\lfloor \frac{j}{2} \right\rfloor \left\lceil \frac{3j+1}{2} \right\rceil \mapsto \{0, 1, 2, 5, 7, 12, 15, 22, \dots\}$$

- For  $1 \leq k \leq n$ , let the correlation component

$$f_{\text{LGF}}[\mathcal{C}](n, k) := \frac{\frac{1}{n} \times [k \in \mathcal{N}_{\text{Pent}}]_\delta \times \mu^2\left(\frac{n}{k}\right) [k|n]_\delta}{\sqrt{2^{\omega(n)} \times \sum_{0 \leq k \leq n} [k \in \mathcal{N}_{\text{Pent}}]_\delta}}.$$

# Visualizing the LGF case – High-level procedure II

- ▶ Then for  $N \gg 1$  (as large as possible, computationally), form the **correlation matrix**

$$\overleftrightarrow{\text{CorrM}}(N) := (f_{\text{LGF}}[\mathcal{C}](n, k) [k \leq n]_{\delta})_{1 \leq n, k \leq N}.$$

- ▶ Pick a clear target image (Tux penguin, below), and partition its pixels into a  $N \times N$  grid
- ▶ Convolve the  $N$ -sized pixels with the prospective correlation matrix,  $\overleftrightarrow{\text{CorrM}}(N)$ . We should observe the following qualitative trends:
  - ▶ Less distortion (e.g., clearer results) indicates good (high) correlation
  - ▶ More distortion (e.g., blurrier results) indicates poor (low) correlation

# Visualizing the LGF case – High-level procedure II

- ▶ Then for  $N \gg 1$  (as large as possible, computationally), form the **correlation matrix**

$$\overleftrightarrow{\text{CorrM}}(N) := (f_{\text{LGF}}[\mathcal{C}](n, k) [k \leq n]_{\delta})_{1 \leq n, k \leq N}.$$

- ▶ Pick a clear target image (**Tux** penguin, below), and partition its pixels into a  $N \times N$  grid
- ▶ Convolve the  $N$ -sized pixels with the prospective correlation matrix,  $\overleftrightarrow{\text{CorrM}}(N)$ . We should observe the following qualitative trends:
  - ▶ Less distortion (e.g., clearer results) indicates good (high) correlation
  - ▶ More distortion (e.g., blurrier results) indicates poor (low) correlation

# Visualizing the LGF case – High-level procedure II

- ▶ Then for  $N \gg 1$  (as large as possible, computationally), form the **correlation matrix**

$$\overleftrightarrow{\text{CorrM}}(N) := (f_{\text{LGF}}[C](n, k) [k \leq n]_{\delta})_{1 \leq n, k \leq N}.$$

- ▶ Pick a clear target image (**Tux** penguin, below), and partition its pixels into a  $N \times N$  grid
- ▶ Convolve the  $N$ -sized pixels with the prospective correlation matrix,  $\overleftrightarrow{\text{CorrM}}(N)$ . We should observe the following qualitative trends:
  - ▶ Less distortion (e.g., clearer results) indicates good (high) correlation
  - ▶ More distortion (e.g., blurrier results) indicates poor (low) correlation

# Visualizing the LGF case – High-level procedure II

- ▶ Then for  $N \gg 1$  (as large as possible, computationally), form the **correlation matrix**

$$\overleftrightarrow{\text{CorrM}}(N) := (f_{\text{LGF}}[C](n, k) [k \leq n]_{\delta})_{1 \leq n, k \leq N}.$$

- ▶ Pick a clear target image (**Tux** penguin, below), and partition its pixels into a  $N \times N$  grid
- ▶ Convolve the  $N$ -sized pixels with the prospective correlation matrix,  $\overleftrightarrow{\text{CorrM}}(N)$ . We should observe the following qualitative trends:
  - ▶ Less distortion (e.g., clearer results) indicates good (high) correlation
  - ▶ More distortion (e.g., blurrier results) indicates poor (low) correlation

# Visualizing the LGF case (cont'd)




(a) *Original image.*



(b)  $\mathcal{C}(q) := (q; q)_{\infty}$ .



(c)  $\mathcal{C}(q) := (q; q)_{\infty}^{-1}$ .



(d)  $\mathcal{C}(q) := (q^2; q^5)_{\infty}$ .



(e)  $\mathcal{C}(q) := (1 - q)^{-\frac{3}{2}}$ .



(f)  $\mathcal{C}(q) := (1 - q)^{-1}$ .



# Generalizing “canonically best” OGF factorizations (cont’d)

- Based on inspection, the visual **Tux** examples for the LGF case, seem to conform to the “*ideal*” case expectation:  
That is, we cannot do better than to choose  $\mathcal{C}(q) := (q; q)_\infty$  (as we had defined to be qualitatively optimal)
- For more general convolution sum types, we seek to **maximize** (**minimize**) the series

$$\text{Corr}(\mathcal{C}, \mathcal{D}) := \sum_{n \geq 1} \frac{\frac{1}{n} \times \sum_{k=1}^n |c_k(\mathcal{C}) \mathcal{D}^{-1}(n, k)|}{\sqrt{\left( \sum_{k=1}^n c_k(\mathcal{C})^2 \right) \left( \sum_{k=1}^n \mathcal{D}^{-1}(n, k)^2 \right)}}$$

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ Based on inspection, the visual **Tux** examples for the LGF case, seem to conform to the “*ideal*” case expectation:  
That is, we cannot do better than to choose  $\mathcal{C}(q) := (q; q)_\infty$  (as we had defined to be qualitatively optimal)
- ▶ For more general convolution sum types, we seek to **maximize** (**minimize**) the series

$$\text{Corr}(\mathcal{C}, \mathcal{D}) := \sum_{n \geq 1} \frac{\frac{1}{n} \times \sum_{k=1}^n |c_k(\mathcal{C}) \mathcal{D}^{-1}(n, k)|}{\sqrt{\left( \sum_{k=1}^n c_k(\mathcal{C})^2 \right) \left( \sum_{k=1}^n \mathcal{D}^{-1}(n, k)^2 \right)}}$$

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ How best to approach finding the optimal OGF? What if we restrict to integer coefficients with  $\mathcal{C}(0) := 1$ ?
- ▶ This is an open topic; Some preliminary conjectures and discussion are given in the last section of the thesis.
- ▶ **Historical notes on correlation statistic tactics:**
  - ▶ There is literature documenting and motivating the use of statistical analysis to study number theoretic objects
  - ▶ Montgomery: Pair correlation to study the non-trivial zeros of  $\zeta(s)$
  - ▶ Hejhal, Rudnick, Sarnak and Odlyzko, respectively, built on HM's work to apply statistical analysis (correlation statistics) to  $L$ -functions, the Gaussian Unitary Ensemble (GUE) and in applications to random matrix theory
  - ▶ See the survey in [9]

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ How best to approach finding the optimal OGF? What if we restrict to integer coefficients with  $\mathcal{C}(0) := 1$ ?
- ▶ This is an open topic; Some preliminary conjectures and discussion are given in the last section of the thesis.
- ▶ **Historical notes on correlation statistic tactics:**
  - ▶ There is literature documenting and motivating the use of statistical analysis to study number theoretic objects
  - ▶ Montgomery: Pair correlation to study the non-trivial zeros of  $\zeta(s)$
  - ▶ Hejhal, Rudnick, Sarnak and Odlyzko, respectively, built on HM's work to apply statistical analysis (correlation statistics) to  $L$ -functions, the Gaussian Unitary Ensemble (GUE) and in applications to random matrix theory
  - ▶ See the survey in [9]

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ How best to approach finding the optimal OGF? What if we restrict to integer coefficients with  $\mathcal{C}(0) := 1$ ?
- ▶ This is an open topic; Some preliminary conjectures and discussion are given in the last section of the thesis.
- ▶ **Historical notes on correlation statistic tactics:**
  - ▶ There is literature documenting and motivating the use of statistical analysis to study number theoretic objects
  - ▶ Montgomery: Pair correlation to study the non-trivial zeros of  $\zeta(s)$
  - ▶ Hejhal, Rudnick, Sarnak and Odlyzko, respectively, built on HM's work to apply statistical analysis (correlation statistics) to  $L$ -functions, the Gaussian Unitary Ensemble (GUE) and in applications to random matrix theory
  - ▶ See the survey in [9]

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ How best to approach finding the optimal OGF? What if we restrict to integer coefficients with  $\mathcal{C}(0) := 1$ ?
- ▶ This is an open topic; Some preliminary conjectures and discussion are given in the last section of the thesis.
- ▶ **Historical notes on correlation statistic tactics:**
  - ▶ There is literature documenting and motivating the use of statistical analysis to study number theoretic objects
  - ▶ Montgomery: Pair correlation to study the non-trivial zeros of  $\zeta(s)$
  - ▶ Hejhal, Rudnick, Sarnak and Odlyzko, respectively, built on HM's work to apply statistical analysis (correlation statistics) to  $L$ -functions, the Gaussian Unitary Ensemble (GUE) and in applications to random matrix theory
  - ▶ See the survey in [9]

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ How best to approach finding the optimal OGF? What if we restrict to integer coefficients with  $\mathcal{C}(0) := 1$ ?
- ▶ This is an open topic; Some preliminary conjectures and discussion are given in the last section of the thesis.
- ▶ **Historical notes on correlation statistic tactics:**
  - ▶ There is literature documenting and motivating the use of statistical analysis to study number theoretic objects
  - ▶ Montgomery: Pair correlation to study the non-trivial zeros of  $\zeta(s)$
  - ▶ Hejhal, Rudnick, Sarnak and Odlyzko, respectively, built on HM’s work to apply statistical analysis (correlation statistics) to  $L$ -functions, the Gaussian Unitary Ensemble (GUE) and in applications to random matrix theory
  - ▶ See the survey in [9]

# Generalizing “canonically best” OGF factorizations (cont’d)

- ▶ How best to approach finding the optimal OGF? What if we restrict to integer coefficients with  $\mathcal{C}(0) := 1$ ?
- ▶ This is an open topic; Some preliminary conjectures and discussion are given in the last section of the thesis.
- ▶ **Historical notes on correlation statistic tactics:**
  - ▶ There is literature documenting and motivating the use of statistical analysis to study number theoretic objects
  - ▶ Montgomery: Pair correlation to study the non-trivial zeros of  $\zeta(s)$
  - ▶ Hejhal, Rudnick, Sarnak and Odlyzko, respectively, built on HM’s work to apply statistical analysis (correlation statistics) to  $L$ -functions, the Gaussian Unitary Ensemble (GUE) and in applications to random matrix theory
  - ▶ See the survey in [9]



# Concluding remarks

The End

Questions?

Comments?

Feedback?

Thank you for your time!

# References I

- [1] G. E. Andrews.  
*The Theory of Partitions*.  
Cambridge University Press, 1984.
- [2] D. E. Knuth.  
Two notes on notation.  
*Amer. Math. Monthly*, 99(5):403–422, 1992.
- [3] M. Merca.  
The Lambert series factorization theorem.  
*Ramanujan Journal*, pages 1–19, 2017.
- [4] M. Merca and M. D. Schmidt.  
Generating special arithmetic functions by Lambert series factorizations.  
*Contributions to Discrete Mathematics*, 2018.
- [5] M. Merca and M. D. Schmidt.  
A partition identity related to Stanley’s theorem.  
*Amer. Math. Monthly*, 125(10):929–933, 2018.

# References II

- [6] M. Merca and M. D. Schmidt.  
Factorization theorems for generalized Lambert series and applications.  
*Ramanujan Journal*, 2019.
- [7] M. Merca and M. D. Schmidt.  
The partition function  $p(n)$  in terms of the classical Möbius function.  
*Ramanujan Journal*, 49:87–96, 2019.
- [8] H. Mousavi and M. D. Schmidt.  
Factorization theorems for relatively prime divisor sums, GCD sums and generalized Ramanujan sums, 2020.
- [9] S. J. Miller O. Barrett, F. W. K. Firk and C. Turnage-Butterbaugh.  
From quantum systems to  $L$ -functions: Pair correlation statistics and beyond.  
*Open Problems in Mathematics*, 2016.  
Available online: [https://web.williams.edu/Mathematics/sjmiller/public\\_html/math/papers/NTandQM\\_BMTB60arxiv.pdf](https://web.williams.edu/Mathematics/sjmiller/public_html/math/papers/NTandQM_BMTB60arxiv.pdf).
- [10] B. Reznick.  
Course notes, Spring 2012, chapter 2, 2012.  
Available online: <https://faculty.math.illinois.edu/~reznick/595-ch2.pdf>.

# References III

- [11] M. D. Schmidt.  
Continued fractions and  $q$ -series generating functions for the generalized sum-of-divisors functions.  
*Journal of Number Theory*, 180:579–605, 2017.
- [12] M. D. Schmidt.  
Jacobi-type continued fractions for the ordinary generating functions of generalized factorial functions.  
*Journal of Integer Sequences*, 20(17.3.4), 2017.
- [13] M. D. Schmidt.  
Zeta series generating function transformations related to polylogarithm functions and the  $k$ -order harmonic numbers.  
*Online journal of analytic combinatorics*, 12(2), 2017.
- [14] M. D. Schmidt.  
Continued fractions for square series generating functions.  
*Ramanujan Journal*, 46:795–820, 2018.

# References IV

- [15] M. D. Schmidt.  
Jacobi-type continued fractions and congruences for binomial coefficients.  
*INTEGERS*, 18(A46), 2018.
- [16] M. D. Schmidt.  
New congruences and finite difference equations for generalized factorial functions.  
*INTEGERS*, 18(A78), 2018.
- [17] M. D. Schmidt.  
New recurrence relations and matrix equations for arithmetic functions generated by Lambert series.  
*Acta Arithmetica*, 181, 2018.
- [18] M. D. Schmidt.  
Zeta series generating function transformations related to generalized Stirling numbers and partial sums of the Hurwitz zeta function.  
*Online journal of analytic combinatorics*, 13(3), 2018.

# References V

- [19] M. D. Schmidt.  
A short note on integral transformations and conversion formulas for sequence generating functions.  
*Axioms*, 8(2), 2019.
- [20] M. D. Schmidt.  
Combinatorial sums and identities involving generalized divisor functions with bounded divisors.  
*INTEGERS*, 20(A85), 2020.
- [21] M. D. Schmidt.  
Exact formulas for the generalized sum-of-divisors functions, 2021.
- [22] M. D. Schmidt.  
Square series generating function transformations.  
2017, 8(2):125–156, *Journal of Inequalities and Special Functions*.
- [23] H. S. Wilf.  
*Generatingfunctionology*.  
Academic Press, 1994.

# Extra slides

## Extra slides and references

# LGFs – Other results (cont'd)

- ▶ The function  $h^{-1}$  is called the **Dirichlet inverse of  $h$**  if  $h * h^{-1} = h^{-1} * h = \varepsilon$  where  $\varepsilon(n) = \delta_{n,1}$  is the multiplicative identity with respect to Dirichlet convolution
- ▶ The function  $h^{-1}$  exists and is unique iff  $h(1) \neq 0$ .
- ▶ When  $h^{-1}$  exists, it is computed recursively via the formula

$$h^{-1}(n) = \begin{cases} \frac{1}{h(1)}, & n = 1; \\ -\frac{1}{h(1)} \times \sum_{\substack{d|n \\ d > 1}} h(d) h^{-1}\left(\frac{n}{d}\right), & n \geq 2. \end{cases}$$



# LGFs – Other results (cont'd)

- ▶ The function  $h^{-1}$  is called the **Dirichlet inverse of  $h$**  if  $h * h^{-1} = h^{-1} * h = \varepsilon$  where  $\varepsilon(n) = \delta_{n,1}$  is the multiplicative identity with respect to Dirichlet convolution
- ▶ The function  $h^{-1}$  exists and is unique iff  $h(1) \neq 0$ .
- ▶ When  $h^{-1}$  exists, it is computed recursively via the formula

$$h^{-1}(n) = \begin{cases} \frac{1}{h(1)}, & n = 1; \\ -\frac{1}{h(1)} \times \sum_{\substack{d|n \\ d > 1}} h(d) h^{-1}\left(\frac{n}{d}\right), & n \geq 2. \end{cases}$$

# LGFs – Other results (cont'd)

- ▶ The function  $h^{-1}$  is called the **Dirichlet inverse of  $h$**  if  $h * h^{-1} = h^{-1} * h = \varepsilon$  where  $\varepsilon(n) = \delta_{n,1}$  is the multiplicative identity with respect to Dirichlet convolution
- ▶ The function  $h^{-1}$  exists and is unique iff  $h(1) \neq 0$ .
- ▶ When  $h^{-1}$  exists, it is computed recursively via the formula

$$h^{-1}(n) = \begin{cases} \frac{1}{h(1)}, & n = 1; \\ -\frac{1}{h(1)} \times \sum_{\substack{d|n \\ d > 1}} h(d) h^{-1}\left(\frac{n}{d}\right), & n \geq 2. \end{cases}$$

# LGFs – Other results (cont'd)

n	$h^{-1}(n)$	n	$h^{-1}(n)$	n	$h^{-1}(n)$
1	$\frac{1}{h(1)}$	4	$-\frac{h(1)h(4) - h(2)^2}{h(1)^3}$	7	$-\frac{h(7)}{h(1)^2}$
2	$-\frac{h(2)}{h(1)^2}$	5	$-\frac{h(5)}{h(1)^2}$	8	$-\frac{h(2)^3 - 2h(1)h(4)h(2) + h(1)^2h(8)}{h(1)^4}$
3	$-\frac{h(3)}{h(1)^2}$	6	$-\frac{h(1)h(6) - 2h(2)h(3)}{h(1)^3}$	9	$-\frac{h(1)h(9) - h(3)^2}{h(1)^3}$

# LGFs – Other results (cont'd)

- For fixed  $f, g$  and any OGF  $\mathcal{C}(q)$  with  $\mathcal{C}(0) \neq 0$ , we define

$$\sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \frac{1}{\mathcal{C}(q)} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k}[\mathcal{C}] f(k) \right) q^n, \quad (\text{i})$$

and let

$$\sum_{n \geq 1} \frac{(f * g)(n)q^n}{1 - q^n} = \frac{1}{\mathcal{C}(q)} \times \sum_{n \geq 1} \left( \sum_{k=1}^n \tilde{s}_{n,k}[\mathcal{C}](g) f(k) \right) q^n. \quad (\text{ii})$$

- We can prove:  $\tilde{s}_{n,k}[\mathcal{C}](g) = \sum_{j=1}^n s_{n,kj}[\mathcal{C}] g(j)$

# LGFs – Other results (cont'd)

- For fixed  $f, g$  and any OGF  $\mathcal{C}(q)$  with  $\mathcal{C}(0) \neq 0$ , we define

$$\sum_{n \geq 1} \frac{f(n)q^n}{1 - q^n} = \frac{1}{\mathcal{C}(q)} \times \sum_{n \geq 1} \left( \sum_{k=1}^n s_{n,k}[\mathcal{C}] f(k) \right) q^n, \quad (\text{i})$$

and let

$$\sum_{n \geq 1} \frac{(f * g)(n)q^n}{1 - q^n} = \frac{1}{\mathcal{C}(q)} \times \sum_{n \geq 1} \left( \sum_{k=1}^n \tilde{s}_{n,k}[\mathcal{C}](g) f(k) \right) q^n. \quad (\text{ii})$$

- We can prove:  $\tilde{s}_{n,k}[\mathcal{C}](g) = \sum_{j=1}^n s_{n,kj}[\mathcal{C}] g(j)$

# LGFs – Other results (cont'd)

Table of the inverse matrices,  $\tilde{s}_{n,k}^{-1}[\mathcal{C}](g)$ :

	1	2	3	4
1	1	0	0	0
2	$-g(2)$	1	0	0
3	$1 - g(3)$	1	1	0
4	$g(2)^2 - g(4) + 2$	$1 - g(2)$	1	1
5	$4 - g(5)$	3	2	1
6	$2g(3)g(2) - g(2) - g(6) + 5$	$-g(2) - g(3) + 3$	$2 - g(2)$	2
7	$10 - g(7)$	7	5	3
8	$-g(2)^3 + 2g(4)g(2) - 2g(2) - g(8) + 12$	$g(2)^2 - g(2) - g(4) + 9$	$6 - g(2)$	$4 - g(2)$
9	$g(3)^2 - g(3) - g(9) + 20$	$14 - g(3)$	$10 - g(3)$	7
10	$2g(5)g(2) - 4g(2) - g(10) + 25$	$-3g(2) - g(5) + 18$	$13 - 2g(2)$	$10 - g(2)$

(Special case where  $g(1) := 1$  for simplicity.)

# Factorization theorems for LGFs – Variants I

- **Notation:** When  $C(q) = (q; q)_\infty$  we write  $s_{n,k}^{-1}[\mathcal{C}](g) \equiv s_{n,k}^{-1}(g)$
- **Notation:** Let the function  $p_k(n) := p(n - k)$
- For  $n \geq 1$ , let

$$f^{-1}(n) := \left( D_{n,f} + \frac{\varepsilon}{f(1)} \right) (n).$$

(The function  $D_{n,f}(n)$  can be defined recursively by partial sums of multiple convolutions of  $f$  with itself.)

- **Theorem:** We can prove that

$$\sum_{d|n} s_{n,k}^{-1}(g) = p_k(n) + (p_k * D_{n,g})(n),$$

# Factorization theorems for LGFs – Variants I

- ▶ **Notation:** When  $C(q) = (q; q)_\infty$  we write  $s_{n,k}^{-1}[\mathcal{C}](g) \equiv s_{n,k}^{-1}(g)$
- ▶ **Notation:** Let the function  $p_k(n) := p(n - k)$
- ▶ For  $n \geq 1$ , let

$$f^{-1}(n) := \left( D_{n,f} + \frac{\varepsilon}{f(1)} \right) (n).$$

(The function  $D_{n,f}(n)$  can be defined recursively by partial sums of multiple convolutions of  $f$  with itself.)

- ▶ **Theorem:** We can prove that

$$\sum_{d|n} s_{n,k}^{-1}(g) = p_k(n) + (p_k * D_{n,g})(n),$$



# Factorization theorems for LGFs – Variants I

- **Notation:** When  $C(q) = (q; q)_\infty$  we write  $s_{n,k}^{-1}[\mathcal{C}](g) \equiv s_{n,k}^{-1}(g)$
- **Notation:** Let the function  $p_k(n) := p(n - k)$
- For  $n \geq 1$ , let

$$f^{-1}(n) := \left( D_{n,f} + \frac{\varepsilon}{f(1)} \right) (n).$$

(The function  $D_{n,f}(n)$  can be defined recursively by partial sums of multiple convolutions of  $f$  with itself.)

- **Theorem:** We can prove that

$$\sum_{d|n} s_{n,k}^{-1}(g) = p_k(n) + (p_k * D_{n,g})(n),$$

# Factorization theorems for LGFs – Variants I

- **Notation:** When  $C(q) = (q; q)_\infty$  we write  $s_{n,k}^{-1}[C](g) \equiv s_{n,k}^{-1}(g)$
- **Notation:** Let the function  $p_k(n) := p(n - k)$
- For  $n \geq 1$ , let

$$f^{-1}(n) := \left( D_{n,f} + \frac{\varepsilon}{f(1)} \right) (n).$$

(The function  $D_{n,f}(n)$  can be defined recursively by partial sums of multiple convolutions of  $f$  with itself.)

- **Theorem:** We can prove that

$$\sum_{d|n} s_{n,k}^{-1}(g) = p_k(n) + (p_k * D_{n,g})(n),$$

# Factorization theorems for LGFs – Variants II

- We also considered factorization theorems for Hadamard products:

$$\sum_{d|n} a_{fg}(d) := \underbrace{\left( \sum_{d|n} f_d \right) \times \left( \sum_{d|n} g_d \right)}_{:=fg(n)}.$$

where

$$\sum_{n \geq 1} \frac{a_{fg}(n) q^n}{1 - q^n} = \frac{1}{(q; q)_{\infty}} \times \sum_{n \geq 1} \sum_{k=1}^n h_{n,k}(f) g_k q^n,$$

# Factorization theorems for LGFs – Variants III

► **Notation:** Let  $\tilde{f}(n) := \sum_{d|n} f_d$

► We prove:

$$h_{n,k}(f) = \tilde{f}(n) [k|n]_{\delta} + \sum_{j=1}^{\left\lfloor \frac{\sqrt{24(n-k)+1}-b}{6} \right\rfloor} (-1)^j \tilde{f}\left(n - \frac{j(3j+b)}{2}\right) \left[k|n - \frac{j(3j+b)}{2}\right]_{\delta}$$

► We prove:  $h_{n,k}^{-1}(f) = \sum_{d|n} \frac{\rho(d-k)}{\tilde{f}(d)} \mu\left(\frac{n}{d}\right)$

# Factorization theorems for LGFs – Variants III

- **Notation:** Let  $\tilde{f}(n) := \sum_{d|n} f_d$
- We prove:

$$h_{n,k}(f) = \tilde{f}(n) [k|n]_{\delta} + \sum_{j=1}^{\left\lfloor \frac{\sqrt{24(n-k)+1}-b}{6} \right\rfloor} (-1)^j \tilde{f}\left(n - \frac{j(3j+b)}{2}\right) \left[ k|n - \frac{j(3j+b)}{2} \right]_{\delta}$$

- We prove:  $h_{n,k}^{-1}(f) = \sum_{d|n} \frac{\rho(d-k)}{\tilde{f}(d)} \mu\left(\frac{n}{d}\right)$

# Factorization theorems for LGFs – Variants III

- **Notation:** Let  $\tilde{f}(n) := \sum_{d|n} f_d$
- We prove:

$$h_{n,k}(f) = \tilde{f}(n) [k|n]_{\delta} + \sum_{j=1}^{\left\lfloor \frac{\sqrt{24(n-k)+1}-b}{6} \right\rfloor} (-1)^j \tilde{f}\left(n - \frac{j(3j+b)}{2}\right) \left[k|n - \frac{j(3j+b)}{2}\right]_{\delta}$$

- We prove:  $h_{n,k}^{-1}(f) = \sum_{d|n} \frac{p(d-k)}{\tilde{f}(d)} \mu\left(\frac{n}{d}\right)$

# Factorization theorems for LGFs – Variants IV

► **Corollaries:** We have so-termed “*exotic*” sums of the form

$$\begin{aligned}\phi(n) &= \sum_{k=1}^n \sum_{d|n} \frac{p(d-k)}{d} \mu\left(\frac{n}{d}\right) \left[ k^2 + \sum_{b=\pm 1} \right. \\ &\quad \left. + \sum_{j=1}^{\lfloor \frac{\sqrt{24k-23}-b}{6} \rfloor} (-1)^j \left( k - \frac{j(3j+b)}{2} \right)^2 \right] \\ n^s &= \sum_{k=1}^n \sum_{d|n} \frac{p(d-k)}{\sigma_t(d)} \mu\left(\frac{n}{d}\right) \left[ \sigma_t(k) \sigma_s(k) \right. \\ &\quad \left. + \sum_{b=\pm 1} \sum_{j=1}^{\lfloor \frac{\sqrt{24k+1}-b}{6} \rfloor} (-1)^j \sigma_t\left(k - \frac{j(3j+b)}{2}\right) \sigma_s\left(k - \frac{j(3j+b)}{2}\right) \right].\end{aligned}$$