





**Quantum  
Phenomena 1**  
xx.06.2023  
Lecture Nr X

Maxim Mai

Office hours: ...  
Room: ...  
Email: ...

# TUNNELING



## Reminder from the last lecture

- Quantum theory is probabilistic:


$$P(x) = |\psi(x)|^2$$

- Wave functions obey Schrödinger equation (time independent):

$$\frac{d^2}{dx^2}\psi(x) = -\frac{2m}{\hbar^2} (E - V(x)) \psi(x)$$

- Wave functions are smooth and continuous.

# Potential barrier



Quantum  
Phenomena 1  
xx.06.2023  
Lecture Nr X

Maxim Mai

Office hours: ...  
Room: ...  
Email: ...

# Potential barrier



Solutions of the Schrödinger equation

$$\Psi_1(x) = A e^{ikx} + B e^{-ikx}$$

$$k = \frac{\sqrt{2mE}}{\hbar}$$




Quantum  
Phenomena 1  
xx.06.2023  
Lecture Nr X

Maxim Mai

Office hours: ...  
Room: ...  
Email: ...

# Potential barrier



Solutions of the Schrödinger equation


$$\Psi_1(x) = A e^{ikx} + B e^{-ikx}$$

$$k = \frac{\sqrt{2mE}}{\hbar}$$

$$\Psi_3(x) = F e^{ikx} + G e^{-ikx}$$


$$k = \frac{\sqrt{2mE}}{\hbar}$$

# Potential barrier




## TODAYS QUIZ:

**A.**



**B.**




Quantum  
Phenomena 1  
xx.06.2023  
Lecture Nr X

Maxim Mai


Office hours: ...  
Room: ...  
Email: ...

# Potential barrier




## TODAYS QUIZ:

**A.**



80%

**B.**



20%




Quantum  
Phenomena 1  
xx.06.2023  
Lecture Nr X

Maxim Mai

Office hours: ...  
Room: ...  
Email: ...



# Potential barrier



Solutions of the Schrödinger equation

$$\Psi_1(x) = A e^{ikx} + B e^{-ikx}$$

$$k = \frac{\sqrt{2mE}}{\hbar}$$


$$\Psi_2(x) = C e^{\kappa x} + D e^{-\kappa x}$$

$$\kappa = \frac{\sqrt{2m(V_0 - E)}}{\hbar}$$

$$\Psi_3(x) = F e^{ikx} + G e^{-ikx}$$

$$k = \frac{\sqrt{2mE}}{\hbar}$$

# Potential barrier




Continuity and smoothness

$$\Psi_1(x) = A e^{ikx} + B e^{-ikx}$$

$$\Psi_2(x) = C e^{\kappa x} + D e^{-\kappa x}$$

$$\Psi_3(x) = F e^{ikx} + G e^{-ikx}$$

# Potential barrier



Continuity and smoothness


$$\Psi_1(x) = A e^{ikx} + B e^{-ikx}$$

$$\Psi_2(x) = C e^{\kappa x} + D e^{-\kappa x}$$

$$\Psi_3(x) = F e^{ikx} + G e^{-ikx}$$

$$\left. \begin{aligned} \Psi_1(x) &= \Psi_2(x) \\ \Psi_1'(x) &= \Psi_2'(x) \end{aligned} \right|_{x=0}$$

# Potential barrier



Continuity and smoothness

$$\Psi_1(x) = A e^{ikx} + B e^{-ikx}$$

$$\Psi_2(x) = C e^{\kappa x} + D e^{-\kappa x}$$


$$\Psi_3(x) = F e^{ikx} + G e^{-ikx}$$

$$\left. \begin{array}{l} \Psi_1(x) = \Psi_2(x) \\ \Psi_1'(x) = \Psi_2'(x) \end{array} \right|_{x=0}$$

$\Downarrow$

$$\left( \begin{array}{l} A + B = C + D \\ ik(A - B) = \kappa(C - D) \end{array} \right)$$

# Potential barrier



Continuity and smoothness

$$\Psi_1(x) = A e^{ikx} + B e^{-ikx}$$

$$\Psi_2(x) = C e^{\kappa x} + D e^{-\kappa x}$$

$$\Psi_3(x) = F e^{ikx} + G e^{-ikx}$$


$$\left. \begin{array}{l} \Psi_1(x) = \Psi_2(x) \\ \Psi'_1(x) = \Psi'_2(x) \end{array} \right|_{x=0}$$

$$\left. \begin{array}{l} \Psi_2(x) = \Psi_3(x) \\ \Psi'_2(x) = \Psi'_3(x) \end{array} \right|_{x=L}$$

$\Downarrow$

$$\left( \begin{array}{l} A + B = C + D \\ ik(A - B) = \kappa(C - D) \end{array} \right)$$

# Potential barrier



Continuity and smoothness

$$\Psi_1(x) = A e^{ikx} + B e^{-ikx}$$

$$\Psi_2(x) = C e^{\kappa x} + D e^{-\kappa x}$$

$$\Psi_3(x) = F e^{ikx} + G e^{-ikx}$$

$$\left. \begin{array}{l} \Psi_1(x) = \Psi_2(x) \\ \Psi'_1(x) = \Psi'_2(x) \end{array} \right|_{x=0}$$

$$\left. \begin{array}{l} \Psi_2(x) = \Psi_3(x) \\ \Psi'_2(x) = \Psi'_3(x) \end{array} \right|_{x=L}$$

$\Downarrow$


$$\left( \begin{array}{l} A + B = C + D \\ ik(A - B) = \kappa(C - D) \end{array} \right)$$

$\Downarrow$

$$\left( \begin{array}{l} \ddots = \ddots \\ \ddots = \ddots \end{array} \right)$$

**Homework**

# Potential barrier




Discussion (non-classical effects)

- Probability for finding particle in the wall

$$P_{\Psi}(x) > 0 \quad \text{for } x \in [0, L)$$

# Potential barrier



Discussion (non-classical effects)

- Probability for “*tunneling*”


$$P_{\Psi}(x) \sim e^{-2\kappa L} > 0 \quad \text{for } x \in [L, \infty)$$



# APPLICATIONS

- $\alpha$ -decay/electronic circuit components/...
- Scanning tunneling microscope (STM)

Binnig and Rohrer, Nobel Prize in Physics in 1986



Quantum  
Phenomena 1  
xx.06.2023  
Lecture Nr X


Maxim Mai

Office hours: ...  
Room: ...  
Email: ...

# APPLICATIONS

- $\alpha$ -decay/electronic circuit components/...
- Scanning tunneling microscope (STM)

Binnig and Rohrer, Nobel Prize in Physics in 1986



Further examples: [https://en.wikipedia.org/wiki/Scanning\\_tunneling\\_microscope](https://en.wikipedia.org/wiki/Scanning_tunneling_microscope)



Quantum  
Phenomena 1  
xx.06.2023  
Lecture Nr X

Maxim Mai

Office hours: ...  
Room: ...  
Email: ...

