

# SPH simulations for space defense

Maximilian Rutz


July 15, 2020

# Roadmap

1 Dart and Hera Missions

2 SPH setup

3 SPH results



# Dart Mission

- Launch in July 2021 on a SpaceX Falcon 9
- Impact in fall 2022
- Impact at 0.07 au to Earth, 29x Earth-Moon, 1/5x Earth-Mars
- Observations with LICIACube and earth based telescopes

# Dart Mission

- Launch in July 2021 on a SpaceX Falcon 9
- Impact in fall 2022
- Impact at 0.07 au to Earth, 29x Earth-Moon, 1/5x Earth-Mars
- Observations with LICIACube and earth based telescopes

# Dart Mission

- Launch in July 2021 on a SpaceX Falcon 9
- Impact in fall 2022
- Impact at 0.07 au to Earth, 29x Earth-Moon, 1/5x Earth-Mars
- Observations with LICIACube and earth based telescopes

# Dart Mission

- Launch in July 2021 on a SpaceX Falcon 9
- Impact in fall 2022
- Impact at 0.07 au to Earth, 29x Earth-Moon, 1/5x Earth-Mars
- Observations with LICIACube and earth based telescopes

# Hera Mission





# Hera Mission

- Launch in 2024
- Arrival in 2026
- Why a second mission?
  - Dust cloud after impact
  - Reduce uncertainty of orbital shift
  - Politics ...

# Hera Mission

- Launch in 2024
- Arrival in 2026
- Why a second mission?
  - Dust cloud after impact
  - Reduce uncertainty of orbital shift
  - Politics ...

# Hera Mission

- Launch in 2024
- Arrival in 2026
- Why a second mission?
  - Dust cloud after impact
  - Reduce uncertainty of orbital shift
  - Politics ...

# Hera Mission

- Launch in 2024
- Arrival in 2026
- Why a second mission?
  - Dust cloud after impact
  - Reduce uncertainty of orbital shift
  - Politics ...

# Hera Mission

- Launch in 2024
- Arrival in 2026
- Why a second mission?
  - Dust cloud after impact
  - Reduce uncertainty of orbital shift
  - Politics ...

# Hera Mission

- Launch in 2024
- Arrival in 2026
- Why a second mission?
  - Dust cloud after impact
  - Reduce uncertainty of orbital shift
  - Politics ...


# Hera Mission

- Launch in 2024
- Arrival in 2026
- Why a second mission?
  - Dust cloud after impact
  - Reduce uncertainty of orbital shift
  - Politics ...



# DART

## Double Asteroid Redirection Test





# Target

<https://www.nasa.gov/planetarydefense/dart>

Dimorphos orbiting Didymos

<https://dart.jhuapl.edu/Gallery/media/graphics/lg/DART>

# Impactor

<https://dart.jhuapl.edu/Mission/Impactor-Spacecraft.php>

- 1.2 1.3 1.3 meters
- artificial viscosity

# Impactor

<https://dart.jhuapl.edu/Mission/Impactor-Spacecraft.php>

- 1.2 1.3 1.3 meters
- artificial viscosity

# Impactor

<https://dart.jhuapl.edu/Mission/Impactor-Spacecraft.php>

- 1.2 1.3 1.3 meters
- artificial viscosity

# SPH method

Smoothed particle hydrodynamics

# Miluphcuda

Smoothed particle hydrodynamics

# Miluphcuda setup

- x3 Kernel function
- Runge Kutta fourth order
- no self gravity
- $p$ - $\alpha$  porosity - micro vs macroporosity

# Miluphcuda setup

- x3 Kernel function
- Runge Kutta fourth order
- no self gravity
- $p-\alpha$  porosity - micro vs macroporosity



# Miluphcuda setup

- x3 Kernel function
- Runge Kutta fourth order
- no self gravity
- $p-\alpha$  porosity - micro vs macroporosity

# Miluphcuda setup

- x3 Kernel function
- Runge Kutta fourth order
- no self gravity
- $p$ - $\alpha$  porosity - micro vs macroporosity

# Miluphcuda setup

- x3 Kernel function
- Runge Kutta fourth order
- no self gravity
- $p$ - $\alpha$  porosity - micro vs macroporosity

# Initial conditions setup

- x3 Kernel function
- artificial viscosity
- Runge Kutta Fourth order
- no self gravity
- $p-\alpha$  porosity

# Initial conditions setup

- x3 Kernel function
- artificial viscosity
- Runge Kutta Fourth order
- no self gravity
- $p-\alpha$  porosity

# Initial conditions setup

- x3 Kernel function
- artificial viscosity
- Runge Kutta Fourth order
- no self gravity
- $p-\alpha$  porosity

# Initial conditions setup

- x3 Kernel function
- artificial viscosity
- Runge Kutta Fourth order
- no self gravity
- $p-\alpha$  porosity

# Initial conditions setup

- x3 Kernel function
- artificial viscosity
- Runge Kutta Fourth order
- no self gravity
- $p-\alpha$  porosity



# smoothing length

- how can resolution be locally increased with SPH method (different radii and sml) - limit of sml  $\rightarrow 0$  is normal hydrodynamics??

# Goal of simulations

Compare numerical results with observations to:

- 1 test numerical codes
- 2 identify target material and properties

# Goal of simulations

Compare numerical results with observations to:

- ① test numerical codes
- ② identify target material and properties

# Beta factor


Momentum change because of ejecta:  $\beta = 1 + \frac{P_{\text{ejecta}}}{P_{\text{impactor}}}$

# Beta factor

Momentum change because of ejecta:  $\beta = 1 + \frac{p_{ejecta}}{p_{impactor}}$

The DART impact into different targets can produce the same  $\beta$ , but different craters.

Measurements of **both**  $\beta$  and crater size/morphology **together** can be diagnostic of target properties.



# Impact angle

Not seen in 2d grid codes Raducan

# Personal observations about SPH

- A lot of individual physics implementable
- Many different codes available
- Many parameters needed to reproduce results of others