Ordering the cusp(AdS-CFT)

Maximilian Weinhold

November 2026

1 If only I had come up with this sooner

Three is less than four because the program we associate with three is shorter. The 'shortness' function is 3's superpower. With the operation -1, initializing on three we would recover almost a whole negative chain: $2, 1, 0, -1, \cdots$. To make the likelihood net zero, we give the whole expansion the expectation: $< 2, 1, 0, -1, \cdots >_3 = 0$

And so also 'expect' to have most instances as positive numbers, given the long possible tail in the negative numbers. So, to keep on the same page, the negative numbers are rare and stretching and the positive numbers are common.

But are the positive numbers the same?

Imagine storing each entry uniquely:

$(, \cdots, 2, 2, 2, 2, 2, 2, 2, 2, 2, \cdots)$

Treated since we begin our recording receiving and stretching information. And since I kept thinking about 2 eventually the rigid precision must be relaxed, so the held-in-mem object for the system is continued with floating point.

But of course, it isn't just 2, since 1 ALSO EXISTS.

Then, the resulting string, for example, $1, 2, 2, 2, 1, 2, 2, 1, 1, 1, 1, 1, 2, 1, 1, 1, 2, , 1, -8, 2, 1, 1, 2$

Provides some information about the statistics and so treats 2 like an ensemble, measuring any floating point against the developing distribution. Saving compute-in-mem/

-8 is only significant if it finds pairs. Nearby pairs may increase the likelihood of spontaneous recombination.

1.1 Talking about the thing

Let us assume that a prime is a more-ordered object, so has a higher cost-to-compute. Then, easily, we would seek to order all other numbers. Kolmogorov complexity seems to work well here, so in a (2,3) language I can express eight by 2: 2^3, 6 the same, two symbols: $2 * 3$, and maybe even 9. This degeneracy is just a property of the redundancy of the language.

Notably for this toy language if we give a commutative construction an additive property, then even 5 is two symbols (and an associated operation).

It seems a negative addition came along for the ride, so with an even and an odd number we can tile the integers. To make the picture interesting, what if we make $2, 2$ unlikely, $3, 3$ likely, and $2, 2, 2, 2$ likely.

Immediately, great towers erupt. All the threes fold and 9s multiply and raise and pretty soon we have to imagine up arrow notation. The 2s behave interesting, enjoying lucky 8 in addition, proper 2^{16} by exponentiation, and the base 16.

Since the author's is assumed to be base-10 we would look to see all the numbers composable. Since no rule is given for 3 and 2 it is assumed they may combine to 1 and the curious 5 as well as 6. 8 is demonstrated with lucky addition, 9 quite easy. The ghosts, 4 and 7 might remain increasingly undiscovered.

Since base 10 gives $0 \equiv 10 \ mod \ 10$ we complete the undergrowth and note that elements two and three give us base 10 with murky four and seven (author noted) under repulsive recombination rules.

Completing the physical analogy, we would create the 'negative' tail as describing expansion in $deSitter$, let the expectation 0 describe the influence of mass, and provoke $AdS-CFT$ to share all of the boring 2 and 1 as a wrapping-within probability with a floating point imprecision.

Conjectured by Max Weinhold, 13/12/26