

Given a **1-indexed** array of integers `numbers` that is already ***sorted in non-decreasing order***, find two numbers such that they add up to a specific `target` number. Let these two numbers be `numbers[index1]` and `numbers[index2]` where `1 <= index1 < index2 <= numbers.length`.

Return *the indices of the two numbers*, `index1` and `index2`, ***added by one*** as an integer array `[index1, index2]` of length 2.

The tests are generated such that there is **exactly one solution**. You **may not** use the same element twice.

Example 1:

Input: `numbers = [2,7,11,15]`, `target = 9`

Output: `[1,2]`

Explanation: The sum of 2 and 7 is 9. Therefore, `index1 = 1`, `index2 = 2`. We return `[1, 2]`.

Example 2:

Input: `numbers = [2,3,4]`, `target = 6`

Output: `[1,3]`

Explanation: The sum of 2 and 4 is 6. Therefore `index1 = 1`, `index2 = 3`. We return `[1, 3]`.

Example 3:

Input: `numbers = [-1,0]`, `target = -1`

Output: `[1,2]`

Explanation: The sum of -1 and 0 is -1. Therefore `index1 = 1`, `index2 = 2`. We return `[1, 2]`.

Constraints:

- `2 <= numbers.length <= 3 * 104`
- `-1000 <= numbers[i] <= 1000`
- `numbers` is sorted in **non-decreasing order**.
- `-1000 <= target <= 1000`
- The tests are generated such that there is **exactly one solution**.