

Consistency and asymptotic normality of Maximum Likelihood Estimator (independent but not necessarily identically distributed)

Max Leung

August 26, 2024

Newey and McFadden (1994) only includes theorems for MLE with i.i.d. assumption. I try to extend them with independent but not necessarily identically distributed assumption. The same asymptotic distribution is derived by Hoadley (1971), but not from Extremum estimator approach.

1 Consistency

Theorem 1 (Consistency of Extremum Estimator with compact parameter space)

Let $Q_0(\boldsymbol{\theta})$ be the limiting objective function and $\hat{\boldsymbol{\theta}} := \operatorname{argmax}_{\boldsymbol{\theta} \in \Theta} \hat{Q}_n(\boldsymbol{\theta})$ i.e., the maximizer of the sample objective function. If (i) $\boldsymbol{\theta}_0$ is the unique maximizer of $Q_0(\boldsymbol{\theta})$. (ii) $\boldsymbol{\theta}_0 \in \Theta$ which is compact. (iii) $Q_0(\boldsymbol{\theta})$ is a continuous function of $\boldsymbol{\theta}$. (iv) $\sup_{\boldsymbol{\theta} \in \Theta} |\hat{Q}_n(\boldsymbol{\theta}) - Q_0(\boldsymbol{\theta})| \rightarrow_p 0$. Then, $\hat{\boldsymbol{\theta}} \rightarrow \boldsymbol{\theta}_0$.

Proof: See Theorem 2.1 of Newey and McFadden (1994).

Q.E.D.

Lemma 2 (Information Inequality; Identification Conditions for MLE)

Let $f_i(y|\mathbf{x}; \boldsymbol{\theta})$ be the conditional density and $\hat{Q}_n(\boldsymbol{\theta}) = n^{-1} \sum_{i=1}^n \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})$. If (i) $f_i(y|\mathbf{x}; \boldsymbol{\theta}) \neq f_i(y|\mathbf{x}; \boldsymbol{\theta}_0)$ for $\forall \boldsymbol{\theta} \in \Theta$ and $\boldsymbol{\theta} \neq \boldsymbol{\theta}_0$. (ii) $\mathbb{E}[|\ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})|] < \infty$ for $\forall i$. (iii) $n^{-1} \sum_{i=1}^n \mathbb{E}[\ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})] := n^{-1} \sum_{i=1}^n Q_{0,i}(\boldsymbol{\theta}) \rightarrow Q_0(\boldsymbol{\theta})$. Then, $\boldsymbol{\theta}_0$ is the unique maximizer of $Q_{0,i}(\boldsymbol{\theta})$ and thus $Q_0(\boldsymbol{\theta})$.

Proof: Replace f with f_i in the Lemma 2.2 of Newey and McFadden (1994).

Q.E.D.

Theorem 3 (Uniform LLN for independent but not necessarily identically distributed case)

If (i) $y_i|\mathbf{x}_i$ is independent but not necessarily identically distributed. (ii) parameter space Θ is compact. (iii) $a_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})$ is a continuous function of $\boldsymbol{\theta}$ for all $y_i|\mathbf{x}_i$ with probability one. (iv) For at least one $\delta > 0$, there exists $d_i(y_i|\mathbf{x}_i)$ such that $|a_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})|^{1+\delta} \leq d_i(y_i|\mathbf{x}_i)$ for $\forall \boldsymbol{\theta} \in \Theta$ and $\mathbb{E}[d_i(y_i|\mathbf{x}_i)] < \infty$. Then, $\sup_{\boldsymbol{\theta} \in \Theta} |n^{-1} \sum_{i=1}^n a_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}) - n^{-1} \sum_{i=1}^n \mathbb{E}[a_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})]| \rightarrow_p 0$ and $n^{-1} \sum_{i=1}^n \mathbb{E}[a_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})]$ is a continuous function of $\boldsymbol{\theta}$.

Reference: Theorem 4.2.2 of Amemiya (1985).

Theorem 4 (Consistency of MLE with compact parameter space)

If (i) $y_i|\mathbf{x}_i$ is independent but not necessarily identically distributed with conditional density $f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)$. (ii) $f_i(y|\mathbf{x}; \boldsymbol{\theta}) \neq f_i(y|\mathbf{x}; \boldsymbol{\theta}_0)$ for $\forall \boldsymbol{\theta} \in \Theta$ and $\boldsymbol{\theta} \neq \boldsymbol{\theta}_0$. (iii) $\boldsymbol{\theta}_0 \in \Theta$ which is compact. (iv) $f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)$ is a continuous function of $\boldsymbol{\theta}$ for all

$y_i|\mathbf{x}_i$ with probability one. (v) $\mathbb{E}[\sup_{\boldsymbol{\theta} \in \Theta} |\ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)|^{1+\delta}] < \infty$ for at least one $\delta > 0$. (vi) $n^{-1} \sum_{i=1}^n \mathbb{E}[\ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})] := n^{-1} \sum_{i=1}^n Q_{0,i}(\boldsymbol{\theta}) \rightarrow Q_0(\boldsymbol{\theta})$. Then, $\hat{\boldsymbol{\theta}} \rightarrow_p \boldsymbol{\theta}_0$.

Proof: (ii), (v) and (vi) implies Theorem 1's (i) by using Lemma 2 because (v) implies Lemma 2's (ii). (iii) is Theorem 1's (ii). Set $a_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}) = \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})$ and $d_i(y_i|\mathbf{x}_i) = \sup_{\boldsymbol{\theta} \in \Theta} |a_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})|^{1+\delta}$ in Theorem 3. (i), (iii), (iv), and (v) is or implies Theorem 3's (i), (ii), (iii), and (iv) respectively. Thus, Theorem 3 implies Theorem 1's (iv) and (iii) since $n^{-1} \sum_{i=1}^n \mathbb{E}[\ln f(y_i|\mathbf{x}_i; \boldsymbol{\theta})] \rightarrow Q_0(\boldsymbol{\theta})$ by (vi). Q.E.D.

2 Asymptotic Normality

Theorem 5 (Asymptotic Normality of Extremum Estimator)

Let $\hat{\boldsymbol{\theta}} := \operatorname{argmax}_{\boldsymbol{\theta} \in \Theta} \hat{Q}_n(\boldsymbol{\theta}) \rightarrow_p \boldsymbol{\theta}_0$. (i) $\boldsymbol{\theta}_0$ is not at the boundary of Θ . (ii) $\nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \hat{Q}_n(\boldsymbol{\theta})$ exists and continuous in N where $\boldsymbol{\theta}_0 \in N$. (iii) $\sqrt{n} \nabla_{\boldsymbol{\theta}} \hat{Q}_n(\boldsymbol{\theta}_0) \rightarrow_d N(\mathbf{0}, \boldsymbol{\Sigma})$. (iv) there exists $\mathbf{H}(\boldsymbol{\theta})$ such that it is continuous at $\boldsymbol{\theta}_0$ and $\sup_{\boldsymbol{\theta} \in N} \|\nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \hat{Q}_n(\boldsymbol{\theta}) - \mathbf{H}(\boldsymbol{\theta})\| \rightarrow_p 0$. (v) $\mathbf{H}(\boldsymbol{\theta}_0)^{-1}$ exists. Then, $\sqrt{n}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \rightarrow_d N(\mathbf{0}, \mathbf{H}^{-1} \boldsymbol{\Sigma} \mathbf{H}^{-1})$.

Proof: See Theorem 3.1 of Newey and McFadden (1994). Q.E.D.

Theorem 6 (Liapounov Central Limit Theorem)

Let y_i be independent. If $\lim_{n \rightarrow \infty} [\sum_{i=1}^n \operatorname{Var}(y_i)]^{-1/2} (\sum_{i=1}^n \mathbb{E}[|y_i - \mathbb{E}[y_i]|^3])^{1/3} = 0$. Then, $\sqrt{n}(n^{-1} \sum_{i=1}^n y_i - n^{-1} \sum_{i=1}^n \mathbb{E}[y_i]) \rightarrow_d N(0, n^{-1} \sum_{i=1}^n \operatorname{Var}(y_i))$.

Reference: Theorem 3.3.5 of Amemiya (1985).

Theorem 7 (Asymptotic Normality of MLE with compact parameter space)

If (i) $y_i|\mathbf{x}_i$ is independent but not necessarily identically distributed with conditional density $f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)$ and other assumptions in Theorem 4 holds. (ii) $\boldsymbol{\theta}_0$ is not at the boundary of Θ . (iii) $f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}) > 0$ is twice continuously differentiable in N neighborhood of $\boldsymbol{\theta}_0$. (iv) Sufficient conditions for interchanging integration and differentiation. (v) $\mathbf{J}_i = \mathbb{E}[\nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0) \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)']$ is invertible. (vi) $\mathbb{E}[\sup_{\boldsymbol{\theta} \in N} \|\nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})\|^{1+\delta}] < \infty$ for at least one $\delta > 0$. (vii) $\lim_{n \rightarrow \infty} [\sum_{i=1}^n \mathbf{J}_i]^{-1/2} (\sum_{i=1}^n \mathbb{E}[\|\nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)\|^3])^{1/3} = 0$. Then, $\sqrt{n}(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}_0) \rightarrow_d N(\mathbf{0}, (n^{-1} \sum_{i=1}^n \mathbf{J}_i)^{-1})$.

Proof: (ii) is Theorem 5's (i). $\nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \hat{Q}_n(\boldsymbol{\theta}) = \nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} n^{-1} \sum_{i=1}^n \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}) = n^{-1} \sum_{i=1}^n \nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})$ and thus (iii) implies Theorem 5's (ii). $\nabla_{\boldsymbol{\theta}} \hat{Q}_n(\boldsymbol{\theta}_0) = \nabla_{\boldsymbol{\theta}} n^{-1} \sum_{i=1}^n \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0) = n^{-1} \sum_{i=1}^n \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)$. $\mathbb{E}[\nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)] = \nabla_{\boldsymbol{\theta}} \mathbb{E}[\ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)] = \nabla_{\boldsymbol{\theta}} Q_{0,i}(\boldsymbol{\theta}_0)$ by (iv). $\nabla_{\boldsymbol{\theta}} Q_{0,i}(\boldsymbol{\theta}_0) = 0$ because Theorem 4's (ii) and (v) implies $\boldsymbol{\theta}_0$ is the unique maximizer of $\mathbb{E}[\ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})] = Q_{0,i}(\boldsymbol{\theta})$. Thus, $\operatorname{Var}(\nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)) = \mathbf{J}_i$ by (v). With (i) and (vii), we apply Liapounov Central Limit Theorem, and have $\sqrt{n} \nabla_{\boldsymbol{\theta}} \hat{Q}_n(\boldsymbol{\theta}_0) = \sqrt{n}(n^{-1} \sum_{i=1}^n \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0) - 0) \rightarrow_d N(\mathbf{0}, n^{-1} \sum_{i=1}^n \mathbf{J}_i)$. Set $a_i(\cdot) = \nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})$ and $d_i(\cdot) = \sup_{\boldsymbol{\theta} \in N} \|\nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})\|^{1+\delta}$ in Theorem 3. Theorem 4's (i), (iii), and this theorem's (iii) and (vi) implies $\sup_{\boldsymbol{\theta} \in N} \|n^{-1} \sum_{i=1}^n \nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}) - n^{-1} \sum_{i=1}^n \mathbb{E}[\nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})]\| \rightarrow_p 0$ and $n^{-1} \sum_{i=1}^n \mathbf{H}_i(\boldsymbol{\theta}) := n^{-1} \sum_{i=1}^n \mathbb{E}[\nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})] = \nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} n^{-1} \sum_{i=1}^n \mathbb{E}[\ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta})] = \nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} n^{-1} \sum_{i=1}^n Q_{0,i}(\boldsymbol{\theta}) \rightarrow \nabla_{\boldsymbol{\theta}} \nabla_{\boldsymbol{\theta}} Q_0(\boldsymbol{\theta}) := \mathbf{H}(\boldsymbol{\theta})$ is a continuous function of $\boldsymbol{\theta}$ and thus $\mathbf{H}(\boldsymbol{\theta})$ is continuous at $\boldsymbol{\theta}_0$. (iv) implies Information Matrix equality i.e., $\mathbf{J}_i = -\mathbf{H}_i$. So, $\mathbf{J} := n^{-1} \sum_{i=1}^n \mathbf{J}_i = -n^{-1} \sum_{i=1}^n \mathbf{H}_i := -\mathbf{H}$. Therefore, (v) implies Theorem 5's (v). Finally, $\mathbf{H}^{-1} \mathbf{J} \mathbf{H}^{-1} = \mathbf{J}^{-1} \mathbf{J} \mathbf{J}^{-1} = \mathbf{J}^{-1} = (n^{-1} \sum_{i=1}^n \mathbf{J}_i)^{-1}$. Q.E.D.

Remark: It is easy to check that the conditions of Theorem 7 implies the conditions of Theorem 2 in Hoadley (1971) for the asymptotic normality. For instance, N5 in Hoadley (1971) is $\mathbb{E}[\nabla_{\boldsymbol{\theta}} \ln f_i(y_i|\mathbf{x}_i; \boldsymbol{\theta}_0)] = 0$ in our proof. N6 is Information Matrix equality, which is implied by (iv) of Theorem 7. N7 is $-n^{-1} \sum_{i=1}^n \mathbf{H}_i(\boldsymbol{\theta}) \rightarrow -\mathbf{H}(\boldsymbol{\theta})$ in our proof. N8 is Liapounov condition, which is Theorem 7's (vii). N9 is dominance condition, which is (vi) of Theorem 7. The asymptotic distribution is exactly the same. However, the conditions in Theorem 7 (I believe) is more primitive i.e., easier to interpret.

3 Application

Binary outcome model with $Pr(y_i = 1|\mathbf{x}_i) = F_i(\mathbf{x}_i'\boldsymbol{\beta}_0)$

$y_i|\mathbf{x}_i$ is independent and Bernoulli distributed with parameter $p_i = Pr(y_i = 1|\mathbf{x}_i) = F_i(\mathbf{x}_i'\boldsymbol{\beta}_0)$. Thus, it is not identically distributed since different i can have different $F_i(\cdot)$ and thus p_i . By specifying conditions which implies the conditions in Theorem 4 and Theorem 7, $\widehat{\boldsymbol{\beta}}$ is consistent and asymptotic normal. In practice, $F_i(\cdot)$ is usually chosen to be Logistic function or standard normal distribution function for $\forall i$. This model relaxes this assumption.

4 Reference

Amemiya, T. (1985). *Advanced Econometrics*, Harvard University Press.

Hoadley, B. (1971). Asymptotic Properties of Maximum Likelihood Estimators for the Independent Not Identically Distributed Case. *The Annals of Mathematical Statistics*, 42(6), 1977–1991.

Newey, W., & McFadden, D. (1986). Large sample estimation and hypothesis testing. *Handbook of Econometrics*, 4, 2111-2245.