

Manuscript

Commented [1]: <https://www.nature.com/natecoleval/about/content>

Larger but younger fish when growth compensates for higher mortality in heated ecosystem

Commented [2]: maximum title length is 100 characters (including spaces)

Deleted: warmed

Max Lindmark^{a,b,1}, Malin Karlsson^a, Anna Gårdmark^c

^a Swedish University of Agricultural Sciences, Department of Aquatic Resources, Institute of Coastal Research, Skolgatan 6, 742 42 Öregrund, Sweden

^b Swedish University of Agricultural Sciences, Department of Aquatic Resources, Institute of Marine Research, Turistgatan 5, 453 30 Lysekil, Sweden

^c Swedish University of Agricultural Sciences, Department of Aquatic Resources, Skolgatan 6, SE-742 42 Öregrund, Sweden

¹ Author to whom correspondence should be addressed. Current address:

Max Lindmark, Swedish University of Agricultural Sciences, Department of Aquatic Resources, Institute of Marine Research, Turistgatan 5, 453 30 Lysekil, Sweden, Tel.: +46(0)104784137, email: max.lindmark@slu.se

Keywords: body growth, size-structure, size-spectrum, mortality, climate change

Abstract

Ectotherms are often predicted to “shrink” with global warming, in line with general growth models and the temperature-size rule (TSR), which predict smaller adult sizes with warming. However, they also predict faster juvenile growth rates, leading to larger size-at-age. Hence, the result of warming on population size-structure also depends on mortality rates and how much adult size changes. We used data from an artificially heated (+8C) bay in comparison with an unheated area, to analyze how warming has affected body growth, mortality rates and population size-structure of Eurasian perch (*Perca fluviatilis*). In the heated bay, body size was larger for all ages and growth faster for all sizes, resulting in larger size-spectrum exponent (greater proportion of large fish) – despite higher mortality. Hence, to understand warming-induced changes in the size-structure of species, which affect ecological interactions and dynamics, it is critical to account for changes in both mortality and growth rate.

B.

Commented [3]: 150 words max (151 nu)

Commented [AG4R3]: nu 150

Deleted: due to smaller adult body sizes, in line with the temperature-size rule (TSR). However, tStill, he TSR also predicts faster juvenile growth rate

Deleted: s.

Commented [5]: Jag har lite svårt att hänga med här. Storleksfördelningen vid högre temperaturer beror väl både på tillväxt (och när i livet den är högre/längre än vid normala temperaturer) och på mortalitet?

Commented [MOU6R5]: Mhm, se om det passar bättre nu kanske!

Deleted: The net effect of warming on population, however, depends also on mortality

Deleted: warm

Commented [MOU7]: Eftersom vi inte använder en regressionsbaserad metod så kallar vi detta numera exponent, eftersom det är den vi skattar direkt!

Deleted: slope

Deleted: Combining

Deleted: temperature effects on growth with those on mortality is necessary for understanding warming-induced changes in size-distributions

Commented [AG8]: this is a general view point, but not something that this study demonstrates (as we don't study how these differences in population size structure affect stability). Better here to have a conclusion that links more to your actual results, and move this type of statements to the end of the discussion instead

Se kommentarer på slut-stycket i diskussionen. Om du köper det upplägget (som jag kan skriva ett utkast på, om du vill), så passar det bättre med en liknande slutsats i abstract.

Commented [MOU9R8]: Var inte heller superförtjust i de två sista meningarna. Vad tror du om detta? En enda slutsats och implications som är mer i linje med vad vi faktiskt visar? Tänker att det ffa är data som är tyngden i studien.

Deleted: ecause ecological dynamics emerge from size-dependent processes and functions, addressing how warming affects size-distributions through multiple processes is essential for predicting the effects of climate change.

Introduction

Ectotherm species, constituting 99% of species globally (Atkinson & Sibly, 1997; Wilson, 1992), are commonly predicted to shrink in a warming world (Cheung et al., 2013; Gardner et al., 2011; Sheridan & Bickford, 2011). Mean body size responses to temperature may however be uninformative, as the size-distribution of many species spans several orders of magnitude. Warming can even shift size-distributions without altering mean size; for example, if increases in juvenile size-at-age outweigh the decline in size-at-age in adults, which is consistent with the temperature size-rule, TSR (Atkinson, 1994). Resolving how warming induces changes in population size-distributions may thus be more instructive (Fritschie & Olden, 2016), especially for inferring warming effects on species' ecological role, biomass production, or energy fluxes (Yvon-Durocher et al., 2011). This is because key traits such as metabolism, feeding, growth, mortality scale with body size (Andersen, 2020; Blanchard et al., 2017; Brown et al., 2004; Pauly, 1980; Thorson et al., 2017; Ursin, 1973). Hence, as the value of these traits at mean body size is not the same as the mean population trait value (Bernhardt et al., 2018), the size-distribution within a population thus matters for its dynamics under warming.

The population size distribution can be represented as a size-spectrum, which generally is the frequency distribution of individual body sizes (A. M. Edwards et al., 2017). It is often described in terms of the spectrum slope (slope of individuals or biomass of a size class over the mean size of that class on log-log scale (A. M. Edwards et al., 2017; Sheldon et al., 1973; White et al., 2007)) or simply the exponent of the power law individual size-distribution (A. M. Edwards et al., 2017). The size-spectrum implicitly captures temperature-dependent ecological processes such as body growth, mortality and recruitment (Blanchard et al., 2017; Heneghan et al., 2019). Despite its rich theoretical foundation (Andersen, 2019) and usefulness as an ecological indicator (Blanchard et al., 2005), few studies have evaluated warming-effects on the species size-spectrum in larger bodied species (but see Blanchard et al. (2005)). There

Commented [10]: The main text (excluding introductory paragraph, Methods section, references and figure legends) is limited to 3,500 words

Commented [11]: An introduction (without heading) of up to 500 words of referenced text expands on the background of the work (some overlap with the summary is acceptable)...

Commented [12R11]: Vi är uppe i 616 nu

Commented [AG13R11]: nu nere i 580 ☺ (och det är ju med utskrivna referenser; blir ju färre sen när du byter till Nature-stil på referenserna)

Commented [MOU14R11]: 523 nu med nature reffar. Det kanske duger?

Deleted: Global warming is commonly predicted to cause e

Deleted: –

Commented [AG15]: undetected implies that there is a change in the mean but that it is hidden. But here you're after that there may be no change in the mean, aren't you? (because changes in one end outweighs those in the other end of the size spectrum)

Deleted: growth

Deleted: growth by

Deleted: as

Commented [AG16]: "scaling with size allometrically" is tautological

Deleted: . T

Deleted: diversity

Deleted: is

Deleted: the number of individuals in different body-size classes related to the mean size of each class.

Deleted: The

Deleted: size-

Deleted: (γ)

Deleted:

Deleted: is the often-linear relationship between these variables on a

Deleted: Hence, t

Deleted: slope

Commented [17]: Probably because they tend to focus on fishing effects on the size spectrum

are numerous [paths](#) by which the species size-spectrum could change with warming (Heneghan et al., 2019). For instance, the clearest consequence of TSR for the size-spectrum is a decline in maximum size. However, unless warming also alters the relative abundances of juveniles and adults, the same number of adults will accumulate in a smaller size-range, resulting in a less steep [slope](#) of the size-spectrum. Warming can also lead to elevated mortality, which truncates the age-distribution towards younger individuals (Barnett et al., 2017). Increased growth rates can however counter the effects of mortality on abundance-at-size, unless only small individuals benefit from warming (Daufresne et al., 2009; Lindmark et al., n.d.). Growth rates can increase due to physiological responses to higher temperatures, or reduced density-dependence following warming-induced mortality, or both. Hence, the effect of warming on the size-spectrum depends on several interlinked processes affecting abundance-at-size and size-at-age. Yet, while warming effects on ectotherm body growth has been thoroughly studied (Atkinson, 1994; Brett et al., 1969; Lindmark et al., n.d.), those on mortality in wild populations have not, nor their joint consequences for population size-spectra in warming environments.

We used data from a unique, large-scale 23-year-long heating-experiment of a coastal ecosystem to quantify how warming changed fish body growth, mortality, and the size structure in an unexploited population of Eurasian perch (*Perca fluviatilis*, 'perch'). We compare fish from this enclosed bay exposed to temperatures 5-10 °C above normal ('[heated area](#)') with fish from a reference area ('cold area') in the adjacent archipelago (Fig. 1). Using hierarchical Bayesian models, we quantify differences in key individual- and population level parameters, such as body growth, asymptotic size, mortality rates, and size-spectra between the [heated](#) and [reference](#) coastal area.

Results

Commented [MOU19R18]: yes paths better!

Deleted: ways

Deleted: slope

Commented [AG20]: from this paragraph it isn't really clear what the main knowledge gap is. worth reiterating here to make it clear.

Commented [AG21]: completely unknown to your readers; need proper introduction

Commented [AG22]: I would go for 'heated area' instead of 'warm', as this signals that it is caused and artificial rather than a natural difference warm-cold

Commented [MOU23R22]: ok!

Deleted: warm

Deleted: warm

Commented [AG24]: see previous comment; suggest you go for 'heated' and 'reference', which clarifies that it is an experimental heating; but since you started with warm cold I continued it here

Commented [MOU25R24]: will do that

Deleted: cold

Commented [26]: ... and is followed by a concise, focused account of the findings (headed 'Results')...

Commented [27R26]: Det star visserligen "concise", men vi är ganska långt ifrån word-limit (och intron är begränsad till 500 ord, och diskussionen till 2 paragrafer, så det är bara här vi kan öka på. Har även kollat i andra nature eco evo articular och jag tycker att många lägger in lite tolkningar och friare meningar här.

Analysis of fish (perch) size-at-age using the von Bertalanffy growth equation (VBGE) revealed that fish cohorts (year classes) in the heated area both grew faster initially (larger size-at-age and VBGE K parameter) and reached larger predicted asymptotic sizes than those in the reference area (Fig. 2). The model with area-specific VBGE parameters (L_{∞} , K and t_0) had the largest expected log pointwise predictive density for a new observation (Table S1), and there is a clear difference in both the estimated values for fish asymptotic length (L_{∞}) and growth rate (K) between the heated and reference area (Fig. 2B-E). For instance, the distribution of differences between the heated and reference area of the posterior samples for L_{∞} and K only had 11% and 2%, respectively, of the density below 0, illustrating that it is unlikely that the parameters are larger in the reference area or similar in the two areas (Fig. 2C, E). We estimated that fish in the heated area had an asymptotic length that was 16% greater than fish in the reference area ($L_{\infty\text{heat}} = 45.7[36.8, 56.3]$, $L_{\infty\text{ref}} = 39.4[35.4, 43.9]$, given as posterior median with 95% credible interval) and had 27% larger growth coefficients, i.e. K parameters ($K_{\text{heat}} = 0.19[0.15, 0.23]$, $K_{\text{ref}} = 0.15[0.12, 0.17]$). Corresponding estimates of the third parameter in the VBGE were $t_{0\text{heat}} = -0.16[-0.21, -0.11]$, and $t_{0\text{ref}} = -0.44[-0.56, -0.33]$.

In addition, we found that the initial growth rate (at small size) was higher in the heated area and that the decline in growth with length was steeper in the reference area (Fig. 3). The best model for growth ($G = \alpha L^{\theta}$) had area-specific parameters (Table S2), and we found even stronger support for differences in growth parameters between the areas (Fig. 3). Initial growth was estimated to be 18% faster in the heated than in the reference area ($\alpha_{\text{heat}} = 509.7[460.1, 563.5]$, $\alpha_{\text{ref}} = 433.5[413.3, 454.1]$), and growth of fish in the heated area had a 4% lower decline with length than in the reference area ($\theta_{\text{heat}} = -1.13[-1.16, -1.11]$, $\theta_{\text{ref}} = -1.18[-1.19, -1.16]$). The distribution of differences of the posterior samples for α

Commented [AG28]: do readers understand this word do we think?

Commented [MOU29R28]: this ok?

Commented [AG30]: needs to be translated for the general reader into something clarifying what this means

Commented [MOU31R30]: erm, will be difficult.

Commented [MOU32]: Vi borde inte kalla detta growth rate eftersom den bara har dimensionen tid

Deleted: grew with

Deleted: faster

Deleted: growth rate

Deleted: warm

Deleted: a

Commented [AG33]: behöver vi så här fin noggrannhet, räcker det inte med en decimal? (eller inga?)

Commented [MOU34R33]: Räcker gott med en

and θ both only had 0.3% of the density below 0 (Fig. 3C, E), indicating high probability that growth rates have differentiated.

By analyzing the decline in catch-per-unit-effort over age, we found that the instantaneous mortality rate Z (rate at which log abundance declines with age) is higher in the heated area (Fig. 4). The overlap with zero is 0.1% for the distribution of differences of posterior samples of Z_{heat} and Z_{ref} (Fig. 4C). We estimated Z_{heat} to be $0.76[0.63,0.88]$ and Z_{ref} to be $0.63[0.58,0.69]$, which correspond to annual mortality rates of 53% in the heated area and 47% in the reference area.

Lastly, analysis of the size-structure in the two areas revealed that the faster growth rates for fish of all sizes, which lead to a larger size-at-age, outweigh the higher mortality rates in the heated area, such that the size-spectrum exponent is larger there (Fig. 5A). However, in contrast to the results on size-at-age, growth and mortality, the differentiation is not as strong statistically, as the 95% confidence intervals overlap. That said, because of this faster growth, the heated area both has the largest individuals and an overall larger proportion of large individuals (Fig. 5C). ↓

Discussion

Our study provides strong evidence for warming-induced differentiation in growth, mortality, and size-structure in a natural population of a non-exploited, temperate fish species exposed to 5-10 °C above normal temperatures for more than two decades. While it is a study on only a single species, these features make it a unique climate change experiment, as most warming studies on natural populations of fish to date are on commercially exploited fish species (and fisheries exploitation affects size-structure both directly and indirectly by selecting for fast growing individuals). While factors other than temperature could have contributed to the elevated growth and mortality, the temperature contrast is unusually large for natural systems

Deleted: that

Commented [MOU35]: Föredrar nog att skriva såhär, att sannolikheten är hög för olika tillväxt än låg för lika tillväxt

Deleted: identical growth in the warm and the cold area is highly unlikely

Deleted: population

Deleted: s

Deleted:

Deleted: ing

Deleted: slope

Deleted: less steep in the heated area

Deleted: As a consequence of

Deleted: have

Deleted: as well as

Deleted: 5D

Deleted: This is evident in the analyses of size-spectrum slopes over time, where we found that the intercept (α) is larger in the heated area (Fig. 5B; where year is centered so such that the intercept corresponds to the first year, i.e., 1987, and the model without an interaction between year and area was favored). The overlap with zero was 2% for the distribution of differences of posterior samples of the size-spectrum intercepts of the heated and the reference area, α_{warm} and α_{cold} (Fig. 5C).

Commented [38]: Hade gärna fått in hur tillväxttöken som TSR innebär ofta överskuggas pga focus på shrinking.

Samt att TSR-bevis är rätt ovanliga på större organismer i naturliga miljöer (mycket mer studier och teori på mindre organismer)

Commented [39]: Två paragrafer discussion lämnar ju mycket osagt... Just nu väldigt lite om mortalitet!

Commented [40]: ... and one or two short paragraphs of discussion (headed 'Discussion').

(i.e., 5-10 °C, which can be compared to the 1.35 °C change in the Baltic Sea between 1982 and 2006 (Belkin, 2009)). Moreover, heating occurred at the scale of a whole ecosystem, which makes the findings highly relevant in the context of global warming. Interestingly, our findings contrast with both broader predictions about declining mean body sizes based on the GOLT hypothesis (Cheung et al., 2013; Pauly, 2021), and with intraspecific patterns such as the TSR (temperature-size rule; Atkinson (1994)). The contrasts lie in that both asymptotic size and size-at-age of mature individuals, as well as the proportion of larger individuals were larger and higher in the heated area – despite the elevated mortality rates. Since optimum growth temperatures decline with size within species generally under food satiation in experimental studies (Lindmark et al., n.d.), the finding that the asymptotic size and the largest individuals were found in the heated area was unexpected. This suggests that growth dynamics under food satiation may not be directly proportional to those under natural feeding conditions (Railsback, 2022). Our results suggest that growth (and mortality) changes emerge not only from direct physiological responses to increased temperatures, but also from warming-induced changes in the food web, e.g., prey productivity, diet composition and trophic transfer efficiencies (Gårdmark & Huss, 2020).

A key question for understanding the implications of warming on ectotherm populations is if larger individuals in a population become rarer or smaller (Ohlberger 2013; Ohlberger et al. 2018) – i.e., how the size- and age-distribution change rather than the mean size. Not only for evaluating population-level changes in functions and ecological roles due to the allometric scaling of vital rates (Fritschie & Olden 2016; Audzijonyte et al. 2020), but also because age-truncated populations tend to have less stable dynamics (Anderson et al. 2008). Our study contributes to the literature revealing large variations across species in terms of the warming effects on life history traits and demographic parameters. A key challenge is to account for this variation in projections on the impacts of climate change on natural populations.

Commented [AG42]: refs needed here on other studies in natural systems

Commented [MOU43R42]: Funkar detta?

Commented [MOU41]: gillar dessa edits!

Deleted: (

Deleted:)

Deleted: and, more specifically, on

Deleted: responses

Deleted:

Deleted: ref

Deleted: (Lindmark et al. n.d.)

Deleted: even

Commented [AG44]: we need to say something of what could have caused this; it is likely that heating at the ecosystem scale results in different responses than pure direct effects of warming on individual organisms, in that warming could increase both physiological rates and food productivity.

Commented [MOU45R44]: tycker dina edits blev bra! förkortade sista meningen bara

Deleted: inhabited

Deleted: that organisms respond differently to warming of whole ecosystems than to direct warming of individuals, and

Commented [AG46]: det här stycket är så kort så här finns utrymme att expandera diskussionen (när vi nu är så långt från word limit)

Tänker att vi borde diskutera uppvärmningseffekter på mortalitet, t ex utifrån
Rodgers et al 18:
<https://link.springer.com/content/pdf/10.1007/s00227-018-3333-z.pdf>
Biro et al 07:
<https://www.pnas.org/content/pnas/104/23/9715.full.pdf>
Pershing et al 15:
<https://www.science.org/doi/pdf/10.1126/science.aac9819>

Innan jag ger mig på att skriva ett helt annat slutstycke än det du har nu kanske vi skall diskutera det, så att vi är överens om vad det skall innehålla?
(Jag tänker mig att man börjar med mortalitetseffekter av uppvärmning, och vad de skulle kunna bero på, för att sedan peka på vikten av att studera mortalitet & tillväxt samtidigt för att se sluteffekten på storleksfördelningar (vilket är viktigt för ...etc).)

Commented [MOU47R46]: Jättebra, du får gärna göra det! Och bra reffar också, har inte läst någon av dem faktiskt... Men om du skriver om, kan vi behålla första meningen? Blev så nöjd med den!

Commented [AG48]: refs needed here

Commented [AG49]: this refers to between-species variation in your previous sentence – isn't it the within-species variation that you should highlight here, as it is the focus of your study

Materials and Methods

Data

We use [size-at-age](#) data from perch sampled annually from the heated enclosed bay ('the Biotest lake') and its reference area with natural temperatures in the years after the onset of warming (first cohort is 1981, [and first](#) and last catch year is [1987 and 2003, respectively](#)) to omit transient dynamics and acute responses, and to ensure we use cohorts that only experience one of the thermal environments during its life. A grid at the outlet of the heated area (Fig. 1) prevented fish larger than 10 cm from migrating between the areas (Adill et al., 2013; Huss et al., 2019), and genetic studies confirm the reproductive isolation between the two populations during the time-period (Björklund et al., 2015). However, since the grid removal in 2004, fish growing up in the heated Biotest lake can easily swim out, meaning we cannot be sure fish in the reference area did not recently arrive from the Biotest lake. Hence, we use data up until [2003](#). This resulted in 12658 length-at-age measurements from 2426 individuals in 256 nets.

We use data from fishing events using survey-gillnets that took place in October in the heated Biotest lake and in August in the reference area when temperatures are most comparable between the two areas (Huss et al., 2019), because temperature affect catchability in static gears. The catch was recorded by 2.5 cm length classes during 1987-2000, and into 1 cm length groups between 2001-2003. To express lengths in a common length standard, 1 cm intervals were converted into 2.5 cm intervals. The unit of catch data is hence the number of fish caught per 2.5 cm size class per night per net (i.e., a catch-per-unit-effort [CPUE] variable). All data from fishing events with disturbance affecting the catch (e.g., seal damage, strong algal growth on the gears, clogging by drifting algae) were removed (years 1996 and 1999 from the heated area in the catch data).

Commented [50]: Asta's paper has 2500 words

Deleted: trait

Commented [51]: 2002 for growth data because of the potential change in age estimation

Age and length-at-age was reconstructed for a semi-random length-stratified subset of individuals each year. This was done using annuli rings on the operculum bones (with control counts done on otoliths), and an established power-law relationship between the distance of annual rings and fish length: $L = \kappa R^s$, where L is the length of the fish, R the operculum radius, κ the intercept, and s the slope of the line for the regression of log-fish length on log-operculum radius from a large reference data set for perch (Thoresson, 1996). Back-calculated length-at-age were obtained from the relationship $L_a = L_s \left(\frac{r_a}{R}\right)^s$, where L_a is the back-calculated body length at age a , L_s is the final body length (body length at catch), r_a is the distance from the center to the annual ring corresponding to age a and $s = 0.861$ for perch (Thoresson, 1996). Since perch exhibits sexual size-dimorphism, and age-determination together with back calculation of growth was not done for males in all years, we only used females for our analyses.

Statistical Analysis

The differences in size-at-age, growth, mortality, and size structure between the heated and the reference area were quantified using hierarchical linear and non-linear regression models fitted in a Bayesian framework. First, we describe each statistical model and then provide details of model fitting, model diagnostics and comparison.

We fit the von Bertalanffy growth equation (VBGE) on a *log* scale, describing length as a function of age to evaluate differences in size-at-age and asymptotic size: $\log(L_t) = \log(L_\infty(1 - e^{(-K(t-t_0))}))$, where L_t is the length at age (t , years), L_∞ is the asymptotic size, K is the Brody coefficient (yr^{-1}) and t_0 is the age when the average length was zero.. We used only age- and size-at-catch as the response variables (i.e., not back-calculated length-at-age). This was to have a simpler model and not have to account for parameters varying within individuals as well as cohorts, as mean sample size per individual was only ~ 5 . We let

Commented [AG52]: this was only started after 2000, if I recall Martina's instructions correctly.

Commented [MOU53R52]: ok! do we need that level of detail though? it's not wrong as it is now! Would you be able to ask about it, if you want to be more specific here, or should I do it?

Deleted: θ

Commented [MOU55]: kan vi kalla exponent för s istället för theta, som vi redan använder i tillväxtmodellen?

Deleted: θ

Commented [AG56]: this is only true for the CPUE analyses, right? not back-calculated growth and the growth models? or? if it is true for all our data sets, then we just state this at the very beginning of the methods (where it currently says 1981)

Deleted: .

To focus on the long-term differentiation in mortality between the heated and reference area we avoided the initial years of warming (when mortality rates increased only to decline again (Sandström *et al.* 1995)) and used data from 1987-2003...

Commented [MOU57R56]: Good point, we kind of already say that actually

Commented [AG58]: already said earlier

parameters vary among cohorts rather than year of catch, because individuals within cohorts share similar environmental conditions and density dependence (Morrongiello & Thresher, 2015). Eight models in total were fitted (with area being dummy-coded), with different combinations of shared and area-specific parameters. We evaluated if models with area-specific parameters led to better fit and quantified the differences in area-specific parameters. The model with all area-specific parameter can be written as:

$$L_i \sim \text{Student-}t(v, \mu_i, \sigma) \quad (1)$$

$$\log(\mu_i) = A_R \log \left[L_{\infty Rj[t]} \left(1 - e^{(-K_{Rj[t]}(t - t_{0Rj[t]}))} \right) \right] + A_H \log \left[L_{\infty Hj[t]} \left(1 - e^{(-K_{Hj[t]}(t - t_{0Hj[t]}))} \right) \right] \quad (2)$$

$$\begin{bmatrix} L_{\infty Rj} \\ L_{\infty Hj} \\ K_{Rj} \\ K_{Hj} \end{bmatrix} \sim \text{MVNormal} \left(\begin{bmatrix} \mu_{L_{\infty R}} \\ \mu_{L_{\infty H}} \\ \mu_{K_R} \\ \mu_{K_H} \end{bmatrix}, \begin{bmatrix} \sigma_{L_{\infty R}} & 0 & 0 & 0 \\ 0 & \sigma_{L_{\infty H}} & 0 & 0 \\ 0 & 0 & \sigma_{K_R} & 0 \\ 0 & 0 & 0 & \sigma_{K_H} \end{bmatrix} \right) \quad (3)$$

where lengths are *Student* – *t* distributed to account for extreme observations, v , μ and ϕ represent the degrees of freedom, mean and the scale parameter, respectively. Henceforth, subscripts H and R are used for the heated and reference area, respectively (except in figures and main text where subscripts are spelled out for clarity). A_R and A_H are dummy variables such that $A_R = 1$ and $A_H = 0$ if it is the reference area, and vice. The multivariate normal distribution in Eq. 3 is the prior for the cohort-varying parameters $L_{\infty Rj}$, $L_{\infty Hj}$, K_{Rj} and K_{Hj} (for cohorts $j = 1981, \dots, 1997$) (note that cohorts extend further back in time than the catch data), with hyper-parameters $\mu_{L_{\infty R}}$, $\mu_{L_{\infty H}}$, μ_{K_R} , μ_{K_H} describing the non-varying population means and a covariance matrix with the between-cohort variation along the diagonal (note we did not model a correlation between the parameters, hence off-diagonals are 0). The other seven models include some or all parameters as parameters common for the two areas, e.g., substituting $L_{\infty Rj}$ and $L_{\infty Hj}$ with $L_{\infty j}$. To aid convergence of this non-linear model, we used informative priors chosen after visualizing draws from prior predictive distributions (Wesner

Commented [AG59]: I suggest you spell it out everywhere – conveys the message better

Commented [MOU60R59]: i think the subscripts become to long then, not good for readability either!

& Pomeranz, 2021) using probable parameter values (*Supporting Information*, Fig. S1, S6). We used the same prior distribution for each parameter class for both areas to not introduce any other sources of differences in parameter estimates between areas. We used the following priors for the VBGE model: $\mu_{L_{\infty R,H}} \sim N(45, 20)$, $\mu_{K_{R,H}} \sim N(0.2, 0.1)$, $t_{0 R,H} \sim N(-0.5, 1)$ and $v \sim \text{gamma}(2, 0.1)$. σ , $\mu_{L_{\infty R}}$, $\mu_{L_{\infty H}}$, μ_{K_R} , μ_{K_H} were given a *Student - t*(3, 0, 2.5) prior.

We also compared how growth scales with size (in contrast to length vs age) in the two areas, by fitting allometric growth models describing how specific growth rate scales with length: $G = \alpha L^\theta$, where G , the specific growth, is defined as: $G = 100 \times (\log(L_2) - \log(L_1)) / (t_2 - t_1)$ and L is the geometric mean length: $L = (L_1 \times L_2)^{0.5}$. Here we also used back-calculated length-at-age, resulting in multiple observations for each individual. As with the VBGE model, we dummy coded area to compare models with different combinations of common and shared parameters. We assumed growth rates were *Student - t* distributed, and the full model can be written as:

$$\begin{aligned} L_i &\sim \text{Student} - t(v, \mu_i, \sigma) & (4) \\ \mu_i &= A_R(\alpha_{Rj[i],k[i]} L^{\theta_R}) + A_H(\alpha_{Hj[i],k[i]} L^{\theta_H}) & (5) \\ \alpha_{R,Hj} &\sim N(\mu_{\alpha_{R,Hj}}, \sigma_{\alpha_{R,Hj}}) & (6) \\ \alpha_{R,Hj} &\sim N(\mu_{\alpha_{R,Hj}}, \sigma_{\alpha_{R,Hj}}) & (7) \\ \theta_{R,Hj} &\sim N(\mu_{\theta_{R,Hj}}, \sigma_{\theta_{R,Hj}}) & (8) \\ \theta_{R,Hj} &\sim N(\mu_{\theta_{R,Hj}}, \sigma_{\theta_{R,Hj}}) & (9) \end{aligned}$$

We assumed only α varied across individuals j within cohorts k , and compared two models: one with θ common for the heated and reference area, and one with an area-specific θ . We used the following priors, after visual exploration of the prior predictive distribution (*Supporting Information*, Fig. S7, S9): $\alpha_{R,H} \sim N(500, 100)$, $\theta_{R,H} \sim N(-1.2, 0.3)$ and $v \sim \text{gamma}(2, 0.1)$. σ , $\sigma_{id:cohort}$ and σ_{cohort} were all given a *Student - t*(3, 0, 13.3) prior.

We estimated total mortality by fitting linear models to the natural log of catch (CPUE) as a function of age (catch curve regression), under the assumption that in a closed population,

Commented [AG61]: varför kursiv? ser ut som en variabel

Commented [MOU62R61]: minns ej

the exponential decline can be described as $N_t = N_0 e^{-Zt}$, where N_t is the population at time t , N_0 is the initial population size and Z is the instantaneous mortality rate. This equation can be rewritten as a linear equation: $\log(C_t) = \log(vN_0) - Zt$, where C_t is catch-at-age t , if catch is assumed proportional to the number of fish (i.e., $C_t = vN_t$). Hence, the negative of the slope of the regression is Z . To get catch-at-age data, we constructed area-specific age-length keys using the sub-sample of the total catch that was age-determined. Age length-keys describe the age-proportions of each length-category (i.e., a matrix with length category as rows, ages as columns). Age is then estimated for the total catch based on the “probability” of fish in each length-category being a certain age. With fit this model with and without an *age-area*-interaction, and the former can be written as:

$$\log(CPUE_i) \sim \text{Student-}t(v, \mu_i, \sigma) \quad (10)$$

$$\mu_i = \beta_{0j[i]}(\text{area}_R) + \beta_{1j[i]}(\text{area}_H) + \beta_{2j[i]} \text{age} + \beta_{3j[i]}(\text{age} \times \text{area}_H) \quad (11)$$

$$\begin{bmatrix} \beta_{0j} \\ \beta_{1j} \\ \beta_{2j} \\ \beta_{3j} \end{bmatrix} \sim \text{MVNormal} \left(\begin{bmatrix} \mu_{\beta_0} \\ \mu_{\beta_1} \\ \mu_{\beta_2} \\ \mu_{\beta_3} \end{bmatrix}, \begin{bmatrix} \sigma_{\beta_0} & 0 & 0 & 0 \\ 0 & \sigma_{\beta_1} & 0 & 0 \\ 0 & 0 & \sigma_{\beta_2} & 0 \\ 0 & 0 & 0 & \sigma_{\beta_3} \end{bmatrix} \right) \quad (12)$$

where β_{0j} and β_{1j} are the intercepts for the reference and heated areas, respectively, β_{2j} is the age slope for the reference area and β_{3j} is the interaction between *age* and *area*. All parameters vary by cohort (for cohort $j = 1981, \dots, 2000$) and their correlation is set to 0 (Eq. 12). We use the following (vague) priors: $\mu_{\beta_0, \dots, \beta_3} \sim N(0, 10)$ (where μ_{β_2} is the population-level estimate for $-Z_R$ and $\mu_{\beta_2} + \mu_{\beta_3}$ is the population-level estimate for $-Z_H$) and $v \sim \text{gamma}(2, 0.1)$. σ and $\sigma_{\beta_0, \dots, \beta_3}$ were given a *Student - t*(3, 0, 2.5) prior.

Lastly, we quantified differences in the size-distributions between the areas using size-spectrum [exponents](#). We estimate [biomass the size-spectrum exponent \$\gamma\$ directly](#), using the likelihood approach for binned data, i.e., the *MLEbin* method in the R package *sizeSpectra* (A. Edwards, 2020; A. M. Edwards et al., 2017, 2020). This method explicitly accounts for

Commented [MOU63]: The reason I don't use α_W , α_C , is because the interaction term does not correspond to the warm slope directly

Deleted: slopes

uncertainty in body masses *within* size-classes (bins) in the data and has been shown to be less biased than regression-based methods or the likelihood method based on bin-midpoints (A. M. Edwards et al., 2017, 2020). [We pooled all years to ensure negative relationships between biomass and size in the size-classes.](#)

All analyses were done using R (R Core Team, 2020)_version 4.0.2 [with R Studio \(2021.09.1\)](#). The packages within the *tidyverse* (Wickham et al., 2019) collection were used to processes and visualize data. Models were fit using the R package *brms* (Bürkner, 2017). When priors were not chosen based on the prior predictive distributions, we used the default priors from *brms* as written above. We used 3 chains and 4000 iterations in total per chain. Models were compared by evaluating their expected predictive accuracy (expected log pointwise predictive density) using leave-one-out cross-validation (LOO-CV) (Vehtari et al., 2017) while ensuring pareto k values < 0.7 , in the R package *loo* (Vehtari et al., 2020). Results of the model comparison can be found in the *Supporting Information*, Table S1-S4. We used *bayesplot* (Gabry et al., 2019) and *tidybayes* (Kay, 2019) to process and visualize model diagnostics and posteriors. Model convergence and fit was assessed by ensuring potential scale reduction factors (\hat{R}) were less than 1.1, suggesting all three chains converged to a common distribution (Gelman et al., 2003), and by visually inspecting trace plots, residuals QQ-plots and with posterior predictive checks (*Supporting Information*, Fig. S2, S8, S10, S12).

Acknowledgements

We thank all staff involved in data collection.

Code and Data Availability

All **data** and R code can be downloaded from a GitHub repository (https://github.com/maxlindmark/warm_life_history) and will be archived on Zenodo upon publication.

Commented [64]: Här menar jag städad data som går rakt in i analysen

Author Contributions

ML conceived the idea and designed the study and the statistical analysis. Data-processing, initial statistical analyses, and initial writing was done by MK and ML. AG contributed critically to all mentioned parts of the paper. All authors contributed to the manuscript writing and gave final approval for publication.

Commented [65]: Låter detta OK? Det som är lite klurigt är väl att beskriva hur det bygger på ditt jobb Malin utan att det ska låta som att de slutgiltiga skripten och analyserna skrevs inom exjobbet, samt hur vi ska få till din input Anna som sträcker sig över allt, dock utan att ha en ledande roll på någon av delarna jag nämnt.

Commented [66]: Här menar jag då uppsatsen samt första draften av ms'et som ändå är ganska olika varandra.

Commented [67]: Jag tycker att det låter bra!

References

- Adill, A., Mo, K., Sevastik, A., Olsson, J., & Bergström, L. (2013). *Biologisk recipientkontroll vid Forsmarks kärnkraftverk* (Rapport 2013:19; Issue 2013:19). <https://pub.epsilon.slu.se/11349/>
- Andersen, K. H. (2019). *Fish Ecology, Evolution, and Exploitation: A New Theoretical Synthesis*. Princeton University Press.
- Andersen, K. H. (2020). Size-based theory for fisheries advice. *ICES Journal of Marine Science*, 77(7–8), 2445–2455. <https://doi.org/10.1093/icesjms/fsaa157>
- Atkinson, D. (1994). Temperature and organism size—A biological law for ectotherms? In *Advances in Ecological Research* (Vol. 25, pp. 1–58). Elsevier. [https://doi.org/10.1016/S0065-2504\(08\)60212-3](https://doi.org/10.1016/S0065-2504(08)60212-3)
- Atkinson, D., & Sibly, R. M. (1997). Why are organisms usually bigger in colder environments? Making sense of a life history puzzle. *Trends in Ecology & Evolution*, 12(6), 235–239.
- Barnett, L. A. K., Branch, T. A., Ranasinghe, R. A., & Essington, T. E. (2017). Old-Growth Fishes Become Scarce under Fishing. *Current Biology*, 27(18), 2843–2848.e2. <https://doi.org/10.1016/j.cub.2017.07.069>
- Belkin, I. M. (2009). Rapid warming of large marine ecosystems. *Progress in Oceanography*, 81(1), 207–213.
- Bernhardt, J. R., Sunday, J. M., Thompson, P. L., & O'Connor, M. I. (2018). Nonlinear averaging of thermal experience predicts population growth rates in a thermally variable environment. *Proceedings of the Royal Society B: Biological Sciences*, 285(1886), 20181076. <https://doi.org/10.1098/rspb.2018.1076>
- Björklund, M., Aho, T., & Behrmann-Godel, J. (2015). Isolation over 35 years in a heated biotest basin causes selection on MHC class IIβ genes in the European perch (*Perca fluviatilis* L.). *Ecology and Evolution*, 5(7), 1440–1455. <https://doi.org/10.1002/ece3.1426>

- Blanchard, J. L., Dulvy, N., Jennings, S., Ellis, J., Pinnegar, J., Tidd, A., & Kell, L. (2005). Do climate and fishing influence size-based indicators of Celtic Sea fish community structure? *ICES Journal of Marine Science*, 62(3), 405–411. <https://doi.org/10.1016/j.icesjms.2005.01.006>
- Blanchard, J. L., Heneghan, R. F., Everett, J. D., Trebilco, R., & Richardson, A. J. (2017). From bacteria to whales: Using functional size spectra to model marine ecosystems. *Trends in Ecology & Evolution*, 32(3), 174–186. <https://doi.org/10.1016/j.tree.2016.12.003>
- Brett, J. R., Shelbourn, J. E., & Shoop, C. T. (1969). Growth rate and body composition of fingerling sockeye salmon, *Oncorhynchus nerka*, in relation to temperature and ration size. *Journal of the Fisheries Research Board of Canada*, 26(9), 2363–2394. <https://doi.org/10.1139/f69-230>
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a metabolic theory of ecology. *Ecology*, 85(7), 1771–1789. <https://doi.org/10.1890/03-9000>
- Bürkner, P.-C. (2017). **brms**: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, 80(1). <https://doi.org/10.18637/jss.v080.i01>
- Cheung, W. W. L., Sarmiento, J. L., Dunne, J., Frölicher, T. L., Lam, V. W. Y., Deng Palomares, M. L., Watson, R., & Pauly, D. (2013). Shrinking of fishes exacerbates impacts of global ocean changes on marine ecosystems. *Nature Climate Change*, 3(3), 254–258. <https://doi.org/10.1038/nclimate1691>
- Daufresne, M., Lengfellner, K., & Sommer, U. (2009). Global warming benefits the small in aquatic ecosystems. *Proceedings of the National Academy of Sciences, USA*, 106(31), 12788–12793. <https://doi.org/10.1073/pnas.0902080106>
- Edwards, A. (2020). *sizeSpectra: Fitting Size Spectra to Ecological Data Using Maximum Likelihood* (R package version 1.0.0.0) [Computer software]. <https://github.com/andrew-edwards/sizeSpectra>
- Edwards, A. M., Robinson, J. P. W., Blanchard, J. L., Baum, J. K., & Plank, M. J. (2020). Accounting for the bin structure of data removes bias when fitting size spectra. *Marine Ecology Progress Series*, 636, 19–33. <https://doi.org/10.3354/meps13230>
- Edwards, A. M., Robinson, J. P. W., Plank, M. J., Baum, J. K., & Blanchard, J. L. (2017). Testing and recommending methods for fitting size spectra to data. *Methods in Ecology and Evolution*, 8(1), 57–67. <https://doi.org/10.1111/2041-210X.12641>
- Fritschie, K. J., & Olden, J. D. (2016). Disentangling the influences of mean body size and size structure on ecosystem functioning: An example of nutrient recycling by a non-native crayfish. *Ecology and Evolution*, 6(1), 159–169. <https://doi.org/10.1002/ece3.1852>
- Gabry, J., Simpson, D., Vehtari, A., Betancourt, M., & Gelman, A. (2019). Visualization in Bayesian workflow. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 182(2), 389–402. <https://doi.org/10.1111/rssa.12378>
- Gårdmark, A., & Huss, M. (2020). Individual variation and interactions explain food web responses to global warming. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 375(1814), 20190449. <https://doi.org/10.1098/rstb.2019.0449>
- Gardner, J. L., Peters, A., Kearney, M. R., Joseph, L., & Heinsohn, R. (2011). Declining body size: A third universal response to warming? *Trends in Ecology & Evolution*, 26(6), 285–291. <https://doi.org/10.1016/j.tree.2011.03.005>
- Gelman, A., Carlin, J., Stern, H., & Rubin, D. (2003). *Bayesian Data Analysis. 2nd edition*. Chapman and Hall/CRC.

- Heneghan, R. F., Hatton, I. A., & Galbraith, E. D. (2019). Climate change impacts on marine ecosystems through the lens of the size spectrum. *Emerging Topics in Life Sciences*, 3(2), 233–243. <https://doi.org/10.1042/ETLS20190042>
- Huss, M., Lindmark, M., Jacobson, P., van Dorst, R. M., & Gårdmark, A. (2019). Experimental evidence of gradual size-dependent shifts in body size and growth of fish in response to warming. *Global Change Biology*, 25(7), 2285–2295. <https://doi.org/10.1111/gcb.14637>
- Kay, M. (2019). *tidybayes: Tidy Data and Geoms for Bayesian Models* (R package version 1.1.0) [Computer software]. <http://doi.org/10.5281/zenodo.1308151>
- Lindmark, M., Ohlberger, J., & Gårdmark, A. (n.d.). Optimum growth temperature declines with body size within fish species. *Global Change Biology*, n/a(n/a). <https://doi.org/10.1111/gcb.16067>
- Morrongiello, J. R., & Thresher, R. E. (2015). A statistical framework to explore ontogenetic growth variation among individuals and populations: A marine fish example. *Ecological Monographs*, 85(1), 93–115. <https://doi.org/10.1890/13-2355.1>
- Pauly, D. (1980). On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *ICES Journal of Marine Science*, 39(2), 175–192. <https://doi.org/10.1093/icesjms/39.2.175>
- Pauly, D. (2021). The gill-oxygen limitation theory (GOLT) and its critics. *Science Advances*, 7(2), eabc6050. <https://doi.org/10.1126/sciadv.abc6050>
- R Core Team. (2020). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Railsback, S. F. (2022). What We Don't Know About the Effects of Temperature on Salmonid Growth. *Transactions of the American Fisheries Society*, 151(1), 3–12. <https://doi.org/10.1002/tafs.10338>
- Sheldon, R. W., Sutcliffe, W. H., & Prakash, A. (1973). The Production of Particles in the Surface Waters of the Ocean with Particular Reference to the Sargasso Sea. *Limnology and Oceanography*, 18(5), 719–733. <https://doi.org/10.4319/lo.1973.18.5.0719>
- Sheridan, J. A., & Bickford, D. (2011). Shrinking body size as an ecological response to climate change. *Nature Climate Change*, 1(8), 401–406. <https://doi.org/10.1038/nclimate1259>
- Thoreson, G. (1996). *Metoder för övervakning av kustfiskbestånd* (No. 3; Kustrapport). Kustlaboratoriet, Fiskeriverket. <http://urn.kb.se/resolve?urn=urn:nbn:se:havochvatten:diva-317>
- Thorson, J. T., Munch, S. B., Cope, J. M., & Gao, J. (2017). Predicting life history parameters for all fishes worldwide. *Ecological Applications*, 27(8), 2262–2276. <https://doi.org/10.1002/eap.1606>
- Ursin, E. (1973). On the prey size preferences of cod and dab. *Meddelelser Fra Danmarks Fiskeri-Og Havun-Dersgelser*, 7:8598.
- Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., Bürkner, P., Paananen, T., & Gelman, A. (2020). *loo: Efficient leave-one-out cross-validation and WAIC for Bayesian models*. (R package version 2.3.1) [Computer software]. <https://mc-stan.org/loo>
- Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. *Statistics and Computing*, 27(5), 1413–1432. <https://doi.org/10.1007/s11222-016-9696-4>
- Wesner, J. S., & Pomeranz, J. P. F. (2021). Choosing priors in Bayesian ecological models by simulating from the prior predictive distribution. *Ecosphere*, 12(9), e03739. <https://doi.org/10.1002/ecs2.3739>

- White, E. P., Ernest, S. K. M., Kerkhoff, A. J., & Enquist, B. J. (2007). Relationships between body size and abundance in ecology. *Trends in Ecology & Evolution*, 22(6), 323–330.
- Wickham, H., Averick, M., Bryan, J., Chang, W., D'Agostino McGowan, L., François, R., Golemund, G., & Alex, H. (2019). *Welcome to the tidyverse*. 1686. <https://doi.org/10.21105/joss.01686>
- Wilson, E. O. (1992). *The Diversity of Life*. Harvard University Press.
- Yvon-Durocher, G., Montoya, J. M., Trimmer, M., & Woodward, G. (2011). Warming alters the size spectrum and shifts the distribution of biomass in freshwater ecosystems. *Global Change Biology*, 17(4), 1681–1694. <https://doi.org/10.1111/j.1365-2486.2010.02321.x>

Figures




Fig. 1. Map of the area with the unique ecosystem warming experiment from which perch in this study was sampled. Inset shows the 1 km² enclosed coastal bay that has been artificially heated for 23 years, the adjacent reference area with natural temperatures, and locations of the cooling water intake and where the heated water outlet from nuclear power plants enters the heated coastal area.

Commented [AG68]: need to use the same wordings as in the main text 'warm area' vs 'cold area' (or, if you go for my suggestion: 'heated area' vs 'reference area'); it's irrelevant that it is called the biotest lake

should we perhaps make this figure more informative, by combining it with a photo of a gill lid, showing the growth rings?

arrow for the warm water outlet needs to be moved to the actual outlet (inside the biotest! in the 'fork')

föreslår också att du gör texten 'Sweden' aningens mindre, så den inte ligger emot några andra linjer

Commented [MOU69R68]: har du en bra gällöcksbild vi kan använda?

Deleted: the

Commented [AG70]: if you stick with 'warm and cold area' those terms should appear here in the text as well (suggest you change to heated and reference area throughout the manuscript)




Fig. 2. The average length-at-age is larger for fish of all ages in the heated (red) compared to the reference (blue) coastal area. Points in panel (A) depicts individual-level length-at-age and lines show the global posterior prediction (both exponentiated) without group-level effects (i.e., cohort) from the von Bertalanffy growth model with area-specific coefficients. The shaded areas correspond to 50% and 90% credible intervals. Panel (B) shows the posterior distributions for growth coefficient (parameters K_{heat} (red) and K_{ref} (blue)) and (C) the distribution of their difference. Panel (D) shows the posterior distributions for asymptotic length (parameters $L_{\infty heat}$ and $L_{\infty ref}$), and (E) the distribution of their difference.

Deleted: rate




Fig. 3. The faster growth rates in the heated area (red) compared to the reference (blue) are maintained as fish grow. The points illustrate specific growth estimated from back-calculated length-at-age (within individuals) as a function of length (expressed as the geometric mean of the length at the start and end of the time interval). Lines show the global posterior prediction without group-level effects (i.e., individual within cohort) from the allometric growth model with area-specific coefficients. The shaded areas correspond to the 90% credible interval. The equation uses mean parameter estimates. Panel (B) shows the posterior distributions for initial growth (α_{heat} (red) and α_{ref} (blue)), and (C) the distribution of their difference. Panel (D) shows the posterior distributions for the allometric decline in growth with length (θ_{heat} and θ_{ref}), and (E) the distribution of their difference.

Commented [AG71]: in contrast to fig 4-5, in figs 2-3 you don't have a color legend for the posterior distributions; is there a reason for why you don't do the same in all figs 2-5?




Fig. 4. The instantaneous mortality rate (Z) is higher in the heated area (red) than in the reference (blue). Panel (A) shows the $\log(\text{CPUE})$ as a function of age , where the slope corresponds to the global $-Z$. Lines show the posterior prediction without group-level effects (i.e., cohort) and the shaded areas correspond to the 50% and 90% credible intervals. The equation uses mean parameter estimates. Panel (B) shows the posterior distributions for mortality rate (Z_{heat} and Z_{ref}), and (C) the distribution of their difference.




Fig. 5. The heated area (red) has a larger proportion of large fish than the reference area (blue), illustrated both in terms of the biomass size-spectrum (A-B), and histograms of proportions (C). Panel (A) shows the size distribution and MLEbins fit (red and blue solid curve for the heated and reference area, respectively) with 95% confidence intervals indicated by dashed lines. The vertical span of rectangles illustrates the possible range of the number of individuals with body mass \geq the body mass of individuals in that bin. Panel (B) shows the estimate of the size-spectrum exponent, γ , and vertical lines depict the 95% confidence interval. Panel (C) illustrates histograms of length groups in the heated and reference area as proportions (for all years pooled).

Deleted: as

Deleted: B

Deleted:

Deleted: and

Deleted: A

Deleted: the generally smaller size-spectrum slopes (which also increased over time in both areas)

Deleted: shows mean annual size-spectrum slopes as points and their 95% confidence interval, estimated using maximum likelihood. Lines and shaded areas correspond to the median and 90% credible interval. The equation uses mean parameter estimates.

Deleted: posterior

Deleted: distributions for the reference and heated intercepts (i.e., β_0 and β_1 in the model shown in panel (A) of size-spectrum slopes as a function of year), and (C) the distribution of their difference.

Deleted: D