

Hypoxic and Hypercapnic Respiratory Failure

Respiratory failure is a life-threatening condition characterized by the inability to maintain adequate oxygenation (hypoxic) or ventilation (hypercapnic). This document provides a comprehensive overview for advanced practice providers (APPs) in a hospital setting.

Definition and Classification

- **Definition:** Respiratory failure occurs when the respiratory system fails to maintain adequate gas exchange, resulting in hypoxemia ($\text{PaO}_2 < 60$ mmHg on room air) and/or hypercapnia ($\text{PaCO}_2 > 50$ mmHg) with acidosis ($\text{pH} < 7.35$).
- **Classification:**
 - Type 1: Hypoxic Respiratory Failure: $\text{PaO}_2 < 60$ mmHg with normal or low PaCO_2 . Caused by impaired oxygenation (e.g., V/Q mismatch, shunt, diffusion limitation).
 - Type 2: Hypercapnic Respiratory Failure: $\text{PaCO}_2 > 50$ mmHg with $\text{pH} < 7.35$, often with hypoxemia. Caused by inadequate ventilation (e.g., hypoventilation, increased dead space).
 - Type 3: Perioperative Respiratory Failure: Hypoxemia due to atelectasis or hypoventilation in the perioperative setting (e.g., post-anesthesia).
 - Type 4: Shock-Related Respiratory Failure: Hypoxemia in shock states (e.g., cardiogenic, septic) due to impaired oxygen delivery and utilization.

Physiology and V/Q Ratios

- **Ventilation-Perfusion (V/Q) Ratio:**
 - V/Q ratio = alveolar ventilation (V) / pulmonary blood flow (Q). Normal V/Q ≈ 0.8 (ventilation 4 L/min, perfusion 5 L/min).
 - V/Q Mismatch: Imbalance between ventilation and perfusion leads to hypoxemia or hypercapnia.
 - Low V/Q (Shunt-Like): Ventilation $<$ perfusion (e.g., pneumonia, ARDS). Causes hypoxemia; supplemental O₂ may improve PaO_2 but not fully correct it.
 - High V/Q (Dead Space-Like): Ventilation $>$ perfusion (e.g., pulmonary embolism). Causes hypercapnia; supplemental O₂ does not correct PaCO_2 .

- Shunt: $V/Q = 0$ (no ventilation, perfusion present; e.g., atelectasis, intrapulmonary shunt in ARDS). Severe hypoxemia, minimally responsive to O_2 .
- Dead Space: $V/Q = \infty$ (ventilation, no perfusion; e.g., PE). Increases $PaCO_2$ due to wasted ventilation.
- **Hypoxic (Type 1) Physiology:**
- Mechanisms:
 - V/Q Mismatch: Most common (e.g., pneumonia, pulmonary edema). Low V/Q regions → Hypoxemia.
 - Shunt: Blood bypasses ventilated alveoli (e.g., ARDS, atelectasis). Severe hypoxemia, poor response to O_2 .
 - Diffusion Limitation: Impaired O_2 transfer across alveolar-capillary membrane (e.g., ILD, pulmonary fibrosis). Hypoxemia worsens with exercise.
 - Low FiO_2 : Reduced inspired O_2 (e.g., high altitude, smoke inhalation).
 - Hypoventilation: Contributes to hypoxemia if severe (e.g., opioid overdose), though primarily causes hypercapnia.
 - A-a Gradient: Elevated in V/Q mismatch, shunt, and diffusion limitation (normal <15 mmHg on room air, age-adjusted). Normal in hypoventilation or low FiO_2 .
 - $PaCO_2$: Normal or low (compensatory hyperventilation to improve oxygenation).
- **Hypercapnic (Type 2) Physiology:**
- Mechanisms:
 - Hypoventilation: Decreased minute ventilation (e.g., COPD exacerbation, opioid overdose, neuromuscular disease). $\uparrow PaCO_2$, $\downarrow PaO_2$.
 - Increased Dead Space: Wasted ventilation (e.g., COPD with bullae, PE). $\uparrow PaCO_2$ due to ineffective CO_2 elimination.
 - Increased CO_2 Production: Rarely the primary cause (e.g., sepsis, fever, hyperthyroidism), but can exacerbate hypercapnia in hypoventilation states.
 - V/Q Mismatch: High V/Q regions (e.g., COPD with emphysema) contribute to hypercapnia by increasing dead space.
 - A-a Gradient: Normal in pure hypoventilation; elevated if V/Q mismatch or shunt present.
 - PaO_2 : Often decreased due to hypoventilation (alveolar gas equation: $PaO_2 = FiO_2 \times (P_{atm} - P_{H_2O}) - (PaCO_2/RQ)$; $\uparrow PaCO_2 \rightarrow \downarrow PaO_2$).

Categories of Hypoxic Respiratory Failure

Category	Common Causes	V/Q Ratio	Notes
V/Q Mismatch	Pneumonia, pulmonary edema (CHF), asthma, atelectasis, early ARDS	Low V/Q (<0.8)	Most common cause of hypoxemia; responds partially to O ₂ .
Shunt	ARDS, atelectasis, pulmonary AV malformation, intracardiac shunt (e.g., PFO)	V/Q = 0	Severe hypoxemia; minimal response to O ₂ ; common in ARDS.
Diffusion Limitation	Interstitial lung disease (ILD), pulmonary fibrosis, pulmonary edema	Normal or low V/Q	Worsens with exercise; responds to O ₂ .
Low FiO ₂	High altitude, smoke inhalation, fire (CO poisoning)	Normal V/Q	Normal A-a gradient; corrects with O ₂ .
Hypoventilation	Opioid overdose, CNS depression, neuromuscular disease (contributes to hypoxemia)	Normal V/Q	Primary cause of hypercapnia; hypoxemia corrects with O ₂ .
Mixed	ARDS with hypoventilation, COPD with pneumonia	Variable	Combination of mechanisms; treat both components.

Categories of Hypercapnic Respiratory Failure

Category	Common Causes	V/Q Ratio	Notes
Hypoventilation	COPD exacerbation, opioid overdose, obesity hypoventilation syndrome (OHS), neuromuscular disease (e.g., ALS, GBS, myasthenia gravis), CNS depression (e.g., stroke, sedative overdose)	Normal V/Q	Primary mechanism; normal A-a gradient; BiPAP first-line.
Increased Dead Space	COPD (emphysema, bullae), pulmonary embolism (PE), severe ARDS	High V/Q (>0.8)	Wasted ventilation; elevated A-a gradient if shunt present.
Increased CO ₂ Production	Sepsis, fever, hyperthyroidism, burns (rare primary cause)	Normal or high V/Q	Exacerbates hypoventilation; treat underlying cause.
Mixed	COPD with pneumonia, OHS with OSA, ARDS with sedative overdose	Variable	Combination of mechanisms; address all components.

Clinical Presentation

- **Hypoxic (Type 1):**
 - Symptoms: Dyspnea, tachypnea, cyanosis, confusion (severe hypoxemia), fatigue.

- Exam: Hypoxemia ($\text{SpO}_2 < 90\%$ on room air), tachycardia, accessory muscle use, crackles (pulmonary edema), wheezing (asthma), diminished breath sounds (pneumothorax).
- Red Flags: $\text{SpO}_2 < 88\%$ despite O_2 , altered mental status, hemodynamic instability → Consider ARDS, massive PE.
- **Hypercapnic (Type 2):**
 - Symptoms: Dyspnea, somnolence, confusion, headache (CO_2 narcosis), asterixis (CO_2 retention).
 - Exam: Hypoxemia (often present), respiratory acidosis ($\text{pH} < 7.35$), shallow breathing (hypoventilation), wheezing (COPD), weakness (neuromuscular disease).
 - Red Flags: $\text{pH} < 7.25$, lethargy/coma, respiratory arrest → Urgent intervention (e.g., BiPAP, intubation).

Diagnostic Workup

• Initial Tests:

- Arterial Blood Gas (ABG):
 - Hypoxic: $\text{PaO}_2 < 60$ mmHg, normal/low PaCO_2 .
 - Hypercapnic: $\text{PaCO}_2 > 50$ mmHg, $\text{pH} < 7.35$, often with $\text{PaO}_2 < 60$ mmHg.
 - Calculate A-a gradient: $A-a = [\text{FiO}_2 \times (\text{Patm} - \text{PH}_2\text{O}) - (\text{PaCO}_2/0.8)] - \text{PaO}_2$. Elevated (> 15 mmHg) in V/Q mismatch, shunt, diffusion limitation.
- Pulse Oximetry: $\text{SpO}_2 < 90\%$ on room air indicates hypoxemia; unreliable for CO_2 assessment.
- Chest X-Ray (CXR): Identify structural causes (e.g., pneumonia, pneumothorax, pulmonary edema, COPD hyperinflation).
- Labs: CBC (anemia, infection), BMP (electrolytes, renal function), lactate (sepsis, shock), D-dimer (if PE suspected), BNP (heart failure).

• Specific Tests:

- V/Q Scan or CT Pulmonary Angiogram (CTPA): For PE (V/Q mismatch, high V/Q).
- Echocardiogram: For pulmonary edema (LV dysfunction), right heart strain (PE).
- Pulmonary Function Tests (PFTs): For chronic diseases (e.g., COPD, ILD); not feasible in acute setting.
- Toxicology Screen: If hypoventilation suspected (e.g., opioids, benzodiazepines).
- Neuromuscular Assessment: Maximal inspiratory pressure (MIP), vital capacity (VC) for neuromuscular disease (e.g., ALS, myasthenia gravis).

- **Key Tips:**

- ABG first to classify type (hypoxic vs. hypercapnic).
- A-a gradient: Normal in hypoventilation or low FiO₂; elevated in V/Q mismatch, shunt, diffusion limitation.
- CXR in all; CTPA if PE suspected (D-dimer + Wells score).

Diagnostic Flowsheet: Respiratory Failure

Step	Description
Step 1: ABG	PaO ₂ <60 mmHg? PaCO ₂ >50 mmHg? pH <7.35?
Step 2: Classify Type	Hypoxic (Type 1): PaO ₂ <60 mmHg, normal/low PaCO ₂ . Hypercapnic (Type 2): PaCO ₂ >50 mmHg, pH <7.35.
Step 3: A-a Gradient	Calculate: $A-a = [FiO_2 \times (P_{atm} - P_{H_2O}) - (PaCO_2/0.8)] - PaO_2$. Elevated (>15 mmHg): V/Q mismatch, shunt, diffusion limitation. Normal: Hypoventilation, low FiO ₂ .
Step 4: Identify Cause	Hypoxic: Pneumonia, ARDS, PE, pulmonary edema, ILD. Hypercapnic: COPD, opioid overdose, OHS, neuromuscular disease.
Step 5: Confirm with Imaging/ Labs	CXR: Pneumonia, ARDS, pneumothorax, edema. CTPA: PE. Echo: CHF, right heart strain. Tox screen: Opioid overdose.

Treatment

- **General Principles:** Correct hypoxemia and hypercapnia, treat the underlying cause, and prevent progression to respiratory arrest.
- **Supportive Care:**
 - Oxygen Therapy:
 - Hypoxic: Titrate O₂ to SpO₂ 88-92% (or 92-96% if no COPD risk). Use nasal cannula, Venturi mask, or high-flow nasal cannula (HFNC).
 - Hypercapnic: Cautious O₂ titration (SpO₂ 88-92%) to avoid suppressing hypoxic drive in COPD patients.
 - Non-Invasive Ventilation (NIV):
 - BiPAP: First-line for hypercapnic failure (e.g., COPD, OHS). Settings: IPAP 10-20 cmH₂O, EPAP 4-8 cmH₂O, titrate to improve pH and PaCO₂.
 - CPAP: For hypoxic failure with pulmonary edema (e.g., CHF). Settings: 5-10 cmH₂O.

- Intubation/Mechanical Ventilation:
 - Indications: Failure of NIV (pH <7.25, PaCO₂ rising), severe hypoxemia (PaO₂/FiO₂ <150 despite O₂), respiratory arrest, inability to protect airway.
 - Settings: ARDS → Low tidal volume (6 mL/kg IBW), PEEP 5-15 cmH₂O; COPD → Avoid overventilation (target pH >7.25, not PaCO₂ normalization).
 - Positioning: Prone positioning for ARDS (PaO₂/FiO₂ <150); improves V/Q matching.
 - Monitoring: Repeat ABG q2-4h, continuous SpO₂, respiratory rate, mental status.
- **Specific Therapy:**
 - **Pneumonia:** Antibiotics (e.g., ceftriaxone 1 g IV + azithromycin 500 mg IV for CAP), oxygen, consider ICU if septic shock.
 - **ARDS:** Low tidal volume ventilation (6 mL/kg IBW), PEEP titration (ARDSnet protocol), prone positioning, fluid restriction, treat underlying cause (e.g., sepsis).
 - **Pulmonary Edema (CHF):** Diuretics (furosemide 40 mg IV), nitroglycerin (if BP allows), CPAP/BiPAP, inotropes (e.g., dobutamine) if cardiogenic shock.
 - **Pulmonary Embolism (PE):** Anticoagulation (heparin 80 units/kg IV bolus, then 18 units/kg/h infusion), thrombolytics (e.g., alteplase 100 mg IV over 2h) for massive PE, oxygen.
 - **Pneumothorax:** Chest tube (if tension or large), oxygen, analgesia.
 - **Asthma/COPD Exacerbation:**
 - Bronchodilators Albuterol 2.5 mg nebulized q20min x 3, then q1-4h; ipratropium 0.5 mg nebulized q20min
 - Steroids: Prednisone 40-60 mg PO or methylprednisolone 60 mg IV daily x 5 days.
 - BiPAP: For hypercapnic failure in COPD (pH <7.35, PaCO₂ >50 mmHg).
 - Magnesium sulfate: 2 g IV over 20 min (severe asthma).
 - **Opioid Overdose:** Naloxone 0.4-2 mg IV q2-3min (titrate to respiratory effort), supportive care, monitor for recurrence (long-acting opioids).
 - **Neuromuscular Disease** (e.g., Myasthenia Gravis, GBS): BiPAP, plasmapheresis/IVIG (myasthenia), monitor VC/MIP (intubate if VC <15 mL/kg or MIP >-20 cmH₂O).
 - **Obesity Hypoventilation Syndrome (OHS):** BiPAP (IPAP 12-20 cmH₂O, EPAP 6-10 cmH₂O), weight loss counseling, treat OSA if present.
 - **Interstitial Lung Disease (ILD):** Oxygen, steroids (e.g., prednisone 1 mg/kg/day) for acute exacerbation, consider antifibrotics (e.g., nintedanib) for IPF.

- **Key Tips:**

- Avoid over-oxygenation in COPD (risk of CO₂ retention); target SpO₂ 88-92%.
- BiPAP contraindicated in pneumothorax, altered mental status, or inability to protect airway → Intubate.
- Prone positioning in ARDS: 16h/day, improves mortality (PROSEVA trial, NEJM 2013).

Examples

1. Case 1: Hypoxic Respiratory Failure (Pneumonia)

- Presentation: 65 y/o M, fever, cough, SpO₂ 85% on room air, CXR shows right lower lobe consolidation, ABG: PaO₂ 55 mmHg, PaCO₂ 35 mmHg, pH 7.42.
- Interpretation: Type 1 (hypoxic) respiratory failure, V/Q mismatch (low V/Q due to pneumonia), elevated A-a gradient.
- Management: O₂ via nasal cannula (titrate to SpO₂ 92-96%), ceftriaxone 1 g IV + azithromycin 500 mg IV, monitor ABG q4h, consider ICU if no improvement.

2. Case 2: Hypercapnic Respiratory Failure (COPD Exacerbation)

- Presentation: 70 y/o F, known COPD, dyspnea, wheezing, SpO₂ 87%, ABG: PaO₂ 50 mmHg, PaCO₂ 70 mmHg, pH 7.28.
- Interpretation: Type 2 (hypercapnic) respiratory failure, hypoventilation + increased dead space (high V/Q), respiratory acidosis.
- Management: BiPAP (IPAP 15 cmH₂O, EPAP 5 cmH₂O), albuterol 2.5 mg + ipratropium 0.5 mg nebulized, methylprednisolone 60 mg IV, target SpO₂ 88-92%, repeat ABG in 1h.

3. Case 3: Hypoxic Respiratory Failure (ARDS)

- Presentation: 40 y/o M, sepsis from pancreatitis, SpO₂ 80% on 100% FiO₂, CXR bilateral infiltrates, ABG: PaO₂ 60 mmHg, PaCO₂ 40 mmHg, pH 7.38, PaO₂/FiO₂ 60.
- Interpretation: Type 1 (hypoxic) respiratory failure, shunt (V/Q = 0), severe ARDS (PaO₂/FiO₂ <100).
- Management: Intubate (low tidal volume 6 mL/kg IBW, PEEP 10 cmH₂O), prone positioning, treat sepsis (antibiotics, fluids), monitor plateau pressure (<30 cmH₂O).

4. Case 4: Hypercapnic Respiratory Failure (Opioid Overdose)

- Presentation: 30 y/o M, found unresponsive, pinpoint pupils, RR 6/min, SpO₂ 82%, ABG: PaO₂ 45 mmHg, PaCO₂ 65 mmHg, pH 7.20.
- Interpretation: Type 2 (hypercapnic) respiratory failure, hypoventilation, normal A-a gradient.
- Management: Naloxone 0.4 mg IV q2min (titrate to RR >12), BiPAP if slow recovery, monitor for recurrence, supportive care.

5. Case 5: Hypoxic Respiratory Failure (Pulmonary Embolism)

- Presentation: 55 y/o F, sudden dyspnea, chest pain, SpO₂ 88%, D-dimer elevated, CTPA shows saddle PE, ABG: PaO₂ 58 mmHg, PaCO₂ 32 mmHg, pH 7.45.
- Interpretation: Type 1 (hypoxic) respiratory failure, V/Q mismatch (high V/Q, dead space), elevated A-a gradient.
- Management: Heparin 80 units/kg IV bolus, then 18 units/kg/h infusion, oxygen (SpO₂ 92-96%), consider thrombolytics if hemodynamic instability, monitor for bleeding.

Complications

- **Acute:**
 - Hypoxemia: Organ dysfunction (e.g., brain, heart), arrhythmias, death.
 - Hypercapnia: CO₂ narcosis (lethargy, coma), acidosis → Cardiovascular collapse.
 - Barotrauma: From mechanical ventilation (e.g., pneumothorax, pneumomediastinum).
 - Ventilator-Associated Pneumonia (VAP): Risk in intubated patients (>48h on vent).
- **Long-Term:**
 - Chronic Respiratory Failure: In progressive diseases (e.g., COPD, ILD, neuromuscular disease).
 - Pulmonary Hypertension: From chronic hypoxemia (e.g., COPD, OHS).
 - Cognitive Impairment: From prolonged hypoxemia or CO₂ retention.

Prognosis

- **Mortality:**
 - ARDS: 30-40% in-hospital mortality (higher if PaO₂/FiO₂ <100).
 - COPD Exacerbation: 10-20% mortality if requiring NIV; 20-40% if intubated.

- Massive PE: 30-50% mortality if untreated; 10-20% with anticoagulation.
- Opioid Overdose: <5% with naloxone and supportive care; higher if delayed.
- **Recovery:**
 - Pneumonia: 70-90% recover with antibiotics and O2; worse in elderly.
 - Neuromuscular Disease: Variable; depends on disease progression (e.g., ALS poor, GBS better with treatment).
 - OHS: Good with BiPAP and weight loss; high recurrence without lifestyle change.
- **Key Factors:**
 - Early NIV in hypercapnic failure reduces intubation rates (e.g., COPD).
 - Underlying comorbidities (e.g., heart failure, malignancy) worsen outcomes.
 - Rapid correction of underlying cause (e.g., PE, pneumonia) improves survival.

Key Pearls

- **ABG first:** PaO2 <60 mmHg → Hypoxic; PaCO2 >50 mmHg + pH <7.35 → Hypercapnic.
- **A-a gradient:** Elevated in V/Q mismatch, shunt, diffusion limitation; normal in hypoventilation.
- **Hypoxic:** Focus on oxygenation (O2, CPAP, intubation); treat cause (e.g., antibiotics for pneumonia, thrombolytics for PE).
- **Hypercapnic:** Prioritize ventilation (BiPAP, intubate if severe); cautious O2 in COPD (SpO2 88-92%).
- **ARDS:** Low tidal volume ventilation, prone positioning, PEEP titration (ARDSnet protocol).
- **COPD:** BiPAP first, avoid overventilation on vent (target pH >7.25, not PaCO2 normalization).

References

- **UpToDate:** "Acute Respiratory Failure" (2025).
- **ATS Guidelines:** "Management of Acute Exacerbations of COPD" (2023).
- **NEJM:** "Prone Positioning in Severe ARDS (PROSEVA Trial)" (2013).
- **CHEST:** "Non-Invasive Ventilation in Acute Respiratory Failure" (2022).
- **ERS/ATS:** "ARDS Definition and Management" (2024).

Visit: medcheatsheets.com for more education, fun resources and 10 category 1 AAPA CME credit!

© Hospital Medicine Cheat Sheets (medcheatsheets.com). For educational purposes only. Do not redistribute or sell. Neither the author nor the company is liable for realworld implications. AI was used in development