

# Generalized Hospital Medicine

## Pearls for Optimal Inpatient Care

### Definition and Overview

**Prevalence Inpatient complications are common:** delirium affects 20-50% of elderly inpatients, constipation ~30-40%, sleep disturbances ~50%, and delayed mobility increases length of stay (LOS) by 1-2 days. Falls occur in 3-5% of admissions, and 30% of discharges face readmission within 30 days. Risk Factors Advanced age, polypharmacy, comorbidities (e.g., dementia, CHF), prolonged hospitalization, immobility. Rare Demographics Pediatric inpatients (e.g., post-surgical), bariatric patients, culturally diverse groups with unique needs.

### Pathophysiology

- **Mechanisms:**

- **Poor sleep** disrupts circadian rhythms, increasing delirium risk via melatonin dysregulation.
- **Constipation** results from opioid use, dehydration, and immobility, altering gut motility.
- **Delirium** stems from neuroinflammation (IL-6, TNF- $\alpha$ ) and cholinergic deficiency.
- **Immobility** causes muscle atrophy, VTE, and deconditioning.
- **Inadequate discharge** planning leads to medication errors and readmissions.

Effects These issues prolong LOS, increase morbidity (e.g., delirium mortality 10-20%), and reduce quality of life. Falls cause fractures in 1-2% of cases, and malnutrition worsens recovery.

- **Molecular Pathways**

- **Sleep:** Reduced melatonin impairs GABA signaling.
- **Constipation:** Opioid  $\mu$ -receptor activation slows peristalsis.
- **Delirium:** IL-6 upregulates microglial activation.
- **Mobility:** Myostatin inhibits muscle protein synthesis.
- **Nutrition:** Low albumin reflects NLRP3 inflammasome activation.
- **Key Pathway:** Inpatient stressors → Physiological disruption (inflammation, immobility) → Complications → Prolonged recovery.

## Hospital Medicine Pearls

Issue	Strategy	Implementation	Notes
Inpatient Sleep Medicine	Optimize sleep hygiene	Dim lights post-8 PM, minimize noise (earplugs), avoid non-urgent vitals 11 PM-5 AM. Melatonin 3-5 mg . Trazodone 50mg	Avoid sedatives (e.g., zolpidem); improves delirium risk by 20%
Constipation Prevention	Proactive bowel regimen	Dulcolax 5mg PO daily, senna 8.6 mg PO qHS, polyethylene glycol 17 g PO daily. movement, hydration	Start with opioids; monitor BMs q24h; avoid in C. diff
Delirium Avoidance	Non-pharmacologic interventions	CAM-ICU q12h, reorient q4h, family presence, avoid restraints. Haloperidol 0.5 mg IV q6h PRN (severe)	Reduce anticholinergics (e.g., diphenhydramine); 4AT score for screening
Early Mobility	Promote ambulation	PT/OT consult if appropriate, ambulate 3x/day, chair positioning. Goal: 200-300 ft/ day	Reduces LOS by 1-2 days, VTE risk by 30%; use walkers if fall risk
Discharge Planning	Structured process	START criteria day 1, med reconciliation, teach-back, follow-up within 7 days. Social work consult	Reduces 30-day readmissions by 15%; include caregiver education
Nutrition Optimization	Address malnutrition	Dietitian consult, high-protein diet (1.2-1.5 g/kg/day), oral supplements (Ensure 240 mL BID). NG if NPO >3 days	Malnutrition Universal Screening Tool (MUST); reduces complications by 20%
Fall Prevention	Risk assessment, mitigation	Morse Fall Scale on admission, bed alarms, non-slip socks, lower bed height. Trazodone 25 mg PO qHS if insomnia	Avoid benzodiazepines; 50% of falls in elderly occur at night
Pain Management	Multimodal approach	Acetaminophen 650 mg PO q6h, lidocaine patches, gabapentin 300 mg PO qHS. Opioids (morphine 2 mg IV PRN) if severe	Non-opioid first; reduces delirium, constipation risk

## Clinical Presentation

### • Symptoms

- **Sleep** Disturbances Insomnia, frequent waking, daytime fatigue
- **Constipation** Abdominal discomfort, infrequent BMs (<3/week)
- **Delirium** Confusion, agitation, inattention
- **Immobility** Weakness, orthostasis, pressure ulcers
- **Discharge** Issues Medication non-adherence, missed follow-ups

- **Rare:** Hallucinations (delirium), fecal impaction (constipation), aspiration (malnutrition)
- **Exam**
  - **Sleep** Fatigue, irritability
  - **Constipation** Distended abdomen, hypoactive bowel sounds
  - **Delirium** Waxing/waning consciousness, CAM-ICU positive
  - **Immobility** Muscle atrophy, DVT signs
  - **Discharge** Poor teach-back, social barriers
  - **Rare** Trousseau's sign (delirium hypocalcemia), skin breakdown (immobility)
- **Red Flags** Delirium: (CAM-ICU positive), no BM >3 days, Morse score >45, MUST >2, readmission risk

## Labs and Studies

- **Labs**
  - CMP Na<sup>+</sup> (delirium), Cr (renal function), glucose (hypoglycemia)
  - CBC Anemia (malnutrition), leukocytosis (infection)
  - Electrolytes K<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup> (delirium, constipation)
  - Advanced Albumin (<3 g/dL, malnutrition), prealbumin (nutrition status), CRP (inflammation)
- **Imaging**
  - CXR Pneumonia (delirium trigger), aspiration (malnutrition)
  - CT Head Rule out stroke, ICH in delirium
  - abdominal X-ray (constipation impaction)

## Management Strategies

General Principles Implement proactive, multidisciplinary strategies to prevent complications, enhance recovery, and ensure smooth transitions, involving medicine, nursing, PT/OT, and social work.

- Supportive Care
  - Monitoring Vitals q4h, daily weights, BM log
- Environment
  - Quiet rooms, low lighting, family involvement
- Patient Education
  - Teach-back for discharge, mobility goals
- Specific Interventions
  - Sleep Earplugs, melatonin, avoid caffeine post-noon
  - Constipation Bowel regimen with opioids, hydration

- Delirium Reorient, remove catheters, avoid restraints Mobility PT/OT, ambulation goals, fall precautions
- Discharge Med reconciliation, follow-up appointments
- Nutrition High-protein diet, supplements, NG if needed
- Falls Bed alarms, assist devices, night rounds
- Pain Non-opioids, regional blocks, minimize opioids
- Weekly nutrition labs (albumin, prealbumin) Discharge readiness (START criteria)

## Complications

### • Acute

- **Delirium:** Prolonged LOS, mortality 10-20%
- **Constipation:** Impaction, obstruction (<1%)
- **Falls:** Fractures (1-2%), head injury
- **Immobility** VTE (2-5%), pressure ulcers (5-10%), ICU myopathy
  - Rare
    - Aspiration pneumonia (malnutrition),
    - torsades (delirium drugs)

### • Long-Term

- Readmission 30% within 30 days without planning
- Functional Decline 20-30% post-immobility
- Malnutrition Chronic wounds, infections

## Clinical Scenarios

### Case 1 Delirium in Elderly

Presentation 80 y/o F with CHF, admitted for pneumonia, develops confusion. Vitals BP 130/80, HR 90, SpO2 94%, RR 18. Exam Disoriented, CAM-ICU positive.

Labs/Studies Na+ 132 mEq/L, Cr 1.5 mg/dL, WBC 12K. CXR RLL infiltrate. I

nterpretation Delirium, infection-triggered.

Management Reorient q4h, family presence, haloperidol 0.5 mg IV PRN, ceftriaxone 2 g IV q24h. Delirium resolves by day 4.

### Case 2 Constipation Post-Surgery

- **Presentation** 55 y/o M post-colectomy, no BM x 3 days, abdominal pain. Vitals BP 120/80, HR 85, SpO2 96%, RR 16.
- **Exam** Distended abdomen.

- **Labs/Studies** Normal CMP, K<sup>+</sup> 4 mEq/L. Abdominal X-ray Air-fluid levels.
- **Interpretation** Opioid-induced constipation.
- **Management** Dulcolax 5mg daily, senna 8.6 mg PO qHS, polyethylene glycol 17 g PO. BM by day 2, pain improves.
  - NOTE: Colace ineffective for constipation, do not use

### Case 3 Delayed Discharge

- **Presentation** 70 y/o M with COPD, ready for discharge but no follow-up plan. Vitals BP 140/80, HR 80, SpO<sub>2</sub> 95%, RR 14.
- **Exam** Stable, poor teach-back.
- **Labs/Studies** Normal CMP, O<sub>2</sub> sat 95% on RA. CXR Stable emphysema.
- **Interpretation** High readmission risk, poor planning.
- **Management** Social work consult, med reconciliation, pulmonology follow-up in 7 days. Discharge day 6, no readmission.

### Expert Tips

- Screen for delirium with CAM-ICU q12h; avoid benzodiazepines
- Start bowel regimen with opioids; polyethylene glycol for refractory constipation
- Ambulate within 24h of admission; PT/OT consult day 1
- Use START criteria for discharge; teach-back reduces errors by 20%
- Assess MUST score on admission; high-protein diet prevents deconditioning
- Pitfall Missing delirium triggers; check infections, electrolytes
- Advanced Actigraphy for sleep optimization; wearable sensors for mobility tracking

### Key Pearls

**Delirium prevention:** Reorient, minimize sedatives, CAM-ICU q12h

**Constipation:** Proactive bowel regimen, hydrate, monitor BMs

**Early mobility:** Ambulate 3x/day, reduces LOS, VTE risk

**Discharge planning:** START criteria, follow-up within 7 days

**Nutrition, falls, pain:** MUST score, Morse scale, non-opioid analgesia

### References

[UpToDate](#) "Hospital Medicine: Patient Safety Strategies" (2025)

[NEJM](#) "Delirium Prevention in Hospitalized Patients" (2024)

[JAMA](#) "Early Mobility in Inpatient Care" (2024)

[Lancet](#) "Nutrition and Outcomes in Hospitalized Patients" (2023)

[J Hosp Med](#) "Discharge Planning: Reducing Readmissions" (2024)

**Visit [medcheatsheets.com](https://medcheatsheets.com) for more education, fun resources and 10 category 1 AAPA CME credit!**

© Hospital Medicine Cheat Sheets ([medcheatsheets.com](https://medcheatsheets.com)). For educational purposes only. Do not redistribute or sell. Neither the author nor the company is liable for real-world implications. AI was used in development.