



Bloch's theorem


Recap of previous lectures

- Crystals and Bravais lattices
- Bragg diffraction, Reciprocal space, Brillouin zones
- Electrons in lattices and the independent electron approximation



Electrons in lattices

- **Independent electron approx:** Neglect e^-e^- interactions
- Each electron experiences a potential $V(\mathbf{r})$ from the ions



$$\left(\frac{-\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) \right) \psi(\mathbf{r}) = E \psi(\mathbf{r})$$

- Potential has the same periodicity as the lattice

$$V(\mathbf{r} + \mathbf{a}) = V(\mathbf{r})$$


for primitive lattice vector \mathbf{a}

- **Bloch's theorem:**
Special structure of eigenstates due to periodicity of V

One dimension

$$\left(\frac{-\hbar^2}{2m} \frac{d^2}{dx^2} + V(x) \right) \psi(x) = E\psi(x)$$

$$V(x + a) = V(x)$$




$$V(x) = \sum_{n=-\infty}^{\infty} V_n e^{inGx}$$


$$G = 2\pi/a$$

Coupling plane wave states

- A periodic lattice potential $V(x)$ only generates superpositions of plane wave states with wavevectors related by multiples of the reciprocal lattice constant $G = 2\pi / a$



$$V(x + a) = V(x)$$
$$\hat{V} |k\rangle = \sum_n V_n |k + nG\rangle$$



X-Ray diffraction,
von Laue, 1912

The structure of eigenstates

- A general form for eigenstates:
$$\psi_q(x) = \sum_n b_n e^{i(q+nG)x}$$

- Bloch's theorem (1D)**


Eigenstates of the Schrödinger equation with a periodic potential have the form

$$\psi_q(x) = e^{iqx} u_q(x)$$

Where q is a wavevector in the first Brillouin zone $[-\pi, \pi]$ and $u_q(x)$ is a periodic function of x

The “Bloch wavefunction”
(periodic)

$$u_q(x) = \sum_n b_n e^{inGx}$$



Three dimensions

- The same ideas all hold in three dimensions

$$V(x) = \sum_{n=-\infty}^{\infty} V_n e^{inGx}$$



$$\psi_q(x) = e^{iqx} \sum_n b_n e^{inGx}$$

$$V(\mathbf{r}) = \sum_{\mathbf{G}} V_{\mathbf{G}} e^{i\mathbf{G} \cdot \mathbf{r}}$$



$\mathbf{G} \in$ reciprocal lattice



$$\begin{aligned} \psi_{\mathbf{q}}(\mathbf{r}) &= e^{i\mathbf{q} \cdot \mathbf{r}} \sum_{\mathbf{G}} b_{\mathbf{G}} e^{i\mathbf{G} \cdot \mathbf{r}} \\ &= e^{i\mathbf{q} \cdot \mathbf{r}} u_{\mathbf{q}}(\mathbf{r}) \end{aligned}$$

Summary

- Electrons in a periodic lattice potential
- Work in basis of plane waves
- Periodic potentials couple plane waves related by reciprocal lattice vectors
- Bloch's theorem: Wavefunctions are product of a periodic part $u_{\mathbf{q}}(\mathbf{r})$ (the “Bloch state”) and a phase factor $e^{i\mathbf{q}\cdot\mathbf{r}}$



Further reading

- Section 7 of Kittel 'Introduction to solid state physics'
- P. 134 of Ashcroft and Mermin 'Solid state physics': Alternative derivation based on symmetry operators and group theory
- My lecture notes:

