

# Cornu Spirals

From Euler to Ferrari

Max Mussavian

Arcana, May 2022

- Definitions: Curves, Lengths and Curvature
- Relating the Curve Length to its Curvature
- The Euler Spiral
- Other Fun Curves
- “Curvature determines the Curve”
- Application 1: Diffraction and Fresnel integrals
- Application 2: Designing Roads and Railways
- Application 3: The Racing Line



# What is a Curve?

Before we begin note that everything we look at here is in 2 dimensions  $\mathbb{R}^2$

## Definition (A Parametric Curve)

A parametric curve is a twice differentiable function that has the form  $x = g(t)$  and  $y = h(t)$  defined on an open interval  $(a, b)$ . The set of points traced out by the curve is called the trace.



TONBRIDGE  
SCHOOL

# How Long is a Curve?

## Definition (Arc Length)

The length of a curve  $s$  is given by  $s = \int_{t_1}^{t_2} \sqrt{x'^2 + y'^2} dt$

Intuition: The velocity at time  $t$  is  $\begin{pmatrix} x'(t) \\ y'(t) \end{pmatrix}$  and hence the speed at time  $t$  is  $\sqrt{x'(t)^2 + y'(t)^2}$ . Distance travelled (arc length) is the integral of speed with respect to time.

Alternatively approximate the trace by line segments. The length of line segments converges to the arc length as the segments get smaller and smaller.



TONBRIDGE  
SCHOOL

# Some Simple Examples

- Circle: We can define a circle with radius  $r$  as  $x(t) = r \cos(t)$  and  $y(t) = r \sin(t)$ . The arc length is

$$s = \int_0^{2\pi} \sqrt{r^2 \sin^2 t + r^2 \cos^2 t} dt = [r]_0^{2\pi} = 2\pi r$$

- Parabola: We can define a parabola as  $x(t) = t$  and  $y(t) = t^2$ . The Arc Length between 0 and 1 is

$$\begin{aligned} s &= \int_0^1 \sqrt{1 + 4t^2} dt \\ &= \left[ \frac{1}{2} t \sqrt{1 + 4t^2} + \frac{1}{4} \ln \left( 2t + \sqrt{1 + 4t^2} \right) \right]_0^1 \\ &\approx 1.48 \end{aligned}$$



TONBRIDGE  
SCHOOL

# How Curved is a Curve?

The curvature at a point on the curve is the reciprocal of the radius of the circle that approximates the curve.

## Definition (Curvature)


The curvature of curve is  $\kappa$  given by  $\kappa = \frac{x'y'' - y'x''}{(x'^2 + y'^2)^{3/2}}$

Put in explanation here



TONBRIDGE  
SCHOOL

# Explaining Definition for Curvature



For any circle with radius  $r$  we have  
 $s = r\theta$ .

Therefore for “kissing” circle  $\frac{ds}{dt} = r \frac{d\theta}{dt}$   
 But  $\tan \theta = \frac{y'(t)}{x'(t)}$  and taking derivatives  
 w.r.t.  $t$  we have  $\sec^2 \theta \frac{d\theta}{dt} = \frac{x'y'' - y'x''}{x'^2}$ .

Further  $\frac{1}{\cos^2 \theta} = \frac{x'^2 + y'^2}{x'^2}$  therefore  
 $\frac{d\theta}{dt} = \frac{x'y'' - y'x''}{x'^2 + y'^2}$

From above  $\frac{ds}{dt} = \sqrt{x'^2 + y'^2}$ .

Combining gives us

$$r = \frac{ds/dt}{d\theta/dt} = \frac{(x'^2 + y'^2)^{3/2}}{x'y'' - y'x''}$$

But  $\kappa = \frac{1}{r}$



TONBRIDGE  
SCHOOL

# Some Simple Examples

- Circle: Recall  $x(t) = r \cos(t)$  and  $y(t) = r \sin(t)$

$$\kappa = \frac{r^2 \sin^2 t + r^2 \cos^2 t}{(r^2 \sin^2 t + r^2 \cos^2 t)^{\frac{3}{2}}} = \frac{1}{r}$$

- Parabola: Recall  $x(t) = t$  and  $y(t) = t^2$ .

$$\kappa = \frac{2}{(1 + 4t^2)^{\frac{3}{2}}}$$



TONBRIDGE  
SCHOOL



# Relating Curve Length to Curvature

Let's try the following parametrisation for  $x$  and  $y$

$$\begin{aligned}x(t) &= \int_0^t \cos f(u) du \\y(t) &= \int_0^t \sin f(u) du\end{aligned}$$

This gives us

$$\begin{aligned}x' &= x'(t) = \cos f(t) & \text{and} & & x'' &= -f'(t) \sin f(t) \\y' &= y'(t) = \sin f(t) & \text{and} & & y'' &= f'(t) \cos f(t)\end{aligned}$$



TONBRIDGE  
SCHOOL

# Relating Curve Length to Curvature

This gives us the following for slope, arc length and curvature

$$\frac{dy}{dx} = \frac{\sin f(t)}{\cos f(t)} = \tan f(t)$$

$$s = \int_0^t \sqrt{\cos^2 f(u) + \sin^2 f(u)} du = t$$

$$\kappa = \frac{f'(t) \cos^2 f(t) + f'(t) \sin^2 f(t)}{\cos^2 f(t) + \sin^2 f(t)} = f'(t)$$



TONBRIDGE  
SCHOOL

# Define the Curve by Curve Length and Curvature

We can get replace “time” variable  $t$  by the curve length  $s$ .  
And the curvature at point  $t$  is  $f'(t)$ . Which means

$$f(t) = \int \kappa(t) dt$$

Thus the equations for the curve become

$$\begin{aligned}x = x(s) &= \int_0^s \cos \left( \int_0^u \kappa(t) dt \right) du \\y = y(s) &= \int_0^s \sin \left( \int_0^u \kappa(t) dt \right) du\end{aligned}$$

Hence the curve is defined by curve length and curvature alone.



TONBRIDGE  
SCHOOL

# A Very Simple Example

We can make the curvature  $\kappa$  constant and equal to 1. Then  $\int_0^u \kappa(t) dt = u$  and

$$\begin{aligned}x = x(s) &= \int_0^s \cos u du = \sin s \\y = y(s) &= \int_0^s \sin u du = -\cos s + 1\end{aligned}$$

which is the parametric curve for a circle with centre  $(0, 1)$  and radius 1.



TONBRIDGE  
SCHOOL

# The Euler Spiral

A more interesting example is the curvature equal to the arc length.

$$\kappa(s) = s$$

Then  $\int_0^u \kappa(t) dt = \frac{u^2}{2}$  and

$$\begin{aligned}x = x(s) &= \int_0^s \cos \frac{u^2}{2} du \\y = y(s) &= \int_0^s \sin \frac{u^2}{2} du\end{aligned}$$

These integrals are called “Fresnel Integrals”.

They cannot be solved analytically. But what do they look like numerically?



TONBRIDGE  
SCHOOL

# Making a Plotter in Python

```
def x_func(upper, lower, func):  
    def integrand(t):  
        return np.cos(func(t))  
    result, _ = integrate.quad(integrand, upper, lower)  
    return result
```

```
def y_func(upper, lower, func):  
    def integrand(t):  
        return np.sin(func(t))  
    result, _ = integrate.quad(integrand, upper, lower)  
    return result
```

Euler Spiral

```
[7] k = s  
    spy.integrate(k, s)  
  
     $\frac{s^2}{2}$ 
```

```
[8] def func_euler(x):  
    return 0.5 * np.power(x, 2)
```

```
[9] arc_length, x_coord, y_coord = x_y_coordinates(x_func, y_func, func_euler, s_steps)
```


First write functions to calculate  
x and y coordinates More words  
here



TONBRIDGE  
SCHOOL

# The Euler Spiral


Figure: The Euler Spiral aka Cornu Curve



TONBRIDGE  
SCHOOL

# The Euler Spiral - x y Coordinates

Figure: Fresnel Integrals with arguments  $\frac{u^2}{2}$



These converge to  $\pm \frac{\sqrt{\pi}}{2} \approx 0.8862$ .




TONBRIDGE  
SCHOOL



# Other Fun Curves: Even Powers of $s$


Figure:  $\kappa(s) = s^2$



TONBRIDGE  
SCHOOL

# More Fun Curves: Mix in a Bit of a Circle


Figure:  $\kappa(s) = s^2 - 2.19$



TONBRIDGE  
SCHOOL

# More Fun Curves: Polynomials


Figure:  $\kappa(s) = 5s^4 - 18s^2 + 5$



TONBRIDGE  
SCHOOL

# More Fun Curves: Trigonometric Functions


Figure:  $\kappa(s) = \cos(s) - s \sin(s)$



TONBRIDGE  
SCHOOL

# More Fun Curves: Hyperbolic Functions

Figure:  $\kappa(s) = \sinh(s) - 5.19$



TONBRIDGE  
SCHOOL

# "Curvature determines the Curve"



TONBRIDGE  
SCHOOL

# Diffraction and Fresnel Integrals



TONBRIDGE  
SCHOOL



# Designing Roads and Railways

- 1 Transition curves are used to link straight sections of motorways or railways.
- 2 They are designed to give passengers a smooth ride.
- 3 In particular so sudden changes in acceleration.

Figure: Cloverleaf Motorway Interchange





# Why Transition Curves are Euler Spirals

- The acceleration along the transition is given by

$$a = s''(t)\vec{T} + \kappa s'(t)^2\vec{N}$$

Where  $\vec{T}$  is the unit tangent vector and  $\vec{N}$  is the unit normal vector.

- If the car/train is going round the curve at constant speed  $s'(t) = \text{constant}$  and  $s''(t) = 0$ .
- The acceleration at constant speed only depends on the curvature  $\kappa$  and speed  $s'(t)$  in the direction of the normal vector.



TONBRIDGE  
SCHOOL

# The Racing Line



TONBRIDGE  
SCHOOL